

**On global smooth solution of Cauchy problem for a class
of quasilinear parabolic systems in several spaces
variables**

O. Bennouna

► **To cite this version:**

O. Bennouna. On global smooth solution of Cauchy problem for a class of quasilinear parabolic systems in several spaces variables. RR-0487, INRIA. 1986. <inria-00076067>

HAL Id: inria-00076067

<https://hal.inria.fr/inria-00076067>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IRIA

CENTRE DE ROCQUENCOURT

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
B.P.105
78153 Le Chesnay Cedex
France
Tél. : (1) 39 63 55 11

Rapports de Recherche

N° 487

**ON GLOBAL SMOOTH SOLUTION
OF CAUCHY PROBLEM
FOR A CLASS OF QUASILINEAR
PARABOLIC SYSTEMS
IN SEVERAL SPACES VARIABLES**

Omar BENNOUNA

Février 1986

ON GLOBAL SMOOTH SOLUTION OF CAUCHY
PROBLEM FOR A CLASS OF QUASILINEAR
PARABOLIC SYSTEMS IN SEVERAL SPACES
VARIABLES

Omar BENNOUNA^(*)

(*) BEL Laboratories, 38 allée des Roses - Fès - MAROC.

Abstract- In this paper we consider a class of quasilinear parabolic systems in several spaces variables :

$$(*) \left\{ \begin{aligned} \frac{\partial u}{\partial t} + \sum_{i=1}^N B_i(t, x, u) \frac{\partial u}{\partial x_i} + C(t, x, u) &= \mu \Delta u \text{ in } \pi_T \\ u(x, 0) &= u_0(x) \text{ in } \mathbb{R}^N \end{aligned} \right.$$

Where B_i , C and u_0 are given data, $\mu > 0$ fixed and $\pi_T =]0, T[\times \mathbb{R}^N$, $u = (u^1, u^2, \dots, u^M)$.

Using probabilistic methods we introduce a notion of generalized solution of (*) and we prove existence and uniqueness results.

Key words : quasilinear - generalized solution

Résumé- Dans ce papier on considère une classe de systèmes quasilinéaires paraboliques avec plusieurs variables d'espace :

$$(*) \left\{ \begin{aligned} \frac{\partial u}{\partial t} + \sum_{i=1}^N B_i(t, x, u) \frac{\partial u}{\partial x_i} + C(t, x, u) &= \mu \Delta u \text{ dans } \pi_T \\ u(x, 0) &= u_0(x) \text{ dans } \mathbb{R}^N \end{aligned} \right.$$

où B_i , C et u_0 sont des données, $\mu > 0$ fixé et $\pi_T =]0, T[\times \mathbb{R}^N$, $u = (u^1, u^2, \dots, u^M)$.

On introduit une notion de solution généralisée pour (*) pour laquelle on a un résultat d'existence et d'unicité ceci à l'aide de techniques probabilistes.

Mots clefs : quasilinéaire - solution généralisée

ON GLOBAL SMOOTH SOLUTION OF CAUCHY
PROBLEM FOR A CLASS OF QUASILINEAR
PARABOLIC SYSTEMS IN SEVERAL SPACES
VARIABLES

Omar BENNOUNA (*)

INTRODUCTION

We consider in this paper the problem of existence and uniqueness of global smooth solutions for the Cauchy problem

$$(*) \quad \frac{\partial u}{\partial t} + \sum_{i=1}^N B_i(t, x, u) \frac{\partial u}{\partial x_i} + C(t, x, u) = \mu \Delta u \quad \text{in } \Pi_T$$
$$u(x, 0) = u_0(x) \quad \text{in } \mathbb{R}^N$$

where B_i , C and u_0 are given data, $\mu > 0$ fixed, and $\Pi_T =]0, T[\times \mathbb{R}^N$,

$$u = (u^1, u^2, \dots, u^M).$$

Using probabilistic methods we introduce a notion of generalized solution of (*). When this generalized solution is smooth enough then it is a classical solution. In the proof of the existence result we use a successive approximation process.

The paper is organized as follows : the part I contains definitions and some usefull lemmas, and the part II is concerned with the existence and uniqueness results.

This paper generalizes the result of D.W. STROOCK [2] on the probabilistic representation of the solution of Cauchy problem for a class of linear parabolic systems.

(*) BEL Laboratories, 38 allée des Roses - Fès - MAROC.

I. DEFINITIONS AND PRELIMINARY RESULTS

I.1. Notations and assumptions

In the following we assume that :

- (1) $B_i^{k\ell}(t,x,v)$, $C^{k\ell}(t,x,v)$ are bounded and have bounded derivatives with respect to v
- (2) $u_0^k(x)$ is a bounded measurable function.

Let us denote

$$(3) \quad G(t,x,v) = \int_0^1 \frac{\partial C}{\partial u}(t,x,\theta v) d\theta \quad , \quad \forall (t,x,v) \in \Pi_T \times \mathbb{R}^M$$

then

$$C(t,x,v) = G(t,x,v)v + C(t,x,0) .$$

In the sequel, for convenience, we take

$$C(t,x,0) \equiv 0 .$$

Let $\Omega = C([0,\infty), \mathbb{R}^N)$. Given $t \geq 0$ and $\omega \in \Omega$ denote by $y(t) \equiv y(t,\omega)$ the value of ω at t . For $0 \leq s \leq t$, define

$$F_t^s = \mathcal{B}[y(\lambda) : s \leq \lambda \leq t]$$

$$F^s = \mathcal{B}[y(\lambda) : \lambda \geq s] .$$

Then it was shown in [1] that for each $s \geq 0$ and $x \in \mathbb{R}^N$ there is a unique probability measure $P_{s,x}$ on (Ω, F^s) such that

$$(4) \quad y(t) = x + \sqrt{2\mu} [w(t) - w(s)] \quad \text{a.s. } P_{s,x}$$

where $w(t)$ is the standard Wiener process.

Let $u(t,x)$ be a bounded measurable vector function and denote by $X_u^s(t)$, $t \geq s$ the solution to the stochastic integral equation

$$(5) \quad X_u^s(t) = I + \frac{1}{\sqrt{2\mu}} \int_s^t X_u^s(\lambda) B_i(\lambda, y(\lambda), u(\lambda, y(\lambda))) dw_i(\lambda) + \int_s^t X_u^s(\lambda) G(\lambda, y(\lambda), u(\lambda, y(\lambda))) d\lambda \quad \text{a.s. } P_{s,x}.$$

For convenience we consider problem (*) with final condition.

I.2. Definitions and lemmas

We now give the definition of a generalized solution.

Definition

A bounded measurable vector function $u(t,x)$ is a generalized solution of (*) if $u(t,x)$ satisfies the integral equation :

$$(6) \quad u(t,x) = E_{t,x} X_u^t(T) u_0(y(T)) \quad , \quad \forall (t,x) \in \Pi_T.$$

Remark 1

From the weak continuity of $P_{s,x}$ we have that $u \in C(\Pi_T)$.

Remark 2

When $u \in C_b^{2,1}(\Pi_T)$, using the Itô formula we see that u is a classical solution.

Let us give some usefull results :

Lemma 1 : Under conditions (1), (2), we have the estimate

$$(7) \quad \|u(t)\|_\infty^2 \leq M \|u_0\|_\infty^2 \exp\left[\frac{(T-t)}{2\mu} (\|B\|_\infty^2 + 4\mu \|G\|_\infty)\right]$$

$$\forall t \in [0, T].$$

Proof : We have

$$(8) \quad |u(t,x)| \leq \|u_0\|_{\infty} (E_{t,x} \|X_u^t(T)\|^2)^{1/2}.$$

On the other hand we see that

$$(9) \quad \begin{aligned} \|X_u^t(T)\|^2 &= M + \frac{1}{\sqrt{2\mu}} \sum_i \int_t^T \text{Tr}[X_u^t(s) B_i(s, y(s), u(s, y(s))) X_u^t(s)^*] dw_i(s) + \\ &+ \frac{1}{2\mu} \sum_i \int_t^T \text{Tr}[X_u^t(s) B_i(s, y(s), u(s, y(s))) \\ &\quad B_i^*(s, y(s), u(s, y(s))) X_u^t(s)^*] ds + \\ &+ 2 \int_t^T \text{Tr}[X_u^t(s) G(s, y(s), u(s, y(s))) X_u^t(s)^*] ds. \end{aligned}$$

Then, using condition (3) we get that

$$(10) \quad E \|X_u^t(T)\|^2 \leq M + \left[\frac{1}{2\mu} \|B\|_{\infty}^2 + 4\mu \|G\|_{\infty} \right] \int_t^T E \|X_u^t(s)\|^2 ds$$

which implies

$$E \|X_u^t(T)\|^2 \leq M \exp \left[\frac{(T-t)}{2\mu} (\|B\|_{\infty}^2 + 4\mu \|G\|_{\infty}) \right]$$

and then (7).

Remark 3

If we assume that : $\text{Tr}[AG(t,x,v)A^*] \leq -\alpha \|A\|^2$ for any $(t,x,v) \in \Pi_T \times \mathbb{R}^M$ and any $M \times M$ real valued matrix A we obtain

$$(11) \quad E \|X_u^t(s)\|^2 \leq M \exp \left[\frac{(s-t)}{2\mu} (\|B\|_{\infty}^2 - 4\alpha\mu) \right] \quad \forall s, 0 \leq t \leq s \leq T.$$

We assume in the following that

$$(12) \quad ||G(t,x,v_1) - G(t,x,v_2)|| \leq K ||v_1 - v_2|| \quad \forall (t,x) \in \Pi_T$$

Lemma 2 : Under conditions (1), (2) and (12) we have

$$(13) \quad E ||X_u^t(s) - X_v^t(s)||^2 \leq M_0(s-t) \int_t^s ||u(\theta) - v(\theta)||_\infty^2 d\theta$$

where

$$(14) \quad M_0(s-t) = 4M \cdot \exp\left[\frac{4}{u} (s-t) ||B||_\infty^2\right] \exp\left[4(s-t) ||G||_\infty (1+(s-t) ||G||_\infty)\right] \\ \cdot \left[||\frac{\partial G}{\partial u}||_\infty^2 (s-t) + \frac{1}{2u} ||\frac{\partial B}{\partial u}||_\infty^2 \right] (*)$$

Proof :

$$(15) \quad X_u^t(s) - X_v^t(s) = \frac{1}{\sqrt{2u}} \int_t^s X_u^t(\theta) [B_i(\theta, y(\theta), u(\theta, y(\theta))) - \\ - B_i(\theta, y(\theta), v(\theta, y(\theta)))] dw_i(\theta) + \\ + \frac{1}{\sqrt{2u}} \int_t^s [X_u^t(\theta) - X_v^t(\theta)] B_i(\theta, y(\theta), v(\theta, y(\theta))) dw_i(\theta) + \\ + \int_t^s X_u^t(\theta) [G(\theta, y(\theta), u(\theta, y(\theta))) - G(\theta, y(\theta), v(\theta, y(\theta)))] d\theta \\ + \int_t^s [X_u^t(\theta) - X_v^t(\theta)] G(\theta, y(\theta), v(\theta, y(\theta))) d\theta \\ = I + II + III + IV$$

We have

(*) $||\frac{\partial f}{\partial u}||_\infty$ is the smallest Lipschitz constant.

$$(16) \quad E(|I|^2) \leq \frac{1}{2\mu} E \int_t^s \|X_u^t(\theta)[B_i(u(\theta, y(\theta))) - B_i(v(\theta, y(\theta)))]\|^2 d\theta$$

$$\leq \frac{1}{2\mu} \|\frac{\partial B}{\partial u}\|_\infty^2 \sup_{t \leq \theta \leq s} (E\|X_u^t(\theta)\|^2) \int_t^s \|u(\theta) - v(\theta)\|^2 d\theta$$

$$(17) \quad E(|II|^2) \leq \frac{1}{2\mu} E \int_t^s \|[X_u^t(\theta) - X_v^t(\theta)]B_i(v(\theta, y(\theta)))\|^2 d\theta$$

$$\leq \frac{1}{2\mu} \|B\|_\infty^2 \int_t^s E\|X_u^t(\theta) - X_v^t(\theta)\|^2 d\theta$$

$$(18) \quad E(|III|^2) \leq \|\frac{\partial G}{\partial u}\|_\infty^2 \sup_{t \leq \theta \leq s} (E\|X_u^t(\theta)\|^2) \int_t^s \|u(\theta) - v(\theta)\|^2 d\theta (s-t)$$

$$(19) \quad E(|IV|^2) \leq \|G\|_\infty^2 \int_t^s E\|X_u^t(\theta) - X_v^t(\theta)\|^2 d\theta (s-t)$$

Hence, we get that

$$(20) \quad E\|X_u^t(s) - X_v^t(s)\|^2 \leq 4\left(\frac{1}{2\mu} \|B\|_\infty^2 + \|G\|_\infty^2 (s-t)\right) \int_t^s E\|X_u^t(\theta) - X_v^t(\theta)\|^2 d\theta +$$

$$+ 4 \sup_{t \leq \theta \leq s} E\|X_u^t(\theta)\|^2 \left[\|\frac{\partial G}{\partial u}\|_\infty^2 (s-t) + \frac{1}{2\mu} \|\frac{\partial B}{\partial u}\|_\infty^2 \int_t^s \|u(\theta) - v(\theta)\|^2 d\theta \right]$$

and then, using (11) and the Gronwall lemma,

$$(21) \quad E\|X_u^t(s) - X_v^t(s)\|^2 \leq 4M(\exp[2(s-t)] \left(\frac{1}{\mu} \|B\|_\infty^2 + 2\|G\|_\infty \right))$$

$$\left[\|\frac{\partial G}{\partial u}\|_\infty^2 (s-t) + \frac{1}{2\mu} \|\frac{\partial B}{\partial u}\|_\infty^2 \int_t^s \|u(\theta) - v(\theta)\|^2 d\theta \right]$$

$$\left(\exp[4(s-t)] \left(\frac{1}{2\mu} \|B\|_\infty^2 + \|G\|_\infty^2 (s-t) \right) \right)$$

$$\int_t^s \|u(\theta) - v(\theta)\|^2 d\theta$$

which completes the proof.

II. EXISTENCE AND UNIQUENESS RESULTS

Let us consider the following process of successive approximations :

$$(22) \quad \begin{aligned} u^0(t,x) &= u_0(x) \\ u^n(t,x) &= E_{t,x} X_{u^{n-1}}^t(T) u_0(y(T)) \quad , \quad \forall n, n \geq 1 \end{aligned}$$

where $X_{u^{n-1}}^t(s)$ is the solution of

$$(23) \quad \begin{aligned} X_{u^{n-1}}^t(s) &= I + \frac{1}{\sqrt{2\mu}} \int_t^s X_{u^{n-1}}^t(\theta) B_i(\theta, y(\theta), u^{n-1}(\theta, y(\theta))) dw_i(\theta) \\ &\quad + \int_t^s X_{u^{n-1}}^t(\theta) G(\theta, y(\theta), u^{n-1}(\theta, y(\theta))) d\theta \end{aligned}$$

then, we have

Theorem 1 : Under conditions (1), (2) and (12) we have

$$(24) \quad \sup_{t \in [0, T]} \| |u^{n+1}(t) - u^n(t)| \|_\infty^2 \leq \frac{(M_1(T) \cdot T)^n}{n!} \sup_{t \in [0, T]} \| |u^1(t) - u^0(t)| \|_\infty^2$$

where

$$(25) \quad M_1(T) = \| |u_0| \|_\infty^2 M_0(T) .$$

Proof : Indeed, we have for $n \geq 1$

$$(26) \quad u^{n+1}(t,x) - u^n(t,x) = E_{t,x} X_{u^n}^t(T) u_0(y(T)) - E_{t,x} X_{u^{n-1}}^t(T) u_0(y(T)) .$$

Thus

$$\begin{aligned} \| |u^{n+1}(t,x) - u^n(t,x)| \| &\leq E_{t,x} \| |u_0(y(T))| \| \| |X_{u^n}^t(T) - X_{u^{n-1}}^t(T)| \| \\ &\leq \| |u_0| \|_\infty E_{t,x} \| |X_{u^n}^t(T) - X_{u^{n-1}}^t(T)| \| . \end{aligned}$$

From (13), we get that

$$(27) \quad \|u^{n+1}(t) - u^n(t)\|_\infty^2 \leq \|u_0\|_\infty^2 M_0(T-t) \int_t^T \|u^n(\theta) - u^{n-1}(\theta)\|_\infty^2 d\theta .$$

Hence

$$(28) \quad \sup_{t \in [0, T]} \|u^{n+1}(t) - u^n(t)\|_\infty^2 \leq \left(\|u_0\|_\infty^2 \right)^n \frac{(M_0(T) \cdot T)^n}{n!} \sup_{t \in [0, T]} \|u^1(t) - u^0(t)\|_\infty^2$$

which completes the proof.

Now from (24) we have the uniform convergence of the sequence $u^n(t, x)$ on $\Pi_{T, R}$ (*) to a vector valued function $u(t, x)$.

On the other hand, there is a matrix-valued function $X_u^s(t)$ such that

$$(29) \quad X_u^s(t) = I + \frac{1}{\sqrt{2\mu}} \int_s^t X_u^s(\theta) B_i(\theta, y(\theta), u(\theta, y(\theta))) d\omega_i(\theta) + \\ + \int_s^t X_u^s(\theta) G(\theta, y(\theta), u(\theta, y(\theta))) d\theta \quad , \quad \text{a.s. } P_{s, x}$$

and $u(t, x)$ satisfies

$$(30) \quad u(t, x) = E_{t, x} X_u^t(T) u_0(y(T)) .$$

Thus $u(t, x)$ is a generalized solution of (*).

To obtain the uniqueness result, let $(X_v^s(t), v(t, x))$ be another generalized solution of (*).

We have

$$(31) \quad u(t, x) - v(t, x) = E_{t, x} X_u^t(T) u_0(y(T)) - E_{t, x} X_v^t(T) u_0(y(T))$$

$$(*) \quad \Pi_{T, R} =]0, T[\times B(0, R) .$$

hence,

$$\|u(t) - v(t)\|_{\infty}^2 \leq \|u_0\|_{\infty}^2 E_{t,x} \|X_u^t(T) - X_v^t(T)\|^2.$$

From this and (13) we obtain that

$$(32) \quad \|u(t) - v(t)\|_{\infty}^2 \leq \|u_0\|_{\infty}^2 M_0(T) \cdot \int_t^T \|u(\theta) - v(\theta)\|_{\infty}^2 d\theta.$$

Thus, by Gronwall's inequality we have that

$$(33) \quad \|u(t) - v(t)\|_{\infty}^2 \equiv 0, \quad \forall t \in [0, T]$$

i.e. $u(t, x) = v(t, x)$ in $\bar{\Pi}_T$.

Remark 4

When $B_i^{k\ell}(t, x, v)$, $G(t, x, v)$ and $u_0(x)$ have two continuous derivatives in x and v then we can obtain a priori estimate of derivatives of the generalized solution.

Remark 5

The results of this paper remain valid for the more general parabolic systems with Cauchy condition :

$$\frac{\partial u^k}{\partial t} + \sum_{i=1}^N \sum_{\ell=1}^M B_i^{k\ell}(t, x, u) \frac{\partial u^{\ell}}{\partial x_i} + C^k(t, x, v) = \sum_{i,j=1}^N a_{ij}(t, x) \frac{\partial^2 u^k}{\partial x_i \partial x_j}$$

in Π_T

$$u^k(0, x) = u_0^k(x) \text{ in } \mathbb{R}^N, \quad 1 \leq k \leq M.$$

REFERENCE

- [1] D. STROOCK, S.R.S. VARADHAN, Diffusion processes with continuous coefficients I, II, Comm. Pure Appl. Math. Vol. 22, 1969.

- [2] D.W. STROOCK, On certain systems of parabolic equations,
Comm. on Pure and Applied Math., Vol. 23 (1970).

Imprimé en France

par

l'Institut National de Recherche en Informatique et en Automatique

