



On strongly coupled elliptic systems: existence,uniqueness and representation results

O. Bennouna

► To cite this version:

O. Bennouna. On strongly coupled elliptic systems: existence,uniqueness and representation results. RR-0453, INRIA. 1985. inria-00076102

HAL Id: inria-00076102

<https://inria.hal.science/inria-00076102>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



CENTRE DE ROCQUENCOURT

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
BP 105
78153 Le Chesnay Cedex
France

Tél. (1) 39 63 55 11

Rapports de Recherche

N° 453

ON STRONGLY COUPLED ELLIPTIC SYSTEMS: EXISTENCE, UNIQUENESS AND REPRESENTATION RESULTS

Omar BENNOUNA

Novembre 1985

ON STRONGLY COUPLED ELLIPTIC SYSTEMS :

EXISTENCE, UNIQUENESS AND REPRESENTATION

RESULTS

Omar BENNOUNA (*)

(*) BEL Laboratories, 38 Allée des Roses - FES - MAROC.

RESUME

Dans ce papier on considère une classe d'équations et de systèmes elliptiques qui peuvent être réduits à la forme

$$(*) \quad -\sum_{i,j} a_{ij} \frac{\partial^2 u_k}{\partial x_i \partial x_j} - \sum_l \sum_i b_{kl,i} \frac{\partial u_l}{\partial x_i} + \sum_l c_{kl} u_l = f_k \quad \text{dans } \mathcal{O}$$

où a_{ij} , $b_{kl,i}$, c_{kl} , f_k sont des données et \mathcal{O} est un ouvert borné de \mathbb{R}^N . On donne un résultat du type principe du maximum et on démontre un résultat d'existence et d'unicité. D'autre part on donne une représentation de la solution du type Feynman-Kac.

MOTS-CLEFS

Elliptique - Principe du maximum - Représentation du type Feynman-Kac.

ABSTRACT

In this paper we consider a class of elliptic equations and systems which can be reduced to the form :

$$(*) \quad -\sum_{i,j} a_{ij} \frac{\partial^2 u_k}{\partial x_i \partial x_j} - \sum_l \sum_i b_{kl,i} \frac{\partial u_l}{\partial x_i} + \sum_l c_{kl} u_l = f_k \quad \text{in } \mathcal{O}$$

where a_{ij} , $b_{kl,i}$, c_{kl} , f_k are given data and \mathcal{O} is an open bounded subset of \mathbb{R}^N . We give a maximum principle type result for this class and we prove existence and uniqueness results.

On the other hand, we give a Feynman-Kac type representation of the solution of (*).

KEY WORDS

Elliptic - Maximum principle - Feynman-Kac representation.

ON STRONGLY COUPLED ELLIPTIC SYSTEMS :
EXISTENCE, UNIQUENESS AND REPRESENTATION
RESULTS

O. Bennouna

INTRODUCTION

We consider a class of elliptic equations and systems which can be reduced to the form :

$$\left\{ \begin{array}{l} - \sum_{i,j} a_{ij} \frac{\partial^2 u_k}{\partial x_i \partial x_j} - \sum_l \sum_i b_{kl,i} \frac{\partial u_l}{\partial x_i} + \sum_l c_{kl} u_l = f_k \quad \text{in } \Omega \\ u_k|_{\partial\Omega} = 0 \quad 1 \leq k \leq M \end{array} \right. (*)$$

where a_{ij} , $b_{kl,i}$, c_{kl} , f_k are given data and Ω is an open bounded subset of \mathbb{R}^N with smooth boundary $\partial\Omega$.

In this note we give first a maximum principle type result for this class. To obtain the existence and uniqueness result we use a perturbation technic of weakly coupled system of second order elliptic partial differential equations to which we apply accretive operators methods.

The parabolic case is studied in [2] by the method of parabolic singular integrals.

On the other hand, we give a Feynman-Kac type representation of the solution of (*) using the notion of Itô random evolution. In the parabolic

case this question is solved in [1]. This type of result is made possible in the elliptic case under a stability condition of (*) which means, roughly speaking, that the dissipation term must be large enough.

Let us give an example of forth order equation which can be reduced to the form (*).

We consider

$$\left\{ \begin{array}{l} -\Delta^2 u + b\Delta u + cu = f \text{ in } \Omega \\ u|_{\partial\Omega} = 0, \quad \Delta u|_{\partial\Omega} = 0 \end{array} \right.$$

when b , c and f are given smooth data.

We denote $v = \Delta u$, then we obtain the second order elliptic system :

$$\left\{ \begin{array}{l} -\Delta v + bv + cu = f \\ -\Delta u + v = 0 \\ u|_{\partial\Omega} = 0, \quad v|_{\partial\Omega} = 0. \end{array} \right.$$

As we shall see below this problem is well posed, and we have a probabilistic representation of u , under some natural conditions on the data.

The paper is organized as follows : the part I contains analytic results and the part II is concerned with the Feynman-Kac type result.

I - THE ANALYTIC RESULTS

I.1. Assumptions and notations

Let Ω be an open bounded smooth domain in \mathbb{R}^N . We assume that :

1) Regularity conditions :

$$\phi = a_{ij}, b_{kl}, i, c_{kl}, f_k ; \quad \phi \in C(\bar{\Omega}) .$$

2) Ellipticity condition :

$$\sum_{i,j} a_{ij} \xi_i \xi_j \geq \gamma |\xi|^2 , \quad \gamma > 0 , \quad \forall \xi \in \mathbb{R}^N .$$

3) Stability condition :

$$\sum_{k,l} c_{kl} v_k v_l \geq \beta |v|^2 , \quad \beta > 0 \text{ large enough} \\ \forall v \in \mathbb{R}^M$$

(as we shall see in remark 2.2 it is sufficient to assume that

$$\sum_{k,l} c_{kl} v_k v_l \geq 0) .$$

I.2. A maximum principle for (*)

In this section we shall give a maximum principle type result.

We have

Theorem 1.1 - Assume that conditions (1), (2), (3) hold and

$$(10) \quad 4\gamma\beta > ||b||_{\infty}^2$$

then we have for $u, \bar{u} \in C(\bar{\Omega}) \cap W^{2,p}(\Omega)$
solutions of (*) corresponding respectively to f, \bar{f}

$$(11) \quad ||u - \bar{u}||_{\infty} \leq \frac{1}{\beta - \frac{||b||_{\infty}^2}{4\gamma}} ||f - \bar{f}||_{\infty}$$

where

$$||b||_{\infty}^2 = \sum_i ||b_i||_{\infty}^2 ,$$

Proof : It is enough to prove (11) for $u_k \in C(\bar{\Omega}) \cap C^2(\Omega)$. Hence, multiplying (*) by u_k and then summing up from 1 to M yields

$$(12) \quad \begin{aligned} & - \sum_{i,j} \frac{a_{ij}}{2} \frac{\partial^2 |u|^2}{\partial x_i \partial x_j} + \sum_{i,j,k} a_{ij} \frac{\partial u_k}{\partial x_i} \frac{\partial u_k}{\partial x_j} - \\ & - \sum_{k,l} b_{kl} \frac{\partial u_l}{\partial x_i} u_k + \sum_{k,l} c_{kl} u_l u_k = \sum_k f_k u_k . \end{aligned}$$

Let x_0 be a point in $\bar{\Omega}$ such that $|u|^2 \equiv \sum_k |u_k|^2$ attains a maximum at x_0 . If $|u|(x_0) = 0$ then

$$(13) \quad \begin{aligned} ||u||_{\infty} & \leq \frac{1}{\beta - \frac{||b||_{\infty}^2}{4\gamma}} ||f||_{\infty} . \end{aligned}$$

If not, we have

$$(14) \quad \sum_{i,j} \frac{a_{ij}}{2}(x_0) \frac{\partial^2 |u|}{\partial x_i \partial x_j}(x_0) \leq 0$$

and using conditions 1), 3) we get that

$$(15) \quad \gamma \left| \frac{\partial u}{\partial x} \right|^2(x_0) - \sum_i \sum_{k,l} b_{kl,i}(x_0) \frac{\partial u_l}{\partial x_i}(x_0) u_k(x_0) + \beta |u|^2(x_0) \\ \leq |f|(x_0) |u|(x_0).$$

Remarking that

$$(16) \quad \left| \sum_i \sum_{k,l} b_{kl,i}(x) \frac{\partial u_l}{\partial x_i} u_k \right| \leq \|b\|_\infty \left| \frac{\partial u}{\partial x} \right| |u|$$

we obtain from (15) and (16)

$$(17) \quad \gamma \left| \frac{\partial u}{\partial x} \right|^2(x_0) - \|b\|_\infty \left| \frac{\partial u}{\partial x} \right| |u|(x_0) + \beta |u|^2(x_0) \\ \leq |f|(x_0) |u|(x_0).$$

Next, noting that

$$\gamma \left| \frac{\partial u}{\partial x} \right|^2 - \|b\|_\infty \left| \frac{\partial u}{\partial x} \right| |u| + \frac{\|b\|_\infty^2}{4\gamma} |u|^2 \geq 0$$

we get finally

$$(18) \quad |u|(x_0) \leq \frac{1}{\frac{\|b\|_\infty^2}{4\gamma}} \|f\|_\infty$$

which implies (11).

1.3. Existence result

Let us now give the existence result. The proof is made in several steps and use accretive operators methods (see [3]).

In the following we need some additional assumptions on the data :

$$2') \quad a_{ij} \in C^1(\bar{\Omega}) \quad , \quad \partial a \in C^2$$

$$3') \quad c_{kk} \geq \beta > 0 \quad \forall k, 1 \leq k \leq M.$$

Let us denote

$$Au = - \sum_{i,j} \frac{\partial}{\partial x_i} (a_{ij} \frac{\partial u}{\partial x_j})$$

we have :

Theorem 1.2 - Assume that 1), 2), 2'), 3) 3') hold. Then there exists a unique solution of (*)

$$(19) \quad u_k \in W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega) \quad , \quad 2 \leq p < +\infty.$$

Proof :

. Step 1 : Approximation of (*)

Let $\beta_\eta(t)$ be a smooth function defined by

$$(20) \quad \beta_\eta(t) = \begin{cases} t & \text{if } |t| \leq 1/\eta - 1 \\ 1/\eta & \text{if } |t| > 1/\eta \end{cases}$$

$$0 \leq \beta_\eta \leq 1 \quad , \quad 0 < \eta < 1 \quad , \quad \eta \text{ fixed.}$$

Then we consider the problem $(*)_\eta$

$$(*)_n \quad \begin{cases} Au_k^n + \sum_l c_{kl} u_l^n = f_k - \beta_n (-\sum_l \sum_i b_{kl,i} \frac{\partial u_l^n}{\partial x_i}) & \text{a.e. in } \Omega \\ u_k^n|_{\partial\Omega} = 0 \end{cases}$$

where

$$\tilde{b}_{kl,i} = \begin{cases} b_{kl,i} & \text{for } k \neq l \\ b_{kk,i} - \sum_j \frac{\partial a_{ij}}{\partial x_j} & \text{for } k = l \end{cases}$$

We have that the operator A with domain

$$\mathcal{D}(A) = \{u \in [L^2(\Omega)]^M \mid u_k \in W_0^{1,2}(\Omega) \text{ with } Au_k \in L^2(\Omega)\}$$

is m -accretive in $[L^2(\Omega)]^M$. Indeed, we have

$$(21) \quad \begin{aligned} [u - \bar{u}, Au - A\bar{u}]_+ &= \sum_k [u_k - \bar{u}_k, Au_k - A\bar{u}_k]_+ \\ &= \sum_k \int_0 \text{sign}(u_k - \bar{u}_k) |u_k - \bar{u}_k| \left(-\sum_{i,j} \frac{\partial}{\partial x_i} (a_{ij} \frac{\partial u_k - \bar{u}_k}{\partial x_j}) \right) dx \\ &= \sum_k \int_0 \sum_{i,j} a_{ij} \frac{\partial u_k - \bar{u}_k}{\partial x_i} \frac{\partial u_k - \bar{u}_k}{\partial x_j} dx \geq 0 \end{aligned}$$

and $R(I + \lambda A) = L^2(\Omega)$ for any $\lambda > 0$.

On the other hand, let

$$(22) \quad Cu \equiv (C_1 u, C_2 u, \dots, C_M u)$$

$$C_k u \equiv \sum_l c_{kl} u_l$$

we now prove that C is an accretive operator in $[L^2(\Omega)]^M$.

Indeed C is everywhere defined, Lipschitz continuous operator with range in $[L^2(\Omega)]^M$ and

$$(23) \quad \begin{aligned} [u - \bar{u}, Cu - C\bar{u}]_+ &= \sum_{k=1}^M [u_k - \bar{u}_k, c_k u - c_k \bar{u}]_+ \\ &= \sum_k \int_0 \text{sign}(u_k - \bar{u}_k) |u_k - \bar{u}_k| (\sum_l c_{kl} (u_l - \bar{u}_l)) dx \\ &= \int_0 \sum_{k,l} c_{kl} (u_k - \bar{u}_k) (u_l - \bar{u}_l) dx \geq 0 \end{aligned}$$

by condition 3).

Now from known results on the perturbation of m -accretive operator ([3]) we have that

$$(24) \quad A + C \text{ is } m\text{-accretive in } [L^2(\Omega)]^M.$$

. Step 2 : Existence result for (*)

We define a mapping from $[W_0^{1,p}(\Omega)]^M$ into itself by

$$(25) \quad S_\eta v = u^\eta$$

where u_k^η is the unique solution in $W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$, $2 \leq p < +\infty$ of the problem

$$(26) \quad \begin{cases} Au_k^\eta + \sum_l c_{kl} u_l^\eta = f_k - \beta_\eta (-\sum_l \sum_i \tilde{b}_{kl,i} \frac{\partial v}{\partial x_i}) \\ u_k^\eta|_{\partial\Omega} = 0. \end{cases}$$

Indeed, since $A+C$ is m -accretive in $[L^2(\Omega)]^M$ there exists a unique solution of (26) in $W_0^{2,1}(\Omega)$ with $Au_k \in L^2(\Omega)$. On the other hand, we have

$$(27) \quad |\beta_\eta| \leq 1/\eta$$

multiplying (26)a) by $u_k^\eta |u_k^\eta|^{p-2}$ and integrating over Ω , give us

$$(28) \quad \beta \frac{||u_k^\eta||}{L^p} \leq C \frac{||u_k^\eta||}{L^p} + C_0(\eta, p) \quad (*).$$

But $u_k^\eta \in W^{2,2}(\Omega)$ implies that $u_k^\eta \in W^{1,\bar{p}}(\Omega)$ for some $\bar{p} > 2$.

Therefore the right hand of (26)a) is in $L^{p \wedge \bar{p}}(\Omega)$ and from regularity results of elliptic equations

$$(29) \quad u_k^\eta \in W^{2,p \wedge \bar{p}}(\Omega).$$

Now by a bootstrap argument we get that

$$(30) \quad u_k^\eta \in W^{2,p}(\Omega) \text{ for each } p : 2 \leq p < +\infty$$

and

$$(31) \quad \frac{||u_k^\eta||}{W^{2,p}(\Omega)} \leq C_1(\eta, p)$$

(the constants C_i do not depend on v).

Next consider the set

$$(32) \quad D_\eta = \{v \in [W_0^{1,p}(\Omega)]^M \mid ||v||_{W^{2,p}} \leq C_1(\eta, p), \quad 2 \leq p < +\infty\}$$

which is a convex compact subset of $[W_0^{1,p}(\Omega)]^M$. D_η is mapped into itself under S_η and the smoothness of β_η implies that S_η is continuous. Therefore, it follows from Schauder's theorem that S_η has a fixed point.

(*) we take β large enough.

. Step 3 : L^∞ -a priori estimates

Using the same technic as in the proof of theorem 1.1 and the fact that

$$(33) \quad |\beta_\eta(t)| \leq |t|$$

we get that

$$(34) \quad \|\mathbf{u}_\eta\|_{L^\infty} \leq \frac{1}{\beta - \frac{\|\mathbf{b}\|_\infty^2 + 4\gamma C}{4\gamma}} \|\mathbf{f}\|_{L^\infty} .$$

. Step 4 : $W^{2,p}$ - a priori estimates

From the first equation in $(*)_\eta$ we get that

$$(35) \quad \|A\mathbf{u}^\eta\|_{L^p} \leq \|\mathbf{f}\|_{L^p} + C_1(\|\mathbf{u}^\eta\|_{W^{1,p}} + \|\mathbf{u}^\eta\|_{L^p})$$

and using the Sobolev interpolation inequality

$$(36) \quad \|\mathbf{u}^\eta\|_{W^{1,p}} \leq C_2(\delta \|\mathbf{u}^\eta\|_{W^{2,p}} + \frac{1}{\delta} \|\mathbf{u}^\eta\|_{L^p}), \quad \delta > 0$$

we obtain

$$(37) \quad \|A\mathbf{u}^\eta\|_{L^p} \leq \|\mathbf{f}\|_{L^p} + C_3 \delta \|\mathbf{u}^\eta\|_{W^{2,p}} + \frac{C_3}{\delta} \|\mathbf{u}^\eta\|_{L^p} .$$

Now, using the estimate (34) we get

$$(38) \quad \|A\mathbf{u}^\eta\|_{L^p} \leq \frac{C_4}{\delta} \|\mathbf{f}\|_{L^\infty} + C_3 \delta \|\mathbf{u}^\eta\|_{W^{2,p}} .$$

On the other hand, from the standard elliptic theory we have

$$(39) \quad \|\mathbf{u}^\eta\|_{W^{2,p}} \leq C_5(\|A\mathbf{u}^\eta\|_{L^p} + \|\mathbf{u}^\eta\|_{L^p})$$

Therefore, combining (38) and (39) we get

$$(40) \quad ||u^n||_{W^{2,p}} \leq \frac{C_6}{\delta} ||f||_{L^\infty} + C_6 \delta ||u^n||_{W^{2,p}}$$

and for δ well chosen we get

$$(41) \quad ||u^n||_{W^{2,p}} \leq C ||f||_{L^\infty}$$

where C does not depend on n .

. Step 5 : Passage to limit

Now, from (41) there exists a subsequence u_n and a function u such that

$$u_n \xrightarrow{\text{weakly}} u_k \text{ in } W^{2,p}(\Omega) \text{ and in } C(\bar{\Omega})$$

(42)

$$\frac{\partial u_n}{\partial x} \xrightarrow{\text{weakly}} \frac{\partial u_k}{\partial x} \text{ in } C(\bar{\Omega}).$$

Next, set $\eta = \eta_n$ and let $\eta_n \rightarrow 0$ in $(*)_n$ yields the existence result for $(*)$.

. Step 6 : Uniqueness

Let $\tilde{u} \in W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$ be another solution of $(*)$. Then we get that

$$(43) \quad - \sum_{i,j} a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} (u_k - \tilde{u}_k) - \sum_{l,i} b_{kl,i} \frac{\partial}{\partial x_i} (u_l - \tilde{u}_l) + \sum_l c_{kl} (u_l - \tilde{u}_l) = 0 \quad \text{a.e. in } \Omega$$

and by the maximum principle result we deduce that

$$(44) \quad u - \tilde{u} \equiv 0.$$

which completes the proof.

Remark 1.1 : From this proof we see that when $p = 2$ it is enough to assume condition 3). But as we shall see below condition 3') is not a restriction at least when Ω is bounded.

II - THE PROBABILISTIC RESULTS

II.1. The Itô Random evolution

Let $\Omega = C([0, \infty); \mathbb{R}^N)$ be the space of continuous functions on $[0, \infty)$ into \mathbb{R}^N .

Given $t \geq 0$, $\omega \in \Omega$, let $y(t, \omega)$ denotes the position of ω at time t .

For $t \geq s \geq 0$, we consider the σ -fields

$$\mu_t^s = \mathcal{B}(y(\lambda); s \leq \lambda \leq t)$$

$$\mu^s = \mu_\infty^s .$$

We assume that 1) holds and

$$4) \quad b_{kl,i}, c_{kl} \in L^\infty(\mathbb{R}^N) ; a_{ij} \in W^{1,\infty}(\mathbb{R}^N)$$

$$5) \quad a_{ij} = a_{ji} \quad (\text{which is not a restriction since } a_{ij} \in W^{1,\infty}(\mathbb{R}^N)).$$

Then in [4] it was shown that, for x, s fixed, there exists a unique probability measure $Q_{x,s}$ on (Ω, μ^s) such that

i) $Q_{x,s}(y(s) = x) = 1$

ii) $\phi(y(t)) + \int_s^t A\phi(y(\lambda))d\lambda$ is a $Q_{x,s}$ -martingale
 $\forall \phi \in \mathcal{D}(\mathbb{R}^N)$,

where

$$(45) \quad A\phi(x) = - \sum_{i,j} a_{ij}(x) \frac{\partial^2 \phi}{\partial x_i \partial x_j}(x).$$

Moreover, there exists a standard Wiener process $w(t)$ such that

$$y(t) = x + \int_s^t \sigma(y(\lambda))dw(\lambda), \text{ a.s. } - Q_{x,s}, \quad t \geq s$$

where $\sigma(x)$ is such that $a = \frac{1}{2}\sigma^2$.

Let us consider the system $F = (\Omega, \mu_t^s, Q_{x,s}, y(t))$, then we have the definition, for $z \in \mathbb{R}^M$ fixed :

Definition 2.1 - A stochastic process $\xi_z(t)$ is called the Itô random evolution associated to F with respect to the coefficients $b_i, c, 1 \leq i \leq N$, if

1) $\xi_z(t)$ is μ_t^s -adapted and continuous in t with probability one

2) $Q_{x,s}(\xi_z(s)=z) = 1$

$$3) d\xi_z(t) = -\xi_z(t)c(y(t))dt + \frac{1}{2} \sum_{i,j=1}^N \xi_z(t)(a^{-1})_{ij}(y(t)) b_j(y(t))dy_i(t)$$

a.s. $Q_{x,s}$, $t \geq s$.

we have the result

Theorem 2.1 - Under conditions 1), 4), 5) there exists a unique, in the a.s.- $Q_{s,x}$ sense, Itô random evolution.

The proof of this result is similar to the one given in [1].

Under some additional condition we shall prove that $\xi_z(t)$ decreases exponentially as $t \rightarrow \infty$. More precisely, let us assume that

$$3) \quad \sum_{k,\ell} c_{k,\ell} z_k z_\ell \geq \beta |z|^2, \quad \forall z \in \mathbb{R}^M$$

then we have

Theorem 2.2 - Assume that 1), 2), 3), 4), 5) hold. Then we have the estimate

$$(46) \quad E_{x,s} |\xi_z(t)|^2 \leq |z|^2 \exp[-(\beta - \frac{\|b\|_\infty^2}{4\gamma})(t-s)]$$

$\forall t \geq s \text{ and } z \in \mathbb{R}^M.$

Proof : Indeed, we have

$$(47) \quad d\xi_z(t)\xi_z^*(t) = \frac{1}{2} \xi_z(t) \left[\sum_{i,j} (a^{-1})_{ij} (y(t)) b_j^*(y(t)) \xi_z^*(t) \right] dy_i(t) +$$

$$+ \frac{1}{2} \left[\sum_{i,j} (a^{-1})_{ij} (y(t)) \xi_z(t) b_j(y(t)) \right] \xi_z^*(t) dy_i(t) -$$

$$- \xi_z(t) c^*(y(t)) \xi_z^*(t) dt - \xi_z(t) c(y(t)) \xi_z^*(t) dt +$$

$$+ \frac{1}{4} \sum_{i,j} (a^{-1})_{ij} \xi_z(t) b_j(y(t)) b_i^*(y(t)) \xi_z^*(t) dt.$$

Hence, as $a^{-1} \leq \frac{1}{\gamma}$ we get that

$$(48) \quad E_{x,s} |\xi_z(t)|^2 \leq |z|^2 - E_{x,s} \int_s^t \xi_z(\lambda) [c(y(\lambda) + c^*(y(\lambda))] \xi_z^*(\lambda) d\lambda + \\ + \frac{1}{4\gamma} \|b\|_\infty^2 E_{x,s} \int_s^t |\xi_z(\lambda)|^2 d\lambda.$$

Using condition (3) we get

$$(49) \quad E_{x,s} |\xi_z(t)|^2 \leq |z|^2 + (\frac{1}{4\gamma} \|b\|_\infty^2 - \beta) \int_s^t E_{x,s} |\xi_z(\lambda)|^2 d\lambda.$$

By Gronwall's lemma we obtain (46).

II.2. A representation result

Our objective here is to give a probabilistic interpretation of the functions u_k .

To this end we will rely on the results of the analytic part.

We will be interested only in the case of $s = 0$ and denote $Q_x = Q_{x,0}$.

Let τ be the first exit time of the process $y(t)$ from the domain O .

$$(50) \quad \tau = \inf\{t \geq 0 : y(t) \notin O\}$$

which is an μ_t^0 -stopping time.

Our aim is to prove that

$$(51) \quad u_k(x) = E_x \int_0^\tau \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda$$

where e_k is the unit vector in the direction coordinate x_k .

We have :

Theorem 2.3 - Assume that 1), 2), 2'), 3), 3'), 4), 5) hold. Then we have

$$(52) \quad u_k^n(x) = E_x \int_0^T \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda, \quad 1 \leq k \leq M.$$

Proof : Let u_k^n be functions belonging in $\mathcal{D}(\bar{\Omega})$ such that

$$(53) \quad u_k^n \rightarrow u_k \text{ in } W^{2,p}(\bar{\Omega}).$$

Since $p > N$, we have

$$(54) \quad Du_k^n \rightarrow Du_k \text{ in } C(\bar{\Omega})$$

$$u_k^n \rightarrow u_k \text{ in } C(\bar{\Omega}).$$

By defining u_k^n outside $\bar{\Omega}$ in a convenient way, we can assume that $u_k^n \in \mathcal{D}(\mathbb{R}^N)$.

We have the vector version of Itô's formula which is derived from component-wise considerations,

$$(55) \quad du^n(y(t)) = \sum_i \frac{\partial u^n}{\partial x_i}(y(t)) dy_i(t) + \sum_{i,j} a_{ij}(y(t)) \frac{\partial^2 u^n}{\partial x_i \partial x_j}(y(t)) dt$$

a.s. Q_x

On the other hand,

$$(56) \quad d\xi_{e_k}(t) = -\xi_{e_k}(t) c(y(t)) dt + \frac{1}{2} \sum_{i,j} \xi_z(t) (a^{-1})_{ij}(y(t)) b_j(y(t))$$

$$dy_i(t) \quad \text{a.s. } Q_x.$$

Therefore, we have

$$\begin{aligned}
 (57) \quad d\xi_{e_k}(t)u^n(y(t)) &= \sum_{i,j} \xi_{e_k}(t)((a^{-1})_{ij}(y(t))b_j(y(t)) + \\
 &\quad + \frac{\partial u^n}{\partial x_i}(y(t)))dy_i(t) + \\
 &\quad + \xi_{e_k}(t) [\sum_{i,j} a_{ij}(y(t)) \frac{\partial^2 u^n}{\partial x_i \partial x_j}(y(t)) + \\
 &\quad + \sum_i b_i(y(t)) \frac{\partial u^n}{\partial x_i}(y(t)) - c(y, t)u^n(y(t))]dt.
 \end{aligned}$$

Hence, we get that for $T > 0$ fixed

$$\begin{aligned}
 (58) \quad E_x \xi_{e_k}(T\wedge\tau)u^n(y(T\wedge\tau)) &= u_k^n(x) + E_x \int_0^{T\wedge\tau} \xi_{e_k}(\lambda) \\
 &\quad [\sum_{i,j} a_{ij}(y(\lambda)) \frac{\partial^2 u^n}{\partial x_i \partial x_j}(y(\lambda)) + \sum_i b_i(y(\lambda)) \frac{\partial u^n}{\partial x_i}(y(\lambda)) - \\
 &\quad - c(y(\lambda))u^n(y(\lambda))]d\lambda.
 \end{aligned}$$

Remark that

$$\begin{aligned}
 (59) \quad E_x \int_0^{T\wedge\tau} |\xi_{e_k}(\lambda)| |Lu^n(y(\lambda)) - Lu(y(\lambda))| d\lambda &\leq \\
 &\leq (E_x [\int_0^T |\xi_{e_k}(\lambda)|^q d\lambda])^{1/q} (E_x [\int_0^T |L(u^n - u)(y(\lambda))|^p d\lambda])^{1/p}
 \end{aligned}$$

with $1/q + 1/p = 1$. Using the L_p -estimate we get that

$$(60) \quad E_x [\int_0^T |L(u^n - u)(y(\lambda))|^p d\lambda] \leq C_{T,p} ||L(u^n - u)||_{L^p} \rightarrow 0$$

as $n \rightarrow \infty$

and as we have by elementary calculation

$$(61) \quad E_x \left[\int_0^T |\xi_{e_k}(\lambda)|^q d\lambda \right] \leq T \exp\left(\frac{T}{2}q(q-1)[||\sigma b||_\infty^2 + ||c||_\infty]\right).$$

we can let $n \rightarrow \infty$ in (58) and we get that

$$(62) \quad E_x \xi_{e_k}(T \wedge \tau) u(y(T \wedge \tau)) = u_k(x) + E_x \left(\int_0^{T \wedge \tau} \xi_{e_k}(\lambda) \right. \\ \left[\sum_{i,j} a_{ij}(y(\lambda)) \frac{\partial^2 u}{\partial x_i \partial x_j}(y(\lambda)) + \sum_i b_i(y(\lambda)) \frac{\partial u}{\partial x_i}(y(\lambda)) - \right. \\ \left. - c(y(\lambda)) u(y(\lambda)) d\lambda \right].$$

Using equation (*), we obtain

$$(63) \quad E_x \xi_{e_k}(T \wedge \tau) u(y(T \wedge \tau)) = u_k(x) - E_x \int_0^{T \wedge \tau} \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda.$$

Furthermore as u, f are bounded by deterministic constant, we can let $T \rightarrow \infty$ in (63), thus.

$$(64) \quad E_x \int_0^{T \wedge \tau} \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda \rightarrow E_x \int_0^T \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda \text{ as } T \rightarrow \infty$$

and using the estimate (46) we obtain that

$$(65) \quad E_x \xi_{e_k}(T \wedge \tau) u(y(T \wedge \tau)) = E_x \xi_{e_k}(T) u(y(T)) \chi_{T < \tau} \rightarrow 0 \text{ as } T \rightarrow \infty.$$

Therefore it follows that

$$(66) \quad u_k(x) = E_x \int_0^T \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda$$

which completes the proof.

Remark 2.1 : When $O \in \mathbb{R}^N$ we have the representation of the solution of (*) given by

$$(67) \quad u_k(x) = E_x \int_0^\infty \xi_{e_k}(\lambda) f(y(\lambda)) d\lambda$$

which makes sense using the estimate (46).

Remark 2.2 : When Ω is bounded, condition (3) is not a restriction. Indeed, making the following change of unknown functions

$$(68) \quad u_k = \omega v_k$$

with

$$(69) \quad \omega(x) = 2 - \exp[-\alpha(|x-x_0|^2 + 1)^{1/2}]$$

where $x_0 \notin \bar{\Omega}$ and α is a constant to be chosen sufficiently large.

Hence we obtain the following system

$$(70) \quad \begin{aligned} & - \sum_{i,j} a_{ij} \frac{\partial^2 v_k}{\partial x_i \partial x_j} - \sum_{i,l} b_{kl,i} \frac{\partial v_l}{\partial x_i} - \sum_{i,j} \frac{2}{\omega} \frac{\partial \omega}{\partial x_j} a_{ij} \frac{\partial v_k}{\partial x_i} + \\ & + \sum_l (c_{kl} v_l - \frac{1}{\omega} \sum_{i,j} a_{ij} \frac{\partial^2 \omega}{\partial x_i \partial x_j} v_k + \frac{1}{\omega} \sum_{i,j} b_{kl,i} \frac{\partial \omega}{\partial x_i} v_l) \\ & = \frac{1}{\omega} f_k \end{aligned}$$

which verifies condition (3) for adequate α .

REFERENCES

- [1] D.W. STROOCK, On certain systems of parabolic equations, Comm. on Pure and Applied Math., Vol. 23 (1970).
- [2] E. FABES, N. RIVIERE, Systems of parabolic equations with uniformly continuous coefficients, J. Anal. Math., Vol. 17 (1966).
- [3] V. BARBU, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff Int. Publishing, 1976.
- [4] D.W. STROOCK, S.R.S. VARADHAN, Diffusion processes with continuous coefficients I,II, Comm. Pure and Appl. Math., Vol. 22 (1969).

Imprimé en France
par
l'Institut National de Recherche en Informatique et en Automatique

