

# Hyperbolic systems in diagonal form with dissipation, in several spaces variables: existence and uniqueness results

O. Bennouna

► **To cite this version:**

O. Bennouna. Hyperbolic systems in diagonal form with dissipation, in several spaces variables: existence and uniqueness results. RR-0452, INRIA. 1985. inria-00076103

**HAL Id: inria-00076103**

**<https://hal.inria.fr/inria-00076103>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# IRIA

CENTRE DE ROCQUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP 105  
78153 Le Chesnay Cedex  
France

Tel. (1) 39 63 55 11

Rapports de Recherche

N° 452

**HYPERBOLIC SYSTEMS IN  
DIAGONAL FORM WITH  
DISSIPATION, IN SEVERAL  
SPACES VARIABLES:  
EXISTENCE AND UNIQUENESS  
RESULTS**

Omar BENNOUNA

Novembre 1985

HYPERBOLIC SYSTEMS IN DIAGONAL FORM  
WITH DISSIPATION, IN SEVERAL SPACES VARIABLES :  
- EXISTENCE AND UNIQUENESS RESULTS

Omar BENNOUNA (\*)

(\*) BEL Laboratories, 38 allée des Roses - Fes - Maroc

## Abstract

In this paper we consider a class of hyperbolic systems in diagonal form with dissipation. We study the Cauchy problem for which we obtain the existence and uniqueness results of generalized solution i.e. solution in  $W^{1,\infty}$ ; assuming the dissipation term large enough.

Key words : Hyperbolic - Dissipation - Generalized solution;

## Résumé

Dans ce papier on considère une classe de systèmes hyperboliques sous forme diagonale avec terme de dissipation. On établit pour le problème de Cauchy l'existence et l'unicité de solution généralisée, c'est-à-dire de solution dans  $W^{1,\infty}$ ; ceci sous la condition que le terme de dissipation est assez grand.

Mots clefs : Hyperbolique - Dissipation - Solution généralisée.

HYPERBOLIC SYSTEMS IN DIAGONAL FORM  
WITH DISSIPATION, IN SEVERAL SPACES VARIABLES :  
EXISTENCE AND UNIQUENESS RESULTS

O. Bennouna

Introduction

It is well known that for the Cauchy problem of first order hyperbolic systems in diagonal form :

$$(*) \quad \left\{ \begin{array}{l} \frac{\partial u_k}{\partial t} + \sum_{i=1}^N \lambda_k^i(t, x, u) \frac{\partial u_k}{\partial x_i} + c_k(t, x, u) u_k = 0 \quad \text{in } \Pi_T \\ u_k(0, x) = v_k(x) \quad \text{in } \mathbb{R}^N, \quad 1 \leq k \leq M \end{array} \right.$$

where

$$u = (u_1, \dots, u_M)$$

$$\Pi_T = ]0, T[ \times \mathbb{R}^N,$$

$v_k(x)$ ,  $\lambda_k^i(t, x, v)$ ,  $c_k(t, x, v)$  are given smooth data; a smooth solution exists only locally in time and singularities i.e. formation of shocks develop in a finite time.

In this note we prove that the addition of dissipation term permit us to obtain the existence and uniqueness of global generalized solution i.e. a Lipschitz continuous solution which satisfies (\*) in the almost everywhere sense.

To obtain the necessary a priori estimates we use a successive approximation method.

For  $N = 1$ , similar results are obtained under other types of dissipation condition in [1], [2].

## I. PRELIMINARY RESULTS

### I.1. Notations, assumptions and definitions

We make the following assumptions about the data :

- (1) The functions  $\lambda_k^i(t, x, v)$ ,  $c_k(t, x, v)$  are bounded and continuous and have bounded derivatives with respect to  $x$  and  $v$ .
- (2)  $v_k(x)$  is bounded continuous and has bounded derivatives.
- (3)  $c_k(t, x, v) \leq -\alpha < 0 \quad \forall (t, x, v) \in \Pi_T \times \mathbb{R}^M$ .

Let us give now the definition of a generalized solution of (\*).

Définition 1.1 :  $u$  is a generalized solution of (\*) if  $u \in [W^{1, \infty}(\Pi_T)]^M$  and satisfies (\*) in the almost everywhere sense.

Let  $y_{x,k}^t(\tau)$  be the  $k$ -th characteristic curve passing through  $(t, x)$ , which satisfies

$$(4) \quad y_{x,k}^t(\tau) = x - \int_{\tau}^t \lambda_k(s, y_{x,k}^t(s), u(s, y_{x,k}^t(s))) ds$$
$$\forall \tau, \quad 0 \leq \tau \leq t \leq T.$$

We shall prove some useful lemma concerning  $y_{x,k}^t(\tau)$ .

### I.2. Some lemma

We have :

Lemma 1 : Assume that (1) holds and let  $u \in [W^{1,\infty}(\Pi_T)]^M$ .

Then

$$(5) \quad \|y_{x,k}^t(\tau) - y_{x+h,k}^t(\tau)\| \leq \|h\| \exp[(t-\tau)\omega_{\lambda_k}(u)] \quad \forall h \in \mathbb{R}^N$$

where

$$\omega_{\lambda_k}(u) = \left\| \frac{\partial \lambda_k}{\partial x} \right\|_{\infty} + M \left\| \frac{\partial \lambda_k}{\partial u} \right\|_{\infty} \left\| \frac{\partial u}{\partial x} \right\|_{\infty}$$

$$\left\| \frac{\partial \lambda_k}{\partial x} \right\|_{\infty} = \max_{s \in [0, T]} \sup_{v \in \mathbb{R}^M} \sup_{x, x' \in \mathbb{R}^N} (|\lambda_k(s, x, v) - \lambda_k(s, x', v)| / \|x - x'\|)$$

$$\left\| \frac{\partial \lambda_k}{\partial u} \right\|_{\infty} = \max_{1 \leq \ell \leq M} \left\| \frac{\partial \lambda_k}{\partial u_{\ell}} \right\|_{\infty}$$

$$\begin{aligned} \left\| \frac{\partial \lambda_k}{\partial u_{\ell}} \right\|_{\infty} &= \max_{s \in [0, T]} \sup_{x \in \mathbb{R}^N} \sup_{v_{\ell}, v'_{\ell} \in \mathbb{R}} (|\lambda_k(s, x, u_1, \dots, v_{\ell}, u_{\ell+1}, \dots) \\ &\quad - \lambda_k(s, x, u_1, \dots, v'_{\ell}, u_{\ell+1}, \dots)| / |v_{\ell} - v'_{\ell}|) \end{aligned}$$

$$\left\| \frac{\partial u}{\partial x} \right\|_{\infty} = \max_{1 \leq \ell \leq M} \left\| \frac{\partial u_{\ell}}{\partial x} \right\|_{\infty}$$

Proof : We consider the function  $\|x\|^2$  on the curve  $y_{x,k}^t(\tau) - y_{x+h,k}^t(\tau)$ , thus we get by integration by part that

$$\begin{aligned} (6) \quad \|y_x(\tau) - y_{x+h}(\tau)\|^2 &= 2 \int_{\tau}^t (y_{x+h}(s) - y_x(s)) \cdot [\lambda_k(s, y_x(s), u(s, y_x(s))) \\ &\quad - \lambda_k(s, y_{x+h}(s), u(s, y_{x+h}(s)))] ds + \|h\|^2 \\ &= 2 \int_{\tau}^t (y_{x+h}(s) - y_x(s)) \cdot [\lambda_k(s, y_x(s), u(s, y_x(s))) \\ &\quad - \lambda_k(s, y_{x+h}(s), u(s, y_x(s)))] ds \end{aligned}$$

$$\begin{aligned}
 & + 2 \int_{\tau}^t (y_{x+h}(s) - y_x(s)) \cdot [\lambda_k(s, y_{x+h}(s), \\
 & \quad u(s, y_x(s))) - \lambda_k(s, y_{x+h}(s), u(s, y_{x+h}(s)))] \\
 & \quad ds + \|h\|^2 \\
 & = I + II + \|h\|^2 .
 \end{aligned}$$

We estimate separately I and II. We have

$$(7) \quad I \leq 2 \left\| \frac{\partial \lambda_k}{\partial x} \right\|_{\infty} \int_{\tau}^t \|y_{x,k}(s) - y_{x+h,k}(s)\|^2 ds$$

$$\begin{aligned}
 (8) \quad II & \leq 2 \sum_{\ell} \left\| \frac{\partial \lambda_k}{\partial u_{\ell}} \right\|_{\infty} \int_{\tau}^t \|y_{x,k}(s) - y_{x+h,k}(s)\| \\
 & \quad |u_{\ell}(s, y_x(s)) - u_{\ell}(s, y_{x+h}(s))| ds \\
 & \leq 2M \left\| \frac{\partial \lambda_k}{\partial u} \right\|_{\infty} \left\| \frac{\partial u}{\partial x} \right\|_{\infty} \int_{\tau}^t \|y_{x,k}(s) - y_{x+h,k}(s)\|^2 ds .
 \end{aligned}$$

Combining (7) and (8) we get that

$$\begin{aligned}
 & \|y_x(\tau) - y_{x+h}(\tau)\|^2 \leq \\
 & \|h\|^2 \exp[2(t-\tau) (\left\| \frac{\partial \lambda_k}{\partial x} \right\|_{\infty} + M \left\| \frac{\partial \lambda_k}{\partial u} \right\|_{\infty} \left\| \frac{\partial u}{\partial x} \right\|_{\infty})]
 \end{aligned}$$

which completes the proof.

Let us denote

$$(9) \quad X_{x,k}^t(\tau) = \exp \int_{\tau}^t c_k(s, y_{x,k}^t(s), u(s, y_{x,k}^t(s))) ds$$

which satisfies the integral equation

$$(10) \quad X_{x,k}^t(\tau) = 1 + \int_{\tau}^t X_{x,k}^t(s) c_k(s, y_{x,k}^t(s), u(s, y_{x,k}^t(s))) ds$$

$$\forall \tau, \quad 0 \leq \tau \leq t \leq T .$$

We have :



Lemma 2 : Assume that (1), (3), (5) hold. Then we have

$$(11) \quad |X_{x,k}^t(\tau) - X_{x+h,k}^t(\tau)| \leq \exp[-(t-\tau)\alpha] \int_{\tau}^t |c_k(s, y_x(s), u(s, y_x(s))) - c_k(s, y_{x+h}(s), u(s, y_{x+h}(s)))| ds$$

$$\forall h \in \mathbb{R}^N.$$

The proof of this result is easy using the fact that

$$|\exp a - \exp b| \leq \max(\exp a, \exp b) |a-b|$$

$$\forall a, b \in \mathbb{R}.$$

## II. EXISTENCE RESULT

To prove the existence of a generalized solution of (\*) we shall use the successive approximations method.

For  $w \in [W^{1,\infty}(\Pi_T)]^M$  let us denote

$$(Tw)_k(t, x) = X_{x,k}^t(0) v(y_{x,k}^t(0)) \quad , \quad 1 \leq k \leq M,$$

where  $y_{x,k}^t(\tau)$  is the solution of the integral equation

$$y_{x,k}^t(\tau) = x - \int_{\tau}^t \lambda_k(s, y_{x,k}^t(s), w(s, y_{x,k}^t(s))) ds$$

$$\forall \tau, \quad 0 \leq \tau \leq t \leq T.$$

We define the successive approximations as follows :

$$(12) \quad \left| \begin{array}{l} (S^0(t)v)_k = v_k(x) \\ (S^{n+1}(t)v)_k(x) = (TS^n(\cdot)v)_k(t,x) \quad , \quad n \geq 0 . \end{array} \right.$$

We shall establish some a priori estimates, more precisely we have :

Lemma 3 : Under assumptions (1), (2), (3) and

$$(13) \quad \alpha \geq \gamma_1 + \gamma_2 M \|v\|_{W^{1,\infty}}$$

we have

$$(14) \quad \frac{1}{\|h\|} |S^{n+1}(t)v(x) - S^{n+1}(t)v(x+h)| \leq \|v\|_{W^{1,\infty}}$$

$$\forall n \geq 0 \quad , \quad \forall t \in [0, T] \quad , \quad \forall h, x \in \mathbb{R}^N ,$$

where

$$\gamma_1 = \max\left(\left\|\frac{\partial \lambda}{\partial x}\right\|_{\infty} , \left\|\frac{\partial c}{\partial x}\right\|_{\infty}\right)$$

$$\gamma_2 = \max\left(\left\|\frac{\partial \lambda}{\partial u}\right\|_{\infty} , \left\|\frac{\partial c}{\partial u}\right\|_{\infty}\right)$$

$$|x| = \max_{1 \leq i \leq N} |x_i| \quad \forall x \in \mathbb{R}^N .$$

Proof : Indeed we have

$$(15) \quad \begin{aligned} & |(S(t)^{n+1}v)_k(x) - (S(t)^{n+1}v)_k(x+h)| = \\ & |X_{x,k,n}^t v_k(y_{x,k,n}^t) - X_{x+h,k,n}^t v_k(y_{x+h,k,n}^t)| \\ & \leq |(X_{x,k,n}^t - X_{x+h,k,n}^t) v_k(y_{x,k,n}^t)| + \\ & |(v_k(y_{x,k,n}^t) - v_k(y_{x+h,k,n}^t)) X_{x+h,k,n}^t| \\ & = I + II . \end{aligned}$$

Then, using lemma 1, 2, we get that

$$\begin{aligned}
 I &\leq \|v\|_{\infty} \exp(-\alpha t) \cdot \int_0^t |c_k(s, y_x(s), v(s, y_x(s)) \\
 &\quad - c_k(s, y_{x+h}(s), v(s, y_{x+h}(s)))| ds \\
 &\leq \|v\|_{\infty} \exp(-\alpha t) \cdot \left\| \frac{\partial c}{\partial x} \right\|_{\infty} \int_0^t \|y_{x,k}(s) - y_{x+h,k}(s)\| ds \\
 &\quad + M \|v\|_{\infty} \exp(-\alpha t) \cdot \left\| \frac{\partial c}{\partial u} \right\|_{\infty} \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty} \cdot \\
 &\quad \int_0^t \|y_{x,k}(s) - y_{x+h,k}(s)\| ds .
 \end{aligned}$$

Thus,

$$\begin{aligned}
 (16) \quad I &\leq \|v\|_{\infty} \exp(-\alpha t) \cdot \left( \left\| \frac{\partial c}{\partial x} \right\|_{\infty} + M \left\| \frac{\partial c}{\partial u} \right\|_{\infty} \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty} \right) \cdot \\
 &\quad \int_0^t \|y_{x,k}(s) - y_{x+h,k}(s)\| ds .
 \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
 (17) \quad &\int_0^t \|y_{x,k}(s) - y_{x+h,k}(s)\| ds \leq \\
 &\|h\| \int_0^t \exp[(t-s) \left( \left\| \frac{\partial \lambda_k}{\partial x} \right\|_{\infty} + M \left\| \frac{\partial \lambda_k}{\partial u} \right\|_{\infty} \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty} \right)] ds \leq \\
 &\|h\| \int_0^t \exp[s(\gamma_1 + M\gamma_2 \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty})] ds .
 \end{aligned}$$

Hence,

$$\begin{aligned}
 (18) \quad &\int_0^t \|y_{x,k}(s) - y_{x+h,k}(s)\| ds \leq \|h\| (\gamma_1 + M\gamma_2 \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty})^{-1} \cdot \\
 &\exp[t(\gamma_1 + M\gamma_2 \left\| \frac{\partial S(\cdot)^n v}{\partial x} \right\|_{\infty})]
 \end{aligned}$$

and then, we get

$$(19) \quad I \leq \|h\| \|v\|_{\infty} \exp[(-\alpha + \gamma_1 + M\gamma_2 \|\frac{\partial S(\cdot)^n v}{\partial x}\|_{\infty})t] .$$

We also have, in a similar way

$$(20) \quad II \leq \|\frac{\partial v}{\partial x}\|_{\infty} \|h\| \exp[(-\alpha + \gamma_1 + M\gamma_2 \|\frac{\partial S(\cdot)^n v}{\partial x}\|_{\infty})t] .$$

Combining (19) and (20) we obtain finally

$$\begin{aligned} & |S^{n+1}(t)v(x) - S^{n+1}(t)v(x+h)| \leq \\ & \|h\| \|v\|_{W^{1,\infty}} \exp[(-\alpha + \gamma_1 + M\gamma_2 \|\frac{\partial S^n(\cdot)v}{\partial x}\|_{\infty})t] . \end{aligned}$$

Let us denote

$$(21) \quad \text{and} \quad \left\{ \begin{aligned} L_n(t) &= \|\frac{\partial S^n(t)v}{\partial x}\|_{\infty} \\ \hat{L}_n &= \sup_{0 \leq t \leq T} L_n(t) \end{aligned} \right.$$

thus we have proved that

$$L_{n+1}(t) \leq \|v\|_{W^{1,\infty}} \exp[(-\alpha + \gamma_1 + M\gamma_2 \hat{L}_n)t] .$$

On the other hand, we have

$$L_0(t) = \|\frac{\partial v}{\partial x}\|_{\infty} \leq \|v\|_{W^{1,\infty}}$$

$$\hat{L}_0 \leq \|v\|_{W^{1,\infty}}$$

since (13) holds, we have

$$-\alpha + \gamma_1 + M\gamma_2 \hat{L}_0 \leq 0$$

and then

$$\hat{L}_1 \leq \|v\|_{W^{1,\infty}}.$$

By induction we see that

$$L_n(t) \leq \|v\|_{W^{1,\infty}}$$

which is exactly (14).

We have :

Lemma 4 : Under conditions (1), (2), (3) and (13), we have the estimate

$$(22) \quad \frac{1}{|\theta|} |S^{n+1}(t+\theta)v(x) - S^{n+1}(t)v(x)| \leq \|v\|_{W^{1,\infty}} \max(\|\lambda\|_{\infty}, \|c\|_{\infty}) \exp(-\alpha t)$$

$$\forall t, \theta \in [0, T], \quad n \geq 0, \quad \forall x \in \mathbb{R}^N.$$

Proof :

$$(23) \quad (S^{n+1}(t+\theta)v)_k(x) - (S^{n+1}(t)v)_k(x)$$

$$= X_{x,k,n}^{t+\theta}(0) v_k(y_{x,k,n}^{t+\theta}(0)) - X_{x,k,n}^t(0) v_k(y_{x,k,n}^t(0))$$

$$= \int_t^{t+\theta} X_{x,k,n}^s(0) \left[ \frac{\partial v_k}{\partial x}(y_{x,k,n}^s(0)) \cdot \lambda_k(s, y_{x,k,n}^s(0), (S^n(s)v)(y_{x,k,n}^s(0))) \right. \\ \left. + c_k(s, y_{x,k,n}^s(0), (S^n(s)v)(y_{x,k,n}^s(0))) v_k(y_{x,k,n}^s(0)) \right] ds.$$

Thus,

$$|(S^{n+1}(t+\theta)v)_k(x) - (S^{n+1}(t)v)_k(x)| \leq |\theta| \left( \left\| \frac{\partial v_k}{\partial x} \right\|_{\infty} \|\lambda_k\|_{\infty} + \|v_k\|_{\infty} \|c_k\|_{\infty} \right) \exp(-\alpha t)$$

which implies (22).

Now, from the above a priori estimates we obtain the following existence result.

Theorem 1 : Assume that (1), (2), (3) and (13) hold, then there exists a generalized solution to (\*) with the initial condition  $v(x)$  given by

$$(24) \quad u(t,x) = \lim_{n \rightarrow \infty} S^n(t)v(x)$$

uniformly on any compact subset of  $\mathbb{R}^N$  and uniformly with respect to  $t$ .

Proof : Indeed, from the estimates (14), (22) and the uniform boundedness of the sequence  $S^n(t)v(x)$ , we get that, for some subsequence, there exists

$$u(t,x) \in C^0(\Pi_T)$$

such that (24) holds.

After some calculation, we have

$$(25) \quad \begin{aligned} & \| |y_{x,k,n}^t(\tau) - y_{x,k,m}^t(\tau)| \| \leq \\ & C_1 T (\exp(C_2 T)) \| |S^n(\cdot)v - S^m(\cdot)v| \|_{\infty} \\ & \forall \tau, 0 \leq \tau \leq t \leq T \end{aligned}$$

hence,

$$(26) \quad \lim_{m,n \rightarrow \infty} \| |y_{x,k,m}^t(\tau) - y_{x,k,n}^t(\tau)| \| = 0 \text{ uniformly in } (\tau,x).$$

Thus, there exists  $y_{x,k}^t(\tau)$  such that

$$(27) \quad y_{x,k}^t(\tau) = x - \int_{\tau}^t \lambda_k(s, y_{x,k}^t(s), u(s, y_{x,k}^t(s))) ds$$

and we obtain that

$$(28) \quad u_k(t, x) = X_{x,k}^t(0) v_k(y_{x,k}^t(0)), \quad 1 \leq k \leq M.$$

Finally from (28) and the fact that  $u(t, x)$  is Lipschitz continuous we see that  $u(t, x)$  is a generalized solution of (\*).

### III. UNIQUENESS RESULTS

We shall prove now that  $u(t, x)$  is the unique generalized solution of (\*).

Indeed, let  $v(t, x)$  be another generalized solution of (\*) and let

$$(29) \quad y_{x,k,v}^t(\tau) = x - \int_{\tau}^t \lambda_k(s, y_{x,k,v}^t(s), v(s, y_{x,k,v}^t(s))) ds$$

then

$$(30) \quad v_k(t, x) = X_{x,k,v}^t(0) v_k(y_{x,k,v}^t(0)) + \int_0^t X_{x,k,v}^s(0) \left[ \frac{\partial v_k}{\partial t}(s, y_{x,k,v}^s(0)) + \lambda_k(s, y_{x,k,v}^s(0), v(s, y_{x,k,v}^s(0))) \frac{\partial v_k}{\partial x}(s, y_{x,k,v}^s(0)) + c_k(s, y_{x,k,v}^s(0), v(s, y_{x,k,v}^s(0))) v_k(s, y_{x,k,v}^s(0)) \right] ds$$

$$= X_{x,k,v}^t(0)$$

On the other hand, we have

$$(31) \quad \|u(t) - v(t)\|_{\infty} \leq c_1 \int_0^t \|y_{x,k,u}^t(s) - y_{x,k,v}^t(s)\| ds + c_2 \int_0^t \|u(s) - v(s)\|_{\infty} ds$$

and

$$(32) \quad \begin{aligned} & \|y_{x,k,u}^t(\tau) - y_{x,k,v}^t(\tau)\| \leq \\ & c_3 \int_{\tau}^t \|u(s) - v(s)\|_{\infty} + c_4 \int_{\tau}^t \|y_{x,k,u}^t(s) - y_{x,k,v}^t(s)\| ds . \end{aligned}$$

Hence

$$(33) \quad \|u(t) - v(t)\|_{\infty} \leq c_1 e^{c_2 t} \int_0^t \|y_{x,k,u}^t(s) - y_{x,k,v}^t(s)\| ds$$

and

$$(34) \quad \|y_{x,k,u}^t(\tau) - y_{x,k,v}^t(\tau)\| \leq c_3 e^{c_2 t} \int_{\tau}^t \|u(s) - v(s)\|_{\infty} ds .$$

Combining (33) and (34) yields

$$\|u(t) - v(t)\|_{\infty} \leq c_1 c_3 e^{(c_2 + c_4)t} \int_0^t \int_s^t \|u(\bar{s}) - v(\bar{s})\|_{\infty} d\bar{s} ds$$

and then,

$$\|u(t) - v(t)\|_{\infty} \leq c_5 t e^{c_6 t} \int_0^t \|u(s) - v(s)\|_{\infty} ds$$

this yields

$$(35) \quad \|u(t) - v(t)\|_{\infty} = 0 \quad \forall t \in [0, T]$$

which completes the proof.

Remark 1 : We can prove similar results for the problems :

$$(36) \quad \left| \begin{aligned} & \frac{\partial u_k}{\partial t} + \sum_{i=1}^N \lambda_k^i(t, x, u) \frac{\partial u_k}{\partial x_i} + \sum_{\ell=1}^M c_{k\ell} u_{\ell} = 0 \\ & u_k(0, x) = v_k(x) \end{aligned} \right.$$



under the supplementary condition :

$$(37) \quad \text{the matrix } (c_{kl}) \text{ is commuting with the diagonal matrices } (\lambda_k^i), \\ 1 \leq i \leq N.$$

Then using the same technic as above yields the result.

Remark 2 : Let us consider the quasilinear hyperbolic system

$$(38) \quad \begin{cases} \frac{\partial u_1}{\partial t} - \frac{\partial u_2}{\partial x} = 0 \\ \frac{\partial u_2}{\partial t} - \frac{\partial \sigma(u_1)}{\partial x} = 0 \end{cases}$$

which arises in fluid mechanics, nonlinear elasticity, and other fields of continuum mechanics. With the change of variables

$$(39) \quad u_2 = \frac{\partial w}{\partial t}, \quad u_1 = \frac{\partial w}{\partial x}$$

(38) is equivalent to the second-order nonlinear wave equation

$$\frac{\partial^2 w}{\partial t^2} - \frac{\partial}{\partial x} \sigma\left(\frac{\partial w}{\partial x}\right) = 0.$$

For (38) it is known that there exist functions  $r$  and  $s$  called Riemann invariants (see [3], [4])

$$(40) \quad \begin{cases} r = u_2 + \int_0^{u_1} \sigma'(v)^{1/2} dv \\ s = u_2 - \int_0^{u_1} \sigma'(v)^{1/2} dv. \end{cases} \quad (*)$$

As we can see,  $r$  and  $s$  satisfy

(\*) we assume  $\sigma' > 0$ .

$$(41) \quad \left| \begin{array}{l} \frac{\partial r}{\partial t} - \sigma'(u_1)^{1/2} \frac{\partial r}{\partial x} = 0 \\ \frac{\partial s}{\partial t} + \sigma'(u_1)^{1/2} \frac{\partial s}{\partial x} = 0 \end{array} \right.$$

Hence if we consider the perturbed problem

$$(42) \quad \left| \begin{array}{l} \frac{\partial r_\varepsilon}{\partial t} - \sigma'(u_{1\varepsilon})^{1/2} \frac{\partial r_\varepsilon}{\partial x} - \varepsilon r_\varepsilon = 0 \\ \frac{\partial s_\varepsilon}{\partial t} + \sigma'(u_{1\varepsilon})^{1/2} \frac{\partial s_\varepsilon}{\partial x} - \varepsilon s_\varepsilon = 0 \end{array} \right.$$

then theorem 1 implies the existence and uniqueness of a generalized solution of (42) with smooth Cauchy data for  $\varepsilon$  large enough.

#### REFERENCES

- [1] T. NISHIDA, "Global smooth solutions for the second-order quasilinear wave equations with first-order dissipation", To appear.
- [2] TA-TSIEN LI, LING HSIAO, "Global smooth solution of the Cauchy problem for a class of quasilinear hyperbolic systems", Rapport de Recherche INRIA, n° 83, juin 1981.
- [3] R. COURANT, K.O. FRIEDRICHS, "Supersonic flow and shock waves", New York, Interscience, (1948).
- [4] P.D. LAX, "Hyperbolic systems of conservation laws II", Comm. Pure Appl. Math., 10 (1957).

Imprimé en France  
par  
l'Institut National de Recherche en Informatique et en Automatique

