

**Matrix-geometric stationary distribution for the
PH/PH/1R queue**

P.P. Bocharov, V.A. Naumov

► **To cite this version:**

P.P. Bocharov, V.A. Naumov. Matrix-geometric stationary distribution for the PH/PH/1R queue. RR-0304, INRIA. 1984. <inria-00076253>

HAL Id: inria-00076253

<https://hal.inria.fr/inria-00076253>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IRIA

CENTRE DE ROCQUENCOURT

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
BP 105
78153 Le Chesnay Cedex
France
Tél. (3) 954 90 20

Rapports de Recherche

N° 304

**MATRIX-GEOMETRIC
STATIONARY DISTRIBUTION
FOR THE PH/PH/1/r QUEUE**

**P. P. BOCHAROV
V. A. NAUMOV**

Mai 1984

**MATRIX-GEOMETIC STATIONARY DISTRIBUTION
FOR THE PH/PH/1/r QUEUE**

P.P. BOCHAROV and V.A. NAUMOV

Department of Mathematics

University of Friendship Patrice Lumumba

Moscow

USSR

ABSTRACT

This paper is devoted to the study of the PH/PH/1/r queueing systems. Using matrix calculus together with the stochastic interpretation of distributions of PH-type, we obtain the steady-state probabilities in a matrix-geometric form. Numerical examples are also given.

RESUME

Cet article analyse la file d'attente PH/PH/1/r. En utilisant le calcul matriciel ainsi que l'interprétation probabiliste de la distribution de type PH, nous obtenons les probabilités stationnaires sous forme "géométrique matricielle". Des exemples numériques sont présentés.

N.B.

Ce travail a été terminé pendant la visite du professeur P.P. BOCHAROV à l'INRIA du 9/3/84 au 9/4/84, dans le projet "Modélisation et évaluation des systèmes informatiques" - responsable scientifique G. FAYOLLE.

1 - INTRODUCTION

A single server queueing system (QS) with a buffer of finite capacity r ($1 \leq r < \infty$) is studied. The customers arrive according to a renewal process with probability distribution function (P.D.F.) $A(x)$. The service times of customers are i.i.d. random variables with P.D.F. $B(x)$. An arriving customer finding a full buffer is lost. Using Kendall's notation, this QS is denoted GI/GI/1/r.

A huge literature has been devoted to special cases of this QS. In the present paper we shall consider a situation, where steady-state distribution is given in a matrix-geometric form [1-6]. In the past, such a form has been obtained by Basharin [1,2] for M/HM/1/r $\leq \infty$ (HM \equiv hyperexponential). The system M/HK/1/r has been analysed in [4] (K \equiv coxian, i.e. rational Laplace transform). The matrix-geometric solution for steady-state probabilities in the QSs M/PH/1/r and PH/M/c/r was given by Neuts [5]. The symbols "PH" denote a distribution of phase type [7]. Recent results have been obtained in [6] for the QS HM/HM/1/r. They rely on some arguments developed in [3], where the arrival process is considered as one server QS of type HM.

In the present paper we analyse the system PH/PH/1/r. The main result tells that the steady-state probabilities have a matrix-geometric form. In section 2 we briefly mention the notations and some properties of the PH-distribution, which are crucial in the proof of the main theorem of section 3. Numerical examples are also given.

2 - PH-distribution

A P.D.F. $F(x)$ of a non-negative random variable is called a PH-distribution [7] if it can be written as

$$F(x) = \bar{f}^T e^{Gx} \bar{1}, \quad x \geq 0, \quad (1)$$

where:

- the m -dimensional row vector \bar{f}^T satisfies

$$\sum_{j=1}^m f_j \leq 1, f_j \geq 0, j = \overline{1, m};$$

- G is an $(m \times m)$ -matrix such that

$$\sum_{j=1}^m G_{ij} \leq 0, G_{ij} \geq 0, G_{ii} < 0, i, j = \overline{1, m};$$

- $\bar{1}$ denotes the column vector having all its components equal to one.

The pair (\bar{f}, G) is then called the PH-representation of order m of the P.D.F. $F(x)$.

A P.D.F. $F(x)$ of the PH-type has a probabilistic interpretation which we show right now.

Let v_1, \dots, v_m be real numbers,

$$v_i \geq -G_{ii}, i = \overline{1, m}.$$

Define $\theta_{ij}, i, j = \overline{1, m}$,

$$\theta_{ij} = \begin{cases} 1 + \frac{G_{ii}}{v_i}, & i = j, \\ \frac{G_{ij}}{v_i}, & i \neq j. \end{cases}$$

Then

$$\sum_{j=1}^m \theta_{ij} \leq 1, \theta_{ij} \geq 0, i, j = \overline{1, m}.$$

Consider an open network with m nodes as shown on Figure 1.




Figure 1

An arriving customer enters node i , $i = \overline{1, m}$, with probability f_i , or is rejected with probability

$$f_0 \stackrel{\text{def}}{=} 1 - \sum_{j=1}^m f_j.$$

His sojourn time at node i has an exponential distribution with parameter ν_i . After completing service at node i , this customer enters node j with probability θ_{ij} and with probability

$$\theta_{i0} \stackrel{\text{def}}{=} 1 - \sum_{j=1}^m \theta_{ij}$$

he leaves the network. Moreover we assume that there is always at most one single customer in this network and τ will denote his sojourn time. Let $\xi(t)$ be the index of the node visited by this customer at time t . The process $\xi(t)$ defined only for $t \in [0, \tau)$ is an homogeneous

terminating Markov process with generator G, where

$$G_{ij} = \begin{cases} v_i(\theta_{ii} - 1), & i = j, \\ v_i\theta_{ij}, & i \neq j. \end{cases}$$

Then, $F(x)$ is the P.D.F. of the time until the process $\xi(t)$ terminates.

A PH-representation (\bar{f}, G) is called irreducible, if $f_0 = 1 - \bar{f}^T \bar{1} \neq 1$ and

$$A = G - \frac{1}{1 - f_0} G \bar{1} \bar{f}^T$$

is irreducible.

This case may be interpreted as a random walk of one customer in the closed network shown on Figure 2.




Figure 2

Lemma 1. Let (\bar{F}, G) be an irreducible PH-representation. Then the matrix $H = -G$ is an invertible M-matrix.

Proof. It is known [8] that, if a matrix B satisfies the condition

$$B_{ij} \leq 0, i \neq j, i, j = \overline{1, m},$$

and if there exists a vector \bar{x} with positive components so that

$$\sum_{j=1}^m B_{ij} x_j \geq 0, i = \overline{1, m},$$

then B is a M-matrix. Such a vector for the matrix H is the vector $\bar{x} = \bar{1}$. Furthermore, from [7], we know that the matrix B of an irreducible PH-representation is invertible. This concludes the proof.



As the real parts of the eigenvalues of an invertible matrix are positive, we have the

Corollary. Let (\bar{f}, G) be an irreducible PH-representation. Then the matrix G is stable.*)

This follows from the fact that a PH-representation is of "exponential type" as introduced in [9].



All characteristics of a P.D.F. $F(x)$ with an irreducible PH-representation (\bar{f}, G) can be obtained from the vector \bar{f} and the matrix G. In particular the Laplace-Stieltjes transforms (LST) of the P.D.F. $F(x)$ is given by

$$F^*(s) = f_0^{-T} (sI - G)^{-1} G \bar{1} = \bar{1}^{-T} (sI - G)^{-1} \bar{1}, \text{ Re } s \geq 0. \quad (2)$$

*) A matrix A is stable if all its eigenvalues λ_i satisfy $\text{Re } \lambda_i < 0$. A is substable if $\text{Re } \lambda_i \leq 0$.

The k'th moment is

$$\int_0^{\infty} x^k dF(x) = (-1)^k k! \bar{f}^T G^{-k} \bar{1}, \quad k = 1, 2, \dots \quad (3)$$

In the sequel we shall also use the following

Lemma 2. Let (\bar{f}, G) be an irreducible PH-representation of order m of the P.D.F. $F(x)$, $\bar{f}^T \bar{1} = 1$. If the $(n \times n)$ -matrix H is stable, then the $(nm \times nm)$ -matrix

$$\tilde{G} = G \otimes I + (I - \bar{1} \bar{f}^T) \otimes H$$

is invertible. *)

Proof. First we shall show that the matrix $G^{-1}(I - \bar{1} \bar{f}^T)$ is substable. Therefore let γ be an eigenvalue of that matrix and \bar{x} be the corresponding eigenvector

$$G^{-1} \bar{x} - (\bar{f}^T \bar{x}) G^{-1} \bar{1} = \gamma \bar{x}. \quad (4)$$

Assume that $\text{Re } \gamma > 0$. G and G^{-1} being stable, $1/\gamma$ (resp. γ) can not be an eigenvalue of G (resp. G^{-1}). Hence $\bar{f}^T \bar{x} \neq 0$. Furthermore relation (4) is equivalent to

$$\bar{x} = \left(\frac{\bar{f}^T \bar{x}}{\gamma} \right) \left(\frac{1}{\gamma} I - G \right)^{-1} \bar{1}.$$

Then it follows that

$$\bar{f}^T \bar{x} = \frac{(\bar{f}^T \bar{x})}{\gamma} \bar{f}^T \left(\frac{1}{\gamma} I - G \right)^{-1} \bar{1}. \quad (5)$$

Using (2) and (5) we get

$$F^* \left(\frac{1}{\gamma} \right) = 1 - \frac{1}{\gamma} \bar{f}^T \left(\frac{1}{\gamma} I - G \right)^{-1} \bar{1} = 0.$$

*) $A \otimes B$ denotes the usual tensor product of the matrices A and B (see for instance [11]).

But the LST of the P.D.F. of a non-negative random variable is strictly positive in the right-hand plane [10]. Hence the equality (6) contradicts the assumption $\text{Re } \gamma > 0$ and the matrix $G^{-1}(I - \bar{1} \bar{f}^T)$ is therefore substable.

We rewrite now the matrix \tilde{G} as

$$\tilde{G} = (G \otimes I) \Psi (I \otimes H),$$

where

$$\Psi = I \otimes H^{-1} + G^{-1}(I - \bar{1} \bar{f}^T) \otimes I.$$

Each eigenvalue of Ψ is the sum of the eigenvalues of the matrices H^{-1} and $G^{-1}(I - \bar{1} \bar{f}^T)$ [11]. H^{-1} is stable and $G^{-1}(I - \bar{1} \bar{f}^T)$ is substable. This implies that Ψ is stable and invertible. Moreover, the matrices $G \otimes I$ and $I \otimes H$ being also invertible, so is \tilde{G} and the proof of Lemma 2 is concluded. □

3 - QS PH/PH/1/r

Let us assume that P.D.F.s $A(t)$ and $B(t)$ allow the following irreducible PH-representations

$$\begin{aligned} A(x) &= 1 - \frac{\alpha^T}{\bar{1}} e^{Ax} \bar{1}, \quad x \geq 0, \quad \frac{\alpha^T}{\bar{1}} \bar{1} = 1, \\ B(x) &= 1 - \frac{\beta^T}{\bar{1}} e^{Mx} \bar{1}, \quad x \geq 0, \quad \frac{\beta^T}{\bar{1}} \bar{1} = 1, \end{aligned}$$

of order l and m respectively.

Using probabilistic interpretation of a PH-distribution, the QS PH/PH/1/r may be considered as an homogeneous Markov process $\{X(t), t \geq 0\}$ on the state space

$$X = \bigcup_{k=0}^{r+1} X_k,$$

where

$$X_0 = \{(i,0) \mid i = \overline{1,\ell}\},$$

$$X_k = \{(i,k,j) \mid i = \overline{1,\ell}, j = \overline{1,m}\}, k = \overline{1,r+1}.$$

The irreducibility of the PH-representations of $A(x)$ and $B(x)$ entails that the process $X(t)$ is irreducible. We denote the stationary probability of the state $x \in X$ by p_x . Define the vectors

$$\begin{aligned} \overline{p}_0 &= (p_{10}, p_{20}, \dots, p_{\ell 0}), \\ \overline{p}_x &= (p_{1k_1}, p_{1k_2}, \dots, p_{1km}, \dots, p_{\ell k_1}, p_{\ell k_2}, \dots, p_{\ell km}), \\ & k = \overline{1,r+1}. \end{aligned}$$

From the above assumptions, the stationary distribution p_x , $x \in X$, is the unique solution of the steady state equilibrium equations

$$\overline{p}_0^T \Lambda + \overline{p}_1^T (I \otimes \overline{\mu}) = \overline{0}^T, \quad (7)$$

$$\overline{p}_0^T (\overline{\lambda} \overline{\alpha}^T \otimes \overline{\beta}^T) + \overline{p}_1^T (\Lambda \otimes I + I \otimes M) + \overline{p}_2^T (I \otimes \overline{\mu} \overline{\beta}^T) = \overline{0}^T, \quad (8)$$

$$\overline{p}_{k-1}^T (\overline{\lambda} \overline{\alpha}^T \otimes I) + \overline{p}_k^T (\Lambda \otimes I + I \otimes M) + \overline{p}_{k+1}^T (I \otimes \overline{\mu} \overline{\beta}^T) = \overline{0}^T, \quad (9)$$

$k = \overline{2,r};$

$$\overline{p}_r^T (\overline{\lambda} \overline{\alpha}^T \otimes I) + \overline{p}_{r+1}^T ((\Lambda + \overline{\lambda} \overline{\alpha}^T) \otimes I + I \otimes M) = \overline{0}^T, \quad (10)$$

where

$$\overline{\lambda} = -\Lambda \overline{1}, \quad \overline{\mu} = -M \overline{1},$$

with the normalizing condition

$$\sum_{k=0}^{r+1} \overline{p}_k^T \overline{1} = 1. \quad (11)$$

Let us defined matrices

$$\tilde{\Lambda} = (\bar{1} \bar{\alpha}^T - I) \otimes M - \Lambda \otimes I, \quad \tilde{M} = \Lambda \otimes (\bar{1} \bar{\beta}^T - I) - I \otimes M.$$

We shall now study some properties of the stationary distribution $p_x, x \in X$, based on results in [3].

1°) Multiplying on the right both sides of equations (8)-(10) by the matrix $I \times \bar{1}$, yields

$$\bar{p}_0^T \bar{\lambda} \bar{\alpha}^T + \bar{p}_1^T (\Lambda \otimes \bar{1} - I \otimes \bar{\mu}) + \bar{p}_2^T (I \otimes \bar{\mu}) = \bar{0}^T, \quad (12)$$

$$\bar{p}_{k-1}^T (\bar{\lambda} \bar{\alpha}^T \otimes \bar{1}) + \bar{p}_k^T (\Lambda \otimes \bar{1} - I \otimes \bar{\mu}) + \bar{p}_{k+1}^T (I \otimes \bar{\mu}) = \bar{0}^T, \quad (13)$$

$$k = \overline{1, r},$$

$$\bar{p}_r^T (\bar{\lambda} \bar{\alpha}^T \otimes \bar{1}) + \bar{p}_{r+1}^T ((\Lambda \oplus \bar{\lambda} \bar{\alpha}^T) \otimes \bar{1}) - I \otimes \bar{\mu}) = \bar{0}^T. \quad (14)$$

Sum (13) with respect to k and combine the result with (7), (12) and (14), to get, after modest rearrangement,

$$\left[\bar{p}_0^T + \sum_{k=1}^{r+1} \bar{p}_k^T (I \otimes \bar{1}) \right] (\Lambda + \bar{\lambda} \bar{\alpha}^T) = \bar{0}^T. \quad (15)$$

2°) After multiplying on the right both sides of (7), (12) and (13) by the vector $\bar{1}$, we obtain

$$\bar{p}_0^T \bar{\lambda} - \bar{p}_1^T (1 \otimes \bar{\mu}) = 0,$$

$$\bar{p}_0^T \bar{\lambda} - \bar{p}_1^T (\bar{\lambda} \otimes \bar{1} + \bar{1} \otimes \bar{\mu}) + \bar{p}_2^T (\bar{1} \otimes \bar{\mu}) = 0,$$

$$\bar{p}_{k-1}^T (\bar{\lambda} \otimes \bar{1}) - \bar{p}_k^T (\bar{\lambda} \otimes \bar{1} + \bar{1} \otimes \bar{\mu}) + \bar{p}_{k+1}^T (\bar{1} \otimes \bar{\mu}) = 0,$$

$$k = \overline{2, r}.$$

The latter equalities together with (7) and (16) give

$$\bar{P}_1^{-T} \bar{M} = \bar{P}_1^{-T} (I \otimes \bar{\mu} \bar{\beta}^T) = -\bar{P}_0^{-T} (\Lambda \otimes \bar{\beta}^T), \quad (17)$$

$$\begin{aligned} \bar{P}_k^{-T} \bar{M} &= \bar{P}_k^{-T} (I \otimes \bar{\mu} \bar{\beta}^T) + \bar{P}_{k-1}^{-T} (\bar{\lambda} \bar{\alpha}^T \otimes I) - \bar{P}_{k-1}^{-T} (\bar{\lambda} \bar{\alpha}^T \otimes \bar{1} \bar{\beta}^T) = \\ &= \bar{P}_k^{-T} (I \otimes \bar{\mu} \bar{\beta}^T) + \bar{P}_{k-1}^{-T} (\bar{\lambda} \bar{\alpha}^T \otimes I) - \bar{P}_k^{-T} (\bar{1} \bar{\alpha}^T \otimes \bar{\mu} \bar{\beta}^T), \end{aligned} \quad (18)$$

$k = \overline{2, r}.$

4°) After multiplying at the right both sides of (8) and (9) by the matrix $(\bar{1} \bar{\alpha}^T - I) \otimes I$, we have

$$\bar{P}_k^{-T} (\Lambda \otimes I + I \otimes M) [(\bar{1} \bar{\alpha}^T - I) \otimes I] + \bar{P}_{k+1}^{-T} [(\bar{1} \bar{\alpha}^T - I) \otimes \bar{\mu} \bar{\beta}^T] = \bar{0}^T, \quad (18)$$

$k = \overline{1, r}.$

This implies that

$$\bar{P}_k^{-T} \Lambda = \bar{P}_k^{-T} (\bar{\lambda} \bar{\alpha}^T \otimes I) + \bar{P}_{k+1}^{-T} [(I - \bar{1} \bar{\alpha}^T) \otimes \bar{\mu} \bar{\beta}^T], \quad k = \overline{1, r}. \quad (19)$$

5°) Now, from (17)-(19) and (10), it follows that :

$$\bar{P}_1^{-T} \bar{M} = -\bar{P}_0^{-T} (\Lambda \otimes \bar{\beta}^T), \quad (20)$$

$$\bar{P}_k^{-T} \bar{M} = \bar{P}_{k-1}^{-T} \Lambda, \quad k = \overline{2, r}, \quad (21)$$

$$\bar{P}_{r+1}^{-T} [\Lambda + \bar{\lambda} \bar{\alpha}^T \otimes I + I \otimes M] - \bar{P}_r^{-T} (\bar{\lambda} \bar{\alpha}^T \otimes I). \quad (22)$$

We note that the substable and irreducible matrix $\Lambda + \bar{\lambda} \bar{\alpha}^T$ is the generator of an homogeneous Markov process. The eigenvalues of $(\Lambda + \bar{\lambda} \bar{\alpha}^T) \otimes I + I \otimes M$ are the sum of the eigenvalues of the substable matrix $\Lambda + \bar{\lambda} \bar{\alpha}^T$ and of the stable matrix M [11]. Hence $(\Lambda + \bar{\lambda} \bar{\alpha}^T) \otimes I + I \otimes M$ is stable and invertible. Lemma 2) entails that the matrices Λ and \bar{M} are also invertible.

Introduce

$$W_0 = - (\Lambda \otimes \bar{\alpha}^T) \bar{M}^{-1}, \quad W = \bar{\Lambda} \bar{M}^{-1},$$

$$W_r = - (\bar{\lambda} \bar{\alpha}^T \otimes I) [(\Lambda + \bar{\lambda} \bar{\alpha}^T) \otimes I + I \otimes M]^{-1},$$

$$V = I + \left(\sum_{k=0}^{r-1} W_0 W^k + W_0 W^{r-1} W_r \right) (I \otimes \bar{1}).$$

Theorem. Let \bar{u} be a solution of the steady-state system of equations

$$\bar{u}^T (\Lambda + \bar{\lambda} \bar{\alpha}^T) = \bar{0}^T \tag{23}$$

$$\bar{u}^T \bar{1} = 1.$$

Then, the stationary distribution p_x , $x \in X$, has the form

$$p_k^T = \begin{cases} \bar{p}_0^T W_0 W^{k-1}, & k = \overline{1, r}, \\ \bar{p}_0^T W_0 W^{r-1} W_r, & k = r + 1, \end{cases} \tag{24}$$

and the vector \bar{p}_0 satisfies the system

$$\bar{p}_0^T V = \bar{u}^T. \tag{25}$$

Proof. The irreducibility of the matrix $\Lambda + \bar{\lambda} \bar{\alpha}^T$ implies that the system (23) has a unique solution. From the relations (20)-(22), we find the stationary distribution p_x , $x \in X$, in the form (24). Moreover (24) and (15) yield that the vector $\bar{p}_0^T V$ satisfy the system

$$\bar{p}_0^T V (\Lambda + \bar{\lambda} \bar{\alpha}^T) = \bar{0}^T.$$

Therefore $\bar{p}_0^T V = c \bar{u}^T$, where c is a constant. The normalizing condition (11) and relation (24) lead to $c = 1$. The proof is concluded. □

4 - A NUMERICAL EXAMPLE

In order to illustrate the theoretical derivations in this paper, some numerical results for the QSs $H_3M/E_4/1/10$, $E_3/H_4MM/1/10$ and $E_3/E_4/1/10$ are presented. The Erlang and hyperexponential distributions are obvious by particular PH-distributions. In the case of the QS $H_3M/E_4/1/10$ e.g. we have respectively

$$G = \text{diag}(-\lambda_1, -\lambda_2, -\lambda_3), \quad \bar{\alpha}^T = (\alpha_1, \alpha_2, \alpha_3),$$

$$\lambda_i > 0, \alpha_i > 0, i = \overline{1,3}, \quad \sum_{i=1}^3 \alpha_i = 1 \text{ and}$$

$$G = \begin{vmatrix} -\mu & \mu & 0 & 0 \\ 0 & -\mu & \mu & 0 \\ 0 & 0 & -\mu & \mu \\ 0 & 0 & 0 & -\mu \end{vmatrix}, \quad \bar{\beta}^T = (1, 0, 0, 0),$$

with $\mu > 0$.

The calculation have been carried out at the department of mathematics of the University of Friendship Patrice Lumumba. The program was written by S.S. Spesivov in Fortran language.

The distribution parameters were given following numerical values :

$$\underline{H_3M/E_4} : \bar{\lambda}^T = (1.0, 2.0, 3.0), \bar{\alpha}^T = (0.2, 0.3, 0.5); \mu = 12.0;$$

$$\underline{E_3/H_4M} : \lambda = 10.5; \bar{\mu}^T = (1.0, 2.0, 3.0, 4.0), \bar{\beta}^T = (0.1, 0.2, 0.3, 0.4);$$

$$\underline{E_3/E_4} : \lambda = 7.5; \mu = 14.0$$

[λ and μ are respectively the mean arrival rate and the mean service rate.].

In Table 1 the values $p_k \stackrel{\text{def}}{=} \sum_i \sum_j p(i,k,j)$, $k = \overline{0,11}$, are given.

The last row of Table 1 check the normalizing condition (11).

Table 1

$\begin{matrix} \text{QS} \\ \backslash \\ k \end{matrix}$	$H_3 M/E_4 /1/10$	$E_3 /H_4 M/1/10$	$E_3 /E_4 /1/10$
0	0.355551	0.003984	0.285716
1	0.248652	0.007517	0.431506
2	0.162687	0.010527	0.189542
3	0.097829	0.015112	0.063441
4	0.057377	0.021910	0.020310
5	0.033436	0.031891	0.006469
6	0.019456	0.046486	0.002059
7	0.011321	0.067787	0.000658
8	0.006597	0.098770	0.000209
9	0.003856	0.143269	0.000066
10	0.002279	0.204365	0.000021
11	0.000962	0.348383	0.000005
\sum	1.000003	1.000001	1.000002

Aknowledgements : We should like to thank Professor G.P. Basharin for helpful discussions on this paper and Dr. G. Fayolle (INRIA, France) for carefull reading of this manuscript.

REFERENCES

- [1] Basharin, G.P. : A single server with a finite queue and items of several types. Theory of Probability and its applications. Vol.X, n° 2, 1965, pp.261-274. (in russian).
- [2] Basharin, G.P., Harkevich, A.D. and Shneps, M.A. : Queueing theory in telephone systems. Moscow, Izd."Nauka", 1968. (in russian).
- [3] Bocharov, P.P. and Naumov, V.A. : Analysis of hyperexponential two-node queueing system with a limited intermediate buffer. In the collection "Telecommunication networks and their structure", Moscow, Izd."Nauka", 1976, pp.168-181. (in russian).
- [4] Naumov, V.A. : On one-server system with a limited buffer and customers of several types. In the collection "Models of systems of distribution information and their analysis". Moscow, Izd."Nauka", 1982, pp.77-82. (in russian).
- [5] Neuts, M.F. : Explicit steady-state solutions to some elementary queueing models. Operations Research, vol. 30, n° 3, 1982, pp. 480-489.
- [6] Bocharov, P.P. : Single server queueing systems with finite capacity. Moscow. Izd. University of Friendship (forthcoming).(in russian).
- [7] Neuts, P.F. : Matrix-geometric solutions in stochastics models. The Johns Hopkins Univ. Press, Baltimore and London, 1981.
- [8] Berman, A. and Plemmons, P. : Nonnegative matrices in the mathematical sciences. Acad. Press, N.Y., 1979.
- [9] Bocharov, P.P. and Naumov, V.A. : On some queueing system with finite capacity. Problems of information transmission, vol. XIII, 1977, pp.96-104. (in russian).
- [10] Lukacs, E. : Characteristic functions, 2ed. Griffen, London, 1970.

Imprimé en France

par

l'Institut National de Recherche en Informatique et en Automatique

