

Global smooth solution of Cauchy problems for a class of quasilinear hyperbolic systems

L. Hsiao, T.T. Li

► **To cite this version:**

L. Hsiao, T.T. Li. Global smooth solution of Cauchy problems for a class of quasilinear hyperbolic systems. RR-0083, INRIA. 1981. <inria-00076478>

HAL Id: inria-00076478

<https://hal.inria.fr/inria-00076478>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IRIA

CENTRE DE ROCQUENCOURT

Rapports de Recherche

N° 83

**GLOBAL SMOOTH SOLUTION
OF CAUCHY PROBLEMS
FOR A CLASS OF QUASILINEAR
HYPERBOLIC SYSTEMS**

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
B.P.105
78153 Le Chesnay Cedex
France
Tél. 954 90 20

Ling HSIAO
Ta-Tsien LI

Juin 1981

GLOBAL SMOOTH SOLUTION OF CAUCHY PROBLEMS FOR A CLASS OF
QUASILINEAR HYPERBOLIC SYSTEMS

par

Ling HSIAO
Institute of Mathematics
Academia Sinica of China
Peking, China

et

Ta-tsien Li
Département de Mathématiques
Université Fudan
Shanghai, China

et

Professeur invité
Collège de France et INRIA
Paris, France

RESUME : Dans cet article, nous avons démontré que la présence de certains types de termes non homogènes peut rendre globalement régulière en temps la solution de systèmes hyperboliques quasilineaires du premier ordre quand les conditions initiales sont assez petites.

ABSTRACT : In this article we proved that certain kinds of inhomogeneous terms can smoothen the solution for first order quasilinear hyperbolic systems globally in time provided that the initial data are small.

GLOBAL SMOOTH SOLUTION OF CAUCHY PROBLEMS FOR A CLASS OF
QUASILINEAR HYPERBOLIC SYSTEMS

by

Ling Hsiao^{*}
Institute of Mathematics
Academia Sinica of China
Peking, China

and

(Visiting Professor, 1979-82)
Division of Applied Mathematics
Lefschetz Center for Dynamical Systems
Brown University
Providence, R. I. 02912

and

Ta-tsien Li
Fudan University
Shanghai, China

(Visiting Professor)
Collège de France and INRIA
Paris, France

May 1981

* This research has been supported in part by the National Science Foundation under Grant #NSF-Eng. CME80-23824.

GLOBAL SMOOTH SOLUTION OF CAUCHY PROBLEMS FOR A CLASS OF
QUASILINEAR HYPERBOLIC SYSTEMS

by

Ling Hsiao and Ta-tsien Li

ABSTRACT

In this article we prove that certain kinds of inhomogeneous terms can smoothen the solution for first order quasilinear hyperbolic systems globally in time provided that the initial data are small.

1. INTRODUCTION

For the Cauchy problem of first order quasilinear hyperbolic systems it is well-known that even if the initial data are very smooth, in general, a smooth solution exists only locally in time and singularities may appear in a finite time. It was discussed in [1] how the presence of various damping and dissipation mechanisms may influence the smoothness of the solution. In the same direction, we consider here the influence of a special kind of dissipative inhomogeneous term and establish the existence and uniqueness of global smooth solutions.

Consider the following Cauchy problem for a first order system of inhomogeneous hyperbolic balance laws:

$$\left\{ \begin{array}{l} \frac{\partial u}{\partial t} + \frac{\partial f(u)}{\partial x} + g(u) = 0, \quad -\infty < x < \infty, \quad 0 \leq t < \infty \end{array} \right. \quad (1)$$

$$\left\{ \begin{array}{l} u(x,0) = \phi(x), \quad -\infty < x < \infty. \end{array} \right. \quad (2)$$

where $u = (u_1, \dots, u_n)^T$ is the unknown vector function, $f(u)$ and $g(u)$ are given smooth vector functions of u with

$$g(0) = 0 \quad (3)$$

and $\phi(x)$ is smooth.

System (1) is hyperbolic on the domain under consideration if

- 1) The $n \times n$ matrix $\nabla f(u)$ has n real eigenvalues:

$$\lambda_1(u), \lambda_2(u), \dots, \lambda_n(u); \quad (4)$$

- 2) $\nabla f(u)$ is diagonalizable, i.e.,

$$\det \zeta(u) \neq 0. \quad (5)$$

where

$$\zeta(u) = \begin{pmatrix} \zeta_1(u) \\ \zeta_2(u) \\ \vdots \\ \zeta_n(u) \end{pmatrix} \quad (6)$$

and $\zeta_\ell(u) = (\zeta_{\ell 1}(u), \zeta_{\ell 2}(u), \dots, \zeta_{\ell n}(u))$ is a left eigenvector corresponding to λ_ℓ :

$$\zeta_\ell(u) \nabla f(u) = \lambda_\ell(u) \zeta_\ell(u). \quad (7)$$

By (3), $g(u)$ can be rewritten as

$$g(u) = B(u)u \quad (8)$$

where $B(u) = (B_{ij}(u))$ is a $n \times n$ matrix with

$$B(0) = \nabla g(0). \quad (9)$$

Let
$$\lambda(u) = \text{diag}\{\lambda_1(u), \lambda_2(u), \dots, \lambda_n(u)\}. \quad (10)$$

The main results in this paper are the following:

THEOREM 1. Assume that (3) holds and the matrix

$$\mathcal{A} \equiv (a_{ij}) = \zeta(0) \nabla g(0) \zeta^{-1}(0) \quad (11)$$

is row diagonal dominant:

$$a_{\ell\ell} > \sum_{j \neq \ell} |a_{\ell j}| \quad (\ell=1, \dots, n) \quad (12)$$

Furthermore, let $\zeta(u), \lambda(u)$ and $B(u)$ be smooth with bounded C^1 norm and satisfy

$$|\det \zeta(u)| \geq D_0 > 0 \quad (D_0 \text{ is constant}). \quad (13)$$

Then the Cauchy problem (1), (2) admits a unique global smooth solution $u = u(x,t)$ on $t \geq 0$ which decays exponentially with respect to t , provided that the C^1 norm of the initial data $\phi(x)$ is sufficiently small.

THEOREM 2. The conclusion of Theorem 1 holds if (12) is replaced by the following weaker condition that there exists a diagonal matrix γ with $\gamma_\ell \neq 0$ ($\ell=1, \dots, n$) such that the matrix $\gamma \mathcal{A} \gamma^{-1}$ is row diagonal dominant.

The results in this paper generalize and make more precise the results in [4] for the one-dimensional case.

We should note that our assumption (12) or even (15) do not cover, for example, the case of the one-dimensional damped wave equation

$$\begin{cases} u_t - v_x = 0 \\ v_t - \sigma(u)_x + v = 0. \end{cases} \quad (14)$$

For a global existence theorem for this special system see [5].

The proofs of Theorem 1 and 2 will be given in the next section.

2. PROOFS OF THEOREMS

Multiplying (1) by ζ from the left and using (7), we obtain the characteristic form

$$\zeta(u) \left[\frac{\partial u}{\partial t} + \lambda(u) \frac{\partial u}{\partial x} \right] + A(u) \zeta(u) u = 0, \quad (15)$$

where

$$A(u) \equiv (A_{ij}(u)) = \zeta(u)B(u)\zeta^{-1}(u). \quad (16)$$

We note, in view of (9), that $A(0) = \mathcal{A}$, i.e., $A_{ij}(0) = a_{ij}$ ($i, j=1, \dots, n$).

It is now clear that the assertion of Theorem 1 follows easily from the following lemma.

LEMMA. Assume that the matrix $A(0) = \mathcal{A}$ satisfies (12) and that $\zeta(u), \lambda(u)$ and $A(u)$ are smooth with bounded C^1 norms and satisfy (13). Then, if the C^1 norm of the initial data $\phi(x)$ is sufficiently small, there exists a unique global smooth solution $u = u(x, t)$ on $t \geq 0$ for the Cauchy problem (15), (2) and this solution decays exponentially as $t \rightarrow \infty$.

PROOF. In order to obtain the existence of the global smooth solution it suffices to prove that if the initial data $\phi(x)$ have a small C^1 norm then the C^1 norm of the smooth solution $u = u(x, t)$ defined on the domain

$$R(T) = \{(t, x) \mid 0 \leq t \leq T, \quad -\infty < x < \infty\} \quad (17)$$

does not depend on T .

Rewrite (15) in the form

$$\begin{aligned} & \sum_j \zeta_{\ell j}(u) \left(\frac{\partial u_j}{\partial t} + \lambda_\ell(u) \frac{\partial u_j}{\partial x} \right) + \sum_j (A_{\ell j}(u) - A_{\ell j}(0)) \sum_k \zeta_{jk}(u) u_k + \\ & \sum_j a_{\ell j} \sum_k \zeta_{jk}(u) u_k = 0 \quad (\ell=1, \dots, n) \end{aligned} \quad (18)$$

Let

$$\xi = f_\ell(\tau; t, x) \quad (19)$$

be the ℓ -th characteristic curve passing through the point (t, x) , which satisfies

$$\begin{cases} \frac{d\xi}{d\tau} = \lambda_\ell(\tau, f_\ell(\tau; t, x)), \\ f_\ell(t; t, x) = x. \end{cases} \quad (20)$$

Setting

$$U_\ell = \sum_k \zeta_{\ell k}(u) u_k \quad (21)$$

and

$$V_\ell = e^{a_{\ell\ell} t} U_\ell, \quad (22)$$

it is easy to see that

$$\left\{ \begin{aligned} \frac{\partial V_\ell}{\partial t} + \lambda_\ell(u) \frac{\partial V_\ell}{\partial x} &= e^{a_{\ell\ell} t} \left\{ \sum_j \left[\frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} \right] u_j \right. \\ &\quad \left. - \sum_j [A_{\ell j}(u) - A_{\ell j}(0)] \sum_k \zeta_{jk}(u) u_k \right\} - e^{a_{\ell\ell} t} \sum_{j \neq \ell} a_{\ell j} U_j \\ &\quad (\ell=1, \dots, n), \\ V_\ell &= U_\ell^0(x) \equiv \sum_k \zeta_{\ell k}(\phi) \phi_k, \quad t=0, \quad -\infty < x < \infty. \end{aligned} \right. \quad (23)$$

Integrating the ℓ -th equation of (23) along the ℓ -th characteristic curve, we get

$$\begin{aligned} V_\ell(t, x) &= U_\ell^0(f_\ell(0; t, x)) + \int_0^t \left\{ e^{a_{\ell\ell} \tau} \left[\sum_j \left(\frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} \right) u_j + \right. \right. \\ &\quad \left. \left. - \sum_j [A_{\ell j}(u) - A_{\ell j}(0)] \sum_k \zeta_{jk}(u) u_k \right] - e^{a_{\ell\ell} \tau} \sum_{j \neq \ell} a_{\ell j} U_j \right\} d\tau, \end{aligned} \quad (24)$$

then

$$U_\ell(t, x) = e^{-a_{\ell\ell}t} U_\ell^0(f_\ell(0; t, x)) + \int_0^t \{ e^{a_{\ell\ell}(\tau-t)} \left[\sum_j \left(\frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} \right) u_j - \sum_j (A_{\ell j}(u) - A_{\ell j}(0)) \sum_k \zeta_{jk}(u) u_k \right] - e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} a_{\ell j} U_j \} d\tau. \quad (25)$$

in which the integration is carried along the characteristic $(\tau, f_\ell(\tau; t, x))$.

We have

$$\frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} = \sum_k \frac{\partial \zeta_{\ell j}(u)}{\partial u_k} \left(\frac{\partial u_k}{\partial t} + \lambda_\ell(u) \frac{\partial u_k}{\partial x} \right). \quad (26)$$

System (18) together with (21) imply that

$$\frac{\partial u_k}{\partial t} = \sum_{p,m} \zeta^{k,p}(u) \zeta_{p,m}(u) \lambda_p(u) \frac{\partial u_m}{\partial x} - \sum_{p,m} \zeta^{k,p}(u) A_{p,m}(u) U_m, \quad (27)$$

where $\zeta^{k,p}$ ($k, p = 1, \dots, n$) denote the elements of ζ^{-1} . Thus, introducing

$$w_i = \frac{\partial u_i}{\partial x}, \quad \bar{w}_\ell = \sum_j \zeta_{\ell j}(u) w_j, \quad (28)$$

it follows from (26) and (27) that

$$\left\{ \begin{aligned} \frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} &= \sum_s P_{\ell j s}(u) \bar{w}_s + \sum_{p,s} Q_{\ell j p}(u) A_{p,s}(u) U_s, \\ \frac{\partial u_k}{\partial t} &= \sum_s R_{k,s}(u) \bar{w}_s + \sum_{p,s} S_{k,p}(u) A_{p,s}(u) U_s, \end{aligned} \right. \quad (29)$$

in which

$$\left\{ \begin{aligned}
 P_{\ell j s}(u) &= \sum_{k,p,m} \frac{\partial \zeta_{\ell j}(u)}{\partial u_k} \zeta^{k,p}(u) \zeta_{p,m}(u) \lambda_p(u) \zeta^{ms}(u) + \sum_k \frac{\partial \zeta_{\ell j}(u)}{\partial u_k} \lambda_\ell(u) \zeta^{ks}(u), \\
 Q_{\ell j p}(u) &= - \sum_k \frac{\partial \zeta_{\ell j}(u)}{\partial u_k} \zeta^{kp}(u), \\
 R_{k s}(u) &= \sum_{p,m} \zeta^{kp}(u) \zeta_{p,m}(u) \lambda_p(u) \zeta^{ms}(u), \\
 S_{kp}(u) &= - \zeta^{kp}(u).
 \end{aligned} \right. \quad (30)$$

Then, from (25) we have

$$\begin{aligned}
 U_\ell(t, x) &= e^{-a_{\ell\ell}t} U_\ell^0(f_\ell(0; t, x)) + \\
 &\int_0^t \{ e^{a_{\ell\ell}(\tau-t)} [\sum_j (\sum_s^p \ell_{j s}(u) \bar{w}_s + \sum_{p,s} Q_{\ell j p}(u) A_{ps}(u) U_s) \cdot \sum_r \zeta^{jr}(u) U_r \\
 &- \sum_j (A_{\ell j}(u) - A_{\ell j}(0)) U_j] - e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} a_{\ell j} U_j \} d\tau.
 \end{aligned} \quad (31)$$

Moreover, if we set

$$H_\ell = e^{a_{\ell\ell}t} \bar{w}_\ell \quad (32)$$

and differentiate (18) with respect to x , it follows in a similar way that

$$\left\{ \begin{aligned}
 \frac{\partial H_\ell}{\partial t} + \lambda_\ell(u) \frac{\partial H_\ell}{\partial x} &= e^{a_{\ell\ell}t} \cdot \sum_j \left(\frac{\partial \zeta_{\ell j}(u)}{\partial t} + \lambda_\ell(u) \frac{\partial \zeta_{\ell j}(u)}{\partial x} w_j \right. \\
 &- e^{a_{\ell\ell}t} \left(\sum_{j,k} \frac{\partial \zeta_{\ell j}}{\partial u_k} \cdot \frac{\partial u_j}{\partial t} w_k + \sum_{j,k} \frac{\partial(\zeta_{\ell j} \lambda_\ell)}{\partial u_k} w_j w_k + \sum_{j,k,p} \frac{\partial(\Lambda_{\ell j} \zeta_{jk})}{\partial u_p} w_p u_k \right. \\
 &\left. \left. + \sum_j (A_{\ell j}(u) - A_{\ell j}(0)) \bar{w}_j \right) - e^{a_{\ell\ell}t} \sum_{j \neq \ell} a_{\ell j} \bar{w}_j, \right. \\
 H &= \bar{w}_\ell^0(x) \equiv \sum_k \zeta_{\ell k}(\phi) \phi'_k(x), \quad t = 0, \quad -\infty < x < \infty.
 \end{aligned} \right. \quad (33)$$

Integrating the ℓ -th equation along the ℓ -th characteristic curve and using (29), we obtain

$$\begin{aligned}
 W_\ell(t, x) = & e^{-a_{\ell\ell} t} \bar{W}_\ell^0(f_\ell(0; t, x)) \\
 & + \int_0^t \{ e^{a_{\ell\ell}(\tau-t)} [\sum_j (\sum_s P_{\ell js}(u) \bar{W}_s + \sum_{p,s} Q_{\ell jp}(u) A_{ps}(u) U_s) \cdot \sum_r \zeta^{jr} \bar{W}_r \\
 & - \sum_{j,k} \frac{\partial \zeta_{\ell j}}{\partial u_k} (\sum_s R_{ks}(u) \bar{W}_s + \sum_{p,s} S_{kp}(u) A_{ps}(u) U_s) \cdot \sum_r \zeta^{kr} \bar{W}_r \\
 & - \sum_{j,k,s,r} \frac{\partial(\zeta_{\ell j} \lambda_\ell)}{\partial u_k} \zeta^{js} \zeta^{kr} \bar{W}_s \bar{W}_r - \sum_{j,k,p,r,s} \frac{\partial(A_{\ell j} \zeta_{jk})}{\partial u_p} \zeta^{kr} \zeta^{ks} \bar{W}_r U_s \\
 & - \sum_j (A_{\ell j}(u) - A_{\ell j}(0)) \bar{W}_j] - e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} a_{\ell j} \bar{W}_j \} d\tau.
 \end{aligned} \tag{34}$$

Let

$$\begin{aligned}
 U_\ell(t) = \sup_{\substack{0 \leq \tau \leq t \\ -\infty < x < \infty}} |U_\ell(\tau, x)|, \quad U(t) = \sup_\ell U_\ell(t),
 \end{aligned} \tag{35}$$

$$\begin{aligned}
 \bar{W}_\ell(t) = \sup_{\substack{0 \leq \tau \leq t \\ -\infty < x < \infty}} |\bar{W}_\ell(\tau, x)|, \quad \bar{W}(t) = \sup_\ell \bar{W}_\ell(t),
 \end{aligned} \tag{36}$$

and
$$a = \min_\ell \{a_{\ell\ell}\}.$$
 (37)

We can easily obtain from (31) that

$$\begin{aligned}
 U_\ell(t) \leq & e^{-at} C_0 + \int_0^t \{ D_1 e^{a(\tau-t)} [\bar{W}(\tau) U(\tau) + U^2(\tau)] + \\
 & e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} |a_{\ell j}| U_j(\tau) \} d\tau,
 \end{aligned} \tag{38}$$

where

$$C_0 = \sup_{-\infty < x < \infty} |U_\ell^0(x)| = \sup_{-\infty < x < \infty} \left| \sum_k \zeta_{\ell k}(\phi) \phi_k \right| \quad (39)$$

and $D_i (i=1,2,\dots)$ will denote throughout various fixed constants.

Since

$$\int_0^t e^{a_{\ell\ell}(\tau-t)} d\tau = \frac{1}{a_{\ell\ell}} (1 - e^{-a_{\ell\ell}t}),$$

it follows on account of (12),

$$\begin{aligned} & \int_0^t e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} |a_{\ell j}| U_j(\tau) d\tau \\ & \leq \int_0^t e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} |a_{\ell j}| d\tau \cdot U(t) \\ & \leq \frac{\sum_{j \neq \ell} |a_{\ell j}|}{a_{\ell\ell}} U(t) \leq dU(t), \end{aligned} \quad (40)$$

where

$$d = \max_{\ell} \frac{\sum_{j \neq \ell} |a_{\ell j}|}{a_{\ell\ell}} < 1. \quad (41)$$

Then (38) gives

$$U(t) \leq \frac{1}{1-d} e^{-at} C_0 + \int_0^t D_2 e^{a(\tau-t)} (\bar{w}(\tau)U(\tau) + U^2(\tau)) d\tau. \quad (42)$$

In a similar way (34) gives

$$\left\{ \begin{aligned} \bar{w}_\ell(t) & \leq e^{-at} C_1 + \int_0^t \{ D_3 e^{a(\tau-t)} (\bar{w}^2(\tau) + \bar{w}(\tau)U(\tau)) \\ & \quad - e^{a_{\ell\ell}(\tau-t)} \sum_{j \neq \ell} |a_{\ell j}| \bar{w}_j(\tau) \} d\tau \end{aligned} \right. \quad (43)$$

where

$$C_1 = \sup_{\ell} |\bar{W}_{\ell}^0(x)| = \sup_{\ell} \left| \sum_k \zeta_{\ell k}(\phi) \phi_k' \right|. \quad (44)$$

$$-\infty < x < \infty \quad -\infty < x < \infty$$

Therefore,

$$\bar{W}(t) \leq \frac{1}{1-d} e^{-at} C_1 + \int_0^t D_4 e^{a(\tau-t)} (\bar{W}^2(\tau) + \bar{W}(\tau)U(\tau)) d\tau. \quad (45)$$

Let

$$X(t) = U(t) + \bar{W}(t). \quad (46)$$

Combining (42) and (45), it is easily seen that

$$X(t) \leq \frac{1}{1-d} e^{-at} (C_0 + C_1) + \int_0^t D_5 e^{a(\tau-t)} X^2(\tau) d\tau. \quad (47)$$

Letting

$$Y(t) = X(t)e^{at}, \quad (48)$$

(47) can be written as

$$Y(t) \leq \frac{1}{1-d} (C_0 + C_1) + \int_0^t D_5 e^{-a\tau} Y^2(\tau) d\tau. \quad (49)$$

Consider

$$\begin{cases} \frac{dZ(t)}{dt} = D_5 e^{-at} Z^2(t), \\ Z = \frac{1}{1-d} (C_0 + C_1), \quad t = 0. \end{cases} \quad (50)$$

We then have the estimation

$$Y(t) \leq Z(t). \quad (51)$$

But

$$Z(t) = \frac{1}{\frac{1-d}{C_0+C_1} - \frac{D_5}{a}(1-e^{-at})} \quad (52)$$

Therefore, if $C_0 + C_1$ is so small that

$$a > D_5 \cdot \frac{C_0+C_1}{1-d}, \quad (53)$$

then

$$Z(t) \leq \frac{C_0+C_1}{1-d} \quad (54)$$

Hence

$$X(t) \leq \frac{C_0+C_1}{1-d} e^{-at}. \quad (55)$$

Thus, by the definition of $X(t)$, the lemma and thenceforth Theorem 1, is proved. VVV

REMARK 1. It follows from the proof of the theorem that assumption (13) of Theorem 1 can be replaced with the weaker

$$\det \zeta(u) \neq 0. \quad (56)$$

REMARK 2. Assumption (12) in Theorem 1 is similar to the assumption in [2] for establishing the corresponding global existence of discontinuous solutions for the Cauchy problem (1),(2).

By means of the method in [3], multiplying the l -th equation of (18) by a real number $\gamma_l \neq 0$, the matrix \mathcal{A} can be replaced with $\gamma \mathcal{A} \gamma^{-1}$, where

$$\gamma = \text{diag}\{\gamma_1, \gamma_2, \dots, \gamma_n\}. \quad (57)$$

Thus Theorem 2 follows from Theorem 1.

REFERENCES

- [1] C.M. Dafermos, Conservation laws with dissipation, July 1980. (preprint)
- [2] C.M. Dafermos and Ling Hsiao, Hyperbolic system of balance laws with inhomogeneity and dissipation, (to appear).
- [3] Ta-tsien Li and Wen-tju Yu, Some existence theorems for quasilinear hyperbolic systems of partial differential equations in two independent variables (1) Typical boundary value problems, Scientia Sinica, 4(1964), pp. 529-550.
- [4] Gu chao-hao, Some global solutions of quasilinear hyperbolic symmetric systems and their asymptotic behavior, Acta Math. Sinica 21(1978) pp.130-134.
- [5] T. Nishida, "Global smooth solutions for the second-order quasilinear wave equations with the first-order dissipation, (preprint).

Imprimé en France
par
l'Institut National de Recherche en Informatique et en Automatique

