



# Curved finite elements of class C1: implementation and numerical experiments

Michel Bernadou, Jean-Marie Boisserie

## ► To cite this version:

Michel Bernadou, Jean-Marie Boisserie. Curved finite elements of class C1: implementation and numerical experiments. [Research Report] RR-1718, INRIA. 1992. inria-00076956

HAL Id: inria-00076956

<https://inria.hal.science/inria-00076956>

Submitted on 29 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNITE DE RECHERCHE  
INRIA ROCQUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP 105  
78153 Le Chesnay Cedex  
France  
Tel (1) 39635511

# Rapports de Recherche

1992



25<sup>e</sup>  
anniversaire

N° 1718

*Programme 6*  
*Calcul Scientifique, Modélisation et*  
*Logiciels numériques*

## CURVED FINITE ELEMENTS OF CLASS C1 : IMPLEMENTATION AND NUMERICAL EXPERIMENTS

**Part 1 :**  
**Construction and numerical tests**  
**of the interpolation properties**

Michel BERNADOU  
Jean-Marie BOISSERIE

Juin 1992



\* R R - 1 7 1 8 \*

# CURVED FINITE ELEMENTS OF CLASS $\mathcal{C}^1$ : IMPLEMENTATION AND NUMERICAL EXPERIMENTS

(\*)(\*\*)

## Part 1 : Construction and numerical tests of the interpolation properties

Michel BERNADOU

INRIA,

Rocquencourt,

78153 Le Chesnay Cedex, France

Jean-Marie BOISSERIE

EDF/DER,

6 quai Watier,

78400 Chatou, France

### Abstract :

The aim of this work is to show how to implement some of the methods of curved finite elements of class  $\mathcal{C}^1$  introduced and analyzed in a previous work by the first author. These methods use curved finite elements which are  $\mathcal{C}^1$ -compatible with the Argyris triangle. The careful description of the implementation is completed by some numerical experiments which show the very high degree of accuracy of the associated interpolation. Subsequently, in a second part, we will show the efficiency of such methods for solving thin plate or thin shell problems set on curved boundary domains.

## ELEMENTS FINIS COURBES DE CLASSE $\mathcal{C}^1$ : IMPLEMENTATION ET TESTS NUMERIQUES

### Partie 1 : Construction et tests numériques des propriétés d'interpolation

### Résumé :

Le but de ce travail est de montrer comment implémenter certaines des méthodes d'éléments finis courbes de classe  $\mathcal{C}^1$  introduites et analysées dans un travail précédent du premier auteur. Ces méthodes utilisent des éléments finis courbes qui sont  $\mathcal{C}^1$ -compatibles avec le triangle d'Argyris. La description détaillée de l'implémentation est complétée par des tests numériques qui montrent le très haut degré de précision de l'interpolation correspondante. Ultérieurement, dans la seconde partie, nous illustrerons l'efficacité de ces méthodes pour la résolution de problèmes de plaques et de coques minces formulés sur des domaines à frontière curviligne.

---

(\*) This work is part of  
- the Project "Numerical Solution of Partial Differential Equations for Engineering Problems" of the Program  
"INRIA-NSF"  
- the Project "Junctions in Elastic Multi-Structures" of the Program "S.C.I.E.N.C.E." of the Commission of  
the European Communities (contract n° SC1-0473 - C(EDB))

(\*\*) To appear in Comput. Methods Appl. Mech. Engng.

## INTRODUCTION

The conforming approximation of thin plate problems with curved boundaries requires the use of curved  $C^1$ -finite elements which are compatible with some classical  $C^1$ -straight finite elements. A similar situation occurs for the conforming approximation of thin shells or for the conforming approximation of junctions between thin shells (see [1]).


We have introduced and analyzed such curved elements in [2,3] so that they are  $C^1$ -compatible with the ARGYRIS triangle [4,5,6]. These methods are particularly interesting to realize a very accurate approximation of thin plate or thin shell problems [2,3].

In this paper, we record the mains steps of the construction of curved finite elements which are  $C^1$ -compatible with the ARGYRIS triangle. and then we give a very detailed description of how to implement such methods. More precisely, paragraph 1 is dedicated to the construction of the approximated domain  $\Omega_h$  of the given curved boundary domain  $\Omega$  while paragraph 2 contains the main steps of the definition of the considered curved  $C^1$  finite element. Next, in paragraph 3, we detail the matrix decompositions which allow to realize the interpolation modules. Finally, in paragraph 4 we report some numerical experiments which prove the efficiency and the very high accuracy of such interpolation methods.

In addition it is worth to mention that all the relations have been checked by using the formal computational tools provided by Mathematica [7].

### Summary of the construction of these curved $C^1$ -elements

Let  $v$  be the function to be interpolated on the curved triangle  $K$ . The construction of



these curved  $C^1$ -elements takes several steps :

- i) triangulation of the curved boundary domain  $\Omega$
- ii) interpolation of the curved triangular sides located on the boundary
- iii) definition of the mapping  $F_K$  ;
- iv) computation of the set of values of the degrees of freedom  $\Sigma_K(v)$  of function  $v$  ;
- v) from  $\Sigma_K(v)$ , computation of the set of values of the degrees of freedom  $\hat{\Delta}_K(v)$  ;
- vi) from the set  $\hat{\Delta}_K(v)$ , computation of the interpolate function  $\hat{w}$  ;
- vii) computation of the interpolate function  $\pi_K v = w = \hat{w} \circ F_K^{-1}$ .

## Notations

For more details concerning curved  $C^1$  finite elements and for notations we refer to [2,3] and, more generally, finite element notations are those used by [5,6].

## Contents

### INTRODUCTION

### 1 - APPROXIMATION OF THE CURVED BOUNDARY DOMAIN $\Omega$

- 1.1. "Exact" triangulation of the domain  $\Omega$
- 1.2. "Approximate" triangulation of the domain  $\Omega$
- 1.3. Examples

### 2 - DEFINITION OF CURVED FINITE ELEMENTS $C^1$ -COMPATIBLE WITH ARGYRIS TRIANGLE

- 2.1. Basic principles
- 2.2. Definition of curved finite elements  $C^1$ -compatible with ARGYRIS triangle for  $F_K \in (P_3)^2$
- 2.3. Definition of curved finite elements  $C^1$ -compatible with ARGYRIS triangle for  $F_K^* \in (P_5)^2$

### 3 - IMPLEMENTATION OF CURVED FINITE ELEMENTS $C^1$ -COMPATIBLE WITH ARGYRIS TRIANGLE

- 3.1. Interpolation modulus associated to curved finite element  $C^1$ -compatible with ARGYRIS triangle when  $F_K \in (P_3)^2$
- 3.2. Interpolation modulus associated to curved finite element  $C^1$ -compatible with ARGYRIS triangle when  $F_K \in (P_5)^2$
- 3.3. Some numerical tests of interpolation properties

## ANNEX

## REFERENCES

### 1 APPROXIMATION OF THE CURVED BOUNDARY DOMAIN $\Omega$

This paragraph records some useful results obtained in [2,3].

#### 1.1 "Exact" triangulation of the domain $\Omega$

Let  $\Omega$  be a plane domain with a curved boundary  $\Gamma$ . We realize partitions of this domain by using regular families of triangulations  $T_h$  including (see Fig. 1.1.1) :

- i) straight triangles, on the one hand ;
- ii) triangles with one curved side located on  $\Gamma$ , on the other hand.

In this way, triangulations  $T_h$  can be splitted as follows :  $T_h = T_h^1 \cup T_h^2$  where  $T_h^1$  collects all the straight triangles  $K$  while  $T_h^2$  collects all the triangles  $K_c$  with one curved side on  $\Gamma$ .

According to Figure 1.1.1, we note  $\tilde{K}$  the straight triangles associated to a one-curved side triangle  $K_c$ .




Figure 1.1.1 : "Exact" triangulation of the domain  $\Omega$  made up of straight triangles  $K$  and one-curved side triangles  $K_c$ .

## 1.2 "Approximate" triangulation of the domain $\Omega$

Any one-curved side triangle  $K_c$  is approximated by a triangle  $K$ . In this way, the curved side of  $K_c$  is approximated by an arc  $\gamma_h$  (see Figure 1.2.1). Then the union of all straight triangles  $K \in T_h^1$  and of all one-curved side triangles  $K \in T_h^2$  give the approximate domain  $\Omega_h$ .




Figure 1.2.1 : "Exact" and "approximate" one-curved side triangles  $K_c$  and  $K$

Such a construction can be splitted in two steps :

**Step 1 : Interpolation of the curved side  $a_1a_2$  of the triangle  $K_c$ .** Assume that  $\mathbb{R}^2$  is referred to an orthonormal system  $(O, \vec{e}_1, \vec{e}_2)$  whose associate coordinates are noted

$(x_1, x_2)$  and that  $\Omega$  is a bounded domain. Moreover, assume that the boundary  $\Gamma$  can be subdivided in a finite number of arcs, each of them having a sufficiently smooth parametric representation of the following type :

$$(1.2.1) \quad x_1 = \chi_1(s), \quad x_2 = \chi_2(s), \quad \underline{s} \leq s \leq \bar{s}.$$

Subsequently, we also use another parameterization of the arc  $\widehat{a_1 a_2}$  :

$$(1.2.2) \quad x_1 = \psi_1(\hat{x}_2), \quad x_2 = \psi_2(\hat{x}_2), \quad 0 \leq \hat{x}_2 \leq 1,$$

where

$$(1.2.3) \quad \psi_\alpha(\hat{x}_2) = \chi_\alpha(\underline{s} + (\bar{s} - \underline{s})\hat{x}_2), \quad \alpha = 1, 2.$$

Then, every component  $\psi_\alpha(\hat{x}_2)$  is interpolated by a polynomial function  $\psi_{\alpha h}$  of degree  $n \geq 2$ , so that

$$(1.2.4) \quad \psi_{\alpha h}(0) = \psi_\alpha(0), \quad \psi_{\alpha h}(1) = \psi_\alpha(1), \quad \alpha = 1, 2.$$

Thus, for  $n \geq 2$ , we get :

$$(1.2.5) \quad \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 + \hat{x}_2(1 - \hat{x}_2)P_{n-2;\alpha}(\hat{x}_2)$$

where  $P_{n-2;\alpha}(\hat{x}_2)$  refers to polynomials of degree  $n - 2$  with respect to  $\hat{x}_2$ . These relations (1.2.5) define the approximate arc  $\gamma_h$ . With the notations of Fig. 1.2.2, observe that relation (1.2.5) can be geometrically interpreted as follows

$$(1.2.6) \quad \overrightarrow{OP_h} = \overrightarrow{O\tilde{P}} + \overrightarrow{\tilde{P}P_h}$$




Figure 1.2.2 : Construction of the approximate arc  $\gamma_h$

**Step 2 : Definition of the application  $F_K : \hat{K} \longrightarrow K$ .** Let  $\hat{K}$  be a given reference triangle, for instance the unit right-angled triangle  $\hat{a}_1 \hat{a}_2 \hat{a}_3$  with  $\hat{a}_1 = (1, 0)$ ,  $\hat{a}_2 = (0, 1)$ ,  $\hat{a}_3 = (0, 0)$ . Then, to any point  $\hat{M}$  of the reference triangle  $\hat{K}$ , the application  $F_K$  associates the point  $M_h$  :

$$(1.2.7) \quad \overrightarrow{OM_h} = F_{K1}(\hat{x}_1, \hat{x}_2)\vec{e}_1 + F_{K2}(\hat{x}_1, \hat{x}_2)\vec{e}_2$$


where the functions  $F_{K\alpha}$ ,  $\alpha = 1, 2$  are defined for  $n \geq 2$  by the relations

$$(1.2.8) \quad \left\{ \begin{array}{l} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 \\ \quad + \frac{1}{2} \hat{x}_1 \hat{x}_2 [P_{n-2;\alpha}(1 - \hat{x}_1) + P_{n-2;\alpha}(\hat{x}_2)]. \end{array} \right.$$

These components are symmetrical with respect to  $\hat{x}_1$  and  $\hat{x}_2$ , and they can be geometrically interpreted in the vectorial form (see Figure 1.2.3) :

$$(1.2.9) \quad \overrightarrow{OM_h} = \overrightarrow{O\tilde{M}} + \frac{1}{2} \overrightarrow{\tilde{M}M_h^1} + \frac{1}{2} \overrightarrow{\tilde{M}M_h^2}$$

The properties of this application  $F_K : \hat{K} \longrightarrow K$  are detailed in [2].



**Figure 1.2.3 :** Transformation  $\tilde{M} \longrightarrow M_h$  (triangles  $Q^1 \tilde{M} M_h^1$  and  $Q^1 \tilde{P}^1 P_h^1$  are homothetic in the ratio  $\hat{x}_2/(1 - \hat{x}_1)$  and triangles  $Q^2 \tilde{M} M_h^2$  and  $Q^2 \tilde{P}^2 P_h^2$  are homothetic in the ratio  $\hat{x}_1/(1 - \hat{x}_2)$ ).

### 1.3 Examples

Hereunder, we give two examples of applications  $\psi_{\alpha h}$  (see (1.2.5)) which define an approximate arc  $\gamma_h$  interpolating a given arc  $a_1 a_2$ . These examples correspond to interpolations by polynomials of degree  $n = 3$  or  $5$  which are in practice the most interesting. For each case, we indicate the expression of the applications  $\psi_{\alpha h}$  from which we deduce the expression of the components  $F_{K\alpha}$ ,  $\alpha = 1, 2$ . Afterwards, these expressions are permanently used ; in order to make distinction between them, we note  $F_K$  the one which is associated to the example 1.3.1 (i.e.,  $F_K \in (P_3)^2$ ) and  $F_K^*$  the one which corresponds to example 1.3.2 (i.e.,  $F_K^* \in (P_5)^2$ ).


**Example 1.3.1 : Construction of the approximate arc  $\gamma_h$  by using polynomials of degree 3 (Figure 1.3.1)**

Expressions (1.2.5) give

$$(1.3.1) \quad \left\{ \begin{array}{l} \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 + \hat{x}_2(1 - \hat{x}_2)\{[2(x_{\alpha 2} - x_{\alpha 1}) \\ - (\bar{s} - s)(\chi'_{\alpha}(s) + \chi'_{\alpha}(\bar{s}))]\hat{x}_2 + x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - s)\chi'_{\alpha}(s)\} \end{array} \right.$$

so that from (1.2.8), we obtain

$$(1.3.2) \quad \left\{ \begin{array}{l} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 + \frac{1}{2}\hat{x}_1\hat{x}_2\{[2(x_{\alpha 2} - x_{\alpha 1}) \\ - (\bar{s} - s)(\chi'_{\alpha}(s) + \chi'_{\alpha}(\bar{s}))][\hat{x}_2 - \hat{x}_1] + (\bar{s} - s)[\chi'_{\alpha}(s) - \chi'_{\alpha}(\bar{s})]\} \end{array} \right.$$



**Figure 1.3.1 :** Boundary interpolation with polynomials of degree 3

**Example 1.3.2 : Construction of the approximate arc  $\gamma_h^*$  by using polynomials of degree 5**

We interpolate the functions  $\psi_{\alpha}$ ,  $\alpha = 1, 2$ , on the interval  $[0, 1]$  by using Hermite polynomials of degree 5. The degrees of freedom of the interpolation are the values of the functions  $\psi_{\alpha}$ ,  $\psi'_{\alpha}$  and  $\psi''_{\alpha}$  at points  $\hat{x}_2 = 0$  or  $\hat{x}_2 = 1$ , i.e.,

$$(1.3.3) \quad \left\{ \begin{array}{l} x_{\alpha 1} = \psi_{\alpha}(0) = \chi_{\alpha}(s), \quad x_{\alpha 2} = \psi_{\alpha}(1) = \chi_{\alpha}(\bar{s}), \\ \psi_{\alpha}^{(\ell)}(0) = (\bar{s} - s)^{\ell}\chi_{\alpha}^{(\ell)}(s), \quad \psi_{\alpha}^{(\ell)}(1) = (\bar{s} - s)^{\ell}\chi_{\alpha}^{(\ell)}(\bar{s}), \quad \ell = 1, 2. \end{array} \right.$$

From the expression (1.2.5) we obtain (we recall that all the results related to the example 1.3.2 are indexed with a star (\*) in order to make the distinction with similar results related to the example 1.3.1)

$$(1.3.4) \quad \psi_{\alpha h}^*(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 + \hat{x}_2(1 - \hat{x}_2)[\beta_{\alpha 3}(\hat{x}_2)^3 + \beta_{\alpha 2}(\hat{x}_2)^2 + \beta_{\alpha 1}\hat{x}_2 + \beta_{\alpha 0}]$$

where the coefficients  $\beta_{\alpha \ell}$ ,  $\ell = 0, 1, 2, 3$  are given by the relations

$$(1.3.5) \quad \left\{ \begin{array}{l} \beta_{\alpha 0} = x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - s)\chi'_{\alpha}(s) \\ \beta_{\alpha 1} = x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - s)\chi'_{\alpha}(s) + \frac{(\bar{s} - s)^2}{2}\chi''_{\alpha}(s) \\ \beta_{\alpha 2} = 9(x_{\alpha 2} - x_{\alpha 1}) - (\bar{s} - s)[5\chi'_{\alpha}(s) + 4\chi'_{\alpha}(\bar{s})] - \frac{(\bar{s} - s)^2}{2}[2\chi''_{\alpha}(s) - \chi''_{\alpha}(\bar{s})] \\ \beta_{\alpha 3} = 6(x_{\alpha 1} - x_{\alpha 2}) + 3(\bar{s} - s)[\chi'_{\alpha}(s) + \chi'_{\alpha}(\bar{s})] + \frac{(\bar{s} - s)^2}{2}[\chi''_{\alpha}(s) - \chi''_{\alpha}(\bar{s})] \end{array} \right.$$

From the expressions (1.2.8) and (1.3.4), we deduce

$$(1.3.6) \quad \left\{ \begin{array}{l} F_{K\alpha}^*(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 \\ \quad + \frac{1}{2}\hat{x}_1\hat{x}_2[\beta_{\alpha 3}(\hat{x}_2)^3 + \beta_{\alpha 2}(\hat{x}_2)^2 + \beta_{\alpha 1}\hat{x}_2 + \beta_{\alpha 0} \\ \quad + \tilde{\beta}_{\alpha 3}(\hat{x}_1)^3 + \tilde{\beta}_{\alpha 2}(\hat{x}_1)^2 + \tilde{\beta}_{\alpha 1}\hat{x}_1 + \tilde{\beta}_{\alpha 0}] \end{array} \right.$$

where the coefficients  $\beta_{\alpha \ell}$ ,  $\ell = 0, 1, 2, 3$  are given by relation (1.3.5) and where coefficients  $\tilde{\beta}_{\alpha \ell}$ ,  $\ell = 0, 1, 2, 3$  are given by

$$(1.3.7) \quad \left\{ \begin{array}{l} \tilde{\beta}_{\alpha 0} = x_{\alpha 2} - x_{\alpha 1} - (\bar{s} - s)\chi'_{\alpha}(\bar{s}) \\ \tilde{\beta}_{\alpha 1} = x_{\alpha 2} - x_{\alpha 1} - (\bar{s} - s)\chi'_{\alpha}(\bar{s}) + \frac{(\bar{s} - s)^2}{2}\chi''_{\alpha}(\bar{s}) \\ \tilde{\beta}_{\alpha 2} = 9(x_{\alpha 1} - x_{\alpha 2}) + (\bar{s} - s)[5\chi'_{\alpha}(\bar{s}) + 4\chi'_{\alpha}(s)] - \frac{(\bar{s} - s)^2}{2}[2\chi''_{\alpha}(\bar{s}) - \chi''_{\alpha}(s)] \\ \tilde{\beta}_{\alpha 3} = 6(x_{\alpha 2} - x_{\alpha 1}) - 3(\bar{s} - s)[\chi'_{\alpha}(\bar{s}) + \chi'_{\alpha}(s)] + \frac{(\bar{s} - s)^2}{2}[\chi''_{\alpha}(\bar{s}) - \chi''_{\alpha}(s)] \end{array} \right.$$

## 2 DEFINITION OF CURVED FINITE ELEMENTS, $C^1$ -COMPATIBLE WITH ARGYRIS TRIANGLE

In this paragraph we record the definitions of **two curved finite elements** which have a connection of class  $C^1$  with the classical ARGYRIS triangle.

i) the **first** corresponds to the interpolation of the boundary considered in the example 1.3.1. This interpolation is realized by using polynomials of degree 3 and it turns out to be sufficient for the approximation of fourth-order problems with homogeneous Dirichlet boundary conditions (see [3]) ;

ii) the **second** corresponds to the interpolation of the boundary described in the example 1.3.2. This interpolation is realized by using polynomials of degree 5 and it can be used for more general boundary conditions (see [3, Remark 3.1]).

Both constructions are fully detailed in [2,3] where, in addition, one can find necessary justifications and corresponding interpolation error estimates. The implementation of these methods is detailed in matrix form in paragraph 3.

## 2.1 Basic principles


We consider separately **essential** and **desirable** conditions.

### Essential conditions :

According to Figure 1.2.1, the connections between ARGYRIS triangles and curved elements are made along the straight sides  $a_3a_1$  and  $a_3a_2$ . Let  $(K, P_K, \Sigma_K)$  be the approximate curved triangle, its associated functional space and its corresponding set of degrees of freedom. To obtain a connection of class  $\mathcal{C}^1$ , it is essential to satisfy (see Figure 2.1.1) :

$$(2.1.1) \quad \left\{ \begin{array}{l} \text{the degrees of freedom of the curved finite elements related to the sides} \\ a_3a_1 \text{ and } a_3a_2 \text{ are identical to those of the ARGYRIS triangle ;} \end{array} \right.$$

$$(2.1.2) \quad \left\{ \begin{array}{l} \text{the traces } p|_{[a_3,a_\alpha]}, \alpha = 1, 2 (\text{resp. } Dp(\cdot)(a_2 - c_2)|_{[a_3,a_1]}; Dp(\cdot)(a_1 - c_1)|_{[a_3,a_2]}) \\ \text{of the functions } p \in P_K \text{ defined on the curved triangle } K, \text{ are one-variable} \\ \text{polynomial of degree 5 (resp. 4), entirely determined by the degrees of} \\ \text{freedom related to the sides } a_3a_\alpha, \alpha = 1, 2. \end{array} \right.$$



**Figure 2.1.1** : The approximate curved triangle  $K$

**Desirable conditions** : The application  $F_K$ , introduced in paragraph 1, associates the curved triangle  $K$  to the reference triangle  $\hat{K}$ .

We use this application  $F_K$  as well to associate at any function  $v$  defined on the triangle  $K$ , one function  $\hat{v}$  defined on the triangle  $\hat{K}$ , i.e.,

$$(2.1.3) \quad v = \hat{v} \circ (F_K)^{-1} ; \quad \hat{v} = v \circ F_K.$$

Consequently, it is "desirable" that the following condition is satisfied :

$$(2.1.4) \quad \left\{ \begin{array}{l} \text{To any function } p \in P_K, \text{ defined on the curved triangle } K, \text{ the} \\ \text{correspondence (2.1.3) associates a polynomial function } \hat{p} = p \circ F_K. \end{array} \right.$$

This condition (2.1.4) is convenient to study the approximation error, to study the effect of numerical integration scheme and to take into account the boundary conditions. But this condition leads to the definition of reference finite elements which are most complicated than those associated to corresponding straight side finite elements. Indeed, with notations of Figure 2.1.1, we obtain for  $a \in [a_3a_1]$  :

$$(2.1.5) \quad \left\{ \begin{array}{l} \frac{\partial \hat{p}}{\partial \hat{x}_2} (\hat{a}) = D\hat{p}(\hat{a})\vec{e}_2 = Dp(a)DF_K(\hat{a})\vec{e}_2 = Dp(a)\frac{\partial F_K}{\partial \hat{x}_2} (\hat{a}) \\ = \left\langle \frac{\partial F_K}{\partial \hat{x}_2} (\hat{a}), \frac{a_1 - a_3}{|a_1 - a_3|^2} Dp(a)(a_1 - a_3) + \frac{a_2 - c_2}{|a_2 - c_2|^2} Dp(a)(a_2 - c_2) \right\rangle \end{array} \right.$$

where  $\langle ., . \rangle$  denotes the usual scalar product in  $\mathbb{R}^2$ . Likewise for  $a \in [a_3a_2]$ , we get :

$$(2.1.6) \quad \frac{\partial \hat{p}}{\partial \hat{x}_1} (\hat{a}) = \left\langle \frac{\partial F_K}{\partial \hat{x}_1} (\hat{a}), \frac{a_2 - a_3}{|a_2 - a_3|^2} Dp(a)(a_2 - a_3) + \frac{a_1 - c_1}{|a_1 - c_1|^2} Dp(a)(a_1 - c_1) \right\rangle.$$

Relations (2.1.5) and (2.1.6) prove that  $\frac{\partial \hat{p}}{\partial \hat{x}_2} (\hat{a})$  (resp.  $\frac{\partial \hat{p}}{\partial \hat{x}_1} (\hat{a})$ ) is a polynomial of degree  $n+3$  with respect to  $\hat{x}_1$  (resp.  $\hat{x}_2$ ) for any  $\hat{a} \in [\hat{a}_3\hat{a}_1]$  (resp.  $\hat{a} \in [\hat{a}_3\hat{a}_2]$ ), with  $n=3$  or  $5$  depending on whether  $F_K$  is of degree  $3$  or  $5$  (see examples 1.3.1 and 1.3.2). Thus to  $\hat{K}$  we have to associate a finite element  $(\hat{K}, \hat{P}, \hat{\Sigma})$  such that

$$(2.1.7) \quad \hat{P}_K \subset P_{n+4} \quad (n=3 \text{ or } 5); \quad \hat{P}_K = \{\hat{p} : \hat{K} \rightarrow \mathbb{R}; \quad \hat{p} = p \circ F_K, \quad p \in P_K\}.$$

## 2.2 Definition of curved finite elements $C^1$ -compatible with AR-GYRIS triangle for $F_K \in (P_3)^2$

In all this section we consider the application  $F_K$  defined in the example 1.3.1. Then, relation (2.1.7) gives  $\hat{P}_K \subset P_7$ .

### 2.2.1 The basic finite element $(\hat{K}, \hat{P}, \hat{\Sigma})$


This finite element is described in Figure 2.2.1 and we refer the reader to [8] for more details. Corresponding basis functions are given by relation (3.1.43).

### 2.2.2 Construction of the interpolating function $v \rightarrow \pi_K v$

Now we use this basic finite element in order to associate an interpolate function  $\pi_K v$  to any function  $v \in C^2(K)$ . This construction takes three steps :

**Step 1 : Definition of the set  $\Sigma_K$  of the degrees of freedom of the curved element.** Set (see Figures 2.2.1 and 2.2.2)

$$(2.2.1) \quad \left\{ \begin{array}{l} a_i = F_K(\hat{a}_i), \quad b_i = F_K(\hat{b}_i), \quad e_i = F_K(\hat{e}_i), \quad i = 1, 2, 3; \\ c_\alpha = \text{orthogonal projection of } a_\alpha \text{ on the side } a_3a_\alpha, \quad \alpha = 1, 2, . \end{array} \right.$$



$\hat{K}$  = unit right-angled triangle ;  $\hat{P} = P_7$  ;  $\dim \hat{P} = 36$

$$\left\{ \begin{array}{l} \hat{\Sigma}(\hat{p}) = [\hat{p}(\hat{a}_i), \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a}_i), \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_1^2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_2^2}(\hat{a}_i), i = 1, 2, 3; \\ - \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{b}_1); - \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{b}_2); \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right)(\hat{b}_3); \hat{p}(\hat{d}_i), i = 1, \dots, 6; \\ - \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{d}_i), i = 1, 2; - \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{d}_i), i = 3, 4; \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right)(\hat{d}_i), i = 5, 6; \\ \hat{p}(\hat{e}_i), i = 1, 2, 3] \end{array} \right.$$

**Figure 2.2.1 :** Basic finite element for the construction of a curved finite element  $C^1$ -compatible with the ARGYRIS triangle ( $F_K \in (P_3)^2$ ).

Then, the set  $\Sigma_K(v)$  of degrees of freedom of function  $v \in \mathcal{C}^2(K)$  is given, in its local version [ $DLLC(v)$ ], by


$$(2.2.2) \quad \left\{ \begin{array}{l} [DLLC(v)]_{1 \times 24} = [v(a_1) \ v(a_2) \ v(a_3) \ Dv(a_1)(a_3 - a_1) \ (\bar{s} - s)Dv(a_1)\vec{\chi}'(s) \\ (\bar{s} - s)Dv(a_2)\vec{\chi}'(\bar{s}) \ Dv(a_2)(a_3 - a_2) \ Dv(a_3)(a_2 - a_3) \ Dv(a_3)(a_1 - a_3) \\ D^2v(a_1)(a_3 - a_1)^2 \ (\bar{s} - s)^2 D^2v(a_1)(\vec{\chi}'(s))^2 \ (\bar{s} - s)^2 D^2v(a_2)(\vec{\chi}'(\bar{s}))^2 \\ D^2v(a_2)(a_3 - a_2)^2 \ D^2v(a_3)(a_2 - a_3)^2 \ D^2v(a_3)(a_1 - a_3)^2 \ D^2v(a_1)(a_2 - a_3)^2 \\ D^2v(a_2)(a_3 - a_1)^2 \ (\bar{s} - s)^2 D^2v(a_3)((\vec{\chi}'(s), \vec{\chi}'(\bar{s})) \ Dv(b_1)(a_1 - c_1) \\ Dv(b_2)(a_2 - c_2) \ Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \ v(e_1) \ v(e_2) \ v(e_3)] \end{array} \right.$$

**Step 2 : Transition from  $\Sigma_K(v)$  to  $\hat{\Delta}_K(v)$  :** Starting from the set  $\Sigma_K(v)$  of 24 elements we introduce the set  $\hat{\Delta}_K(v)$  of 36 values that we need to attribute to the set  $\hat{\Sigma}$  of degrees of freedom in order to obtain a suitable interpolated function  $\hat{w} \in \hat{P}$ . In this way, it is convenient to introduce the following partition of the set  $\hat{\Sigma}$  :

$$(2.2.3) \quad \left\{ \begin{array}{l} \hat{\Sigma}(\hat{p}) = \hat{\Sigma}_1(\hat{p}) \cup \hat{\Sigma}_2(\hat{p}) \cup \hat{\Sigma}_3(\hat{p}), \text{ where} \\ \\ \hat{\Sigma}_1(\hat{p}) = \{(D^\alpha \hat{p}(\hat{a}_i); |\alpha| = 0, 1, 2; i = 1, 2, 3); \hat{p}(\hat{e}_i), i = 1, 2, 3\} \\ \\ \hat{\Sigma}_2(\hat{p}) = \left\{ \hat{p}(\hat{d}_i), i = 1, \dots, 4; -\frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{b}_1); -\frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{d}_i), i = 1, 2; \right. \\ \left. -\frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{b}_2); -\frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{d}_i), i = 3, 4 \right\} \\ \\ \hat{\Sigma}_3(\hat{p}) = \left\{ \hat{p}(\hat{d}_i), i = 5, 6; \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right)(\hat{b}_3); \right. \\ \left. \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right)(\hat{d}_i), i = 5, 6 \right\} \end{array} \right.$$

and the associate partition of  $\hat{\Delta}_K(v)$  :

$$(2.2.4) \quad \hat{\Delta}_K(v) = \hat{\Delta}_{K1}(v) \cup \hat{\Delta}_{K2}(v) \cup \hat{\Delta}_{K3}(v).$$



**Figure 2.2.2 :** The set of degrees of freedom  $\Sigma_K$  for the curved finite elements  $C^1$ -compatible with the ARGYRIS finite elements ( $F_K \in (P_3)^2$ ).

Then, from the knowledge of the set  $\Sigma_K(v)$  of values of the degrees of freedom of  $v$ , we construct the corresponding set of values  $\hat{\Delta}_{K_i}(v)$ ,  $i = 1, 2, 3$  :

**Construction of  $\hat{\Delta}_{K1}(v)$  :** The knowledge of the application  $F_K$  and of the set of values  $\{(D^\alpha v(a_i), |\alpha| = 0, 1, 2; v(e_i)), i = 1, 2, 3\}$  leads immediately to  $\hat{\Delta}_{K1}(v)$ . It suffices to use relations (2.1.3) so that

$$(2.2.5) \quad \hat{\Delta}_{K1}(v) = \{(D^\alpha \hat{v}(\hat{a}_i), |\alpha| = 0, 1, 2 ; i = 1, 2, 3) ; \hat{v}(\hat{e}_i), i = 1, 2, 3\}.$$

This correspondence is detailed in section 3.1.

**Construction of  $\hat{\Delta}_{K2}(v)$**  : The construction of this set of values is much more tricky. Firstly, let us examine the case of the degrees of freedom of  $\hat{\Sigma}_2$  located on the side  $\hat{a}_3\hat{a}_1$ . To verify conditions (2.1.2), the expected interpolating function  $\pi_K v$  has to satisfy :

\* **On the one hand**, its trace  $\pi_K v|_{[a_3, a_1]}$  coincides with the one-variable Hermite polynomial of degree 5, determined by the data of the degrees of freedom

$$(2.2.6) \quad \left\{ \begin{array}{l} \{v(a_1), v(a_3), Dv(a_1)(a_3 - a_1), Dv(a_3)(a_1 - a_3), \\ D^2v(a_1)(a_3 - a_1)^2, D^2v(a_3)(a_1 - a_3)^2\}. \end{array} \right.$$

We will use  $\hat{x}_1$  as a parameterization variable of the side  $a_3a_1$ , i.e.,

$$(2.2.7) \quad x_1 = x_{13} + (x_{11} - x_{13})\hat{x}_1, \quad x_2 = x_{23} + (x_{21} - x_{23})\hat{x}_1$$

where  $x_{\alpha i}$  ( $\alpha = 1, 2$  ;  $i = 1, 2, 3$ ) notes the  $\alpha^{th}$ -coordinate of vertex  $a_i$ . Corresponding Hermite polynomial is named

$$(2.2.8) \quad \hat{f}_1 \text{ (which has to coincide with } (\pi_K v) \circ F_K|_{[\hat{a}_3, \hat{a}_1]})$$

\* **On the other hand**, the trace  $(D\pi_K v(.)(a_2 - c_2)) \circ F_K|_{[\hat{a}_3, \hat{a}_1]}$  of its normal derivative  $D\pi_K v(.)(a_2 - c_2)$  coincides with the one variable Hermite polynomial of degree 4 determined by the data of the degrees of freedom

$$(2.2.9) \quad \{Dv(a_i)(a_2 - c_2), D^2v(a_i)(a_2 - c_2, a_1 - a_3), i \in \{1, 3\} ; Dv(b_2)(a_2 - c_2)\}.$$

In a similar way to (2.2.7) and (2.2.8), the corresponding Hermite polynomial is named

$$(2.2.10) \quad \hat{g}_1 \text{ (which has to coincide with } (D\pi_K v(.)(a_2 - c_2)) \circ F_K|_{[\hat{a}_3, \hat{a}_1]}).$$

By analogy with the above definitions we could define the following Hermite polynomials over the second straight side  $a_3a_2$  of the triangle  $K$ , i.e.,

$$(2.2.11) \quad x_1 = x_{13} + (x_{12} - x_{13})\hat{x}_2, \quad x_2 = x_{23} + (x_{22} - x_{23})\hat{x}_2$$

$$(2.2.12) \quad \hat{f}_2 \text{ (which has to coincide with } (\pi_K v) \circ F_K|_{[\hat{a}_3, \hat{a}_2]})$$

$$(2.2.13) \quad \hat{g}_2 \text{ (which has to coincide with } (D\pi_K v(.)(a_1 - c_1)) \circ F_K|_{[\hat{a}_3, \hat{a}_2]}).$$

Then, the set of values  $\hat{\Delta}_{K2}(v)$  associated to the set of degrees of freedom  $\hat{\Sigma}_2$  (see (2.1.5) (2.1.6) (2.2.3)) is given by

$$(2.2.14) \quad \left\{ \begin{array}{l} \hat{\Delta}_{K2}(v) = [\hat{f}_2(\hat{d}_i), i = 1, 2; \hat{f}_1(\hat{d}_i), i = 3, 4; \\ - < \frac{\partial F_K}{\partial \hat{x}_1}(\hat{b}_1), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d\hat{f}_2}{d\hat{x}_2}(\hat{b}_1) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{g}_2(\hat{b}_1) >; \\ - < \frac{\partial F_K}{\partial \hat{x}_1}(\hat{d}_i), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d\hat{f}_2}{d\hat{x}_2}(\hat{d}_i) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{g}_2(\hat{d}_i) >, i = 1, 2; \\ - < \frac{\partial F_K}{\partial \hat{x}_2}(\hat{b}_2), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{b}_2) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{g}_1(\hat{b}_2) >; \\ - < \frac{\partial F_K}{\partial \hat{x}_2}(\hat{d}_i), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{d}_i) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{g}_1(\hat{d}_i) >, i = 3, 4]. \end{array} \right.$$

Note that the definition of the last six elements uses relations (2.1.6) and (2.1.5), specifications (2.2.12) (2.2.13) (2.2.8) (2.2.10), and :

$$(2.2.15) \quad \left\{ \begin{array}{l} \frac{d\hat{f}_2}{d\hat{x}_2} \text{ which will have to coincide with } (D\pi_K v(\cdot)(a_2 - a_3)) \circ F_K|_{[\hat{a}_3, \hat{a}_2]} \\ \frac{d\hat{f}_1}{d\hat{x}_1} \text{ which will have to coincide with } (D\pi_K v(\cdot)(a_1 - a_3)) \circ F_K|_{[\hat{a}_3, \hat{a}_1]} \end{array} \right.$$

**Construction of  $\hat{\Delta}_{K3}(v)$**  : It remains to compute the values that we have to attribute to the five degrees of freedom of  $\hat{\Sigma}_3$ . Firstly the correspondence  $\hat{v} = v \circ F_K$  leads to

$$(2.2.16) \quad D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3),$$

which is known since this value is among the degrees of freedom of  $\Sigma_K$  (see (2.2.2)).

Next, the values  $D^\alpha \hat{v}(\hat{a}_i)$ ,  $|\alpha| = 0, 1, 2$ ;  $i = 1, 2, 3$  are listed in (2.2.5) while the side  $\hat{a}_1 \hat{a}_2$  of the triangle  $\hat{K}$  can be parameterized by

$$(2.2.17) \quad \hat{x}_1 = \hat{x}_1, \quad \hat{x}_2 = 1 - \hat{x}_1.$$

Then, let  $\hat{f}_3(\hat{x}_1)$  be the Hermite polynomial of degree 5, defined by the data of the degrees of freedom

$$(2.2.18) \quad \left\{ \begin{array}{l} \{\hat{v}(\hat{a}_1), \hat{v}(\hat{a}_2), D\hat{v}(\hat{a}_1)(\hat{a}_2 - \hat{a}_1), D\hat{v}(\hat{a}_2)(\hat{a}_1 - \hat{a}_2), \\ D^2\hat{v}(\hat{a}_1)(\hat{a}_2 - \hat{a}_1)^2, D^2\hat{v}(\hat{a}_2)(\hat{a}_1 - \hat{a}_2)^2\} \end{array} \right.$$

and let  $\hat{g}_3(\hat{x}_1)$  be the Hermite polynomial of degree 4, determined by the data of the degrees of freedom

$$(2.2.19) \quad \left\{ \begin{array}{l} \{D\hat{v}(\hat{a}_\alpha)(\hat{a}_3 - \hat{b}_3), \alpha = 1, 2; D^2\hat{v}(\hat{a}_1)(\hat{a}_3 - \hat{b}_3, \hat{a}_2 - \hat{a}_1), \\ D^2\hat{v}(\hat{a}_2)(\hat{a}_3 - \hat{b}_3, \hat{a}_1 - \hat{a}_2), D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3)\}. \end{array} \right.$$

Then, we set

$$(2.2.20) \quad \hat{\Delta}_{K3}(v) = \left\{ \hat{f}_3\left(\frac{3}{4}\right); \hat{f}_3\left(\frac{1}{4}\right); -\sqrt{2}\hat{g}_3\left(\frac{1}{2}\right); -\sqrt{2}\hat{g}_3\left(\frac{3}{4}\right); -\sqrt{2}\hat{g}_3\left(\frac{1}{4}\right) \right\}.$$

Thus, only from the knowledge of the values  $\Sigma_K(v)$  of the degrees of freedom of the function  $v$ , the relations (2.2.5) (2.2.14) and (2.2.20) assign one and only one value to every degree of freedom of  $\hat{\Sigma}$ . In view of the implementation, we will give a matrix presentation of these correspondences in paragraph 3.

**Step 3 : Transition from the set of values  $\hat{\Delta}_K(v)$  to the interpolate function  $w = \pi_K v$  :** Let  $\hat{w}$  be the function of  $\hat{P}$  which takes the set of values  $\hat{\Delta}_K(v)$  on the set of degrees of freedom  $\hat{\Sigma}$  (see Figure 2.2.1). Then, to  $\hat{w}$  defined on the reference triangle  $\hat{K}$ , relations (2.1.3) associate the function

$$(2.2.21) \quad \pi_K v = w = \hat{w} \circ (F_K)^{-1}$$

where the function  $(F_K)^{-1}$  is the inverse of function  $F_K : \hat{K} \longrightarrow K$  explicit in Example 1.3.1 (see relation (1.3.2)). ■

Finally, according to [2], this function  $w$  is exactly the expected function  $\pi_K v$ , i.e.,  $w = \pi_K v$ , since :

**Theorem 2.2.1 :** The function  $w$ , defined in (2.2.21) is determined in a unique way by the data (2.2.2) of the local version  $[DLIC(v)]$  of the set  $\Sigma_K(v)$  of the values of the degrees of freedom of the function  $v$  to interpolate. Moreover the function  $w$  satisfies

$$(2.2.22) \quad \begin{cases} D^\alpha w(a_i) = D^\alpha v(a_i), \ |\alpha| = 0, 1, 2; i = 1, 2, 3; \\ Dw(b_1)(a_1 - c_1) = Dv(b_1)(a_1 - c_1) \\ Dw(b_2)(a_2 - c_2) = Dv(b_2)(a_2 - c_2) \\ Dw(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \\ w(e_i) = v(e_i), \ i = 1, 2, 3 \end{cases}$$


and the conditions (2.1.1) (2.1.2) and (2.1.4). Thus  $w = \pi_K v$ . ■

## 2.3 Definition of curved finite elements $C^1$ -compatible with AR-GYRIS elements for $F_K^* \in (P_5)^2$

Since the construction is very similar to the case  $F_K \in (P_3)^2$  we only detail the new aspects. Firstly, relation (2.1.7) involves  $\hat{P}_K^* \subset P_9$  (let us record that a star (\*) makes the distinction between the cases  $F_K \in (P_3)^2$  and  $F_K^* \in (P_5)^2$ ).

### 2.3.1 The basic finite element $(\hat{K}, \hat{P}^*, \hat{\Sigma}^*)$

This element is described in Figure 2.3.1 and basis functions are given by (3.2.25).



$$\hat{K} = \text{unit right-angled triangle} ; \quad \hat{P}^* = P_9 ; \quad \dim \hat{P}^* = 55$$

$$\left\{ \begin{aligned} \hat{\Sigma}^*(\hat{p}) = & \left\{ \hat{p}(\hat{a}_i), \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a}_i), \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_1^2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{p}}{\partial \hat{x}_2^2}(\hat{a}_i), i = 1, 2, 3 ; \right. \\ & - \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{b}_1) ; - \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{b}_2) ; \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{b}_3) ; \hat{p}(\hat{d}_i^*), i = 1, \dots, 12 ; \\ & - \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{d}_i^*), i = 1, \dots, 4 ; - \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{d}_i^*), i = 5, \dots, 8 ; \\ & \left. \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{d}_i^*), i = 9, \dots, 12 ; \hat{p}(\hat{e}_i^*), i = 1, \dots, 10 \right\} \end{aligned} \right.$$

**Figure 2.3.1 :** Basic finite element for the construction of a curved finite element  $\mathcal{C}^1$ -compatible with the ARGYRIS triangle ( $F_K^* \in (P_5)^2$ ).

### 2.3.2 Construction of the interpolating function $v \rightarrow \pi_K^* v$


Like in section 2.2, this construction can be splitted in three steps :

**Step 1 : Definition of the set  $\Sigma_K^*$  of degrees of freedom of the curved element.**  
Set (see Figures 2.3.1 and 2.3.2) :

$$(2.3.1) \quad a_i = F_K^*(\hat{a}_i); \quad b_i = F_K^*(\hat{b}_i), \quad i = 1, 2, 3; \quad e_i^* = F_K^*(\hat{e}_i^*), \quad i = 1, \dots, 10$$

while  $c_\alpha$  is still the orthogonal projection of  $a_\alpha$  on the side  $a_3 a_\alpha$ ,  $\alpha = 1, 2$ . Then the set  $\Sigma_K^*(v)$  of degrees of freedom of the function  $v \in C^2(K)$  is given, in its local version [ $DLLC^*(v)$ ], by

$$(2.3.2) \quad \left\{ \begin{array}{l} \Sigma_K^*(v) = [DLLC^*(v)]_{1 \times 31} = [v(a_1) \quad v(a_2) \quad v(a_3) \quad Dv(a_1)(a_3 - a_1) \\ (\bar{s} - \underline{s})Dv(a_1)\vec{\chi}'(\underline{s}) \quad (\underline{s} - \bar{s})Dv(a_2)\vec{\chi}'(\bar{s}) \quad Dv(a_2)(a_3 - a_2) \\ Dv(a_3)(a_2 - a_3) \quad Dv(a_3)(a_1 - a_3) \quad D^2v(a_1)(a_3 - a_1)^2 \\ (\bar{s} - \underline{s})^2D^2v(a_1)(\vec{\chi}'(\underline{s}))^2 \quad (\underline{s} - \bar{s})^2D^2v(a_2)(\vec{\chi}'(\bar{s}))^2 \quad D^2v(a_2)(a_3 - a_2)^2 \\ D^2v(a_3)(a_2 - a_3)^2 \quad D^2v(a_3)(a_1 - a_3)^2 \quad D^2v(a_1)(a_2 - a_3)^2 \\ D^2v(a_2)(a_3 - a_1)^2 \quad (\bar{s} - \underline{s})^2D^2v(a_3)((\vec{\chi}'(\underline{s}), \vec{\chi}'(\bar{s})) \quad Dv(b_1)(a_1 - c_1) \\ Dv(b_2)(a_2 - c_2) \quad Dv(b_3)DF_K^*(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \quad v(e_1^*) \dots v(e_{10}^*)] \end{array} \right.$$



**Figure 2.3.2 :** The set of degrees of freedom  $\Sigma_K$  for the curved finite elements  $C^1$ -compatible with the ARGYRIS finite element ( $F_K^* \in (P_5)^2$ ).

**Step 2 : Transition from  $\Sigma_K^*(v)$  to  $\hat{\Delta}_K^*(v)$  :** Starting from the set  $\Sigma_K^*(v)$  including 31 elements we define the set  $\hat{\Delta}_K^*(v)$  of 55 values that we need to attribute to the set  $\hat{\Sigma}^*$  of degrees of freedom in order to obtain a suitable interpolated function  $\hat{w} \in \hat{P}$ . In this way we consider the following partition of the set  $\hat{\Sigma}^*$  :

$$(2.3.3) \quad \left\{ \begin{array}{l} \hat{\Sigma}^*(\hat{p}) = \hat{\Sigma}_1^*(\hat{p}) \cup \hat{\Sigma}_2^*(\hat{p}) \cup \hat{\Sigma}_3^*(\hat{p}), \text{ where} \\ \\ \hat{\Sigma}_1^*(\hat{p}) = \{(D^\alpha \hat{p}(\hat{a}_i) ; |\alpha| = 0, 1, 2 ; i = 1, 2, 3) ; \hat{p}(\hat{e}_i^*), i = 1, \dots, 10\} \\ \\ \hat{\Sigma}_2^*(\hat{p}) = \left\{ \begin{array}{l} \hat{p}(\hat{d}_i^*), i = 1, \dots, 8 ; - \frac{\partial \hat{p}}{\partial \hat{x}_1} (\hat{b}_1) ; - \frac{\partial \hat{p}}{\partial \hat{x}_1} (\hat{d}_i), i = 1, \dots, 4 ; \\ - \frac{\partial \hat{p}}{\partial \hat{x}_2} (\hat{b}_2) ; - \frac{\partial \hat{p}}{\partial \hat{x}_2} (\hat{d}_i), i = 5, \dots, 8 \end{array} \right\} \\ \\ \hat{\Sigma}_3^*(\hat{p}) = \left\{ \begin{array}{l} \hat{p}(\hat{d}_i^*), i = 9, \dots, 12 ; \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{b}_3) ; \\ \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{d}_i^*), i = 9, \dots, 12 \end{array} \right\} \end{array} \right.$$

Like in section 2.2, to  $\Sigma_K^*(v)$  we associate the following partition of the set  $\hat{\Delta}_K^*(v)$  :

$$(2.3.4) \quad \hat{\Delta}_K^*(v) = \hat{\Delta}_{K1}^*(v) \cup \hat{\Delta}_{K2}^*(v) \cup \hat{\Delta}_{K3}^*(v)$$

where each  $\hat{\Delta}_{Ki}^*$  is associated to the corresponding  $\hat{\Sigma}_i^*$ ,  $i = 1, 2, 3$ . We get successively :

**Construction of  $\hat{\Delta}_{K1}^*(v)$**  : The knowledge of  $F_K^*$ , of the set of values  $\{(D^\alpha v(a_i), |\alpha| = 0, 1, 2 ; i = 1, 2, 3) ; v(e_i^*), i = 1, \dots, 10\}$  and relation  $\hat{v} = v \circ F_K^*$  immediately involve the knowledge of :

$$(2.3.5) \quad \hat{\Delta}_{K1}^*(v) = \{(D^\alpha \hat{v}(\hat{a}_i), |\alpha| = 0, 1, 2 ; i = 1, 2, 3) ; \hat{v}(\hat{e}_i^*), i = 1, \dots, 10\}.$$

**Construction of  $\hat{\Delta}_{K2}^*(v)$**  : The functions  $\hat{f}_1, \hat{g}_1, \hat{f}_2, \hat{g}_2$  determined in section 2.2 play exactly the same role here and, since  $F_K|_{[\hat{a}_3, \hat{a}_1]} \equiv F_K^*|_{[\hat{a}_3, \hat{a}_1]}$  are affine, they are identical. Then, by similarity, we obtain

$$(2.3.6) \quad \left\{ \begin{array}{l} \hat{\Delta}_{K2}^*(v) = [\hat{f}_2(\hat{d}_i^*), i = 1, \dots, 4 ; \hat{f}_1(\hat{d}_i^*), i = 5, \dots, 8 ; \\ \\ - < \frac{\partial F_K^*}{\partial \hat{x}_1} (\hat{b}_1), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d\hat{f}_2}{d\hat{x}_2} (\hat{b}_1) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{g}_2(\hat{b}_1) > ; \\ \\ - < \frac{\partial F_K^*}{\partial \hat{x}_1} (\hat{d}_i^*), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d\hat{f}_2}{d\hat{x}_2} (\hat{d}_i^*) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{g}_2(\hat{d}_i^*) >, i = 1, \dots, 4 ; \\ \\ - < \frac{\partial F_K^*}{\partial \hat{x}_2} (\hat{b}_2), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{f}_1}{d\hat{x}_1} (\hat{b}_2) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{g}_1(\hat{b}_2) > ; \\ \\ - < \frac{\partial F_K^*}{\partial \hat{x}_2} (\hat{d}_i^*), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{f}_1}{d\hat{x}_1} (\hat{d}_i^*) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{g}_1(\hat{d}_i^*) >, i = 5, \dots, 8]. \end{array} \right.$$

**Construction of  $\hat{\Delta}_{K3}^*(v)$**  : Since the restrictions to  $[\hat{a}_1, \hat{a}_2]$  of  $F_K^*$  and  $F_K$  are different, the associated functions  $\hat{f}_3^*$  and  $\hat{g}_3^*$  are generally different from  $\hat{f}_3$  and  $\hat{g}_3$ . We obtain

$$(2.3.7) \quad \hat{\Delta}_{K3}^*(v) = \{\hat{f}_3^*(\hat{d}_i^*), i = 9, \dots, 12; -\sqrt{2}\hat{g}_3^*(\hat{b}_3); -\sqrt{2}\hat{g}_3^*(\hat{d}_i^*), i = 9, \dots, 12\}.$$

**Step 3 : Transition from  $\hat{\Delta}_{K3}^*(v)$  to the function  $w^* = \pi_K^* v$**  : Let  $\hat{w}^* \in \hat{P}^*$  be the polynomial of degree 9 which takes the values  $\hat{\Delta}_{K3}^*(v)$  on the set of degrees of freedom  $\hat{\Sigma}^*$ . Then

$$(2.3.8) \quad \pi_K^* v = w^* = \hat{w}^* \circ (F_K^*)^{-1}. \quad \blacksquare$$

In [2], we have proved that this function  $w^* = \pi_K^* v$  satisfy all the expected results :

**Theorem 2.3.1** : The function  $w^*$ , defined by (2.3.8), is determined in a unique way by the data of the set  $\Sigma_K^*(v)$  of the values of the degrees of freedom of the function  $v$ . Moreover the function  $w^*$  satisfies

$$(2.3.9) \quad \left\{ \begin{array}{l} D^\alpha w^*(a_i) = D^\alpha v(a_i), |\alpha| = 0, 1, 2; i = 1, 2, 3; \\ Dw^*(b_1)(a_1 - c_1) = Dv(b_1)(a_1 - c_1) \\ Dw^*(b_2)(a_2 - c_2) = Dv(b_2)(a_2 - c_2) \\ Dw^*(b_3)DF_K^*(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K^*(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \\ w(e_i^*) = v(e_i^*), i = 1, \dots, 10 \end{array} \right.$$

and the conditions (2.1.1) (2.1.2) and (2.1.4). Thus  $w^* = \pi_K^* v$ . ■

### 3 IMPLEMENTATION OF CURVED FINITE ELEMENTS $\mathcal{C}^1$ -COMPATIBLE WITH ARGYRIS TRIANGLE

In this paragraph we detail the matrix decompositions which allow to easily realize the interpolation modules related to both curved finite elements  $\mathcal{C}^1$  compatible with ARGYRIS triangle.

#### 3.1 Interpolation modules associated to curved finite element $\mathcal{C}^1$ -compatible with ARGYRIS triangle when $F_K \in (P_3)^2$

Let  $v$  be a function of  $\mathcal{C}^2(K)$ . By using the construction detailed in section 2.2, we are able to define its interpolate  $\pi_K v$ , i.e.,

$$(3.1.1) \quad \pi_K v(x) = w(x) = \hat{w}(\hat{x}).$$

Subsequently we will use this function  $\hat{w}$  on  $\hat{K}$  instead of  $w$  on  $K$ . This is usual in finite element implementation : in particular this allows to use numerical integration schemes on  $\hat{K}$ . Nevertheless, note that in [9] we took advantage of the linearity of  $F_K$  (we only considered straight triangles), of the properties of barycentric coordinates and of the eccentricity parameters to work directly on  $K$ .

Now it remains to transcribe in matrix expressions the relation (3.1.1). By similarity with section 2.2, we consider three steps :

**Step 1** :  $v \longrightarrow \Sigma_K(v) = [DLLC(v)]$  : The application  $F_K$  is given in Example 1.3.1 while  $\Sigma_K(v)$  is the following set of values (see (2.2.2)) :

$$(3.1.2) \quad \left\{ \begin{array}{l} [DLLC(v)]_{1 \times 24} = [v(a_1) \ v(a_2) \ v(a_3) \ Dv(a_1)(a_3 - a_1) \ (\bar{s} - \underline{s})Dv(a_1)\vec{\chi}'(\underline{s}) \\ \quad - (\bar{s} - \underline{s})Dv(a_2)\vec{\chi}'(\bar{s}) \ Dv(a_2)(a_3 - a_2) \ Dv(a_3)(a_2 - a_3) \ Dv(a_3)(a_1 - a_3) \\ \quad D^2v(a_1)(a_3 - a_1)^2 \ (\bar{s} - \underline{s})^2 D^2v(a_1)(\vec{\chi}'(\underline{s}))^2 \ (\bar{s} - \underline{s})^2 D^2v(a_2)(\vec{\chi}'(\bar{s}))^2 \\ \quad D^2v(a_2)(a_3 - a_2)^2 \ D^2v(a_3)(a_2 - a_3)^2 \ D^2v(a_3)(a_1 - a_3)^2 \ D^2v(a_1)(a_2 - a_3)^2 \\ \quad D^2v(a_2)(a_3 - a_1)^2 \ (\bar{s} - \underline{s})^2 D^2v(a_3)((\vec{\chi}'(\underline{s}), \vec{\chi}'(\bar{s})) \ Dv(b_1)(a_1 - c_1) \\ \quad Dv(b_2)(a_2 - c_2) \ Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \ v(e_1) \ v(e_2) \ v(e_3)] \end{array} \right.$$

When the side  $a_1a_2$  is straight, note that the first twenty degrees of freedom become identical to the first twenty local degrees of freedom of the ARGYRIS element (see [9, (3.1.18)]) since

$$(3.1.3) \quad \vec{\chi}'(s) = \frac{1}{\bar{s} - \underline{s}} \sum_{\alpha=1}^2 (\chi_\alpha(\bar{s}) - \chi_\alpha(\underline{s})) \vec{e}_\alpha = \frac{1}{\bar{s} - \underline{s}} (a_2 - a_1).$$

To the set (3.1.2), we associate the set of global degrees of freedom :

$$(3.1.4) \quad \left\{ \begin{array}{l} [DLGL(v)]_{1 \times 24} = [v(a_1) \ v(a_2) \ v(a_3) \ \frac{\partial v}{\partial x_1}(a_1) \ \frac{\partial v}{\partial x_2}(a_1) \ \frac{\partial v}{\partial x_1}(a_2) \\ \quad \frac{\partial v}{\partial x_2}(a_2) \ \frac{\partial v}{\partial x_1}(a_3) \ \frac{\partial v}{\partial x_2}(a_3) \ \frac{\partial^2 v}{\partial x_1^2}(a_1) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_1) \ \frac{\partial^2 v}{\partial x_2^2}(a_1) \\ \quad \frac{\partial^2 v}{\partial x_1^2}(a_2) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_2) \ \frac{\partial^2 v}{\partial x_2^2}(a_2) \ \frac{\partial^2 v}{\partial x_1^2}(a_3) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_3) \ \frac{\partial^2 v}{\partial x_2^2}(a_3) \\ \quad \frac{\partial v}{\partial \nu_1}(b_1) \ \frac{\partial v}{\partial \nu_2}(b_2) \ Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \ v(e_1) \ v(e_2) \ v(e_3)] \end{array} \right.$$

Local and global degrees of freedom are linked by the relation

$$(3.1.5) \quad [DLLC(v)]_{1 \times 24} = [DLGL(v)]_{1 \times 24} [\dot{D}]_{24 \times 24}$$

with

$$(3.1.6) \quad \check{D} = \begin{bmatrix} I_3 & & & & & \\ & \check{d}_1 & & & & \\ & & \check{d}_2 & & & \bigcirc \\ & & & \check{d}_3 & & \\ & & & & \check{d}_4 & \\ & \bigcirc & & & & n_1 \\ & & & & & n_2 \\ & & & & & I_4 \end{bmatrix}$$

where, by using relations (1.3.3) and by noting  $a_i(x_{1i}, x_{2i})$ , we have

$$(3.1.7) \quad \check{d}_1 = \begin{bmatrix} X_{31} & \psi'_1(0) \\ Y_{31} & \psi'_2(0) \end{bmatrix}; \quad \check{d}_2 = \begin{bmatrix} -\psi'_1(1) & X_{32} \\ -\psi'_2(1) & Y_{32} \end{bmatrix}; \quad \check{d}_3 = \begin{bmatrix} X_{23} & X_{13} \\ Y_{23} & Y_{13} \end{bmatrix}$$

$$(3.1.8) \quad \check{d}_4 = \begin{bmatrix} (X_{31})^2 & (\psi'_1(0))^2 & 0 & 0 & 0 & 0 & (X_{32})^2 & 0 & 0 \\ 2X_{31}Y_{31} & 2\psi'_1(0)\psi'_2(0) & 0 & 0 & 0 & 0 & 2X_{32}Y_{32} & 0 & 0 \\ (Y_{31})^2 & (\psi'_2(0))^2 & 0 & 0 & 0 & 0 & (Y_{32})^2 & 0 & 0 \\ 0 & 0 & (\psi'_1(1))^2 & (X_{32})^2 & 0 & 0 & 0 & (X_{31})^2 & 0 \\ 0 & 0 & 2\psi'_1(1)\psi'_2(1) & 2X_{32}Y_{32} & 0 & 0 & 0 & 2X_{31}Y_{31} & 0 \\ 0 & 0 & (\psi'_2(1))^2 & (Y_{32})^2 & 0 & 0 & 0 & (Y_{31})^2 & 0 \\ 0 & 0 & 0 & 0 & (X_{32})^2 & (X_{31})^2 & 0 & 0 & \psi'_1(0)\psi'_1(1) \\ 0 & 0 & 0 & 0 & 2X_{32}Y_{32} & 2X_{31}Y_{31} & 0 & 0 & \left\{ \begin{array}{l} \psi'_1(0)\psi'_2(1) \\ +\psi'_2(0)\psi'_1(1) \end{array} \right\} \\ 0 & 0 & 0 & 0 & (Y_{32})^2 & (Y_{31})^2 & 0 & 0 & \psi'_2(0)\psi'_2(1) \end{bmatrix}$$

with  $X_{ij} = x_{1i} - x_{1j}$ ,  $Y_{ij} = x_{2i} - x_{2j}$ ,  $1 \leq i, j \leq 3$ , and  $n_1$  and  $n_2$  as given in [9, (3.1.25)].

**Step 2 : Transition from  $\Sigma_K(v) = [DLLC(v)]$  to  $\hat{\Delta}_K(v) = [DL(\hat{w})]$  :** On the fixed reference triangle  $\hat{K}$ , i.e., the unit right-angled triangle, we introduce only one set of degrees of freedom (see Figure 2.2.1) :

$$(3.1.9) \quad \left\{ \begin{array}{l} [DL(\hat{w})]_{1 \times 36} = [\hat{w}(\hat{a}_1) \ \hat{w}(\hat{a}_2) \ \hat{w}(\hat{a}_3) \ \frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{a}_1) \ \frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{a}_1) \ \frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{a}_2) \ \frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{a}_2) \\ \frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{a}_3) \ \frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{a}_3) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2}(\hat{a}_1) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_1) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2}(\hat{a}_1) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2}(\hat{a}_2) \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_2) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2}(\hat{a}_2) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2}(\hat{a}_3) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_3) \ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2}(\hat{a}_3) \\ -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{b}_1) \ -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{b}_2) \ \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{b}_3) \ \hat{w}(\hat{d}_1) \ \hat{w}(\hat{d}_2) \ \hat{w}(\hat{d}_3) \ \hat{w}(\hat{d}_4) \\ \hat{w}(\hat{d}_5) \ \hat{w}(\hat{d}_6) \ -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{d}_1) \ -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{d}_2) \ -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{d}_3) \ -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{d}_4) \\ \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{d}_5) \ \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{d}_6) \ \hat{w}(\hat{e}_1) \ \hat{w}(\hat{e}_2) \ \hat{w}(\hat{e}_3) \end{array} \right]$$

Then, step 2 of section 2.2 leads to

$$(3.1.10) \quad [DL(\hat{w})]_{1 \times 36} = [DLLC(v)]_{1 \times 24} [B]_{24 \times 36}$$

where the matrix  $B$  is piecewise constructed hereunder. Then relations (3.1.5) and (3.1.10) give

$$(3.1.11) \quad [DL(\hat{w})]_{1 \times 36} = [DLGL(v)]_{1 \times 24} [\tilde{D}]_{24 \times 24} [B]_{24 \times 36}.$$

By consideration of relations (3.1.9) and (3.1.10), it is natural to realize the following partition of matrix  $B$

$$(3.1.12) \quad [B]_{24 \times 36} = [B_1 \ B_2 \ B_3 \ B_4 \ B_5 \ B_6 \ B_7]$$

where submatrices  $B_i$ ,  $i = 1, \dots, 7$ , have 24 lines and respectively 3, 6, 9, 3, 6, 6 and 3 columns, and are determined as follows :

**Construction of submatrix  $B_1$**  : From (2.2.21) and (2.2.22), one gets  $\hat{w}(\hat{a}_i) = v(a_i)$ ,  $i = 1, 2, 3$  so that

$$(3.1.13) \quad {}^t B_1 = [I_3 \ ; \ 0_{3 \times 21}].$$

**Construction of submatrix  $B_2$**  : Since  $\hat{w} = w \circ F_K$ , we obtain

$$\frac{\partial \hat{w}}{\partial \hat{x}_\alpha}(\hat{x}) = D\hat{w}(\hat{x})\vec{\hat{e}}_\alpha = Dw(x)DF_K(\hat{x})\vec{\hat{e}}_\alpha = Dw(x) \frac{\partial F_K}{\partial \hat{x}_\alpha}(\hat{x})$$

so that relation (2.2.22) involves

$$(3.1.14) \quad \frac{\partial \hat{w}}{\partial \hat{x}_\alpha} (\hat{a}_i) = Dv(a_i) \frac{\partial F_K}{\partial \hat{x}_\alpha} (\hat{a}_i), \quad i = 1, 2, 3$$

and with (1.3.2)

$$(3.1.15) \quad \left\{ \begin{array}{l} \frac{\partial \hat{w}}{\partial \hat{x}_1} (\hat{a}_1) = Dv(a_1)(a_1 - a_3); \quad \frac{\partial \hat{w}}{\partial \hat{x}_2} (\hat{a}_1) = Dv(a_1)[a_1 - a_3 + (\bar{s} - \underline{s})\vec{\chi}'(\underline{s})] \\ \frac{\partial \hat{w}}{\partial \hat{x}_1} (\hat{a}_2) = Dv(a_2)[a_2 - a_3 - (\bar{s} - \underline{s})\vec{\chi}'(\bar{s})]; \quad \frac{\partial \hat{w}}{\partial \hat{x}_2} (\hat{a}_2) = Dv(a_2)(a_2 - a_3) \\ \frac{\partial \hat{w}}{\partial \hat{x}_1} (\hat{a}_3) = Dv(a_3)(a_1 - a_3); \quad \frac{\partial \hat{w}}{\partial \hat{x}_2} (\hat{a}_3) = Dv(a_3)(a_2 - a_3) \end{array} \right.$$

From these expressions and relations (3.1.2) (3.1.9) (3.1.10) and (3.1.12) we obtain

$$(3.1.16) \quad {}^t B_2 = [0_{6 \times 3} ; \ {}^t(b_2)_{6 \times 6} ; \ 0_{6 \times 15}]$$

with

$$(3.1.17) \quad b_2 = \begin{bmatrix} -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

**Construction of submatrix  $B_3$**  : Relations  $\hat{w} = w \circ F_K$  and (2.2.22) involve for  $\alpha = 1, 2$  :

$$(3.1.18) \quad \frac{\partial^2 \hat{w}}{\partial \hat{x}_\alpha \partial \hat{x}_\beta} (\hat{x}) = Dw(x) \frac{\partial^2 F_K}{\partial \hat{x}_\alpha \partial \hat{x}_\beta} (\hat{x}) + D^2 w(x) \left( \frac{\partial F_K}{\partial \hat{x}_\alpha} (\hat{x}), \frac{\partial F_K}{\partial \hat{x}_\beta} (\hat{x}) \right)$$

so that with (1.3.2) we get

$$\begin{aligned}
(3.1.19) \quad & \left\{ \begin{array}{ll} \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2} (\hat{a}_1) & = D^2 v(a_1)(a_1 - a_3)^2 \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_1) & = D^2 v(a_1)(a_1 - a_3, a_1 - a_3 + (\bar{s} - s)\vec{\chi}'(s))] \\ & + Dv(a_1)[2(a_1 - a_2) + \frac{1}{2}(\bar{s} - s)(3\vec{\chi}'(s) + \vec{\chi}'(\bar{s}))] \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2} (\hat{a}_1) & = D^2 v(a_1)[a_1 - a_3 + (\bar{s} - s)\vec{\chi}'(s)]^2 \\ & - Dv(a_1)[2(a_1 - a_2) + (\bar{s} - s)(\vec{\chi}'(s) + \vec{\chi}'(\bar{s}))] \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3 - (\bar{s} - s)\vec{\chi}'(\bar{s})]^2 \\ & - Dv(a_2)[2(a_2 - a_1) - (\bar{s} - s)(\vec{\chi}'(s) + \vec{\chi}'(\bar{s}))] \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3 - (\bar{s} - s)\vec{\chi}'(\bar{s}), a_2 - a_3] \\ & + Dv(a_2)[2(a_2 - a_1) - \frac{1}{2}(\bar{s} - s)(3\vec{\chi}'(\bar{s}) + \vec{\chi}'(s))] \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3]^2 \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2} (\hat{a}_3) & = D^2 v(a_3)[a_1 - a_3]^2 \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_3) & = D^2 v(a_3)[a_1 - a_3, a_2 - a_3] - \frac{1}{2}(\bar{s} - s)Dv(a_3)[\vec{\chi}'(\bar{s}) - \vec{\chi}'(s)] \\ \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2} (\hat{a}_3) & = D^2 v(a_3)[a_2 - a_3]^2 \end{array} \right. \end{aligned}$$

From these expressions and relations (3.1.2) (3.1.9) (3.1.10) and (3.1.12), we deduce

$$(3.1.20) \quad {}^t B_3 = [0_{9 \times 3} ; {}^t(b_{31})_{9 \times 6} ; {}^t(b_{32})_{9 \times 9} ; 0_{9 \times 6}]$$

where matrices  $b_{31}$  and  $b_{32}$  are given by relations (3.1.21) (3.1.22).

$$(3.1.21) \quad b_{31} = \begin{bmatrix} 0 & \left(2\tilde{a}^1 + \frac{1}{2}\tilde{\tilde{a}}^1\right) & -(2\tilde{a}^1 + \tilde{\tilde{a}}^1) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \left(\frac{3}{2} + 2\tilde{a}^2 + \frac{1}{2}\tilde{\tilde{a}}^2\right) & -(1 + 2\tilde{a}^2 + \tilde{\tilde{a}}^2) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -(1 + 2\tilde{b}^1 - \tilde{\tilde{b}}^1) & \left(\frac{3}{2} + 2\tilde{b}^1 - \frac{1}{2}\tilde{\tilde{b}}^1\right) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -(2\tilde{b}^2 - \tilde{\tilde{b}}^2) & \left(2\tilde{b}^2 - \frac{1}{2}\tilde{\tilde{b}}^2\right) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{2}(\tilde{c}^1 + \tilde{\tilde{c}}^1) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{2}(\tilde{c}^2 + \tilde{\tilde{c}}^2) & 0 \end{bmatrix}$$

$$(3.1.22) \quad b_{32} = \begin{bmatrix} 1 & \left(1 + \frac{1+\tilde{a}^1}{2\tilde{a}^2}\right) & \left(1 + \frac{1+\tilde{a}^1}{\tilde{a}^2}\right) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{\tilde{a}^2}{2(1+\tilde{a}^1)} & \left(1 + \frac{\tilde{a}^2}{1+\tilde{a}^1}\right) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \left(1 + \frac{\tilde{b}^1}{1+\tilde{b}^2}\right) & \frac{\tilde{b}^1}{2(1+\tilde{b}^2)} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \left(1 + \frac{1+\tilde{b}^2}{\tilde{b}^1}\right) & \left(1 + \frac{1+\tilde{b}^2}{2\tilde{b}^1}\right) & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\frac{\tilde{c}^1\tilde{c}^1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1} & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -\frac{\tilde{c}^2\tilde{c}^2}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1} & 0 \\ 0 & \frac{-1}{2\tilde{a}^2(1+\tilde{a}^1)} & \frac{-1}{\tilde{a}^2(1+\tilde{a}^1)} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{-1}{\tilde{b}^1(1+\tilde{b}^2)} & \frac{-1}{2\tilde{b}^1(1+\tilde{b}^2)} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{-1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1} & 0 \end{bmatrix}$$

It remains to indicate how to obtain the coefficients of the submatrices  $b_{31}$  and  $b_{32}$ . The second and third columns of  $b_{31}$  include terms in  $Dv(a_1)$  that we have to express by using the fourth and fifth degrees of freedom of the matrix (3.1.2). Set

$$\vec{A}_1 = a_3 - a_1 \quad \text{and} \quad \vec{A}_2 = (\bar{s} - s)\vec{\chi}'(s)$$

and assume that these vectors are linearly independent. Then  $\tilde{a}^\alpha$  and  $\tilde{\tilde{a}}^\alpha$  are defined by

$$a_1 - a_2 = \tilde{a}^\alpha \vec{A}_\alpha ; \quad (\bar{s} - s)\vec{\chi}'(\bar{s}) = \tilde{\tilde{a}}^\alpha \vec{A}_\alpha$$

so that, if we denote  $\vec{e}_3 = \vec{e}_1 \times \vec{e}_2$ :

$$\begin{cases} \tilde{a}^1 = \frac{[(a_1 - a_2) \times \vec{A}_2] \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} & ; \quad \tilde{a}^2 = \frac{[\vec{A}_1 \times (a_1 - a_2)] \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} \\ \tilde{\tilde{a}}^1 = (\bar{s} - s) \frac{[\vec{\chi}'(\bar{s}) \times \vec{A}_2] \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} & ; \quad \tilde{\tilde{a}}^2 = (\bar{s} - s) \frac{[\vec{A}_1 \times \vec{\chi}'(\bar{s})] \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3}. \end{cases}$$

Similarly, set

$$\begin{aligned} \vec{B}_1 &= -(\bar{s} - s)\vec{\chi}'(\bar{s}) ; \quad \vec{B}_2 = a_3 - a_2 ; \quad a_2 - a_1 = \tilde{b}^\alpha \vec{B}_\alpha ; \quad (\bar{s} - s)\vec{\chi}'(s) = \tilde{\tilde{b}}^\alpha \vec{B}_\alpha \\ \vec{C}_1 &= a_2 - a_3 ; \quad \vec{C}_2 = a_1 - a_3 ; \quad (\bar{s} - s)\vec{\chi}'(s) = \tilde{c}^\alpha \vec{C}_\alpha ; \quad -(\bar{s} - s)\vec{\chi}'(\bar{s}) = \tilde{\tilde{c}}^\alpha \vec{C}_\alpha. \end{aligned}$$

From these definitions, we finally obtain :

$$(3.1.23) \quad \left\{ \begin{array}{l} \tilde{a}^1 = \frac{(\vec{B}_2 \times \vec{A}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} - 1 \quad ; \quad \tilde{a}^2 = \frac{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} \\ \tilde{a}^1 = - \frac{(\vec{B}_1 \times \vec{A}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} \quad ; \quad \tilde{a}^2 = - \frac{(\vec{A}_1 \times \vec{B}_1) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} \\ \tilde{b}^1 = \frac{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} \quad ; \quad \tilde{b}^2 = \frac{(\vec{B}_1 \times \vec{A}_1) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} - 1 \\ \tilde{b}^1 = \frac{(\vec{A}_2 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} \quad ; \quad \tilde{b}^2 = \frac{(\vec{B}_1 \times \vec{A}_2) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} \\ \tilde{c}^1 = - \frac{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3} = - \frac{1}{\tilde{a}^2} \quad ; \quad \tilde{c}^2 = - \frac{(\vec{A}_2 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3} = - \frac{\tilde{b}^1}{\tilde{b}^1} \\ \tilde{c}^1 = - \frac{(\vec{A}_1 \times \vec{B}_1) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3} = \frac{\tilde{a}^2}{\tilde{a}^2} \quad ; \quad \tilde{c}^2 = - \frac{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{B}_2) \cdot \vec{e}_3} = - \frac{1}{\tilde{b}^1} \end{array} \right.$$

**Construction of submatrices  $B_i$ ,  $i = 4, 5, 6$**  : We have seen in section 2.2 that it is convenient to introduce the functions  $\hat{f}_1$  (see (2.2.8)),  $\hat{f}_2$  (see (2.2.12)),  $\hat{f}_3$  (see (2.2.18)),  $\hat{g}_1$  (see (2.2.10)),  $\hat{g}_2$  (see (2.2.13)) and  $\hat{g}_3$  (see (2.2.19)). Relations (1.3.2) (3.1.2),  $\hat{v} = v \circ F_K$  and some technical computations prove that these functions  $\hat{f}_i$ ,  $\hat{g}_i$ ,  $i = 1, 2, 3$  can be written as

$$(3.1.24) \quad \left\{ \begin{array}{l} [\hat{f}_1(\hat{x}_1) \quad \hat{f}_2(\hat{x}_2) \quad \hat{f}_3(\hat{x}_1)] = [DLLC(v)]_{1 \times 24} \quad [\hat{F}_1(\hat{x}_1) \quad \hat{F}_2(\hat{x}_2) \quad \hat{F}_3(\hat{x}_1)]_{24 \times 3} \\ [\hat{g}_1(\hat{x}_1) \quad \hat{g}_2(\hat{x}_2) \quad \hat{g}_3(\hat{x}_1)] = [DLLC(v)]_{1 \times 24} \quad [\hat{G}_1(\hat{x}_1) \quad \hat{G}_2(\hat{x}_2) \quad \hat{G}_3(\hat{x}_1)]_{24 \times 3} \end{array} \right.$$

with

$$(3.1.25) \quad \left\{ \begin{array}{l} {}^t[\hat{F}_1(\hat{x}_1)] = [\hat{x}_1^3(6\hat{x}_1^2 - 15\hat{x}_1 + 10) ; 0 ; (1 - \hat{x}_1)^3(6\hat{x}_1^2 + 3\hat{x}_1 + 1) ; \\ \hat{x}_1^3(1 - \hat{x}_1)(4 - 3\hat{x}_1) ; 0 0 0 0 ; \hat{x}_1(1 - \hat{x}_1)^3(1 + 3\hat{x}_1) ; \\ \frac{1}{2} \hat{x}_1^3(1 - \hat{x}_1)^2 ; 0 0 0 0 ; \frac{1}{2} \hat{x}_1^2(1 - \hat{x}_1)^3 ; 0 0 0 0 0 0 0 0 0 ] \end{array} \right.$$

$$(3.1.26) \quad \left\{ \begin{array}{l} {}^t[\hat{F}_2(\hat{x}_2)] = [0 ; \hat{x}_2^3(6\hat{x}_2^2 - 15\hat{x}_2 + 10) ; (1 - \hat{x}_2)^3(6\hat{x}_2^2 + 3\hat{x}_2 + 1) ; \\ 0 0 0 ; \hat{x}_2^3(1 - \hat{x}_2)(4 - 3\hat{x}_2) ; \hat{x}_2(1 - \hat{x}_2)^3(1 + 3\hat{x}_2) ; 0 0 0 0 ; \\ \frac{1}{2} \hat{x}_2^3(1 - \hat{x}_2)^2 ; \frac{1}{2} \hat{x}_2^2(1 - \hat{x}_2)^3 ; 0 0 0 0 0 0 0 0 0 ] \end{array} \right.$$

$$\begin{aligned}
(3.1.27) \quad & \left\{ \begin{array}{l} {}^t[\hat{\mathcal{F}}_3(\hat{x}_1)] = [\hat{x}_1^3(6\hat{x}_1^2 - 15\hat{x}_1 + 10); (1 - \hat{x}_1)^3(6\hat{x}_1^2 + 3\hat{x}_1 + 1); 0; \\ \quad - (3\tilde{a}^1 + \tilde{a}^1)\hat{x}_1^3(1 - \hat{x}_1)^2; \hat{x}_1^3(1 - \hat{x}_1)\{2 - \hat{x}_1 - (1 - \hat{x}_1)(3\tilde{a}^2 + \tilde{a}^2)\} \\ \quad + \hat{x}_1(1 - \hat{x}_1)^3\{1 + \hat{x}_1 - \hat{x}_1(3\tilde{b}^1 - \tilde{b}^1)\}; - \hat{x}_1^2(1 - \hat{x}_1)^3(3\tilde{b}^2 - \tilde{b}^2); \\ \quad 0\ 0\ 0; \frac{1}{2}\hat{x}_1^3(1 - \hat{x}_1)^2; \frac{1}{2}\hat{x}_1^2(1 - \hat{x}_1)^3; 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0] \end{array} \right. \\
(3.1.28) \quad & \left\{ \begin{array}{l} {}^t[\hat{\mathcal{G}}_1(\hat{x}_1)] = [0\ 0\ 0; -\frac{1}{2}(1 - \eta_2 + 2\tilde{a}^1)\hat{x}_1^2(2\hat{x}_1 - 1)(5 - 4\hat{x}_1); \\ \quad - \tilde{a}^2\hat{x}_1^2(2\hat{x}_1 - 1)(5 - 4\hat{x}_1); 0\ 0; (1 - \hat{x}_1)^2(1 - 2\hat{x}_1)(1 + 4\hat{x}_1); \\ \quad - \frac{1}{2}(1 + \eta_2)(1 - \hat{x}_1)^2(1 - 2\hat{x}_1)(1 + 4\hat{x}_1); \frac{1}{2}(\tilde{a}^1 - \eta_2)\hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1); \\ \quad - \frac{1}{2}\frac{(\tilde{a}^2)^2}{1 + \tilde{a}^1}\hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1); 0\ 0; \frac{-\tilde{c}^1\tilde{c}^1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1}\hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1); \\ \quad - \left\{ \frac{1}{2}(1 + \eta_2) + \frac{\tilde{c}^2\tilde{c}^2}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1} \right\}\hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1); \frac{1}{2(1 + \tilde{a}^1)}\hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1); \\ \quad 0; \frac{-1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1}\hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1); 0; 16\hat{x}_1^2(1 - \hat{x}_1)^2; 0\ 0\ 0\ 0] \end{array} \right. \\
(3.1.29) \quad & \left\{ \begin{array}{l} {}^t[\hat{\mathcal{G}}_2(\hat{x}_2)] = [0\ 0\ 0\ 0\ 0; -\tilde{b}^1\hat{x}_2^2(2\hat{x}_2 - 1)(5 - 4\hat{x}_2); \\ \quad - \frac{1}{2}(2\tilde{b}^2 + 1 + \eta_1)\hat{x}_2^2(2\hat{x}_2 - 1)(5 - 4\hat{x}_2); -\frac{1}{2}(1 - \eta_1)(1 - \hat{x}_2)^2(1 - 2\hat{x}_2)(1 + 4\hat{x}_2); \\ \quad (1 - \hat{x}_2)^2(1 - 2\hat{x}_2)(1 + 4\hat{x}_2); 0\ 0; -\frac{1}{2}\frac{(\tilde{b}^1)^2}{1 + \tilde{b}^2}\hat{x}_2^2(1 - \hat{x}_2)(1 - 2\hat{x}_2); \\ \quad \frac{1}{2}(\tilde{b}^2 + \eta_1)\hat{x}_2^2(1 - \hat{x}_2)(1 - 2\hat{x}_2); -\frac{1}{2}\left\{ \frac{2\tilde{c}^1\tilde{c}^1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1} + 1 - \eta_1 \right\}\hat{x}_2(1 - \hat{x}_2)^2(1 - 2\hat{x}_2); \\ \quad -\frac{\tilde{c}^2\tilde{c}^2}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1}\hat{x}_2(1 - \hat{x}_2)^2(1 - 2\hat{x}_2); 0; \frac{1}{2}\frac{1}{1 + \tilde{b}^2}\hat{x}_2^2(1 - \hat{x}_2)(1 - 2\hat{x}_2); \\ \quad -\frac{1}{\tilde{c}^1\tilde{c}^2 + \tilde{c}^2\tilde{c}^1}\hat{x}_2(1 - \hat{x}_2)^2(1 - 2\hat{x}_2); 16(1 - \hat{x}_2)^2\hat{x}_2^2; 0\ 0\ 0\ 0\ 0] \end{array} \right. \end{aligned}$$

$$(3.1.30) \quad \left\{ \begin{array}{l} {}^t[\hat{\mathcal{G}}_3(\hat{x}_1)] = [0 \ 0 \ 0 ; \ \hat{x}_1^2(2\hat{x}_1-1) \left\{ 5-4\hat{x}_1+(1-\hat{x}_1) \left( \tilde{a}^1 + \frac{1}{2} \tilde{\tilde{a}}^1 \right) \right\} ; \\ \hat{x}_1^2(2\hat{x}_1-1) \left\{ -2 + \frac{3}{2} \hat{x}_1 + (1-\hat{x}_1) \left( \tilde{a}^2 + \frac{1}{2} \tilde{\tilde{a}}^2 \right) \right\} ; \\ -\frac{1}{2} (1-\hat{x}_1)^2(1-2\hat{x}_1) \left\{ (1+3\hat{x}_1) - (2\tilde{b}^1 - \tilde{\tilde{b}}^1) \hat{x}_1 \right\} ; \\ (1-\hat{x}_1)^2(1-2\hat{x}_1) \left\{ 1+4\hat{x}_1 + \frac{1}{2} (2\tilde{b}^2 - \tilde{\tilde{b}}^2) \hat{x}_1 \right\} \ 0 \ 0 ; \\ \frac{1+\tilde{a}^1}{2\tilde{a}^2} \hat{x}_1^2(1-\hat{x}_1)(1-2\hat{x}_1) ; \ \frac{1}{2} \left( 1 + \frac{\tilde{a}^2}{1+\tilde{a}^1} \right) \hat{x}_1^2(1-\hat{x}_1)(1-2\hat{x}_1) ; \\ -\frac{1}{2} \left( 1 + \frac{\tilde{b}^1}{1+\tilde{b}^2} \right) \hat{x}_1(1-\hat{x}_1)^2(1-2\hat{x}_1) ; \ -\frac{1+\tilde{b}^2}{2\tilde{b}^1} \hat{x}_1(1-\hat{x}_1)^2(1-2\hat{x}_1) ; \\ 0 \ 0 ; \ -\frac{1}{2(1+\tilde{a}^1)\tilde{a}^2} \hat{x}_1^2(1-\hat{x}_1)(1-2\hat{x}_1) ; \\ \frac{1}{2\tilde{b}^1(1+\tilde{b}^2)} \hat{x}_1(1-\hat{x}_1)^2(1-2\hat{x}_1) ; \ 0 \ 0 \ 0 ; \ 16(1-\hat{x}_1)^2\hat{x}_1^2 ; \ 0 \ 0 \ 0 ] \end{array} \right.$$

To express  $\hat{\mathcal{G}}_1(\hat{x}_1)$  and  $\hat{\mathcal{G}}_2(\hat{x}_2)$ , we have used the eccentricity parameters  $\eta_1$  and  $\eta_2$  of the triangle  $a_1a_2a_3$  which are given by (see [9, p. 69]) :

$$a_1 - c_1 = a_1 - a_2 - \frac{1}{2} (1 + \eta_1)(a_3 - a_2) ; \ a_2 - c_2 = a_2 - a_1 - \frac{1}{2} (1 - \eta_2)(a_3 - a_1).$$

Then we are able to construct the submatrices  $B_i$ ,  $i = 4, 5, 6$ .

**Construction of submatrix  $B_4$**  : Relations (2.2.3) and (2.2.14) involve

$$(3.1.31) \quad - \frac{\partial \hat{w}}{\partial \hat{x}_1} (\hat{b}_1) = - < \frac{\partial F_K}{\partial \hat{x}_1} (\hat{b}_1), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d\hat{f}_2}{d\hat{x}_2} \left( \frac{1}{2} \right) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{g}_2 \left( \frac{1}{2} \right) >$$

where  $< . , . >$  notes the usual scalar product in  $\mathbb{R}^2$ . Hence relations (3.1.24) (3.1.26) (3.1.28) and (3.1.31) give

$$(3.1.32) \quad - \frac{\partial \hat{w}}{\partial \hat{x}_1} (\hat{b}_1) = [DLLC(v)]_{1 \times 24} \left[ E^1 \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2 \left( \frac{1}{2} \right) + E^2 \hat{\mathcal{G}}_2 \left( \frac{1}{2} \right) \right]_{24 \times 1}$$

where we have set

$$(3.1.33) \quad \begin{cases} E^1 = < a_3 - a_1 + \frac{1}{4} (a_1 - a_2) + \frac{1}{8} (\bar{s} - s) \{3\vec{\chi}'(\bar{s}) - \vec{\chi}'(s)\}, \frac{a_2 - a_3}{|a_2 - a_3|} > \\ E^2 = < a_3 - a_1 + \frac{1}{4} (a_1 - a_2) + \frac{1}{8} (\bar{s} - s) \{3\vec{\chi}'(\bar{s}) - \vec{\chi}'(s)\}, \frac{a_1 - c_1}{|a_1 - c_1|^2} > \end{cases}$$

since (1.3.2) involves

$$-\frac{\partial F_K}{\partial \hat{x}_1} (\hat{b}_1) = a_3 - a_1 + \frac{1}{4} (a_1 - a_2) + \frac{1}{8} (\bar{s} - s) \{3\vec{\chi}'(\bar{s}) - \vec{\chi}'(s)\}$$

By similarity, we obtain

$$(3.1.34) \quad -\frac{\partial \hat{w}}{\partial \hat{x}_2} (\hat{b}_2) = [DLLC(v)]_{1 \times 24} \left[ F^1 \frac{d}{d\hat{x}_1} \hat{\mathcal{F}}_1 \left( \frac{1}{2} \right) + F^2 \hat{\mathcal{G}}_1 \left( \frac{1}{2} \right) \right]_{24 \times 1}$$

where

$$(3.1.35) \quad \begin{cases} F^1 = < a_3 - a_2 + \frac{1}{4} (a_2 - a_1) - \frac{1}{8} (\bar{s} - s) \{3\vec{\chi}'(s) - \vec{\chi}'(\bar{s})\}, \frac{a_1 - a_3}{|a_1 - a_3|^2} > \\ F^2 = < a_3 - a_2 + \frac{1}{4} (a_2 - a_1) - \frac{1}{8} (\bar{s} - s) \{3\vec{\chi}'(s) - \vec{\chi}'(\bar{s})\}, \frac{a_2 - c_2}{|a_2 - c_2|^2} > \end{cases}$$

Moreover

$$(3.1.36) \quad \begin{cases} \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{b}_3) = -\sqrt{2} D \hat{w} (\hat{b}_3) (\hat{a}_3 - \hat{b}_3) \\ = [DLLC(v)]_{1 \times 24} {}^t [0_{1 \times 20}; -\sqrt{2}; 0 \ 0 \ 0]. \end{cases}$$

Thus, relations (3.1.32) (3.1.34) and (3.1.36) lead to

$$(3.1.37) \quad B_4 = \left[ E^1 \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2 \left( \frac{1}{2} \right) + E^2 \hat{\mathcal{G}}_2 \left( \frac{1}{2} \right); F^1 \frac{d}{d\hat{x}_1} \hat{\mathcal{F}}_1 \left( \frac{1}{2} \right) + F^2 \hat{\mathcal{G}}_1 \left( \frac{1}{2} \right); \begin{pmatrix} 0_{20 \times 1} \\ -\sqrt{2} \\ 0_{3 \times 1} \end{pmatrix} \right].$$

**Construction of submatrix  $B_5$**  : Relations (2.2.3) (2.2.14) (2.2.20) and (3.1.24) give

$$(3.1.38) \quad B_5 = \left[ \hat{\mathcal{F}}_2 \left( \frac{3}{4} \right); \hat{\mathcal{F}}_2 \left( \frac{1}{4} \right); \hat{\mathcal{F}}_1 \left( \frac{1}{4} \right); \hat{\mathcal{F}}_1 \left( \frac{3}{4} \right); \hat{\mathcal{F}}_3 \left( \frac{3}{4} \right); \hat{\mathcal{F}}_3 \left( \frac{1}{4} \right) \right]_{24 \times 6}$$

**Construction of submatrix  $B_6$**  : By similarity with the obtention of (3.1.32) (3.1.34) and (3.1.36) we get

$$(3.1.39) \quad \left\{ \begin{array}{l} B_6 = \left[ G^1 \frac{d\hat{\mathcal{F}}_2}{d\hat{x}_2} \left( \frac{3}{4} \right) + G^2 \hat{\mathcal{G}}_2 \left( \frac{3}{4} \right) ; H^1 \frac{d\hat{\mathcal{F}}_2}{d\hat{x}_2} \left( \frac{1}{4} \right) + H^2 \hat{\mathcal{G}}_2 \left( \frac{1}{4} \right) ; \right. \\ J^1 \frac{d\hat{\mathcal{F}}_1}{d\hat{x}_1} \left( \frac{1}{4} \right) + J^2 \hat{\mathcal{G}}_1 \left( \frac{1}{4} \right) ; K^1 \frac{d\hat{\mathcal{F}}_1}{d\hat{x}_1} \left( \frac{3}{4} \right) + K^2 \hat{\mathcal{G}}_1 \left( \frac{3}{4} \right) ; \\ \left. - \sqrt{2} \hat{\mathcal{G}}_3 \left( \frac{3}{4} \right) ; - \sqrt{2} \hat{\mathcal{G}}_3 \left( \frac{1}{4} \right) \right]_{24 \times 6} \end{array} \right.$$

where we have set

$$(3.1.40) \quad \left\{ \begin{array}{l} G^1 = - < a_1 - a_3 + \frac{9}{16} (a_2 - a_1) + \frac{3}{32} (\bar{s} - \underline{s}) \{ \vec{\chi}'(\underline{s}) - 7\vec{\chi}'(\bar{s}) \}, \frac{a_2 - a_3}{|a_2 - a_3|^2} > \\ G^2 = - < a_1 - a_3 + \frac{9}{16} (a_2 - a_1) + \frac{3}{32} (\bar{s} - \underline{s}) \{ \vec{\chi}'(\underline{s}) - 7\vec{\chi}'(\bar{s}) \}, \frac{a_1 - c_1}{|a_1 - c_1|^2} > \\ H^1 = - < a_1 - a_3 + \frac{1}{16} (a_2 - a_1) + \frac{1}{32} (\bar{s} - \underline{s}) \{ 3\vec{\chi}'(\underline{s}) - 5\vec{\chi}'(\bar{s}) \}, \frac{a_2 - a_3}{|a_2 - a_3|^2} > \\ H^2 = - < a_1 - a_3 + \frac{1}{16} (a_2 - a_1) + \frac{1}{32} (\bar{s} - \underline{s}) \{ 3\vec{\chi}'(\underline{s}) - 5\vec{\chi}'(\bar{s}) \}, \frac{a_1 - c_1}{|a_1 - c_1|^2} > \\ J^1 = - < a_2 - a_3 + \frac{1}{16} (a_1 - a_2) - \frac{1}{32} (\bar{s} - \underline{s}) \{ 3\vec{\chi}'(\bar{s}) - 5\vec{\chi}'(\underline{s}) \}, \frac{a_1 - a_3}{|a_1 - a_3|^2} > \\ J^2 = - < a_2 - a_3 + \frac{1}{16} (a_1 - a_2) - \frac{1}{32} (\bar{s} - \underline{s}) \{ 3\vec{\chi}'(\bar{s}) - 5\vec{\chi}'(\underline{s}) \}, \frac{a_2 - c_2}{|a_2 - c_2|^2} > \\ K^1 = - < a_2 - a_3 + \frac{9}{16} (a_1 - a_2) - \frac{3}{32} (\bar{s} - \underline{s}) \{ \vec{\chi}'(\bar{s}) - 7\vec{\chi}'(\underline{s}) \}, \frac{a_1 - a_3}{|a_1 - a_3|^2} > \\ K^2 = - < a_2 - a_3 + \frac{9}{16} (a_1 - a_2) - \frac{3}{32} (\bar{s} - \underline{s}) \{ \vec{\chi}'(\bar{s}) - 7\vec{\chi}'(\underline{s}) \}, \frac{a_2 - c_2}{|a_2 - c_2|^2} > \end{array} \right.$$

**Construction of submatrix  $B_7$**  : Since  $\hat{w}(\hat{e}_i) = w(e_i) = v(e_i)$ ,  $i = 1, 2, 3$ , we get

$$(3.1.41) \quad {}^t B_7 = [0_{3 \times 21} ; I_{3 \times 3}]$$

**Step 3 : Transition from  $\hat{\Delta}_K(v)$  to the function  $\hat{w}(\hat{x}) = w(x) = \pi_K v(x)$**  : By using the finite element described in Figure 2.2.1, we construct the function  $\hat{w} \in \hat{P} = P_7$  which takes the values  $\hat{\Delta}_K(v)$  on the set of degrees of freedom  $\hat{\Sigma}$  so that, with relation (3.1.9)

$$(3.1.42) \quad \hat{w} = [DL(\hat{w})]_{1 \times 36} [p]_{36 \times 1}$$

where  $[p]$  denotes the column matrix of basis polynomials. They can be written

$$(3.1.43) \quad [p]_{36 \times 1} = [A]_{36 \times 36} [m7]_{36 \times 1}$$

where  $[m7]_{36 \times 1}$  collects the 36 basis monomials of degrees less or equal to 7 with respect to  $\hat{x}_1$  and  $\hat{x}_2$ . From (3.1.42), we have

$$[m7]_{36 \times 1} = [DL(m7)]_{36 \times 36} [p]_{36 \times 1} \text{ so that } [A]_{36 \times 36} = [DL(m7)]^{-1}.$$

These basis polynomials and the matrix  $[A]$  are detailed in the annex of [10].

Finally, we get the interpolating function  $\pi_K v$  of the function  $v$  by collecting relations (3.1.1) (3.1.11) (3.1.42) and (3.1.43) :

$$(3.1.44) \quad \pi_K v(x) = \hat{w}(\hat{x}) = [DLGL(v)]_{1 \times 24} [\tilde{D}]_{24 \times 24} [B]_{24 \times 36} [A]_{36 \times 36} [m7]_{36 \times 1}$$

### 3.2 Interpolation modulus associated to the curved finite element $C^1$ -compatible with ARGYRIS element when $F_K^* \in (P_5)^2$

The matrix decompositions are entirely similar to that of section 3.1 so that we just give the results and we send back to the above section for more details. Now

$$(3.2.1) \quad \pi_K^* v(x) = w^*(x)$$

which is constructed in three steps from the application  $F_K^* \in (P_5)^2$  defined by (1.3.6).

**Step 1 :**  $v \longrightarrow \Sigma_K^*(v) = [DLLC^*(v)]$ . From (2.3.2), we get :

$$(3.2.2) \quad \left\{ \begin{array}{l} [DLLC^*(v)]_{1 \times 31} = [v(a_1) \ v(a_2) \ v(a_3) \ Dv(a_1)(a_3 - a_1) \ (\bar{s} - s)Dv(a_1)\vec{\chi}'(s) \\ \quad - (\bar{s} - s)Dv(a_2)\vec{\chi}'(\bar{s}) \ Dv(a_2)(a_3 - a_2) \ Dv(a_3)(a_2 - a_3) \ Dv(a_3)(a_1 - a_3) \\ \quad D^2v(a_1)(a_3 - a_1)^2 \ (\bar{s} - s)^2 D^2v(a_1)(\vec{\chi}'(s))^2 \ (s - \bar{s})^2 D^2v(a_2)(\vec{\chi}'(\bar{s}))^2 \\ \quad D^2v(a_2)(a_3 - a_2)^2 \ D^2v(a_3)(a_2 - a_3)^2 \ D^2v(a_3)(a_1 - a_3)^2 \ D^2v(a_1)(a_2 - a_3)^2 \\ \quad D^2v(a_2)(a_3 - a_1)^2 \ (\bar{s} - s)^2 D^2v(a_3)((\vec{\chi}'(s), \vec{\chi}'(\bar{s})) \ Dv(b_1)(a_1 - c_1) \\ \quad Dv(b_2)(a_2 - c_2) \ Dv(b_3)DF_K^*(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \ v(e_1^*) \dots v(e_{10}^*)] \end{array} \right.$$

while the set of global degrees of freedom is given by

$$(3.2.3) \quad \left\{ \begin{array}{l} [DLGL^*(v)]_{1 \times 31} = [v(a_1) \ v(a_2) \ v(a_3) \ \frac{\partial v}{\partial x_1}(a_1) \ \frac{\partial v}{\partial x_2}(a_1) \ \frac{\partial v}{\partial x_1}(a_2) \\ \quad \frac{\partial v}{\partial x_2}(a_2) \ \frac{\partial v}{\partial x_1}(a_3) \ \frac{\partial v}{\partial x_2}(a_3) \ \frac{\partial^2 v}{\partial x_1^2}(a_1) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_1) \ \frac{\partial^2 v}{\partial x_2^2}(a_1) \\ \quad \frac{\partial^2 v}{\partial x_1^2}(a_2) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_2) \ \frac{\partial^2 v}{\partial x_2^2}(a_2) \ \frac{\partial^2 v}{\partial x_1^2}(a_3) \ \frac{\partial^2 v}{\partial x_1 \partial x_2}(a_3) \ \frac{\partial^2 v}{\partial x_2^2}(a_3) \\ \quad \frac{\partial v}{\partial \nu_1}(b_1) \ \frac{\partial v}{\partial \nu_2}(b_2) \ Dv(b_3)DF_K^*(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) \ v(e_1^*) \dots v(e_{10}^*)] \end{array} \right.$$

Then, with (3.1.6), we get

$$(3.2.4) \quad [DLLC^*(v)]_{1 \times 31} = [DLGL^*(v)]_{1 \times 31} [\tilde{D}^*]_{31 \times 31}, \quad \tilde{D}^* = \begin{bmatrix} \tilde{D} & 0 \\ 0 & I_7 \end{bmatrix}$$

**Step 2 : Transition from  $\Sigma_K^*(v) = [DLLC^*(v)]$  to  $\hat{\Delta}_K^*(v) = [DL^*(\hat{w}^*)]$  :** According to Figure 2.3.1, we set

$$(3.2.5) \quad \left\{ \begin{array}{l} [DL^*(\hat{w}^*)]_{1 \times 55} = [\hat{w}^*(\hat{a}_1) \quad \hat{w}^*(\hat{a}_2) \quad \hat{w}^*(\hat{a}_3) \quad \frac{\partial \hat{w}^*}{\partial \hat{x}_1}(\hat{a}_1) \quad \frac{\partial \hat{w}^*}{\partial \hat{x}_2}(\hat{a}_1) \\ \frac{\partial \hat{w}^*}{\partial \hat{x}_1}(\hat{a}_2) \quad \frac{\partial \hat{w}^*}{\partial \hat{x}_2}(\hat{a}_2) \quad \frac{\partial \hat{w}^*}{\partial \hat{x}_1}(\hat{a}_3) \quad \frac{\partial \hat{w}^*}{\partial \hat{x}_2}(\hat{a}_3) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2}(\hat{a}_1) \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_1) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2}(\hat{a}_1) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2}(\hat{a}_2) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_2) \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2}(\hat{a}_2) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2}(\hat{a}_3) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_3) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2}(\hat{a}_3) \quad - \frac{\partial \hat{w}^*}{\partial \hat{x}_1}(\hat{b}_1) \\ - \frac{\partial \hat{w}^*}{\partial \hat{x}_2}(\hat{b}_2) \quad \frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}^*}{\partial \hat{x}_1} + \frac{\partial \hat{w}^*}{\partial \hat{x}_2} \right)(\hat{b}_3) \quad \{\hat{w}^*(\hat{d}_i^*), i = 1, \dots, 12\} \\ \{- \frac{\partial \hat{w}^*}{\partial \hat{x}_1}(\hat{d}_i^*), i = 1, \dots, 4\} \quad \{- \frac{\partial \hat{w}^*}{\partial \hat{x}_2}(\hat{d}_i^*), i = 5, \dots, 8\} \\ \{\frac{\sqrt{2}}{2} \left( \frac{\partial \hat{w}^*}{\partial \hat{x}_1} + \frac{\partial \hat{w}^*}{\partial \hat{x}_2} \right)(\hat{d}_i^*), i = 9, \dots, 12\} \quad \{\hat{w}(\hat{e}_i^*), i = 1, \dots, 10\} \end{array} \right.$$

It remains to determine the transition matrix  $[B^*]$ , i.e.,

$$(3.2.6) \quad [DL^*(\hat{w}^*)]_{1 \times 55} = [DLLC^*(v)]_{1 \times 31} [B^*]_{31 \times 55}$$

so that

$$(3.2.7) \quad [DL^*(\hat{w}^*)]_{1 \times 55} = [DLGL^*(v)]_{1 \times 31} [\tilde{D}^*]_{31 \times 31} [B^*]_{31 \times 55}.$$

To specify  $B^*$ , it is convenient to consider the following partition :

$$(3.2.8) \quad [B^*]_{31 \times 55} = [B_1^* \ B_2^* \ B_3^* \ B_4^* \ B_5^* \ B_6^* \ B_7^* \ B_8^* \ B_9^*]$$

where matrices  $B_i^*$ ,  $i = 1, \dots, 9$  have 31 lines and 3, 6, 9, 3, 12, 4, 4, 4, 10 columns.

**Construction of submatrix  $B_1^*$  and  $B_9^*$  :** Relations (2.3.8) and (2.3.9) involve  $\hat{w}^*(\hat{a}_i) = w^*(a_i) = v(a_i)$ ,  $i = 1, \dots, 3$  and  $\hat{w}^*(\hat{e}_i^*) = w^*(e_i) = v(e_i)$ ,  $i = 1, \dots, 10$ , so that

$$(3.2.9) \quad {}^t B_1^* = [I_{3 \times 3} ; 0_{3 \times 28}], \quad {}^t B_9^* = [0_{10 \times 21} ; I_{10 \times 10}].$$

**Construction of submatrix  $B_2^*$**  : Relations (3.1.14) are unchanged, except for the star (\*). Then, from (1.3.6), relation (3.1.16) remains valid so that with (3.1.17), we get :

$$(3.2.10) \quad {}^t B_2^* = [0_{6 \times 3} ; {}^t(b_2)_{6 \times 6} ; 0_{6 \times 22}].$$

**Construction of submatrix  $B_3^*$**  : Relations (2.3.9) and (3.1.18) involve

$$(3.2.11) \quad \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_\alpha \partial \hat{x}_\beta} (\hat{a}_i) = D^2 v(a_i) \left( \frac{\partial F_K^*}{\partial \hat{x}_\alpha} (\hat{a}_i), \frac{\partial F_K^*}{\partial \hat{x}_\beta} (\hat{a}_i) \right) + Dv(a_i) \frac{\partial^2 F_K^*}{\partial \hat{x}_\alpha \partial \hat{x}_\beta} (\hat{a}_i)$$

so that, with (1.3.6),

$$(3.2.12) \quad \left\{ \begin{array}{lcl} \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2} (\hat{a}_1) & = D^2 v(a_1)(a_1 - a_3)^2 \\ \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_1) & = D^2 v(a_1)[a_1 - a_3, a_1 - a_3 + (\bar{s} - s)\vec{\chi}'(s)] \\ & + \frac{1}{2} Dv(a_1) \left[ a_1 - a_2 + (\bar{s} - s)\vec{\chi}'(s) - \frac{1}{2} (\bar{s} - s)^2 \vec{\chi}''(s) \right] \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2} (\hat{a}_1) & = D^2 v(a_1)[a_1 - a_3 + (\bar{s} - s)\vec{\chi}'(s)]^2 \\ & + Dv(a_1) \left[ a_1 - a_2 + (\bar{s} - s)\vec{\chi}'(s) + \frac{(\bar{s} - s)^2}{2} \vec{\chi}''(s) \right] \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3 - (\bar{s} - s)\vec{\chi}'(\bar{s})]^2 \\ & + Dv(a_2) \left[ a_2 - a_1 - (\bar{s} - s)\vec{\chi}'(\bar{s}) + \frac{(\bar{s} - s)^2}{2} \vec{\chi}''(\bar{s}) \right] \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3 - (\bar{s} - s)\vec{\chi}'(\bar{s}), a_2 - a_3] \\ & + \frac{1}{2} Dv(a_2) \left[ a_2 - a_1 - (\bar{s} - s)\vec{\chi}'(\bar{s}) - \frac{1}{2} (\bar{s} - s)^2 \vec{\chi}''(\bar{s}) \right] \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2} (\hat{a}_2) & = D^2 v(a_2)[a_2 - a_3]^2 \\ \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1^2} (\hat{a}_3) & = D^2 v(a_3)[a_1 - a_3]^2 \\ \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_1 \partial \hat{x}_2} (\hat{a}_3) & = D^2 v(a_3)[a_1 - a_3, a_2 - a_3] + \frac{1}{2} (\bar{s} - s) Dv(a_3)[\vec{\chi}'(s) - \vec{\chi}'(\bar{s})] \\ \\ \frac{\partial^2 \hat{w}^*}{\partial \hat{x}_2^2} (\hat{a}_3) & = D^2 v(a_3)[a_2 - a_3]^2 \end{array} \right.$$

Then

$$(3.2.13) \quad {}^t B_3^* = [0_{9 \times 3} ; {}^t(b_{31}^*)_{9 \times 6} ; {}^t(b_{32})_{9 \times 9} ; 0_{9 \times 13}]$$

where matrix  $b_{32}$  is defined by (3.1.22) while  $b_{31}^*$  is given by (3.2.14).

$$(3.2.14) \quad b_{31}^* = \begin{bmatrix} 0 & \frac{1}{4}(2\tilde{a}^1 - \underline{a}^1) & (\tilde{a}^1 + \frac{1}{2}\underline{a}^1) & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4}(2 + 2\tilde{a}^2 - \underline{a}^2) & 1 + \tilde{a}^2 + \frac{1}{2}\underline{a}^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & (1 + \tilde{b}^1 + \frac{1}{2}\underline{b}^1) & \frac{1}{4}(2 + 2\tilde{b}^1 - \underline{b}^1) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & (\tilde{b}^2 + \frac{1}{2}\underline{b}^2) & \frac{1}{4}(2\tilde{b}^2 - \underline{b}^2) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{2}(\tilde{c}^1 + \underline{c}^1) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{2}(\tilde{c}^2 + \underline{c}^2) & 0 \end{bmatrix}$$

In this expression, parameters  $\tilde{a}^\alpha$ ,  $\tilde{c}^\alpha$  and  $\tilde{\tilde{c}}^\alpha$  are given by relations (3.1.29) while parameters  $\underline{a}^\alpha$  and  $\underline{b}^\alpha$  are defined by

$$(3.2.15) \quad \begin{cases} \underline{a}^1 = \frac{(\vec{D}_1 \times \vec{A}_2) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} & ; \quad \underline{a}^2 = \frac{(\vec{A}_1 \times \vec{D}_1) \cdot \vec{e}_3}{(\vec{A}_1 \times \vec{A}_2) \cdot \vec{e}_3} \\ \underline{b}^1 = \frac{(\vec{D}_2 \times \vec{B}_2) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} & ; \quad \underline{b}^2 = \frac{(\vec{B}_1 \times \vec{D}_2) \cdot \vec{e}_3}{(\vec{B}_1 \times \vec{B}_2) \cdot \vec{e}_3} \\ \vec{D}_1 = (\bar{s} - s)^2 \chi''(s) = \underline{a}^\alpha \vec{A}_\alpha & ; \quad \vec{D}_2 = (\bar{s} - s)^2 \chi''(\bar{s}) = \underline{b}^\alpha \vec{B}_\alpha \end{cases}$$

**Construction of submatrix  $B_4^*$**  : Since  $F_K$  and  $F_K^*$  are affine along the sides  $[\hat{a}_3^*, \hat{a}_\alpha^*]$ ,  $\alpha = 1, 2$ , functions  $\hat{f}_\alpha, \hat{g}_\alpha$ ,  $\alpha = 1, 2$  are unchanged. By using (1.3.6), we obtain

$$(3.2.16) \quad \begin{cases} [\hat{f}_1(\hat{x}_1) \quad \hat{f}_2(\hat{x}_2) \quad \hat{f}_3^*(\hat{x}_1)] & = [DLLC^*(v)]_{1 \times 31} [\hat{\mathcal{F}}_1^*(\hat{x}_1) \quad \hat{\mathcal{F}}_2^*(\hat{x}_2) \quad \hat{\mathcal{F}}_3^*(\hat{x}_1)]_{31 \times 3} \\ [\hat{g}_1(\hat{x}_1) \quad \hat{g}_2(\hat{x}_2) \quad \hat{g}_3^*(\hat{x}_1)] & = [DLLC^*(v)]_{1 \times 31} [\hat{\mathcal{G}}_1^*(\hat{x}_1) \quad \hat{\mathcal{G}}_2^*(\hat{x}_2) \quad \hat{\mathcal{G}}_3^*(\hat{x}_1)]_{31 \times 3} \end{cases}$$

with

$$(3.2.17) \quad \left\{ {}^t[\hat{\mathcal{F}}_\alpha^*(\hat{x}_\alpha)] = [{}^t\hat{\mathcal{F}}_\alpha(\hat{x}_\alpha) \ ; \ 0_{1 \times 7}] \ ; \ {}^t[\hat{\mathcal{G}}_\alpha^*(\hat{x}_\alpha)] = [{}^t\hat{\mathcal{G}}_\alpha(\hat{x}_\alpha) \ ; \ 0_{1 \times 7}], \quad \alpha = 1, 2 \right.$$

$$(3.2.18) \quad \left\{ {}^t[\hat{\mathcal{F}}_3^*(\hat{x}_1)] = [(\hat{x}_1)^3 \{6\hat{x}_1^2 - 15\hat{x}_1 + 10\} \ ; \ (1 - \hat{x}_1)^3 \{6\hat{x}_1^2 + 3\hat{x}_1 + 1\}] ; \right.$$

$$0 \ ; \ \frac{1}{2}\hat{x}_1^3(1 - \hat{x}_1)^2\underline{a}^1 \ ; \ \hat{x}_1^3(1 - \hat{x}_1)\{4 - 3\hat{x}_1 + \frac{1}{2}(1 - \hat{x}_1)\underline{a}^2\} ;$$

$$\hat{x}_1(1 - \hat{x}_1)^3 \left\{ 1 + 3\hat{x}_1 + \frac{1}{2}\hat{x}_1\underline{b}^1 \right\} \ ; \ \frac{1}{2}\hat{x}_1^2(1 - \hat{x}_1)^3\underline{b}^2 \ ; \ 0 \ 0 \ 0 ;$$

$$\left. \frac{1}{2}\hat{x}_1^3(1 - \hat{x}_1)^2 \ ; \ \frac{1}{2}\hat{x}_1^2(1 - \hat{x}_1)^3 \ ; \ 0_{1 \times 19} \right]$$

$$(3.2.19) \quad \left\{ \begin{array}{l} {}^t[\hat{\mathcal{G}}_3^*(\hat{x}_1)] = [0 \ 0 \ 0 ; \ \hat{x}_1^2(2\hat{x}_1 - 1) \left\{ 5 - 4\hat{x}_1 - \frac{1}{2}(1 - \hat{x}_1) \left( \tilde{a}^1 + \frac{1}{2}\tilde{a}^1 \right) \right\} ; \\ \hat{x}_1^2(2\hat{x}_1 - 1) \left\{ -3 + \frac{5\hat{x}_1}{2} - \frac{1}{2}(1 - \hat{x}_1) \left( \tilde{a}^2 + \frac{1}{2}\tilde{a}^2 \right) \right\} ; \\ - \frac{1}{2}(1 - \hat{x}_1)^2(1 - 2\hat{x}_1) \left\{ 1 + 5\hat{x}_1 + \hat{x}_1(\tilde{b}^1 + \frac{1}{2}\tilde{b}^1) \right\} ; \\ (1 - \hat{x}_1)^2(1 - 2\hat{x}_1) \left\{ 1 + 4\hat{x}_1 - \frac{1}{2}\hat{x}_1(\tilde{b}^2 + \frac{1}{2}\tilde{b}^2) \right\} ; \ 0 \ 0 ; \\ \frac{1 + \tilde{a}^1}{2\tilde{a}^2} \hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1) ; \ \frac{1}{2} \left( 1 + \frac{\tilde{a}^2}{1 + \tilde{a}^1} \right) \hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1) ; \\ - \frac{1}{2} \left( 1 + \frac{\tilde{b}^1}{1 + \tilde{b}^2} \right) \hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1) ; \ - \frac{1 + \tilde{b}^2}{2\tilde{b}^1(1 + \tilde{b}^2)} \hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1) ; \ 0 \ 0 ; \\ - \frac{1}{2(1 + \tilde{a}^1)\tilde{a}^2} \hat{x}_1^2(1 - \hat{x}_1)(1 - 2\hat{x}_1) ; \ \frac{1}{2\tilde{b}^1(1 + \tilde{b}^2)} \hat{x}_1(1 - \hat{x}_1)^2(1 - 2\hat{x}_1) ; \\ 0 \ 0 \ 0 ; \ 16(1 - \hat{x}_1)^2\hat{x}_1^2 ; \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 ] \end{array} \right.$$

Then, by similarity with relation (3.1.37), we obtain

$$(3.2.20) \quad B_4^* = \left[ \tilde{E}^1 \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2^* \left( \frac{1}{2} \right) + \tilde{E}^2 \hat{\mathcal{G}}_2^* \left( \frac{1}{2} \right) ; \ \tilde{F}^1 \frac{d}{d\hat{x}_1} \hat{\mathcal{F}}_1^* \left( \frac{1}{2} \right) + \tilde{F}^2 \hat{\mathcal{G}}_1^* \left( \frac{1}{2} \right) ; \begin{pmatrix} 0_{20 \times 1} \\ -\sqrt{2} \\ 0_{10 \times 3} \end{pmatrix} \right]_{31 \times 3}$$

with

$$\begin{aligned} \tilde{E}^1 &= \langle a_3 - a_1 - \frac{1}{4}(a_2 - a_1) - \frac{1}{32}(\bar{s} - s)[5\vec{\chi}'(s) - 13\vec{\chi}'(\bar{s})] - \frac{(\bar{s} - s)^2}{64}[\vec{\chi}''(s) + \vec{\chi}''(\bar{s})], \frac{a_2 - a_3}{|a_2 - a_3|^2} \rangle \\ \tilde{E}^2 &= \langle a_3 - a_1 - \frac{1}{4}(a_2 - a_1) - \frac{1}{32}(\bar{s} - s)[5\vec{\chi}'(s) - 13\vec{\chi}'(\bar{s})] - \frac{(\bar{s} - s)^2}{64}[\vec{\chi}''(s) + \vec{\chi}''(\bar{s})], \frac{a_1 - c_1}{|a_1 - c_1|^2} \rangle \\ \tilde{F}^1 &= \langle a_3 - a_2 - \frac{1}{4}(a_1 - a_2) - \frac{1}{32}(\bar{s} - s)(5\vec{\chi}'(\bar{s}) - 13\vec{\chi}'(s)) + \frac{(\bar{s} - s)^2}{64}[\vec{\chi}''(s) + \vec{\chi}''(\bar{s})], \frac{a_1 - a_3}{|a_1 - a_3|^2} \rangle \\ \tilde{F}^2 &= \langle a_3 - a_2 - \frac{1}{4}(a_1 - a_2) - \frac{1}{32}(\bar{s} - s)(5\vec{\chi}'(\bar{s}) - 13\vec{\chi}'(s)) + \frac{(\bar{s} - s)^2}{64}[\vec{\chi}''(s) + \vec{\chi}''(\bar{s})], \frac{a_2 - c_2}{|a_2 - c_2|^2} \rangle. \end{aligned}$$

**Construction of submatrix  $B_5^*$**  : Relations (2.3.3) (2.3.6) (2.3.7) and (3.2.21) give

$$(3.2.21) \quad \left\{ \begin{array}{l} B_5^* = \left[ \hat{\mathcal{F}}_2^* \left( \frac{5}{6} \right) ; \ \hat{\mathcal{F}}_2^* \left( \frac{2}{3} \right) ; \ \hat{\mathcal{F}}_2^* \left( \frac{1}{3} \right) ; \ \hat{\mathcal{F}}_2^* \left( \frac{1}{6} \right) ; \ \hat{\mathcal{F}}_1^* \left( \frac{1}{6} \right) ; \ \hat{\mathcal{F}}_1^* \left( \frac{1}{3} \right) ; \right. \\ \left. \hat{\mathcal{F}}_1^* \left( \frac{2}{3} \right) ; \ \hat{\mathcal{F}}_1^* \left( \frac{5}{6} \right) ; \ \hat{\mathcal{F}}_3^* \left( \frac{5}{6} \right) ; \ \hat{\mathcal{F}}_3^* \left( \frac{2}{3} \right) ; \ \hat{\mathcal{F}}_3^* \left( \frac{1}{3} \right) ; \ \hat{\mathcal{F}}_3^* \left( \frac{1}{6} \right) \right]_{31 \times 12} \end{array} \right.$$

**Construction of submatrices  $B_6^*$  and  $B_7^*$  :** From relations (2.3.3) (2.3.6) (2.3.7) and (3.2.16), we obtain

$$(3.2.22) \quad \left\{ \begin{array}{l} B_6^* = \left[ - < \frac{\partial F_K^*}{\partial \hat{x}_1}(\hat{d}_1^*), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2^* \left( \frac{5}{6} \right) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{\mathcal{G}}_2^* \left( \frac{5}{6} \right) > ; \right. \\ \quad - < \frac{\partial F_K^*}{\partial \hat{x}_1}(\hat{d}_2^*), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2^* \left( \frac{2}{3} \right) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{\mathcal{G}}_2^* \left( \frac{2}{3} \right) > ; \\ \quad - < \frac{\partial F_K^*}{\partial \hat{x}_1}(\hat{d}_3^*), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2^* \left( \frac{1}{3} \right) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{\mathcal{G}}_2^* \left( \frac{1}{3} \right) > ; \\ \quad \left. - < \frac{\partial F_K^*}{\partial \hat{x}_1}(\hat{d}_4^*), \frac{a_2 - a_3}{|a_2 - a_3|^2} \frac{d}{d\hat{x}_2} \hat{\mathcal{F}}_2^* \left( \frac{1}{6} \right) + \frac{a_1 - c_1}{|a_1 - c_1|^2} \hat{\mathcal{G}}_2^* \left( \frac{1}{6} \right) > \right]_{31 \times 4} \end{array} \right.$$

$$(3.2.23) \quad \left\{ \begin{array}{l} B_7^* = \left[ - < \frac{\partial F_K^*}{\partial \hat{x}_2}(\hat{d}_5^*), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{\mathcal{F}}_1^*}{d\hat{x}_1} \left( \frac{1}{6} \right) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{\mathcal{G}}_1^* \left( \frac{1}{6} \right) > ; \right. \\ \quad - < \frac{\partial F_K^*}{\partial \hat{x}_2}(\hat{d}_6^*), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{\mathcal{F}}_1^*}{d\hat{x}_1} \left( \frac{1}{3} \right) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{\mathcal{G}}_1^* \left( \frac{1}{3} \right) > ; \\ \quad - < \frac{\partial F_K^*}{\partial \hat{x}_2}(\hat{d}_7^*), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{\mathcal{F}}_1^*}{d\hat{x}_1} \left( \frac{2}{3} \right) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{\mathcal{G}}_1^* \left( \frac{2}{3} \right) > ; \\ \quad \left. - < \frac{\partial F_K^*}{\partial \hat{x}_2}(\hat{d}_8^*), \frac{a_1 - a_3}{|a_1 - a_3|^2} \frac{d\hat{\mathcal{F}}_1^*}{d\hat{x}_1} \left( \frac{5}{6} \right) + \frac{a_2 - c_2}{|a_2 - c_2|^2} \hat{\mathcal{G}}_1^* \left( \frac{5}{6} \right) > \right] \end{array} \right.$$

$$(3.2.24) \quad B_8^* = \left[ -\sqrt{2}\hat{\mathcal{G}}_3^* \left( \frac{5}{6} \right) ; -\sqrt{2}\hat{\mathcal{G}}_3^* \left( \frac{2}{3} \right) ; -\sqrt{2}\hat{\mathcal{G}}_3^* \left( \frac{1}{3} \right) ; -\sqrt{2}\hat{\mathcal{G}}_3^* \left( \frac{1}{6} \right) \right]$$

**Step 3 : Transition from  $\hat{\Delta}_K^*(v)$  to  $\pi_K^* v(x) = w^*(x) = \hat{w}^*(\hat{x})$  :** Let  $[m9]_{55 \times 1}$  be the 55 basis monomials of degree less or equal to 9 in  $\hat{x}_1$  and  $\hat{x}_2$  and  $[A^*]_{55 \times 55} = [DL^*(m9)]^{-1}$ . Corresponding basis polynomials are detailed in the annex of [10]. Then,

$$(3.2.25) \quad \pi_K^* v(x) = \hat{w}^*(\hat{x}) = [DLGL^*(v)]_{1 \times 31} [\tilde{D}^*]_{31 \times 31} [B^*]_{31 \times 55} [A^*]_{55 \times 55} [m9]_{55 \times 1}.$$

### 3.3 Some numerical tests of interpolation properties

It is important to test numerically the interpolation properties. We will compute the asymptotic order of the interpolation error for the following two cases :

- case of an exact representation of the curved boundary ;
- case of an approximate representation of the curved boundary.

#### 3.3.1 Tests on the asymptotic order of interpolation error (generalities)

From [2, (4.1) and (4.51)], we have to numerically verify that

$$(3.3.1) \quad \left\{ \begin{array}{l} |v - \pi_K v|_{m,K} \leq ch_K^{6-m} \|v\|_{6,K} \\ |v - \pi_K^* v|_{m,K} \leq ch_K^{6-m} \|v\|_{6,K} \end{array} , \quad m = 0, \dots, 6, \quad \forall v \in H^6(K). \right.$$

For simplicity, since it can be proved that the asymptotic error estimates in norms  $\|\cdot\|_{L^2(K)}$  and  $\|\cdot\|_{L^\infty(K)}$  have the same order, we will just check that

$$(3.3.2) \quad |v - \pi_K v|_{0,\infty,K} = O(h_K^6), \quad |v - \pi_K^* v|_{0,\infty,K} = O(h_K^6), \quad \forall v \in W^{6,\infty}(K),$$

or, more easily, that the asymptotic behaviour of  $|v - \pi_K v|$  (resp.  $|v - \pi_K^* v|$ ) at a given point  $F_{K_h}(\hat{x}_0, \hat{x}_0)$  (resp.  $F_{K_h}^*(\hat{x}_0, \hat{x}_0)$ ) is in  $O(h_h^6)$ .

In this way, for each example into consideration (see §§ 3.3.2 and 3.3.3), we successively


- i) consider a series of nested triangles whose sizes decrease as  $h, \frac{h}{2}, \frac{h}{4}, \dots$ ;
- ii) interpolate given regular functions  $v$  ;
- iii) illustrate the results on graphics displaying –  $\log h$  in abscissa and –  $\log |v - \pi_K v|$  (resp. –  $\log |v - \pi_K^* v|$ ) in ordinate and check that

$$(3.3.3) \quad \log |v - \pi_K v| = C + 6 \log h ; \quad \log |v - \pi_K^* v| = C + 6 \log h.$$

### 3.3.2 Numerical tests for an "exact" interpolation of the boundary

Firstly it seems interesting to make some numerical experiments for an exact representation of the curved boundary so that it only remains the interpolation error of the function.

#### Example 3.3.1 : Unit right-angled triangle



**Figure 3.3.1 :** The unit right-angled triangle and the associate sequence of triangles  $K_h$

Of course this is the most simple case which is associated to the following representation of the curved (here straight !) side  $a_1a_2$  :

$$(3.3.4) \quad x_1 = \chi_1(s) = 1 - s, \quad x_2 = \chi_2(s) = s \quad \text{with } \underline{s} = 0 \leq s \leq \bar{s} = 1.$$

This test is interesting since  $F_K = I$  and  $F_K^* = I$  involves


$$(3.3.5) \quad \pi_K v(x) \equiv v(x), \quad \pi_K^* v(x) = v(x), \quad \forall v \in P_5$$

or, equivalently, by using (3.1.44) and (3.2.25)

$$(3.3.6) \quad \begin{cases} x_1^p x_2^q = [DLGL(x_1^p x_2^q)] [\tilde{D}] [B] [A] [m7], & \forall p, q \in \mathbb{N} \text{ and } p + q = 5; \\ x_1^p x_2^q = [DLGL^*(x_1^p x_2^q)] [\tilde{D}^*] [B^*] [A^*] [m9], & \forall p, q \in \mathbb{N} \text{ and } p + q = 5. \end{cases}$$

The numerical results show that identities (3.3.5) are exactly satisfied. We have also checked the estimates (3.3.3); in this way, we have considered a sequence of right-angled triangle deduced from the unit right-angled triangle by an homothetic transformation centered at point  $(\frac{1}{2}, \frac{1}{2})$  and whose ratio is  $h = \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$  (see Figure 3.3.1). More generally, we have successfully checked the same properties for any straight triangle  $K$  (note that the associate mapping  $F_K$  is still affine in this case).

**Example 3.3.2 : A curved unit right-angled triangle (case  $F_K \in (P_3)^2$ )**



**Figure 3.3.2 :** The curved unit right-angled triangle and the associate sequence of triangles  $K_h$

Now the straight side  $a_1 a_2$  is replaced by the curved side whose parametric equations are (see Figure 3.3.2) :

$$(3.3.7) \quad x_1 = \chi_1(s) = 1 - s, \quad x_2 = \chi_2(s) = \frac{1}{2} s[s^2 - 3s + 4], \quad s = 0 \leq s \leq \bar{s} = 1,$$

so that its interpolation by the mapping (1.3.2) is exact. The associate application  $F_K$  is given by (see (1.3.2)) :

$$(3.3.8) \quad F_{K1}(\hat{x}_1, \hat{x}_2) = \hat{x}_1, \quad F_{K2}(\hat{x}_1, \hat{x}_2) = \hat{x}_2 + \frac{1}{4} \hat{x}_1 \hat{x}_2 [3 + \hat{x}_1 - \hat{x}_2].$$

Now, since the mapping  $F_K$  is no longer affine, the identities (3.3.5) are no longer satisfied. In order to test the estimate (3.3.2)<sub>1</sub>, we have to introduce a sequence of triangle  $K_h$  whose diameter  $h$  is decreasing to 0 and whose curved sides coincide (for this particular example) with the curved boundary of the domain. In this way we will use the triangles  $K_h$ .

whose vertices  $(a_{1h}, a_{2h}, a_{3h})$  are given by :

$$(3.3.9) \quad \left\{ \begin{array}{l} \text{Vertex } a_{1h} : s_h = \frac{1}{2} (1-h) \implies (x_{11} = \frac{1}{2} (1+h); x_{21} = \frac{1}{16} (1-h)(11+4h+h^2)) \\ \text{Vertex } a_{2h} : \bar{s}_h = \frac{1}{2} (1+h) \implies (x_{12} = \frac{1}{2} (1-h); x_{22} = \frac{1}{16} (1+h)(11-4h+h^2)) \\ \text{Vertex } a_{3h} : (x_{13} = \frac{1}{2} (1-h); x_{23} = \frac{1}{16} (1-h)(11+4h+h^2)) \end{array} \right.$$


so that the associate application  $F_{K_h}$  is given by relation (1.3.2), i.e.,

$$(3.3.10) \quad \left\{ \begin{array}{l} F_{K1h}(\hat{x}_1, \hat{x}_2) = \frac{1}{2} (1-h) + h\hat{x}_1 \\ F_{K2h}(\hat{x}_1, \hat{x}_2) = \frac{1}{16} (1-h)(11+4h+h^2) + \frac{h}{8} (7+h^2)\hat{x}_2 \\ \quad + \frac{1}{4} \hat{x}_1 \hat{x}_2 h^2 [3 + h(\hat{x}_1 - \hat{x}_2)]. \end{array} \right.$$

Then, to check estimate (3.3.2)<sub>1</sub>, we graphically represent (3.3.3)<sub>1</sub> for given functions  $v$ , for instance  $v(x_1, x_2) = x_1^6 x_2^4$ , and  $v(x_1, x_2) = \cos(3x_1) \sin(2x_2)$ , at the image of point  $(\hat{x}_1 = \frac{3}{8}, \hat{x}_2 = \frac{3}{8})$  by the application  $F_{K_h}$ , i.e.

$$(3.3.11) \quad x_1 = F_{K1h} \left( \frac{3}{8}, \frac{3}{8} \right) = \frac{1}{2} - \frac{h}{8}; \quad x_2 = F_{K2h} \left( \frac{3}{8}, \frac{3}{8} \right) = \frac{11}{16} - \frac{7h}{64} - \frac{21h^2}{256} - \frac{h^3}{64}$$


for  $h = \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$ . Thus, Figure 3.3.3 shows that the slope is effectively 6 according to the theoretical prediction (3.3.3).



**Figure 3.3.3** : Asymptotic behaviour of  $|v - \pi_K v|$  at point  $F_{K_h}(\frac{3}{8}, \frac{3}{8})$  when  $h \rightarrow 0$ .

**Example 3.3.3** : A curved unit right-angled triangle (case  $F_K^* \in (P_5)^2$ ) and the associate sequence of triangles  $K_h$

Consider the curved side  $a_1 a_2$  whose parametric equations are (see Figure 3.3.4)



**Figure 3.3.4 :** The curved unit right-angled triangle and the associate sequence of triangles  $K_h^*$

$$(3.3.12) \quad \begin{cases} x_1 = \chi_1(s) = 1 - s, \quad x_2 = \chi_2(s) = \frac{s}{10} [4s^4 + 13s^2 - 27s + 20], \\ \underline{s} = 0 \leq s \leq \bar{s} = 1 \end{cases}$$

so that its interpolation by the mapping (1.3.6) is exact. The associate application  $F_K^*$  is given by (see (1.3.6)) :

$$(3.3.13) \quad \begin{cases} F_{K1}^*(\hat{x}_1, \hat{x}_2) = \hat{x}_1 \\ F_{K2}^*(\hat{x}_1, \hat{x}_2) = \hat{x}_2 + \frac{1}{20} \hat{x}_1 \hat{x}_2 [-4(\hat{x}_2)^3 - 4(\hat{x}_2)^2 - 17\hat{x}_2 - 5 \\ \quad + 4(\hat{x}_1)^3 - 16(\hat{x}_1)^2 + 37\hat{x}_1] \end{cases}$$


Here again we consider the sequence of triangles  $K_h$  whose vertices are given by

$$(3.3.14) \quad \begin{cases} \text{vertex } a_{1h} : \left( x_{11} = \frac{1}{2} (1+h); x_{21} = \frac{1}{80} (1-h)(40+24h+19h^2-4h^3+h^4) \right) \\ \text{vertex } a_{2h} : \left( x_{12} = \frac{1}{2} (1-h); x_{22} = \frac{1}{80} (1+h)(40-24h+19h^2+4h^3+h^4) \right) \\ \text{vertex } a_{3h} : \left( x_{13} = \frac{1}{2} (1-h); x_{23} = \frac{1}{80} (1-h)(40+24h+19h^2-4h^3+h^4) \right) \end{cases}$$

so that the application  $F_{K_h}^*$  is given by relation (1.3.6) :

$$(3.3.15) \quad \left\{ \begin{array}{l} F_{K1h}^*(\hat{x}_1, \hat{x}_2) = \frac{1}{2} (1 - h) + h\hat{x}_1 \\ F_{K2h}^*(\hat{x}_1, \hat{x}_2) = \frac{1}{80} (1 - h)(40 + 24h + 19h^2 - 4h^3 + h^4) \\ \quad + \frac{h}{40} (16 + 23h^2 + h^4)\hat{x}_2 + \frac{1}{2}\hat{x}_1\hat{x}_2[\frac{h^2}{2}(1 - 2h^2) \\ \quad + \frac{h^3}{10}(23 + 10h + 4h^2)\hat{x}_1 + \frac{h^3}{10}(-23 + 10h - 4h^2)\hat{x}_2 \\ \quad - \frac{h^4}{5}(5 + 3h)(\hat{x}_1)^2 - \frac{h^4}{5}(5 - 3h)(\hat{x}_2)^2 + \frac{2h^5}{5}(\hat{x}_1)^3 - \frac{2h^5}{5}(\hat{x}_2)^3] \end{array} \right.$$

Now we obtain Figure 3.3.5 which shows again an  $O(h^6)$  error estimate.



**Figure 3.3.5 :** Asymptotic behaviour of  $|v - \pi_K^* v|$  at point  $F_{K_h}^*(\frac{1}{4}, \frac{1}{4})$  when  $h \rightarrow 0$ .

### 3.3.3 Numerical tests for an approximate interpolation of the boundary


Previous examples are such that applications  $F_K \in (P_3)^2$  or  $F_K^* \in (P_5)^2$  give an exact representation of the curved side  $a_1 a_2$ . Now let us consider some examples for which the curved side is only approximated by the applications  $F_K$  or  $F_K^*$ .

**Example 3.3.4 :** The  $(P_3)^2$  approximation of the curved unit right-angled triangle studied in the example 3.3.3

In the example 3.3.3 we have considered a sequence of triangles  $K_h^*$  such that the curved side  $a_{1h}a_{2h}$  coincided with the initial boundary. In this way we have used the application  $F_{K_h}^* \in (P_5)^2$  defined by relation (3.3.15). Now, starting from the same sequence of curved triangle  $a_{1h}a_{2h}a_{3h}$  we consider the  $(P_3)^2$ -approximation of the curved side. Thus, by substituting relations (3.3.12) (3.3.14) into relations (1.3.2), we get the application  $F_{K_h}$

$$(3.3.16) \quad \left\{ \begin{array}{l} F_{K1h}(\hat{x}_1, \hat{x}_2) = \frac{1}{2} (1 - h) + h\hat{x}_1 \\ F_{K2h}(\hat{x}_1, \hat{x}_2) = \frac{1}{80} (1 - h)(40 + 24h + 19h^2 - 4h^3 + h^4) \\ \quad + \frac{h}{40} (16 + 23h^2 + h^4)\hat{x}_2 + \frac{h^2}{20} \hat{x}_1\hat{x}_2[5 - 10h^2 + h(23 + 2h^2)(\hat{x}_1 - \hat{x}_2)] \end{array} \right.$$

Here again Figure 3.3.6 shows an  $O(h^6)$  error estimate




**Figure 3.3.6 :** Asymptotic behaviour of  $|v - \pi_K v|$  at point  $F_{K_h}(\frac{3}{8}, \frac{3}{8})$  when  $h \rightarrow 0$ .

### Example 3.3.5 : Case of an elliptic edge


The curved side of triangle  $K$  is represented by (see Figure 3.3.7)

$$(3.3.17) \quad x_1 = \chi_1(s) = \cos s; \quad x_2 = \chi_2(s) = R \sin s, \quad 0 < R < 1, \quad s = \frac{\pi}{6} \leq s \leq \bar{s} = \frac{\pi}{3},$$

and it can be approximated with the help of  $F_K \in (P_3)^2$  or  $F_K^* \in (P_5)^2$ , respectively defined by relations (1.3.2) or (1.3.6).



**Figure 3.3.7 :** The "curved elliptic" triangle  $K$  and the associate sequence of triangles  $K_h$  (or  $K_h^*$ ) when  $R = 0.4$



**Figure 3.3.8 :** Asymptotic behaviour of  $|v - \pi_K v|$  (resp.  $|v - \pi_K^* v|$ ) at point  $F_{K_h}(\frac{3}{8}, \frac{3}{8})$  (resp.  $F_{K_h}^*(\frac{1}{4}, \frac{1}{4})$ ) when  $h \rightarrow 0$ .

Here again we introduce the sequence  $K_h$  (or  $K_h^*$ ) of triangles whose vertices are given by  $(0 < h < \frac{\pi}{12})$  :

$$(3.3.18) \quad \begin{cases} \text{vertex } a_{1h} : \underline{s}_h = \frac{\pi}{4} - h \implies \left( x_1 = \cos \left( \frac{\pi}{4} - h \right); x_2 = R \sin \left( \frac{\pi}{4} - h \right) \right) \\ \text{vertex } a_{2h} : \bar{s}_h = \frac{\pi}{4} + h \implies \left( x_1 = \cos \left( \frac{\pi}{4} + h \right); x_2 = R \sin \left( \frac{\pi}{4} + h \right) \right) \\ \text{vertex } a_{3h} : \left( x_1 = \cos \left( \frac{\pi}{4} + h \right), x_2 = R \sin \left( \frac{\pi}{4} - h \right) \right) \end{cases}$$

while the computation of the error is made at an internal point of triangles  $K_h$  (or  $K_h^*$ ). The associated results are displayed on Figure 3.3.8. Here again, we find an asymptotic error in  $O(h^6)$ .

## Concluding remarks

Thus, these examples show the great efficiency of these curved  $C^1$  elements and they illustrate the relevance of the theoretical study of the order of the approximation. These methods will be used to approximate thin plate and thin shell problems in the second part of this work [10].

## Acknowledgments

The authors are indebted to P.G. CIARLET for suggesting this study and for his constant interest during the investigation. Also we take this opportunity to thank I. BABUSKA for his stimulating incitement to realize the corresponding implementation.

## References

- [1] M. BERNADOU, Variational formulation and approximation of junctions between thin shells in "Proceedings of the Fifth International Symposium on Numerical Methods in Engineering, Vol. 1, pp. 407-414, Edited by R. GRUBER, J. PERIAUX and R.P. SHAW, CMP and Springer-Verlag, 1989.
- [2] M. BERNADOU,  $C^1$ -curved finite elements with numerical integration for thin plate and thin shell problems. Part 1 : Construction and interpolation properties of curved  $C^1$  finite elements (to appear in Comput. Methods Appl. Mech. Engrg.).
- [3] M. BERNADOU,  $C^1$ -curved finite elements with numerical integration for thin plate and thin shell problems. Part 2 : Approximation of thin plate and thin shell problems (to appear in Comput. Methods Appl. Mech. Engrg.).
- [4] J.H. ARGYRIS, I. FRIED, D.W. SCHARPF, The TUBA family of plate elements for the matrix displacement method, The Aeronautical Journal of the Royal Aeronautical Society, 72 (1968), pp.701-709.
- [5] P.G. CIARLET, The Finite Element Method for Elliptic Problems. North-Holland, Amsterdam, 1978.
- [6] P.G. CIARLET, Basic Error Estimates for Elliptic Problems. In : Handbook of Numerical Analysis (P.G. CIARLET and J.L. LIONS editors). Vol. II : Finite Element Methods, Part 1, pp. 17-351, North-Holland, Amsterdam, 1991.
- [7] S. WOLFRAM, Mathematica : A system for Doing Mathematics by Computer, Addison-Wesley Publishing Company, Redwood City, 1988.
- [8] J.M. BOISSERIE, A new  $C^1$  finite element : full heptic, Internat. J. Numer. Meth. Engng. 28, pp. 667-677, 1989.
- [9] M. BERNADOU, J.M. BOISSERIE, The Finite Element Method in Thin Shell Theory : Application to Arch Dam Simulations, Birkhäuser, Boston, 1982.
- [10] M. BERNADOU, J.M. BOISSERIE, Curved finite elements of class  $C^1$  : implementation and numerical experiments. Part 2 : Applications to thin plate and thin shell problems. Rapports de Recherche INRIA (to appear).

## ANNEX : Basis polynomials for $P_7$ and $P_9$ finite elements

In order to construct curved finite elements  $C^1$ -compatible with the Argyris triangle, we have proposed two methods :

(i) The first one is based in the interpolation of the boundary by polynomials of degree 3. In this way, we are led to the basic finite element displayed on Figure 2.2.1. Corresponding basis polynomials are given by relation (3.1.43), e.g.,

$$[p]_{36 \times 1} = [A]_{36 \times 36} [m7]_{36 \times 1}.$$

The matrices  $[m7]$  and  $[A]$  are described on Figure A.1 and A.2.

(ii) The second one is based on the interpolation of the boundary by polynomials of degree 5. Then, we are led to the basic finite element which is shown on Figure 2.3.1. Corresponding basis polynomials are given by relation (3.2.25), e.g.,

$$[p^*]_{55 \times 1} = [A^*]_{55 \times 55} [m9]_{55 \times 1}.$$

The associate matrices  $[m9]$  and  $[A^*]$  are given on Figure A.1 and A.3.

$$[m7] = \left[ \begin{array}{l}
 X1^{**7} \\
 X1^{**6} * X2 \\
 X1^{**5} * X2^{**2} \\
 X1^{**4} * X2^{**3} \\
 X1^{**3} * X2^{**4} \\
 X1^{**2} * X2^{**5} \\
 X1 * X2^{**6} \\
 X2^{**7} \\
 X1^{**6} \\
 X1^{**5} * X2 \\
 X1^{**4} * X2^{**2} \\
 X1^{**3} * X2^{**3} \\
 X1^{**2} * X2^{**4} \\
 X1 * X2^{**5} \\
 X2^{**6} \\
 X1^{**5} \\
 X1^{**4} * X2 \\
 X1^{**3} * X2^{**2} \\
 X1^{**2} * X2^{**3} \\
 X1 * X2^{**4} \\
 X2^{**5} \\
 X1^{**4} \\
 X1^{**3} * X2 \\
 X1^{**2} * X2^{**2} \\
 X1 * X2^{**3} \\
 X2^{**4} \\
 X1^{**3} \\
 X1^{**2} * X2 \\
 X1 * X2^{**2} \\
 X2^{**3} \\
 X1^{**2} \\
 X1 * X2 \\
 X2^{**2} \\
 X1 \\
 X2 \\
 1.
 \end{array} \right]
 \quad
 [m9] = \left[ \begin{array}{l}
 X1^{**9} \\
 X1^{**8} * X2 \\
 X1^{**7} * X2^{**2} \\
 X1^{**6} * X2^{**3} \\
 X1^{**5} * X2^{**4} \\
 X1^{**4} * X2^{**5} \\
 X1^{**3} * X2^{**6} \\
 X1^{**2} * X2^{**7} \\
 X1 * X2^{**8} \\
 X2^{**9} \\
 X1^{**8} \\
 X1^{**7} * X2 \\
 X1^{**6} * X2^{**2} \\
 X1^{**5} * X2^{**3} \\
 X1^{**4} * X2^{**4} \\
 X1^{**3} * X2^{**5} \\
 X1^{**2} * X2^{**6} \\
 X1 * X2^{**7} \\
 X2^{**8} \\
 X1^{**7} \\
 X1^{**6} * X2 \\
 X1^{**5} * X2^{**2} \\
 X1^{**4} * X2^{**3} \\
 X1^{**3} * X2^{**4} \\
 X1^{**2} * X2^{**5} \\
 X1 * X2^{**6} \\
 X2^{**7} \\
 X1^{**6} \\
 X1^{**5} * X2 \\
 X1^{**4} * X2^{**2} \\
 X1^{**3} * X2^{**3} \\
 X1^{**2} * X2^{**4} \\
 X1 * X2^{**5} \\
 X2^{**6} \\
 X1^{**5} \\
 X1^{**4} * X2 \\
 X1^{**3} * X2^{**2} \\
 X1^{**2} * X2^{**3} \\
 X1 * X2^{**4} \\
 X2^{**5} \\
 X1^{**4} \\
 X1^{**3} * X2 \\
 X1^{**2} * X2^{**2} \\
 X1 * X2^{**3} \\
 X2^{**4} \\
 X1^{**3} \\
 X1^{**2} * X2 \\
 X1 * X2^{**2} \\
 X2^{**3} \\
 X1^{**2} \\
 X1 * X2 \\
 X2^{**2} \\
 X1 \\
 X2 \\
 1.
 \end{array} \right]$$

Figure A.1 : Basis monomials of  $P_7$  and  $P_9$

A(1, 1)	-6496/27	A(7, 11)	-1472/9	A(14, 3)	-(-8)/3	A(20, 13)	-64	A(28, 24)	-512/9
A(1, 3)	-55376/27	A(7, 13)	-7448/9	A(14, 4)	-(-80)/3	A(20, 17)	-816	A(28, 25)	-2048/9
A(1, 4)	-15296/3	A(7, 15)	-416/3	A(14, 5)	-(-80)/3	A(20, 18)	-992	A(28, 29)	-(-256)/9
A(1, 5)	-86464/27	A(7, 18)	-(-4337)/18	A(14, 6)	-8/3	A(20, 19)	-176	A(29, 3)	-(-2048)/9
A(1, 6)	-464/3	A(7, 19)	-(-1177)/2	A(14, 7)	-32/3	A(20, 23)	--352	A(29, 4)	-(-8192)/9
A(1, 9)	-(-6896)/9	A(7, 21)	-(-457)/3	A(14, 11)	-32/3	A(20, 24)	--160	A(29, 5)	-(-4096)/3
A(1, 11)	-(-136424)/27	A(7, 24)	-619/6	A(14, 12)	-128/3	A(20, 28)	-48	A(29, 6)	-(-8192)/9
A(1, 12)	-(-187040)/27	A(7, 26)	-607/9	A(14, 14)	-(-80)/3	A(21, 4)	-128*sqrt(2)	A(29, 7)	-(-2048)/9
A(1, 13)	-(-27880)/27	A(7, 30)	-(-28)/3	A(14, 18)	-(-65)/6	A(21, 5)	-128*sqrt(2)	A(29, 11)	-6656/9
A(1, 16)	-7750/9	A(8, 1)	-(-400)/9	A(14, 19)	-(-43)/6	A(21, 11)	-32*sqrt(2)	A(29, 12)	-6656/3
A(1, 18)	-98371/27	A(8, 3)	-1456/9	A(14, 20)	-70/3	A(21, 12)	-192*sqrt(2)	A(29, 13)	-6656/3
A(1, 19)	-41027/27	A(8, 4)	-(-992)/9	A(14, 24)	-11/6	A(21, 13)	-32*sqrt(2)	A(29, 14)	-6656/9
A(1, 22)	-(-10501)/27	A(8, 5)	-(-5248)/9	A(14, 25)	-(-25)/3	A(21, 18)	-40*sqrt(2)	A(29, 18)	-(-7936)/9
A(1, 24)	-(-17323)/27	A(8, 6)	-(-4096)/9	A(14, 29)	-1	A(21, 19)	-40*sqrt(2)	A(29, 19)	-(-15872)/9
A(1, 27)	-490/9	A(8, 7)	-(-896)/9	A(15, 3)	-6/3	A(21, 24)	-8*sqrt(2)	A(29, 20)	-(-7936)/9
A(2, 3)	-464/3	A(8, 9)	-1552/9	A(15, 4)	-24	A(22, 4)	-65536/27	A(29, 24)	-4096/9
A(2, 4)	-86464/27	A(8, 11)	-(-464)/3	A(15, 5)	-40	A(22, 5)	-(-114688)/27	A(29, 25)	-4096/3
A(2, 5)	-15296/3	A(8, 12)	-2464/3	A(15, 6)	-52/3	A(22, 6)	-(-40960)/27	A(29, 29)	-(-256)/3
A(2, 6)	-55376/27	A(8, 13)	-3424/3	A(15, 8)	-8/3	A(22, 8)	-(-8192)/27	A(30, 2)	-(-2048)/9
A(2, 8)	-6496/27	A(8, 14)	-3040/9	A(15, 11)	--8	A(22, 11)	-16384/27	A(30, 3)	-(-8192)/9
A(2, 11)	-(-27880)/27	A(8, 16)	-(-761)/3	A(15, 12)	-(-160)/3	A(22, 12)	-53248/9	A(30, 4)	-(-4096)/3
A(2, 12)	-(-187040)/27	A(8, 18)	-(-1831)/9	A(15, 13)	-(-124)/3	A(22, 13)	-38912/9	A(30, 5)	-(-8192)/9
A(2, 13)	-(-136424)/27	A(8, 19)	-938	A(15, 15)	--8	A(22, 15)	-26624/27	A(30, 6)	-(-2048)/9
A(2, 15)	-(-6896)/9	A(8, 20)	-(-1288)/3	A(15, 18)	-47/4	A(22, 18)	-(-32768)/27	A(30, 10)	-6656/9
A(2, 18)	-41027/27	A(8, 22)	-1528/9	A(15, 19)	-349/12	A(22, 21)	-(-10240)/9	A(30, 11)	-6656/3
A(2, 19)	-98371/27	A(8, 24)	-2266/9	A(15, 21)	-17/2	A(22, 24)	-16384/27	A(30, 13)	-6656/9
A(2, 21)	-7750/9	A(8, 25)	-2210/9	A(15, 24)	-(-61)/12	A(22, 26)	-14336/27	A(30, 17)	-(-7936)/9
A(2, 24)	-(-17323)/27	A(8, 27)	-(-406)/9	A(15, 26)	-(-11)/3	A(22, 30)	-(-2048)/27	A(30, 18)	-(-15872)/9
A(2, 26)	-(-10501)/27	A(8, 29)	-(-499)/9	A(15, 30)	-1/2	A(23, 3)	-53248/27	A(30, 19)	-(-7936)/9
A(2, 30)	-490/9	A(8, 31)	-1	A(16, 11)	-(-8)/3	A(23, 4)	-40960/9	A(30, 23)	-4096/9
A(3, 1)	-(-6496)/27	A(9, 2)	-(-896)/9	A(16, 3)	-56/3	A(23, 5)	-69632/27	A(30, 24)	-4096/9
A(3, 3)	-14176/27	A(9, 3)	-(-4096)/9	A(16, 4)	-80/3	A(23, 6)	-(-8192)/27	A(30, 28)	-(-256)/3
A(3, 4)	-(-2944)/27	A(9, 4)	-(-5248)/9	A(16, 5)	-32/3	A(23, 8)	-8192/27	A(31, 2)	-(-2048)/9
A(3, 5)	-(-2944)/27	A(9, 5)	-(-992)/9	A(16, 9)	-32/3	A(23, 11)	-(-48128)/9	A(31, 3)	-(-4096)/9
A(3, 6)	-14176/27	A(9, 6)	-1456/9	A(16, 11)	-40	A(23, 12)	-(-22528)/3	A(31, 4)	-(-2048)/9
A(3, 8)	-(-6496)/27	A(9, 8)	-(-400)/9	A(16, 12)	-32	A(23, 13)	-1024	A(31, 10)	-5632/9
A(3, 9)	-24784/27	A(9, 10)	-3040/9	A(16, 13)	-(-8)/3	A(23, 15)	-(-10240)/9	A(31, 11)	-7168/9
A(3, 11)	-(-5680)/27	A(9, 11)	-3424/3	A(16, 16)	-(-33)/2	A(23, 18)	-14336/3	A(31, 12)	-512/3
A(3, 12)	-12736/9	A(9, 12)	-2464/3	A(16, 18)	-51/2	A(23, 19)	-8192/3	A(31, 17)	-(-1792)/3
A(3, 13)	-(-5680)/27	A(9, 13)	-(-464)/3	A(16, 19)	-19/3	A(23, 21)	-14336/9	A(31, 18)	-(-3584)/9
A(3, 15)	-24784/27	A(9, 15)	-1552/9	A(16, 22)	-73/6	A(23, 24)	-(-37888)/27	A(31, 19)	-(-256)/9
A(3, 16)	-(-11846)/9	A(9, 17)	-(-1288)/3	A(16, 24)	-(-25)/6	A(23, 26)	-(-26624)/27	A(31, 23)	-2048/9
A(3, 18)	-(-26750)/27	A(9, 18)	-938	A(16, 27)	-(-25)/6	A(23, 30)	-2048/9	A(31, 24)	-512/9
A(3, 19)	-(-26750)/27	A(9, 19)	-(-1831)/9	A(16, 31)	-1/2	A(24, 1)	-8192/27	A(31, 28)	-(-256)/9
A(3, 21)	-(-11846)/9	A(9, 21)	-(-761)/3	A(17, 2)	-(-32)/3	A(24, 3)	-(-8192)/27	A(32, 3)	-1024*sqrt(2)/9
A(3, 22)	-22789/27	A(9, 23)	-2210/9	A(17, 3)	-(-160)/3	A(24, 5)	-40960/9	A(32, 11)	-1792*sqrt(2)/9
A(3, 24)	-18262/27	A(9, 24)	-2266/9	A(17, 4)	-(-320)/3	A(24, 6)	-53248/27	A(32, 12)	-256*sqrt(2)/3
A(3, 26)	-22789/27	A(9, 26)	-1528/9	A(17, 5)	-(-320)/3	A(24, 9)	-(-10240)/9	A(32, 18)	-896*sqrt(2)/9
A(3, 27)	-(-5566)/27	A(9, 28)	-(-499)/9	A(17, 6)	-(-160)/3	A(24, 11)	-1024	A(32, 19)	-128*sqrt(2)/9
A(3, 30)	-(-5566)/27	A(9, 30)	-(-406)/9	A(17, 7)	-(-32)/3	A(24, 12)	-(-22528)/3	A(32, 24)	-128*sqrt(2)/9
A(3, 36)	-1	A(9, 33)	-1	A(17, 10)	-112/3	A(24, 13)	-(-48128)/9	A(33, 5)	-1024*sqrt(2)/9
A(4, 1)	-(-400)/9	A(10, 1)	-8/3	A(17, 11)	-448/3	A(24, 16)	-14336/9	A(33, 6)	-1024*sqrt(2)/9
A(4, 3)	-(-3080)/9	A(10, 3)	-52/3	A(17, 12)	-224	A(24, 18)	-8192/3	A(33, 12)	-256*sqrt(2)/3
A(4, 4)	-816	A(10, 4)	-40	A(17, 13)	-448/3	A(24, 19)	-14336/3	A(33, 13)	-1792*sqrt(2)/9
A(4, 5)	-(-4496)/9	A(10, 5)	-24	A(17, 14)	-112/3	A(24, 22)	-(-26624)/27	A(33, 18)	-128*sqrt(2)/9
A(4, 6)	-(-232)/9	A(10, 6)	-4/3	A(17, 17)	--50	A(24, 24)	-(-37888)/27	A(33, 19)	-896*sqrt(2)/9
A(4, 9)	-416/3	A(10, 9)	--8	A(17, 18)	-150	A(24, 27)	-2048/9	A(33, 24)	-128*sqrt(2)/9
A(4, 11)	-7448/9	A(10, 11)	-(-124)/3	A(17, 19)	-150	A(25, 1)	-(-8192)/27	A(34, 3)	-4096
A(4, 12)	-9872/9	A(10, 12)	-(-160)/3	A(17, 20)	-50	A(25, 3)	-(-40960)/27	A(34, 4)	-8192
A(4, 13)	-1472/9	A(10, 13)	--8	A(17, 23)	-95/3	A(25, 4)	-(-114688)/27	A(34, 5)	-4096
A(4, 16)	-(-457)/3	A(10, 16)	-17/2	A(17, 24)	-190/3	A(25, 5)	-(-65536)/27	A(34, 11)	-9216
A(4, 18)	-(-1177)/2	A(10, 18)	-349/12	A(17, 25)	-95/3	A(25, 9)	-26624/27	A(34, 12)	-10240
A(4, 19)	-(-4337)/18	A(10, 19)	-47/4	A(17, 28)	-(-28)/3	A(25, 11)	-38912/9	A(34, 13)	-1024
A(4, 22)	-507/9	A(10, 22)	-(-11)/3	A(17, 29)	-(-28)/3	A(25, 12)	-53248/9	A(34, 18)	-6144
A(4, 24)	-619/6	A(10, 24)	-(-61)/12	A(17, 32)	-1	A(25, 13)	-16384/27	A(34, 19)	-2048
A(4, 27)	-(-28)/3	A(10, 27)	-1/2	A(18, 4)	-32/3	A(25, 16)	-(-10240)/9	A(34, 24)	-1024
A(5, 2)	-(-896)/9	A(11, 2)	-32/3	A(18, 5)	-80/3	A(25, 18)	-(-10240)/3	A(35, 4)	-4096
A(5, 3)	-(-392)/9	A(11, 3)	-8/3	A(18, 6)	-56/3	A(25, 19)	-(-32768)/27	A(35, 5)	-8192
A(5, 4)	-2384/9	A(11, 4)	-(-80)/3	A(18, 8)	-(-8)/3	A(25, 22)	-14336/27	A(35, 6)	-4096
A(5, 5)	-2512/9	A(11, 5)	-(-80)/3	A(18, 11)	-(-8)/3	A(25, 24)	-16384/27	A(35, 11)	-1024
A(5, 6)	-232/9	A(11, 6)	-(-8)/3	A(18, 12)	-32	A(25, 27)	-(-2048)/27	A(35, 12)	-10240
A(5, 10)	-2336/9	A(11, 10)	-(-80)/3	A(18, 13)	-40	A(26, 3)	-(-71680)/27	A(35, 13)	-9216
A(5, 11)	-(-8)/9	A(11, 11)	-8/3	A(18, 15)	-32/3	A(26, 4)	-(-145408)/27	A(35, 18)	-2048
A(5, 12)	-(-3920)/9	A(11, 12)	-128/3	A(18, 16)	-19/3	A(26, 5)	-(-80641)/27	A(35, 19)	-6144
A(5, 13)	-(-976)/9	A(11, 13)	-32/3	A(18, 19)	-51/2	A(26, 6)	-(-2048)/9	A(35, 24)	-1024
A(5, 17)	-(-2104)/9	A(11, 17)	-70/3	A(18, 21)	-(-33)/2	A(26, 11)	-150016/27	A(36, 3)	-4096
A(5, 18)	-123/2	A(11, 18)	-(-43)/6	A(18, 24)	-(-25)/6	A(26, 12)	-183296/27	A(36, 4)	-12288
A(5, 19)	-1981/18	A(11, 19)	-(-65)/6	A(18, 26)	-73/6	A(26, 13)	-31232/27	A(36, 5)	-12288
A(5, 23)	-766/9	A(11, 23)	-(-25)/3	A(18, 30)	-(-25)/6	A(26, 18)	-(-94208)/27	A(36, 6)	-4096
A(5, 24)	-(-307)/18	A(11, 24)	-11/6	A(18, 33)	-1/2	A(26, 19)	-(-40960)/27	A(36, 11)	-11264
A(5, 25)	-123/2	A(11, 28)	-1	A(19, 4)	-256	A(26, 24)	-15872/27	A(36, 13)	-11264
A(5, 26)	-2384/9	A(12, 4)	-40/3	A(19, 5)	-768	A(27, 3)	-(-2048)/9	A(36, 18)	-10240
A(5, 6)	-(-392)/9	A(12, 5)	-40/3	A(19, 7)	-256	A(27, 4)	-(-8064)/27	A(36, 19)	-10240
A(5, 7)	-(-896)/9	A(12, 6)	-4/3	A(19, 11)	--64	A(27, 5)	-(-145408)/27	A(36, 24)	-3072
A(5, 11)	-(-976)/9	A(12, 11)	-(-28)/3	A(19, 12)	-896	A(27, 6)	-(-71680)/27		
A(6, 12)	-(-3920)/9	A(12, 12)	-(-64)/3	A(19, 13)	-1600	A(27, 11)	-31232/27		
A(6, 13)	-(-3920)/9	A(12, 13)	-(-16)/3	A(19, 14)	-768	A(27, 12)	-183296/27		
A(6, 14)	-2336/9	A(12, 19)	-65/12	A(19, 18)	-176	A(27, 13)	-150016/27		
A(6, 18)	-1981/18	A(12, 24)	-(-17)/12	A(19, 19)	-992	A(27, 18)	-(-40960)/27		
A(6, 19)	-123/2	A(13, 3)	-4/3	A(19, 20)	-816	A(27, 19)	-(-94208)/27		
A(6, 20)	-(-2104)/9	A(13, 4)	-40/3	A(19, 24)	-160	A(27, 24)	-15872/27		
A(6, 24)	-(-307)/18	A(13, 5)	-40/3	A(19, 25)	-352	A(28, 5)	-(-2048)/9		
A(6, 25)	-766/9	A(13, 6)	-4	A(19, 29)	-48	A(28, 6)	-(-4096)/9		
A(6, 29)	-(-31)/3	A(13, 11)	-(-16)/3	A(20, 2)	-256	A(28, 7)	-(-2048)/9		
A(7, 3)	-(-232)/9	A(13, 12)	-(-64)/3	A(20, 3)	-768	A(28, 12)	-512/3		
A(7, 4)	-(-4496)/9	A(13, 13)	-(-28)/3	A(20, 4)	-768	A(28, 13)	-7168/9		
A(7, 5)	-816	A(13, 18)	-65/12	A(20, 5)	-256	A(28, 14)	-5632/9		
A(7, 6)	-(-3080)/9	A(13, 19)	-29/4	A(20, 10)	-768	A(28, 18)	-(-256)/9		
A(7, 8)	-(-400)/9	A(13, 24)	-(-17)/12	A(20					

A(1,1)	-1084509/250	A(4,14)	-1629477/50	A(7,49)	-(-157)/10	A(11,3)	-243/10	A(15,8)	-3969/20
A(1,3)	-62720001/1000	A(4,15)	-853821/100	A(8,1)	-(-11907)/25	A(11,4)	-243/10	A(15,10)	-81/5
A(1,4)	-148003999/1000	A(4,16)	-(-53703)/50	A(8,3)	-106191/25	A(11,5)	-1296/5	A(15,13)	-(-81)/10
A(1,5)	-4098681/50	A(4,17)	-6399/25	A(8,4)	-148959/50	A(11,6)	-1296/5	A(15,14)	-162/5
A(1,6)	-(-4220667)/500	A(4,20)	-(-312687)/100	A(8,5)	-(-320679)/25	A(11,7)	-243/10	A(15,15)	-(-5103)/20
A(1,7)	-(-386299)/1000	A(4,22)	-(-10136979)/400	A(8,6)	-(-1257849)/50	A(11,8)	-(-81)/10	A(15,16)	-(-9801)/10
A(1,8)	-1239057/1000	A(4,23)	-(-1055557)/80	A(8,7)	-(-470934)/25	A(11,12)	-(-1134)/5	A(15,17)	-(-12879)/20
A(1,11)	-(-2228553)/125	A(4,24)	-(-919827)/400	A(8,8)	-(-168318)/25	A(11,13)	-(-324)/5	A(15,19)	-(-324)/5
A(1,13)	-(-20589151)/1000	A(4,25)	-(-59229)/400	A(8,9)	-(-25434)/25	A(11,14)	-(-1701)/10	A(15,22)	-(-387)/80
A(1,22)	-2595969/1000	A(4,28)	-252999/100	A(8,11)	-2349	A(11,15)	-(-2673)/5	A(15,23)	-5481/80
A(1,14)	-(-162127899)/500	A(4,30)	-1367451/100	A(8,13)	-(-1121769)/100	A(11,16)	-(-1701)/10	A(15,24)	-1151/16
A(1,15)	-(-85727727)/1000	A(4,31)	-1240929/200	A(8,14)	-654399/100	A(11,17)	-81/5	A(15,25)	-61677/80
A(1,16)	-1337229/125	A(4,32)	-31797/100	A(8,15)	-5075217/100	A(11,21)	-315	A(15,27)	-2061/20
A(1,17)	-(-1224963)/500	A(4,35)	-(-4251)/4	A(8,16)	-5612409/100	A(11,22)	-3141/40	A(15,30)	-(-387)/40
A(1,20)	-29055969/1000	A(4,37)	-(-1284763)/400	A(8,17)	-632448/25	A(11,23)	-9621/40	A(15,31)	-(-1863)/10
A(1,22)	-994544973/4000	A(4,38)	-(-44513)/80	A(8,18)	-112833/25	A(11,24)	-9621/40	A(15,32)	-(-4149)/10
A(1,23)	-184869399/800	A(4,41)	-21409/100	A(8,20)	-(-47279)/100	A(11,25)	-81/40	A(15,34)	-(-1647)/20
A(1,24)	-92584809/4000	A(4,43)	-51869/200	A(8,22)	-1666503/200	A(11,29)	-(-441)/2	A(15,37)	-1339/80
A(1,25)	-5508243/4000	A(4,46)	-(-157)/10	A(8,23)	-(-1395351)/50	A(11,30)	-(-207)/4	A(15,38)	-778/80
A(1,28)	-(-23717313)/1000	A(5,2)	-(-25434)/25	A(8,24)	-(-12094803)/200	A(11,31)	-(-189)/2	A(15,40)	-137/4
A(1,30)	-(-33656553)/250	A(5,3)	-(-56781)/100	A(8,25)	-(-3747951)/100	A(11,32)	-18	A(15,43)	-(-157)/20
A(1,31)	-(-61872849)/1000	A(5,4)	-(-50301)/100	A(8,26)	-(-410733)/50	A(11,36)	-406/5	A(15,45)	-(-137)/20
A(1,32)	-(-6293133)/2000	A(5,5)	-(-107406)/25	A(8,28)	-255753/50	A(11,37)	-599/40	A(15,49)	-1/2
A(1,35)	-401433/40	A(5,6)	-(-108621)/25	A(8,30)	-225423/200	A(11,38)	-65/8	A(16,11)	-(-81)/5
A(1,37)	-23525283/800	A(5,7)	-(-39609)/100	A(8,31)	-1351359/50	A(11,42)	-(-147)/10	A(16,3)	-1053/5
A(1,38)	-22167939/4000	A(5,8)	-12879/100	A(8,32)	-5469291/200	A(11,43)	-(-5)/4	A(16,4)	-4617/10
A(1,41)	-(-2032683)/1000	A(5,12)	-90639/25	A(8,33)	-157311/20	A(11,47)	-1	A(16,5)	-1863/5
A(1,43)	-(-256161)/100	A(5,13)	-37584/25	A(8,35)	-(-76431)/25	A(12,3)	-81/4	A(16,6)	-1053/10
A(1,46)	-14959/100	A(5,14)	-298647/100	A(8,37)	-(-744541)/200	A(12,4)	-(-243)/20	A(16,11)	-81
A(2,3)	-1239057/1000	A(5,15)	-445581/50	A(8,38)	-(-490793)/50	A(12,5)	-(-64)/5	A(16,13)	-(-13527)/20
A(2,4)	-(-386299)/1000	A(5,16)	-283257/100	A(8,39)	-(-420191)/100	A(12,6)	-(-64)/5	A(16,14)	-(-21303)/20
A(2,5)	-(-4220667)/500	A(5,17)	-(-6399)/25	A(8,41)	-48757/50	A(12,7)	-(-243)/20	A(16,15)	-(-10449)/20
A(2,6)	-4098681/50	A(5,21)	-(-10215)/2	A(8,43)	-276743/200	A(12,8)	-81/20	A(16,16)	-(-1053)/20
A(2,7)	-148009539/1000	A(5,22)	-(-687627)/400	A(8,44)	-120393/100	A(12,13)	-(-243)/5	A(16,20)	-(-3357)/20
A(2,8)	-62720001/1000	A(5,23)	-(-1640007)/400	A(8,46)	-(-13389)/100	A(12,14)	-1701/20	A(16,22)	-32409/40
A(2,10)	-1084509/250	A(5,24)	-(-1602207)/400	A(8,48)	-(-15059)/100	A(12,15)	-2673/10	A(16,23)	-4086/5
A(2,13)	-(-1224963)/500	A(5,25)	-(-18927)/400	A(8,53)	-1	A(12,16)	-1701/20	A(16,24)	-7551/40
A(2,14)	-1337229/125	A(5,29)	-72297/20	A(9,2)	-(-25434)/25	A(12,17)	-(-81)/10	A(16,25)	-117/20
A(2,15)	-(-8572727)/1000	A(5,30)	-8379/8	A(9,3)	-(-168318)/25	A(12,22)	-2979/80	A(16,28)	-927/5
A(2,16)	-(-162127899)/500	A(5,31)	-63837/40	A(9,4)	-(-470934)/25	A(12,23)	-(-9621)/80	A(16,30)	-(-17631)/40
A(2,17)	-(-20589151)/1000	A(5,32)	-12231/40	A(9,5)	-(-1257849)/50	A(12,24)	-(-9621)/80	A(16,31)	-(-4671)/20
A(2,19)	-(-2228553)/125	A(5,36)	-(-671717)/50	A(9,6)	-(-320679)/25	A(12,25)	-(-81)/80	A(16,32)	-(-7471)/40
A(2,22)	-5508243/4000	A(5,37)	-(-115031)/400	A(9,7)	-148959/50	A(12,30)	-(-63)/8	A(16,35)	-(-583)/5
A(2,23)	-92584809/4000	A(5,38)	-(-55323)/400	A(9,8)	-106191/25	A(12,31)	-189/4	A(16,37)	-4191/40
A(2,24)	-184869399/800	A(5,42)	-24449/100	A(9,10)	-(-11907)/25	A(12,32)	-9	A(16,38)	-104/5
A(2,25)	-994544973/4000	A(5,43)	-593/25	A(9,12)	-112833/25	A(12,37)	-(-51)/80	A(16,41)	-411/10
A(2,27)	-29055969/1000	A(5,47)	-(-167)/10	A(9,13)	-632448/25	A(12,38)	-(-65)/16	A(16,43)	-(-339)/40
A(2,30)	-(-6293133)/2000	A(6,3)	-12879/100	A(9,14)	-5612409/100	A(12,43)	-1/8	A(16,46)	-(-147)/20
A(2,31)	-61872849/1000	A(6,4)	-(-39609)/100	A(9,15)	-5075217/100	A(13,3)	-81/20	A(16,50)	-1/2
A(2,32)	-(-33656553)/250	A(6,5)	-(-108621)/25	A(9,16)	-654399/100	A(13,4)	-(-243)/20	A(17,2)	-(-324)/5
A(2,34)	-(-23717313)/1000	A(6,6)	-(-107406)/25	A(9,17)	-(-1121769)/100	A(13,5)	-(-64)/5	A(17,3)	-(-2268)/5
A(2,37)	-22167939/4000	A(6,7)	-(-50301)/100	A(9,19)	-2349	A(13,6)	-(-64)/5	A(17,4)	-(-6804)/5
A(2,38)	-25352583/800	A(6,8)	-(-56781)/100	A(9,21)	-(-410733)/50	A(13,7)	-(-243)/20	A(17,5)	-2268
A(2,40)	-401433/40	A(6,9)	-(-25434)/25	A(9,22)	-(-3747951)/100	A(13,8)	-81/4	A(17,6)	-2268
A(2,43)	-(-256161)/100	A(6,13)	-(-6399)/25	A(9,23)	-(-12094803)/200	A(13,13)	-(-81)/10	A(17,7)	-(-6804)/5
A(2,45)	-(-2032683)/1000	A(6,14)	-283257/100	A(9,24)	-(-1395351)/50	A(13,14)	-1701/20	A(17,8)	-(-2268)/5
A(2,49)	-14959/100	A(6,15)	-445581/50	A(9,25)	-1666503/200	A(13,15)	-2673/10	A(17,9)	-(-324)/5
A(3,1)	-(-1084059)/250	A(6,16)	-298647/100	A(9,27)	-(-47279)/100	A(13,16)	-1701/20	A(17,12)	-1458/5
A(3,3)	-7246017/250	A(6,17)	-37584/25	A(9,29)	-157311/20	A(13,17)	-(-243)/5	A(17,13)	-8748/5
A(3,4)	-(-9056367)/500	A(6,18)	-90639/25	A(9,30)	-5469291/200	A(13,22)	-(-81)/80	A(17,14)	-4374
A(3,5)	-(-80108109)/500	A(6,22)	-(-18927)/400	A(9,31)	-1351359/50	A(13,23)	-(-9621)/80	A(17,15)	-5832
A(3,6)	-(-80108109)/500	A(6,23)	-(-1602207)/400	A(9,32)	-225423/200	A(13,24)	-(-9621)/80	A(17,16)	-4374
A(3,7)	-(-9056367)/500	A(6,24)	-(-1640007)/400	A(9,34)	-255753/50	A(13,25)	-2979/80	A(17,17)	-8740/5
A(3,8)	-7246017/250	A(6,25)	-(-687627)/400	A(9,36)	-(-420191)/100	A(13,30)	-9	A(17,18)	-1458/5
A(3,10)	-(-1084059)/250	A(6,26)	-(-10215)/2	A(9,37)	-(-490793)/50	A(13,31)	-189/4	A(17,21)	-(-2709)/5
A(3,11)	-212139/10	A(6,30)	-12231/40	A(9,38)	-(-744541)/200	A(13,32)	-(-63)/8	A(17,22)	-2709
A(3,13)	-(-61872903)/1000	A(6,31)	-63837/40	A(9,40)	-(-76431)/25	A(13,37)	-(-65)/16	A(17,23)	-5418
A(3,14)	-45764919/250	A(6,32)	-8379/8	A(9,42)	-120393/100	A(13,38)	-(-51)/80	A(17,24)	-5418
A(3,15)	-223718679/500	A(6,33)	-72297/20	A(9,43)	-276743/200	A(13,43)	-1/8	A(17,25)	-2709
A(3,16)	-45764919/250	A(6,37)	-(-55323)/400	A(9,45)	-48757/50	A(14,3)	-(-81)/10	A(17,26)	-(-2709)/5
A(3,17)	-(-61872903)/1000	A(6,38)	-(-115031)/400	A(9,47)	-(-15059)/100	A(14,4)	-243/10	A(17,29)	-1071/2
A(3,19)	-212139/10	A(6,39)	-(-671715)/50	A(9,49)	-(-13389)/100	A(14,5)	-1296/5	A(17,30)	-2142
A(3,20)	-(-42597873)/1000	A(6,43)	-593/25	A(9,54)	-1	A(14,6)	-1296/5	A(17,31)	-3213
A(3,22)	-27182547/2000	A(6,44)	-24449/100	A(10,1)	-81/5	A(14,7)	-243/10	A(17,32)	-2142
A(3,23)	-(-681986169)/2000	A(6,48)	-(-167)/10	A(10,3)	-3969/20	A(14,8)	-243/10	A(17,33)	-1071/2
A(3,24)	-(-681986169)/2000	A(7,3)	-(-12879)/100	A(10,4)	-891/20	A(14,9)	-324/5	A(17,36)	-(-3017)/10
A(3,25)	-27182547/2000	A(7,4)	-39609/100	A(10,5)	-243	A(14,13)	-81/5	A(17,37)	-(-9051)/10
A(3,27)	-(-42597873)/1000	A(7,5)	-41391/50	A(10,6)	-(-243)/10	A(14,14)	-(-1701)/10	A(17,38)	-(-9051)/10
A(3,28)	-22436811/500	A(7,6)	-(-40743)/5	A(10,7)	-(-243)/20	A(14,15)	-(-2673)/5	A(17,39)	-(-3017)/10
A(3,30)	-51302269/1000	A(7,7)	-(-1491777)/100	A(10,8)	-81/20	A(14,16)	-(-1701)/10	A(17,42)	-959/10
A(3,31)	-108928341/500	A(7,8)	-(-646623)/100	A(10,11)	-(-324)/5	A(14,17)	-(-324)/5	A(17,43)	-959/5
A(3,32)	-151382269/1000	A(7,10)	-(-11907)/25	A(10,13)	-(-12879)/20	A(14,18)	-(-1134)/5	A(17,44)	-959/10
A(3,34)	-22436811/500	A(7,13)	-6399/25	A(10,14)	-(-9801)/10	A(14,22)	-81/40	A(17,47)	-(-157)/10
A(3,35)	-(-3263511)/125	A(7,14)	-(-53703)/50	A(10,15)	-(-5103)/20	A(14,23)	-9621/40	A(17,48)	-(-157)/10
A(3,37)	-(-81036633)/2000	A(7,15)	-853821/100	A(10,16)	-162/5	A(14,24)	-9621/40	A(17,51)	-1
A(3,38)	-(-81036633)/2000	A(7,16)	-1629477/50	A(10,17)	-(-81)/10	A(14,25)	-3141/40	A(18,5)	-1053/10
A(3,40)	-(-3263511)/125	A(7,17)	-210673/100	A(10,18)	-2061/20	A(14,26)	-315	A(18,6)	-1863/5
A(3,41)	-3974259/500	A(7,19)	-48438/25	A(10,22)	-61677/80	A(14,30)	-18	A(18,7)	-4617/10
A(3,43)	-8433609/1000	A(7,22)	-(-59229)/400	A(10,23)	-11151/16	A(14,31)	-(-189)/2	A(18,8)	-1053/5
A(3,45)	-3974259/500	A(7,23)	-(-91827)/400	A(10,24)	-5481/80	A(14,32)	-(-207)/4	A(18,14)	-(-81)/5
A(3,46)	-(-993043)/1000	A(7,24)	-(-1055557)/80</						

$A(18, 32) = -(-176311)/40$   
 $A(18, 34) = -927/5$   
 $A(18, 37) = -104/5$   
 $A(18, 38) = -4191/40$   
 $A(18, 40) = -(-583)/5$   
 $A(18, 43) = -(-339)/40$   
 $A(18, 45) = -411/10$   
 $A(18, 49) = -(-147)/20$   
 $A(18, 52) = -1/2$   
 $A(19, 5) = -5184$   
 $A(19, 6) = -20736$   
 $A(19, 7) = -31104$   
 $A(19, 8) = -20736$   
 $A(19, 9) = -5184$   
 $A(19, 14) = -2392$   
 $A(19, 15) = -28512$   
 $A(19, 16) = -69984$   
 $A(19, 17) = -64800$   
 $A(19, 18) = -20736$   
 $A(19, 22) = -288$   
 $A(19, 23) = -10224$   
 $A(19, 24) = -52560$   
 $A(19, 25) = -73600$   
 $A(19, 26) = -32976$   
 $A(19, 30) = -1008$   
 $A(19, 31) = -14832$   
 $A(19, 32) = -40176$   
 $A(19, 33) = -26352$   
 $A(19, 37) = -1312$   
 $A(19, 38) = -9392$   
 $A(19, 39) = -10960$   
 $A(19, 43) = -752$   
 $A(19, 44) = -2192$   
 $A(19, 48) = -160$   
 $A(20, 2) = -5184$   
 $A(20, 3) = -20736$   
 $A(20, 4) = -31104$   
 $A(20, 5) = -20736$   
 $A(20, 6) = -5184$   
 $A(20, 12) = -20736$   
 $A(20, 13) = -64800$   
 $A(20, 14) = -69984$   
 $A(20, 15) = -28512$   
 $A(20, 16) = -2592$   
 $A(20, 21) = -32976$   
 $A(20, 22) = -75600$   
 $A(20, 23) = -52560$   
 $A(20, 24) = -10224$   
 $A(20, 25) = -288$   
 $A(20, 29) = -26352$   
 $A(20, 30) = -40176$   
 $A(20, 31) = -14832$   
 $A(20, 32) = -1008$   
 $A(20, 36) = -10960$   
 $A(20, 37) = -9392$   
 $A(20, 38) = -1312$   
 $A(20, 42) = -2192$   
 $A(20, 43) = -752$   
 $A(20, 47) = -160$   
 $A(21, 5) = -2592^2 \cdot \text{Sqrt}(2)$   
 $A(21, 6) = -2592^2 \cdot \text{Sqrt}(2)$   
 $A(21, 14) = -1296^2 \cdot \text{Sqrt}(2)$   
 $A(21, 15) = -5184^2 \cdot \text{Sqrt}(2)$   
 $A(21, 16) = -1296^2 \cdot \text{Sqrt}(2)$   
 $A(21, 22) = -144^2 \cdot \text{Sqrt}(2)$   
 $A(21, 23) = -2088^2 \cdot \text{Sqrt}(2)$   
 $A(21, 24) = -2088^2 \cdot \text{Sqrt}(2)$   
 $A(21, 25) = -144^2 \cdot \text{Sqrt}(2)$   
 $A(21, 30) = -216^2 \cdot \text{Sqrt}(2)$   
 $A(21, 31) = -792^2 \cdot \text{Sqrt}(2)$   
 $A(21, 32) = -216^2 \cdot \text{Sqrt}(2)$   
 $A(21, 37) = -80^2 \cdot \text{Sqrt}(2)$   
 $A(21, 38) = -80^2 \cdot \text{Sqrt}(2)$   
 $A(21, 43) = -8^2 \cdot \text{Sqrt}(2)$   
 $A(22, 5) = -(-8108120)/125$   
 $A(22, 6) = -(-28793376)/125$   
 $A(22, 7) = -(-31702752)/125$   
 $A(22, 8) = -(-10077696)/125$   
 $A(22, 10) = -(-839808)/125$   
 $A(22, 14) = -4094064/125$   
 $A(22, 15) = -40625712/125$   
 $A(22, 16) = -7582764/125$   
 $A(22, 17) = -35796816/125$   
 $A(22, 19) = -139968/5$   
 $A(22, 22) = -(-454896)/125$   
 $A(22, 23) = -(-14679144)/125$   
 $A(22, 24) = -(-59463072)/125$   
 $A(22, 25) = -(-46381896)/125$   
 $A(22, 27) = -(-5705344)/125$   
 $A(22, 30) = -1452168/125$   
 $A(22, 31) = -17180848/125$   
 $A(22, 32) = -26693064/125$   
 $A(22, 34) = -4790016/125$   
 $A(22, 37) = -(-1539648)/125$   
 $A(22, 38) = -(-6572664)/125$   
 $A(22, 40) = -(-2052864)/125$   
 $A(22, 43) = -542376/125$   
 $A(22, 45) = -419904/125$   
 $A(22, 49) = -(-31104)/125$   
 $A(23, 5) = -255879/4$   
 $A(23, 6) = -347733/2$   
 $A(23, 7) = -597051/4$   
 $A(23, 8) = -72171/2$   
 $A(23, 10) = -6561/2$   
 $A(23, 14) = -(-255879)/8$   
 $A(23, 15) = -(-2119203)/8$   
 $A(23, 16) = -(-3112101)/8$   
 $A(23, 17) = -(-1135053)/8$   
 $A(23, 19) = -(-28431)/2$   
 $A(23, 22) = -28431/8$   
 $A(23, 23) = -1579285/16$   
 $A(23, 24) = -329508$   
 $A(23, 25) = -3227283/16$   
 $A(23, 27) = -196101/8$   
 $A(23, 30) = -(-156193)/16$   
 $A(23, 31) = -(-795825)/8$   
 $A(23, 32) = -(-2012769)/16$   
 $A(23, 34) = -(-84807)/4$   
 $A(23, 37) = -18225/2$   
 $A(23, 38) = -522693/16$   
 $A(23, 40) = -9477$   
 $A(23, 43) = -(-44469)/16$   
 $A(23, 45) = -(-8019)/4$   
 $A(23, 49) = -1215/8$   
 $A(24, 4) = -61236$   
 $A(24, 5) = -(-863065)/4$   
 $A(24, 6) = -(-535815)/2$   
 $A(24, 7) = -(-505197)/4$   
 $A(24, 8) = -(-19683)/2$   
 $A(24, 10) = -(-6561)/2$   
 $A(24, 13) = -10206$   
 $A(24, 14) = -1971945/8$   
 $A(24, 15) = -4680909/8$   
 $A(24, 16) = -3483891/8$   
 $A(24, 17) = -570807/8$   
 $A(24, 19) = -15309$   
 $A(24, 22) = -(-280665)/8$   
 $A(24, 23) = -(-5725323)/16$   
 $A(24, 24) = -(-1011609)/2$   
 $A(24, 25) = -(-2422953)/16$   
 $A(24, 27) = -(-231093)/8$   
 $A(24, 30) = -714015/16$   
 $A(24, 31) = -1773981/8$   
 $A(24, 32) = -2139453/16$   
 $A(24, 34) = -223317/8$   
 $A(24, 37) = -(-49653)/2$   
 $A(24, 38) = -(-781893)/16$   
 $A(24, 40) = -(-114453)/8$   
 $A(24, 43) = -81243/16$   
 $A(24, 45) = -(-28431)/8$   
 $A(24, 49) = -(-1215)/4$   
 $A(25, 3) = -5213808/125$   
 $A(25, 4) = -29183328/125$   
 $A(25, 5) = -58471632/125$   
 $A(25, 6) = -49408704/125$   
 $A(25, 7) = -2387168/125$   
 $A(25, 8) = -(-3592323)/125$   
 $A(25, 10) = -39808/125$   
 $A(25, 13) = -(-21893328)/125$   
 $A(25, 14) = -(-96399696)/125$   
 $A(25, 15) = -(-136492128)/125$   
 $A(25, 16) = -(-63090576)/125$   
 $A(25, 17) = -3254256/125$   
 $A(25, 19) = -(-4059072)/125$   
 $A(25, 22) = -36398484/125$   
 $A(25, 23) = -115928496/125$   
 $A(25, 24) = -98365428/125$   
 $A(25, 25) = -10285704/125$   
 $A(25, 27) = -8024832/125$   
 $A(25, 30) = -(-29893212)/125$   
 $A(25, 31) = -(-59819472)/125$   
 $A(25, 32) = -(-18911556)/125$   
 $A(25, 34) = -(-8273664)/125$   
 $A(25, 37) = -12095244/125$   
 $A(25, 38) = -10651824/125$   
 $A(25, 40) = -16624/5$   
 $A(25, 43) = -(-1920996)/125$   
 $A(25, 45) = -(-1353024)/125$   
 $A(25, 49) = -31104/25$   
 $A(26, 1) = -39808/125$   
 $A(26, 3) = -(-3592323)/125$   
 $A(26, 4) = -12387168/125$   
 $A(26, 5) = -49408704/125$   
 $A(26, 6) = -58471632/125$   
 $A(26, 7) = -29183328/125$   
 $A(26, 8) = -5213808/125$   
 $A(26, 11) = -(-4059072)/125$   
 $A(26, 13) = -3254256/125$   
 $A(26, 14) = -(-63090576)/125$   
 $A(26, 15) = -(-136492128)/125$   
 $A(26, 16) = -(-96099696)/125$   
 $A(26, 17) = -(-21893328)/125$   
 $A(26, 20) = -8024832/125$   
 $A(26, 22) = -10285704/125$   
 $A(26, 23) = -98365428/125$   
 $A(26, 24) = -115928496/125$   
 $A(26, 25) = -36398484/125$   
 $A(26, 28) = -(-8273664)/125$   
 $A(26, 30) = -(-18911556)/125$   
 $A(26, 31) = -(-59819472)/125$   
 $A(26, 32) = -(-29893212)/125$   
 $A(26, 35) = -186624/5$   
 $A(26, 37) = -10651824/125$   
 $A(26, 38) = -12095244/125$   
 $A(26, 41) = -(-1353024)/125$   
 $A(26, 43) = -(-1920996)/125$   
 $A(26, 46) = -31104/25$   
 $A(27, 1) = -(-6561)/2$   
 $A(27, 3) = -(-19683)/2$   
 $A(27, 4) = -(-505197)/4$   
 $A(27, 5) = -(-535815)/2$   
 $A(27, 6) = -(-863865)/4$   
 $A(27, 7) = -61236$   
 $A(27, 11) = -15309$   
 $A(27, 13) = -570807/8$   
 $A(27, 14) = -3483891/8$   
 $A(27, 15) = -4680909/8$   
 $A(27, 16) = -1971945/8$   
 $A(27, 17) = -10206$   
 $A(27, 20) = -(-231093)/8$   
 $A(27, 22) = -(-2422953)/16$   
 $A(27, 23) = -(-1011609)/2$   
 $A(27, 24) = -(-5725323)/16$   
 $A(27, 25) = -(-280665)/8$   
 $A(27, 28) = -223317/8$   
 $A(27, 30) = -2139453/16$   
 $A(27, 31) = -1773981/8$   
 $A(27, 32) = -714015/16$   
 $A(27, 35) = -(-114453)/8$   
 $A(27, 37) = -(-781893)/16$   
 $A(27, 38) = -(-49653)/2$   
 $A(27, 41) = -24431/8$   
 $A(27, 43) = -81243/16$   
 $A(27, 46) = -(-1215)/4$   
 $A(28, 1) = -6561/2$   
 $A(28, 3) = -72171/2$   
 $A(28, 4) = -557051/4$   
 $A(28, 5) = -347733/2$   
 $A(28, 6) = -255879/4$   
 $A(28, 11) = -(-28431)/2$   
 $A(28, 13) = -(-1135053)/8$   
 $A(28, 14) = -(-3112101)/8$   
 $A(28, 15) = -(-2119203)/8$   
 $A(28, 16) = -(-255879)/8$   
 $A(28, 20) = -196101/8$   
 $A(28, 22) = -3227283/16$   
 $A(28, 23) = -329508$   
 $A(28, 24) = -1578205/16$   
 $A(28, 25) = -24431/8$   
 $A(28, 28) = -(-84807)/4$   
 $A(28, 30) = -(-2012769)/16$   
 $A(28, 31) = -(-79525)/8$   
 $A(28, 32) = -(-158193)/16$   
 $A(28, 35) = -9477$   
 $A(28, 37) = -522693/16$   
 $A(28, 38) = -18225/2$   
 $A(28, 41) = -(-8019)/4$   
 $A(28, 43) = -(-44469)/16$   
 $A(28, 46) = -1215/8$   
 $A(29, 1) = -(-839808)/125$   
 $A(29, 3) = -(-10077696)/125$   
 $A(29, 4) = -(-31702752)/125$   
 $A(29, 5) = -(-28973376)/125$   
 $A(29, 6) = -(-8188120)/125$   
 $A(29, 11) = -139968/5$   
 $A(29, 13) = -3579616/125$   
 $A(29, 15) = -15739616/125$   
 $A(29, 19) = -75827664/125$   
 $A(29, 21) = -4026572/125$   
 $A(29, 23) = -(-59463072)/125$   
 $A(29, 24) = -(-14679144)/125$   
 $A(29, 25) = -(-454896)/125$   
 $A(29, 28) = -4790016/125$   
 $A(29, 30) = -26693064/125$   
 $A(29, 31) = -17180848/125$   
 $A(29, 32) = -1452168/125$   
 $A(29, 35) = -(-2052864)/125$   
 $A(29, 37) = -(-6572664)/125$   
 $A(29, 38) = -(-1539648)/125$   
 $A(29, 41) = -419904/125$   
 $A(29, 43) = -542376/125$   
 $A(29, 46) = -(-31104)/125$   
 $A(30, 3) = -(-6071121)/125$   
 $A(30, 4) = -(-192456)/5$   
 $A(30, 5) = -9552016/125$   
 $A(30, 6) = -10252656/125$   
 $A(30, 7) = -857304/125$   
 $A(30, 8) = -(-52488)/25$   
 $A(30, 13) = -653194/5$   
 $A(30, 14) = -3242592/125$   
 $A(30, 15) = -(-20435328)/125$   
 $A(30, 16) = -(-6555160)/125$   
 $A(30, 17) = -513216/125$   
 $A(30, 22) = -(-15723558)/125$   
 $A(30, 23) = -3546342/125$   
 $A(30, 24) = -9193662/125$   
 $A(30, 25) = -(-20435328)/125$   
 $A(30, 26) = -24431/8$   
 $A(30, 30) = -263736/5$   
 $A(30, 31) = -18144$   
 $A(30, 32) = -(-1470961)/25$   
 $A(30, 37) = -(-1212246)/125$   
 $A(30, 38) = -212706/125$   
 $A(30, 43) = -83916/125$   
 $A(31, 3) = -(-32805)/8$   
 $A(31, 4) = -(-618921)/8$   
 $A(31, 5) = -148716$   
 $A(31, 6) = -85293$   
 $A(31, 7) = -(-41553)/8$   
 $A(31, 8) = -10935/8$   
 $A(31, 13) = -92583/4$   
 $A(31, 14) = -1661391/8$   
 $A(31, 15) = -955719/4$   
 $A(31, 16) = -410427/8$   
 $A(31, 17) = -(-5103)/2$   
 $A(31, 22) = -(-1285227)/32$   
 $A(31, 23) = -(-5839047)/32$   
 $A(31, 24) = -(-3059127)/32$   
 $A(31, 25) = -(-94527)/32$   
 $A(31, 30) = -227529/8$   
 $A(31, 31) = -57429$   
 $A(31, 32) = -136971/16$   
 $A(31, 37) = -(-256527)/32$   
 $A(31, 38) = -(-163701)/32$   
 $A(32, 2) = -(-1212246)/32$   
 $A(32, 3) = -(-5103)/2$   
 $A(32, 4) = -(-41553)/8$   
 $A(32, 5) = -85293$   
 $A(32, 6) = -148716$   
 $A(32, 7) = -(-618921)/8$   
 $A(32, 8) = -(-32805)/8$   
 $A(32, 13) = -(-5103)/2$   
 $A(32, 14) = -410427/8$   
 $A(32, 15) = -955719/4$   
 $A(32, 16) = -1661391/8$   
 $A(32, 17) = -92583/4$   
 $A(32, 22) = -(-94527)/32$   
 $A(32, 23) = -(-3059127)/32$   
 $A(32, 24) = -(-5839047)/32$   
 $A(32, 25) = -(-1285227)/32$   
 $A(32, 30) = -136971/16$   
 $A(32, 31) = -57429$   
 $A(32, 32) = -227529/8$   
 $A(32, 37) = -(-163701)/32$   
 $A(32, 38) = -(-256527)/32$   
 $A(32, 43) = -11097/16$   
 $A(32, 46) = -(-2012769)/16$   
 $A(32, 48) = -(-52488)/25$   
 $A(32, 51) = -857304/125$   
 $A(33, 5) = -10252656/125$   
 $A(33, 6) = -9552816/125$   
 $A(33, 7) = -(-192456)/5$   
 $A(33, 8) = -(-607112)/125$   
 $A(33, 13) = -513216/125$   
 $A(33, 14) = -(-6555168)/125$   
 $A(33, 15) = -(-20435320)/125$   
 $A(33, 16) = -3242592/125$   
 $A(33, 17) = -653194/5$   
 $A(33, 22) = -180082/125$   
 $A(33, 23) = -9193662/125$   
 $A(33, 24) = -3546342/125$   
 $A(33, 25) = -(-15723558)/125$   
 $A(33, 30) = -(-1470961)/25$   
 $A(33, 31) = -18144$   
 $A(33, 32) = -263736/5$   
 $A(33, 37) = -212706/125$   
 $A(33, 38) = -(-1212246)/125$   
 $A(33, 43) = -83916/125$   
 $A(33, 46) = -(-69984)/25$   
 $A(33, 49) = -(-6572664)/125$   
 $A(33, 51) = -2387168/125$   
 $A(33, 54) = -(-34773376)/125$   
 $A(33, 57) = -(-1539648)/125$   
 $A(33, 60) = -(-1017361)/25$   
 $A(34, 43) = -2592/25$   
 $A(34, 44) = -18792/25$   
 $A(34, 48) = -(-1296)/25$   
 $A(34, 52) = -(-73872)/5$   
 $A(34, 26) = -(-371304)/25$   
 $A(34, 31) = -648$   
 $A(34, 32) = -33696/5$   
 $A(34, 33) = -53784/5$   
 $A(34, 37) = -(-1296)/25$   
 $A(34, 38) = -(-34992)/25$   
 $A(34, 39) = -(-1017361)/25$   
 $A(34, 43) = -2592/25$   
 $A(34, 44) = -18792/25$   
 $A(34, 48) = -(-1296)/25$   
 $A(35, 6) = -4374$   
 $A(35, 7) = -13122$   
 $A(35, 8) = -13122$   
 $A(35, 9) = -4374$   
 $A(35, 15) = -4374$   
 $A(35, 16) = -25515$   
 $A(35, 17) = -37908$   
 $A(35, 18) = -16767$   
 $A(35, 23) = -2673/2$   
 $A(35, 24) = -32805/2$   
 $A(35, 25) = -80919/2$   
 $A(35, 26) = -50787/2$   
 $A(35, 30) = -(-243)/2$   
 $A(35, 31) = -(-16605)/4$   
 $A(35, 32) = -19602$   
 $A(35, 33) = -(-76069)/4$   
 $A(35, 37) = -1377/4$   
 $A(35, 38) = -8505/2$   
 $A(35, 39) = -30213/4$   
 $A(35, 43) = -324$   
 $A(35, 44) = -(-5751)/4$   
 $A(35, 48) = -405/4$

Figure A.3<sub>2</sub> : Matrix  $[A^*]_{55 \times 55}$

$A(36, 4) = -4374$   
 $A(36, 5) = -21870$   
 $A(36, 6) = -43740$   
 $A(36, 7) = -43740$   
 $A(36, 8) = -43740$   
 $A(36, 9) = -4374$   
 $A(36, 13) = -729$   
 $A(36, 14) = -21141$   
 $A(36, 15) = -77274$   
 $A(36, 16) = -112266$   
 $A(36, 17) = -73629$   
 $A(36, 18) = -18225$   
 $A(36, 22) = -2916$   
 $A(36, 23) = -78489/2$   
 $A(36, 24) = -200475/2$   
 $A(36, 25) = -188811/2$   
 $A(36, 26) = -60993/2$   
 $A(36, 30) = (-18387)/4$   
 $A(36, 31) = (-140697)/4$   
 $A(36, 32) = (-226233)/4$   
 $A(36, 33) = (-103923)/4$   
 $A(36, 37) = -3564$   
 $A(36, 38) = -30537/2$   
 $A(36, 39) = -23409/2$   
 $A(36, 43) = (-5427)/4$   
 $A(36, 44) = (-10267)/4$   
 $A(36, 48) = -405/2$   
 $A(37, 3) = (-69984)/25$   
 $A(37, 4) = (-419904)/25$   
 $A(37, 5) = (-209952)/5$   
 $A(37, 6) = (-279936)/5$   
 $A(37, 7) = (-209952)/5$   
 $A(37, 8) = (-419904)/25$   
 $A(37, 9) = (-69984)/25$   
 $A(37, 13) = -303264/25$   
 $A(37, 14) = -303264/5$   
 $A(37, 15) = -606528/5$   
 $A(37, 16) = -606528/5$   
 $A(37, 17) = -303264/5$   
 $A(37, 18) = -303264/25$   
 $A(37, 22) = -21384$   
 $A(37, 23) = -85536$   
 $A(37, 24) = -128304$   
 $A(37, 25) = -85536$   
 $A(37, 26) = -21384$   
 $A(37, 30) = -97840/5$   
 $A(37, 31) = -293544/5$   
 $A(37, 32) = -293544/5$   
 $A(37, 33) = -97848/5$   
 $A(37, 37) = (-244296)/25$   
 $A(37, 38) = (-488592)/25$   
 $A(37, 39) = (-244296)/25$   
 $A(37, 43) = -62856/25$   
 $A(37, 44) = -62856/25$   
 $A(37, 48) = (-1296)/5$   
 $A(38, 2) = (-69984)/25$   
 $A(38, 3) = (-419904)/25$   
 $A(38, 4) = (-209952)/5$   
 $A(38, 5) = (-279936)/5$   
 $A(38, 6) = (-209952)/5$   
 $A(38, 7) = (-419904)/25$   
 $A(38, 8) = (-69984)/25$   
 $A(38, 12) = -303264/25$   
 $A(38, 13) = -303264/5$   
 $A(38, 14) = -606528/5$   
 $A(38, 15) = -606528/5$   
 $A(38, 16) = -303264/5$   
 $A(38, 17) = -303264/25$   
 $A(38, 21) = -21384$   
 $A(38, 22) = -85536$   
 $A(38, 23) = -128304$   
 $A(38, 24) = -85536$   
 $A(38, 25) = -21384$   
 $A(38, 29) = -97840/5$   
 $A(38, 30) = -293544/5$   
 $A(38, 31) = -293544/5$   
 $A(38, 32) = -97848/5$   
 $A(38, 36) = (-244296)/25$   
 $A(38, 37) = (-488592)/25$   
 $A(38, 38) = (-244296)/25$   
 $A(38, 42) = -62856/25$   
 $A(38, 43) = -62856/25$   
 $A(38, 47) = (-1296)/5$   
 $A(39, 2) = -4374$   
 $A(39, 3) = -21870$   
 $A(39, 4) = -43740$   
 $A(39, 5) = -43740$   
 $A(39, 6) = -21870$   
 $A(39, 7) = -4374$   
 $A(39, 12) = -18225$   
 $A(39, 13) = -73629$   
 $A(39, 14) = -112266$   
 $A(39, 15) = -77274$   
 $A(39, 16) = -21141$   
 $A(39, 17) = -729$   
 $A(39, 21) = -60993/2$   
 $A(39, 22) = -188811/2$   
 $A(39, 23) = -200475/2$   
 $A(39, 24) = -78489/2$   
 $A(39, 25) = -2916$   
 $A(39, 29) = (-103923)/4$   
 $A(39, 30) = (-226233)/4$

$A(39, 31) = (-140697)/4$   
 $A(39, 32) = (-18387)/4$   
 $A(39, 36) = -23409/2$   
 $A(39, 37) = -30537/2$   
 $A(39, 38) = -3564$   
 $A(39, 42) = (-10287)/4$   
 $A(39, 43) = (-5427)/4$   
 $A(39, 47) = -405/2$   
 $A(40, 2) = -4374$   
 $A(40, 3) = -13122$   
 $A(40, 4) = -13122$   
 $A(40, 5) = -4374$   
 $A(40, 12) = -16767$   
 $A(40, 13) = -37908$   
 $A(40, 14) = -23515$   
 $A(40, 15) = -4374$   
 $A(40, 21) = -50787/2$   
 $A(40, 22) = -10319/2$   
 $A(40, 23) = -32805/2$   
 $A(40, 24) = -2673/2$   
 $A(40, 29) = (-76869)/4$   
 $A(40, 30) = -19602$   
 $A(40, 31) = (-16605)/4$   
 $A(40, 32) = (-243)/2$   
 $A(40, 36) = -30213/4$   
 $A(40, 37) = -8505/2$   
 $A(40, 38) = -1377/4$   
 $A(40, 42) = (-5751)/4$   
 $A(40, 43) = -324$   
 $A(40, 47) = -405/4$   
 $A(41, 2) = (-69984)/25$   
 $A(41, 3) = (-139968)/25$   
 $A(41, 4) = (-69984)/25$   
 $A(41, 12) = -256608/25$   
 $A(41, 13) = -373248/25$   
 $A(41, 14) = -23328/5$   
 $A(41, 21) = (-371304)/25$   
 $A(41, 22) = (-73872)/5$   
 $A(41, 23) = (-13608)/5$   
 $A(41, 29) = -53784/5$   
 $A(41, 30) = -33696/5$   
 $A(41, 31) = -648$   
 $A(41, 36) = (-101736)/25$   
 $A(41, 37) = (-34992)/25$   
 $A(41, 38) = (-1296)/25$   
 $A(41, 42) = -18792/25$   
 $A(41, 43) = -2592/25$   
 $A(41, 47) = (-1296)/25$   
 $A(42, 3) = -34992*\text{Sqrt}(2)/25$   
 $A(42, 4) = -34992*\text{Sqrt}(2)/25$   
 $A(42, 13) = -93312*\text{Sqrt}(2)/25$   
 $A(42, 14) = -11664*\text{Sqrt}(2)/25$   
 $A(42, 22) = -18468*\text{Sqrt}(2)/5$   
 $A(42, 23) = -6804*\text{Sqrt}(2)/5$   
 $A(42, 30) = -8424*\text{Sqrt}(2)/5$   
 $A(42, 31) = -324*\text{Sqrt}(2)$   
 $A(42, 37) = -8748*\text{Sqrt}(2)/25$   
 $A(42, 38) = -648*\text{Sqrt}(2)/25$   
 $A(42, 43) = -2519424$   
 $A(42, 44) = -648*\text{Sqrt}(2)/25$   
 $A(42, 45) = -2167*\text{Sqrt}(2)$   
 $A(42, 5) = -2187*\text{Sqrt}(2)$   
 $A(43, 13) = -729*\text{Sqrt}(2)/2$   
 $A(43, 14) = -9477*\text{Sqrt}(2)/2$   
 $A(43, 15) = -2187*\text{Sqrt}(2)$   
 $A(43, 22) = -729*\text{Sqrt}(2)$   
 $A(43, 23) = -12879*\text{Sqrt}(2)/4$   
 $A(43, 24) = -2673*\text{Sqrt}(2)/4$   
 $A(43, 30) = -807*\text{Sqrt}(2)/8$   
 $A(43, 31) = -6723*\text{Sqrt}(2)/8$   
 $A(43, 32) = -243*\text{Sqrt}(2)/4$   
 $A(43, 37) = -243*\text{Sqrt}(2)/2$   
 $A(43, 38) = -567*\text{Sqrt}(2)/8$   
 $A(43, 43) = -81*\text{Sqrt}(2)/8$   
 $A(44, 6) = -2187*\text{Sqrt}(2)$   
 $A(44, 7) = -2187*\text{Sqrt}(2)$   
 $A(44, 15) = -2187*\text{Sqrt}(2)$   
 $A(44, 16) = -9477*\text{Sqrt}(2)/2$   
 $A(44, 17) = -729*\text{Sqrt}(2)/2$   
 $A(44, 23) = -2673*\text{Sqrt}(2)/4$   
 $A(44, 24) = -12879*\text{Sqrt}(2)/4$   
 $A(44, 25) = -729*\text{Sqrt}(2)$   
 $A(44, 30) = -243*\text{Sqrt}(2)/4$   
 $A(44, 31) = -6723*\text{Sqrt}(2)/8$   
 $A(44, 32) = -3807*\text{Sqrt}(2)/8$   
 $A(44, 37) = -567*\text{Sqrt}(2)/8$   
 $A(44, 38) = -243*\text{Sqrt}(2)/2$   
 $A(44, 43) = -81*\text{Sqrt}(2)/8$   
 $A(45, 7) = -34992*\text{Sqrt}(2)/25$   
 $A(45, 8) = -34992*\text{Sqrt}(2)/25$   
 $A(45, 16) = -11664*\text{Sqrt}(2)/5$   
 $A(45, 17) = -93312*\text{Sqrt}(2)/25$   
 $A(45, 24) = -6804*\text{Sqrt}(2)/5$   
 $A(45, 25) = -18468*\text{Sqrt}(2)/5$   
 $A(45, 31) = -324*\text{Sqrt}(2)$   
 $A(45, 32) = -8424*\text{Sqrt}(2)/5$   
 $A(45, 37) = -648*\text{Sqrt}(2)/25$   
 $A(45, 38) = -8748*\text{Sqrt}(2)/25$   
 $A(45, 43) = -648*\text{Sqrt}(2)/25$   
 $A(46, 6) = -104976$   
 $A(46, 7) = -209952$   
 $A(46, 8) = -104976$   
 $A(46, 15) = -104976$

$A(46, 16) = -419904$   
 $A(46, 17) = -314928$   
 $A(46, 23) = -32076$   
 $A(46, 24) = -274104$   
 $A(46, 25) = -347004$   
 $A(46, 30) = -2916$   
 $A(46, 31) = -69984$   
 $A(46, 32) = -172044$   
 $A(46, 37) = -5832$   
 $A(46, 38) = -37908$   
 $A(46, 43) = -2916$   
 $A(46, 5) = -139968$   
 $A(47, 6) = -419904$   
 $A(47, 7) = -419904$   
 $A(47, 8) = -139968$   
 $A(47, 14) = -69984$   
 $A(47, 15) = -606528$   
 $A(47, 22) = -1003104$   
 $A(47, 23) = -320680$   
 $A(47, 24) = -466560$   
 $A(47, 25) = -7776$   
 $A(47, 26) = -221616$   
 $A(47, 27) = -793152$   
 $A(47, 28) = -79056$   
 $A(47, 29) = -579312$   
 $A(47, 30) = -22032$   
 $A(47, 31) = -230680$   
 $A(47, 32) = -325296$   
 $A(47, 33) = -20736$   
 $A(47, 34) = -111456$   
 $A(47, 35) = -139968$   
 $A(47, 36) = -279936$   
 $A(47, 37) = -139968$   
 $A(47, 38) = -2916$   
 $A(47, 39) = -37908$   
 $A(47, 40) = -5832$   
 $A(47, 41) = -139968$   
 $A(47, 42) = -2916$   
 $A(47, 43) = -6480$   
 $A(47, 44) = -139968$   
 $A(47, 45) = -558972$   
 $A(47, 46) = -839808$   
 $A(47, 47) = -558972$   
 $A(47, 48) = -139968$   
 $A(47, 49) = -81648$   
 $A(47, 50) = -882576$   
 $A(47, 51) = -1520208$   
 $A(47, 52) = -719280$   
 $A(47, 53) = -106272$   
 $A(47, 54) = -578016$   
 $A(47, 55) = -471744$   
 $A(47, 56) = -60912$   
 $A(47, 57) = -130672$   
 $A(47, 58) = -12960$   
 $A(47, 59) = -104976$   
 $A(47, 60) = -524880$   
 $A(47, 61) = -1049760$   
 $A(47, 62) = -1049760$   
 $A(47, 63) = -524880$   
 $A(47, 64) = -104976$   
 $A(47, 65) = -167616$   
 $A(47, 66) = -611932$   
 $A(47, 67) = -1985796$   
 $A(47, 68) = -1985796$   
 $A(47, 69) = -661932$   
 $A(47, 70) = -1679616$   
 $A(47, 71) = -419904$   
 $A(47, 72) = -661932$   
 $A(47, 73) = -1985796$   
 $A(47, 74) = -1985796$   
 $A(47, 75) = -661932$   
 $A(47, 76) = -1679616$   
 $A(47, 77) = -419904$   
 $A(47, 78) = -661932$   
 $A(47, 79) = -1985796$   
 $A(47, 80) = -1985796$   
 $A(47, 81) = -661932$   
 $A(47, 82) = -1679616$   
 $A(47, 83) = -419904$   
 $A(47, 84) = -661932$   
 $A(47, 85) = -1985796$   
 $A(47, 86) = -1985796$   
 $A(47, 87) = -661932$   
 $A(47, 88) = -1679616$   
 $A(47, 89) = -419904$   
 $A(47, 90) = -661932$   
 $A(47, 91) = -1985796$   
 $A(47, 92) = -1985796$   
 $A(47, 93) = -661932$   
 $A(47, 94) = -1679616$   
 $A(47, 95) = -419904$   
 $A(47, 96) = -661932$   
 $A(47, 97) = -1985796$   
 $A(47, 98) = -1985796$   
 $A(47, 99) = -661932$   
 $A(47, 100) = -1679616$   
 $A(47, 101) = -419904$   
 $A(47, 102) = -661932$   
 $A(47, 103) = -1985796$   
 $A(47, 104) = -1985796$   
 $A(47, 105) = -661932$   
 $A(47, 106) = -1679616$   
 $A(47, 107) = -419904$   
 $A(47, 108) = -661932$   
 $A(47, 109) = -1985796$   
 $A(47, 110) = -1985796$   
 $A(47, 111) = -661932$   
 $A(47, 112) = -1679616$   
 $A(47, 113) = -419904$   
 $A(47, 114) = -661932$   
 $A(47, 115) = -1985796$   
 $A(47, 116) = -1985796$   
 $A(47, 117) = -661932$   
 $A(47, 118) = -1679616$   
 $A(47, 119) = -419904$   
 $A(47, 120) = -661932$   
 $A(47, 121) = -1985796$   
 $A(47, 122) = -1985796$   
 $A(47, 123) = -661932$   
 $A(47, 124) = -1679616$   
 $A(47, 125) = -419904$   
 $A(47, 126) = -661932$   
 $A(47, 127) = -1985796$   
 $A(47, 128) = -1985796$   
 $A(47, 129) = -661932$   
 $A(47, 130) = -1679616$   
 $A(47, 131) = -419904$   
 $A(47, 132) = -661932$   
 $A(47, 133) = -1985796$   
 $A(47, 134) = -1985796$   
 $A(47, 135) = -661932$   
 $A(47, 136) = -1679616$   
 $A(47, 137) = -419904$   
 $A(47, 138) = -661932$   
 $A(47, 139) = -1985796$   
 $A(47, 140) = -1985796$   
 $A(47, 141) = -661932$   
 $A(47, 142) = -1679616$   
 $A(47, 143) = -419904$   
 $A(47, 144) = -661932$   
 $A(47, 145) = -1985796$   
 $A(47, 146) = -1985796$   
 $A(47, 147) = -661932$   
 $A(47, 148) = -1679616$   
 $A(47, 149) = -419904$   
 $A(47, 150) = -661932$   
 $A(47, 151) = -1985796$   
 $A(47, 152) = -1985796$   
 $A(47, 153) = -661932$   
 $A(47, 154) = -1679616$   
 $A(47, 155) = -419904$   
 $A(47, 156) = -661932$   
 $A(47, 157) = -1985796$   
 $A(47, 158) = -1985796$   
 $A(47, 159) = -661932$   
 $A(47, 160) = -1679616$   
 $A(47, 161) = -419904$   
 $A(47, 162) = -661932$   
 $A(47, 163) = -1985796$   
 $A(47, 164) = -1985796$   
 $A(47, 165) = -661932$   
 $A(47, 166) = -1679616$   
 $A(47, 167) = -419904$   
 $A(47, 168) = -661932$   
 $A(47, 169) = -1985796$   
 $A(47, 170) = -1985796$   
 $A(47, 171) = -661932$   
 $A(47, 172) = -1679616$   
 $A(47, 173) = -419904$   
 $A(47, 174) = -661932$   
 $A(47, 175) = -1985796$   
 $A(47, 176) = -1985796$   
 $A(47, 177) = -661932$   
 $A(47, 178) = -1679616$   
 $A(47, 179) = -419904$   
 $A(47, 180) = -661932$   
 $A(47, 181) = -1985796$   
 $A(47, 182) = -1985796$   
 $A(47, 183) = -661932$   
 $A(47, 184) = -1679616$   
 $A(47, 185) = -419904$   
 $A(47, 186) = -661932$   
 $A(47, 187) = -1985796$   
 $A(47, 188) = -1985796$   
 $A(47, 189) = -661932$   
 $A(47, 190) = -1679616$   
 $A(47, 191) = -419904$   
 $A(47, 192) = -661932$   
 $A(47, 193) = -1985796$   
 $A(47, 194) = -1985796$   
 $A(47, 195) = -661932$   
 $A(47, 196) = -1679616$   
 $A(47, 197) = -419904$   
 $A(47, 198) = -661932$   
 $A(47, 199) = -1985796$   
 $A(47, 200) = -1985796$   
 $A(47, 201) = -661932$   
 $A(47, 202) = -1679616$   
 $A(47, 203) = -419904$   
 $A(47, 204) = -661932$   
 $A(47, 205) = -1985796$   
 $A(47, 206) = -1985796$   
 $A(47, 207) = -661932$   
 $A(47, 208) = -1679616$   
 $A(47, 209) = -419904$   
 $A(47, 210) = -661932$   
 $A(47, 211) = -1985796$   
 $A(47, 212) = -1985796$   
 $A(47, 213) = -661932$   
 $A(47, 214) = -1679616$   
 $A(47, 215) = -419904$   
 $A(47, 216) = -661932$   
 $A(47, 217) = -1985796$   
 $A(47, 218) = -1985796$   
 $A(47, 219) = -661932$   
 $A(47, 220) = -1679616$   
 $A(47, 221) = -419904$   
 $A(47, 222) = -661932$   
 $A(47, 223) = -1985796$   
 $A(47, 224) = -1985796$   
 $A(47, 225) = -661932$   
 $A(47, 226) = -1679616$   
 $A(47, 227) = -419904$   
 $A(47, 228) = -661932$   
 $A(47, 229) = -1985796$   
 $A(47, 230) = -1985796$   
 $A(47, 231) = -661932$   
 $A(47, 232) = -1679616$   
 $A(47, 233) = -419904$   
 $A(47, 234) = -661932$   
 $A(47, 235) = -1985796$   
 $A(47, 236) = -1985796$   
 $A(47, 237) = -661932$   
 $A(47, 238) = -1679616$   
 $A(47, 239) = -419904$   
 $A(47, 240) = -661932$   
 $A(47, 241) = -1985796$   
 $A(47, 242) = -1985796$   
 $A(47, 243) = -661932$   
 $A(47, 244) = -1679616$   
 $A(47, 245) = -419904$   
 $A(47, 246) = -661932$   
 $A(47, 247) = -1985796$   
 $A(47, 248) = -1985796$   
 $A(47, 249) = -661932$   
 $A(47, 250) = -$

**ISSN 0249-6399**