

Explicit factors of some iterated resultants and discriminants

Laurent Busé, Bernard Mourrain

► **To cite this version:**

Laurent Busé, Bernard Mourrain. Explicit factors of some iterated resultants and discriminants. Mathematics of Computation, American Mathematical Society, 2009, 78 (265), pp.345–386. inria-00119287v2

HAL Id: inria-00119287

<https://hal.inria.fr/inria-00119287v2>

Submitted on 15 Oct 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXPLICIT FACTORS OF SOME ITERATED RESULTANTS AND DISCRIMINANTS

LAURENT BUSÉ AND BERNARD MOURRAIN

dedicated to Professor Jean-Pierre Jouanolou

ABSTRACT. In this paper, the result of applying iterative univariate resultant constructions to multivariate polynomials is analyzed. We consider the input polynomials as generic polynomials of a given degree and exhibit explicit decompositions into irreducible factors of several constructions involving two times iterated univariate resultants and discriminants over the integer universal ring of coefficients of the entry polynomials. Cases involving from two to four generic polynomials and resultants or discriminants in one of their variables are treated. The decompositions into irreducible factors we get are obtained by exploiting fundamental properties of the univariate resultants and discriminants and induction on the degree of the polynomials. As a consequence, each irreducible factor can be separately and explicitly computed in terms of a certain multivariate resultant. With this approach, we also obtain as direct corollaries some results conjectured by Collins [9] and McCallum [26, 27] which correspond to the case of polynomials whose coefficients are themselves generic polynomials in other variables. Finally, a geometric interpretation of the algebraic factorization of the iterated discriminant of a single polynomial is detailed.

1. INTRODUCTION

Resultants provide an essential tool in constructive algebra and in equation solving, for projecting the solution of a polynomial system into a space of smaller dimension. In the univariate case, a well-known construction due to J.J. Sylvester (1840) consists in eliminating the monomials $1, X, \dots, X^{m+n-1}$ in the multiples

$$(X^i P(X))_{0 \leq i \leq n-1}, (X^j Q(X))_{0 \leq j \leq m-1},$$

of two given polynomials P, Q of degree m and n , and in taking the determinant of the corresponding $(m+n) \times (m+n)$ matrix. Though the first resultant construction appeared probably in the work of E. Bézout [4] (see also Euler's work), and although contemporary to related works (Jacobi 1835, Richelot 1840, Cauchy 1840, . . .), this method remains well-known as Sylvester's resultant. It is nowadays a fundamental tool used in effective algebra to eliminate a variable between two polynomials.

It is a natural belief that equipped with such a tool which eliminates one variable at a time, one can iteratively eliminate several variables. This approach was actually exploited, for instance in [30], to deduce theoretical results (such as the existence of eliminant polynomials) in several variables. However, if we are interested in structural results as well as practical computations or complexity issues, this

Date: October 15, 2007.

This work was first presented at the conference in honor of Jean-Pierre Jouanolou, held at Luminy, Marseille, 15-19th May 2006.

approach is far from being optimal. This explains the study and development of different types of multivariate resultants, including projective [23, 18], anisotropic [18, 19], toric [14, 13, 12], residual [19, 7, 5], determinantal [8] resultants.

Nevertheless, in some algorithms, such as in Cylindrical Algebraic Decomposition (CAD), an induction is applied on the dimension of the problems and iterated univariate resultants and subresultants are used at many steps of the algorithm [9, 3]. In its original paper on quantifier elimination for real closed fields by cylindrical algebraic decomposition [9], Collins used these iterated resultants in a geometric context and observed certain intriguing factorizations that “suggest some theorems” [9, p.178]. The same year, 1975, Van der Waerden responded to these observations in a handwritten letter where he gave some intuitive hints for some of the phenomena noticed by Collins. More recently, McCallum [26, 27] proved rigorously that certain iterated resultants have some irreducible factors, but he only showed the existence and did not give a way to compute them independently. As pointed out by Jean-Pierre Jouanolou, in 1868 and 1869 Olaus Henrici published two outstanding papers [16, 17] addressing the decomposition of the discriminant of a discriminant. In particular, he gave the expected factorization of such a repeated discriminant. However, he did not prove the irreducibility of these factors which is a more difficult task. One of our goals in this paper is to give the decomposition into irreducible factors of these iterated resultant computations.

From a geometric point of view, we are interested in the solutions of equations depending on some parameters and by analyzing what happens “above”, when we move these parameters. The number of solutions might change if we cross the set of points where a vertical line is tangent to the solution set (that is to the polar variety in the vertical direction). This polar variety projected in one dimension less, might have singularities where the number of solutions changes effectively or which are only due to the superposition of distinct points of this polar variety. These critical points of the polar variety of an algebraic surface $f(x, y, z) = 0$ in \mathbb{R}^3 are used effectively in algorithms for computing the topology of the surface (see e.g. [28]). The projection of the polar curve (say on the (x, y) -plane) is obtained by a discriminant computation and the critical points of this projected curve are again computed by a discriminant. These values are then used to analyze where the topology of a plane section is changing in order to deduce the topology of the whole surface $f(x, y, z) = 0$. Similar projection tools are also implicitly used in higher dimension for the triangulation of hypersurfaces (see e.g. [15]), which leads in the algebraic context to univariate resultant computation (see e.g. [10, 3]).

Another of our objectives is to show how these critical points corresponding generically to folds, double folds or pleats of the surface can be related to explicit factors in iterated resultant constructions. The main results of this paper are complete explicit factorizations of two times iterated univariate resultants and discriminants of generic polynomials of given degree. We actually give the decomposition of these iterated resultants over the integer universal ring of coefficients of polynomials of given degree. Such a formulation has the advantage to allow the pre-computation of a given factor. It has direct applications to the topological computation of algebraic surfaces, which was our starting point.

Our approach is based on the study of these iterated resultants in generic situations. Most of the interesting formulas are obtained by suitable specialization of

this case. These specializations are performed using the formalism of the multivariate resultant as it has been originally introduced and deeply developed by Jouanolou to whom this paper is dedicated in recognition of its outstanding contributions to resultant theory (e.g. [18, 19, 20]). It should be noticed that this approach can be pushed further to study more particular situations (corresponding to other type of multivariate resultants) that we did not consider in this paper, but which could be interesting for specific applications.

The paper is structured as follows. In the next section, we recall the definitions and main properties of resultants and discriminants that we will use. In section 3, we consider the computation of iterated resultants of 4 and then 3 polynomials. In section 4, we analyze the iterated computation of resultants of discriminants, first the resultant of the discriminants of two distinct polynomials P_1, P_2 and next the resultant of the discriminant of P_1 and the resultant of P_1, P_2 . We here extend the previous work [26] and prove some properties conjectured by Collins and McCallum. In section 5, we study the discriminant of a resultant, simplifying the proof and also extending some results of [26, 27]. These developments are used in cascade to provide, in section 6, the complete factorization into irreducible components of a discriminant of a discriminant for a generic polynomial, as conjectured in [27]. These new results have direct corollaries for polynomials whose coefficients are themselves generic polynomials in other parameter variables, which we provide.

2. BACKGROUND MATERIAL AND NOTATION

In this section we give the notation and quickly present the tools, as resultants and discriminants, that we will use all along this paper.

2.1. Resultants.

2.1.1. *The univariate case.* Let \mathbb{S} be a commutative ring (with unity) and consider the two polynomials in $\mathbb{S}[X]$

$$f(X) := a_m X^m + a_{m-1} X^{m-1} + \dots + a_0, \quad g(X) := b_n X^n + b_{n-1} X^{n-1} + \dots + b_0,$$

where $a_i, b_j \in \mathbb{S}$, and m and n are both positive integers. Their resultant (in degrees m, n)¹ that we will denote $\text{Res}_X(f, g)$, is defined as the determinant of the well-known Sylvester matrix

$$\begin{pmatrix} a_m & 0 & \dots & 0 & b_n & 0 & 0 \\ a_{m-1} & a_m & & \vdots & b_{n-1} & \ddots & 0 \\ \vdots & & \ddots & 0 & \vdots & & b_n \\ a_0 & & & a_m & b_1 & & b_{n-1} \\ 0 & a_m & & a_{m-1} & b_0 & & \vdots \\ \vdots & & \ddots & \vdots & 0 & \ddots & b_1 \\ 0 & \dots & 0 & a_0 & 0 & 0 & b_0 \end{pmatrix}.$$

Remark 2.1. We emphasize that the notation $\text{Res}_X(f, g)$ denotes the resultant of f and g with respect to the variable X as polynomials of their *expected* degree, which is here m and n respectively. It is important to keep this in mind since the closed

¹Notice that the dependence on the degrees m, n can be avoided if one considers homogeneous polynomials.

formulas we will prove in this paper are using this convention; see Remark 3.5 for an illustration.

This “eliminant polynomial” has a long history and many known properties. One can learn about it in many places in the literature, for instance [11, 22]; see also [14, chapter 12] and [1] for a detailed exposition. In the sequel we will especially use the following properties:

- $\text{Res}_X(f, g)$ belongs to the ideal $(f, g) \subset \mathbb{S}[X]$,
- $\text{Res}_X(f, g)$ is homogeneous of degree n , resp. m , in the a_i 's, resp. in the b_j 's,
- $\text{Res}_X(f, g)$ is homogeneous of degree mn if we set $\deg(a_i) := m - i$ and $\deg(b_j) := n - j$ for all $i = 0, \dots, m$ and $j = 0, \dots, n$.

We also recall the definition of the principal subresultant of f and g that we will use later on. It is defined as the determinant of the above Sylvester matrix where the two last lines and columns number n and $n + m$ are erased; more precisely

$$\text{SRes}_X^{(1)}(f, g) := \begin{vmatrix} a_m & 0 & \cdots & 0 & 0 & b_n & 0 & 0 & 0 \\ a_{m-1} & a_m & & \vdots & \vdots & b_{n-1} & \ddots & 0 & 0 \\ \vdots & a_{m-1} & \ddots & 0 & 0 & \vdots & & b_n & 0 \\ a_2 & & \ddots & a_m & 0 & \vdots & & b_{n-1} & b_n \\ a_1 & a_2 & & a_{m-1} & a_m & b_2 & & \vdots & b_{n-1} \\ a_0 & a_1 & \ddots & \vdots & a_{m-1} & b_1 & & \vdots & \vdots \\ 0 & \ddots & \ddots & a_2 & \vdots & b_0 & \ddots & b_2 & \vdots \\ \vdots & & \ddots & a_1 & a_2 & 0 & \ddots & b_1 & b_2 \\ 0 & \cdots & 0 & a_0 & a_1 & 0 & 0 & b_0 & b_1 \end{vmatrix}$$

(it is a square matrix of size $(m + n - 2) \times (m + n - 2)$). Note that the subresultants share a lot of properties with the resultants; we refer the interested reader to [1].

2.1.2. *The multivariate case.* All along this paper we will also use resultants of several homogeneous polynomials; we now quickly recall this notion. Although they are usually defined “geometrically” (as equations of certain hypersurfaces obtained by projection of an incidence variety), we will follow the formalism developed by Jouanolou [18] because it easily provides many properties of resultants.

Suppose given an integer $n \geq 1$, and a sequence of positive integers d_1, \dots, d_n . One considers the n “generic” homogeneous polynomials of degree d_1, \dots, d_n , respectively, in the variables X_1, \dots, X_n (all assumed to have weight 1) :

$$f_i(X_1, \dots, X_n) = \sum_{|\alpha|=d_i} U_{i,\alpha} X^\alpha, \quad i = 1, \dots, n.$$

Denoting $\mathbb{U} := \mathbb{Z}[U_{i,\alpha} : i = 1, \dots, n, |\alpha| = d_i]$, the polynomials f_1, \dots, f_n belongs to the ring $C := \mathbb{U}[X_1, \dots, X_n]$. The *ideal of inertia forms* of these polynomials is the ideal of C

$$\text{TF}_m(f_1, \dots, f_n) := \{f \in C : \exists \nu \in \mathbb{N} \ m^\nu f \subset (f_1, \dots, f_n)\} \subset C$$

where $\mathfrak{m} := (X_1, \dots, X_n) \subset C$. It is naturally graded and it turns out that its degree zero graded part, denoted $\text{TF}_m(f_1, \dots, f_n)_0$, is a principal ideal of \mathbb{U} and

has a *unique generator*, denoted $\text{Res}_{X_1, \dots, X_n}$ or simply Res , which satisfies

$$(2.1) \quad \text{Res}(X_1^{d_1}, \dots, X_n^{d_n}) = 1.$$

To define the resultant of any given n -uples of homogeneous polynomials in the variables X_1, \dots, X_n (and also to clarify the left side of the equality (2.1)) one proceeds as follows: Let \mathbb{S} be a commutative ring. For all integers $i \in \{1, \dots, n\}$, suppose given a homogeneous polynomial of degree d_i in the variables X_1, \dots, X_n

$$g_i = \sum_{|\alpha|=d_i} u_{i,\alpha} X^\alpha \in \mathbb{S}[X_1, \dots, X_n]_{d_i}$$

and consider the morphism $\theta : \mathbb{U} \rightarrow \mathbb{S} : U_{j,\alpha} \mapsto u_{j,\alpha}$ which corresponds to the *specialization* of the polynomials f_i to the polynomials g_i . Then, given an inertia form $a \in \text{TF}_m(f_1, \dots, f_n)$ we set $a(g_1, \dots, g_n) := \theta(a)$. In particular, the resultant of g_1, \dots, g_n is nothing but

$$\text{Res}(g_1, \dots, g_n) := \theta(\text{Res}(f_1, \dots, f_n)).$$

Also, if $\mathbb{S} = \mathbb{U}$ and θ is the identity (i.e. $g_i = f_i$ for all i), then we get $a = a(f_1, \dots, f_n)$; this clarifies the notation $\text{Res}(f_1, \dots, f_n)$ for the inertia form $\text{Res} \in \mathbb{U}$.

Resultants have a lot of interesting properties. We recall the ones we will use in the sequel and refer the reader to [18, §5] for the proofs. Mention that many formulas are known to compute explicitly these resultants (e.g. [23], [20], [14], [11] and the reference therein).

Let \mathbb{S} be any commutative ring and suppose given f_1, \dots, f_n homogeneous polynomials in the polynomial ring $\mathbb{S}[X_1, X_2, \dots, X_n]$ of positive degree d_1, \dots, d_n respectively.

- *homogeneity*: for all $i = 1, \dots, n$, $\text{Res}(f_1, \dots, f_n)$ is homogeneous w.r.t. the coefficients of f_i of degree $d_1 \dots d_n / d_i$,
- *isobaricity*: $\text{Res}(f_1, \dots, f_n)$ is isobaric of degree $d_1 \dots d_n$ by giving to each coefficient of the f_i 's the power of its corresponding monomial in the variable X_n ,
- *permutation of variables*:

$$\text{Res}(f_{\sigma(1)}, \dots, f_{\sigma(n)}) = (\mathcal{E}(\sigma))^{d_1 \dots d_n} \text{Res}(f_1, \dots, f_n)$$

for any permutation σ of the set $\{1, \dots, n\}$ ($\mathcal{E}(\sigma)$ denotes the signature of the permutation σ),

- *elementary transformations*:

$$\text{Res}(f_1, \dots, f_i + \sum_{j \neq i} h_j f_j, \dots, f_n) = \text{Res}(f_1, \dots, f_n),$$

- *multiplicativity*:

$$\text{Res}(f_1' f_1'', f_2, \dots, f_n) = \text{Res}(f_1', f_2, \dots, f_n) \text{Res}(f_1'', f_2, \dots, f_n)$$

- *base change formula*: if g_1, \dots, g_n are homogeneous polynomials in $\mathbb{S}[X]$ of the same positive degree d , then

$$(2.2) \quad \text{Res}(f_1(g_1, \dots, g_n), \dots, f_n(g_1, \dots, g_n)) = \text{Res}(g_1, \dots, g_n)^{d_1 \dots d_n} \text{Res}(f_1, \dots, f_n)^{d^{n-1}},$$

- *divisibility*: if g_1, \dots, g_n are homogeneous polynomials in $\mathbb{S}[X]$ such that for all $i = 1, \dots, n$ there exists an integer μ_i such that $f_i \in (g_1, \dots, g_n)^{\mu_i}$, then $\text{Res}(g_1, \dots, g_n)^{\mu_1 \cdots \mu_n}$ divides $\text{Res}(f_1, \dots, f_n)$ in \mathbb{S} .

In the following, we are going to consider resultant computation for eliminating a subset X_{i_1}, \dots, X_{i_k} of the variables. Such a resultant, obtained by considering the homogenization of the polynomials with respect to this variable subset, will be denoted hereafter $\text{Res}_{X_{i_1}, \dots, X_{i_k}}$. With this notation, for homogeneous polynomials $f_1, \dots, f_n \in \mathbb{S}[X]$, we have

$$\text{Res}_{X_1, \dots, X_n}(f_1, \dots, f_n) = \text{Res}_{X_2, \dots, X_n}(f_1(1, X_2, \dots, X_n), \dots, f_n(1, X_2, \dots, X_n)).$$

2.2. Discriminants. Given a polynomial

$$P(X) := a_0 X^n + a_1 X^{n-1} + \cdots + a_{n-1} X + a_n \in \mathbb{S}[X]$$

where $n \geq 1$ and \mathbb{S} is any commutative ring, we recall that the discriminant of P , denoted $\text{Disc}_X(P)$, satisfies the equality

$$(2.3) \quad a_0 \text{Disc}_X(P(X)) = \text{Res}_X(P(X), \partial_X P(X)) = \text{Res}_X(\partial_X P(X), P(X))$$

where ∂_X stands for the derivative with respect to the variable X , i.e. $\partial/\partial X$. Its properties follow immediately from the ones of the resultant. In particular, its degree in the coefficients of P is $2(d-1)$. Note that in our notation the polynomial P is seen as a polynomial of its expected degree, here n , as we did for the resultant. Indeed, in (2.3) the considered resultants are in degrees $(n, n-1)$ and $(n-1, n)$ respectively.

Given a homogeneous polynomial $P(X_1, X_2, X_3) = \sum_{|\alpha|=d} U_\alpha X_1^{\alpha_1} X_2^{\alpha_2} X_3^{\alpha_3}$ of degree d , its discriminant is the polynomial in the universal ring of coefficients $\mathbb{U} := \mathbb{Z}[U_\alpha; |\alpha|=d]$, denoted $\text{Disc}_{X_1: X_2: X_3}(P)$ or simply $\text{Disc}(P)$, which satisfies

$$(2.4) \quad d^{d^2-3d+3} \text{Disc}_{X_1: X_2: X_3}(P) = \text{Res}_{X_1: X_2: X_3}(\partial_1 P, \partial_2 P, \partial_3 P)$$

where ∂_i stands for $\partial/\partial X_i$ for all i . It is an irreducible polynomial in \mathbb{U} which is homogeneous of degree $3(d-1)^2$ in the U_α 's. Note that we also have the following equality in \mathbb{U} :

$$\begin{aligned} \text{Res}_{X_1: X_2: X_3}(\partial_1 P, \partial_2 P, P) &= \text{Disc}_{X_1: X_2: X_3}(P) \text{Disc}_{X_2: X_3}(P(0, X_2, X_3)) \\ &= \text{Disc}_{X_1: X_2: X_3}(P) \text{Disc}_{X_3}(P(0, 1, X_3)). \end{aligned}$$

Finally, we also recall that being given two homogeneous polynomials P_1 and P_2 in the variables X_1, X_2, X_3 of degree d_1, d_2 respectively, their discriminant is defined by the formula [21]

$$\text{Disc}_{X_1: X_2: X_3}(P_1, P_2) = \frac{\text{Res}_{X_1: X_2: X_3}(P_1, P_2, \partial_2 P_1 \partial_3 P_2 - \partial_2 P_2 \partial_3 P_1)}{\text{Res}_{X_1: X_2: X_3}(P_1, P_2, X_1)} \in \mathbb{U}$$

where \mathbb{U} denotes the universal ring of coefficients of P_1 and P_2 over \mathbb{Z} . It is an irreducible polynomial in \mathbb{U} which is homogeneous with respect to the coefficients of P_1 of degree $d_2(2(d_1-1)+d_2-1)$ and homogeneous with respect to the coefficients of P_2 of degree $d_1(2(d_2-1)+d_1-1)$.

2.3. Notations and derivatives. Let \mathbb{S} be a commutative ring and suppose given a polynomial $P(X) \in \mathbb{S}[X]$. We denote by $\partial_X P$ the formal derivative of P with respect to the variable X . More precisely, the map

$$\partial_X : \mathbb{S}[X] \rightarrow \mathbb{S}[X] : P \mapsto \partial_X P$$

is \mathbb{S} -linear and for all integers $k \geq 0$ we have $\partial_X(X^k) = kX^{k-1}$. For any integer $i \geq 1$ we define ∂_X^i as the composition of ∂_X with itself i times; for instance $\partial_X^i(X^k)$ is either equal to 0 if $k < i$ or either equal to $k!/(k-i)!X^{k-i}$ if $k \geq i$.

Now, suppose given an element $a \in \mathbb{S}$ and consider the \mathbb{S} -linear map

$$\delta_a : \mathbb{S}[X] \rightarrow \mathbb{S}[X] : P \mapsto \delta_a P := \frac{P(X) - P(a)}{X - a}.$$

For instance, for all integers $k \geq 1$ we have $\delta_a(X^k) = \sum_{j=0}^{k-1} a^{k-1-j} X^j$. As above, for any integer $i \geq 1$ we define δ_a^i as the composition of δ_a with itself i times. It follows that for all integers $n \geq 1$

(2.5)

$$P(X) = P(a) + (X - a)\delta_a(P)(a) + (X - a)^2\delta_a^2(P)(a) + \cdots + (X - a)^n\delta_a^n(P)(X).$$

Indeed, we have $P(X) = P(a) + (X - a)\delta_a(P)(X)$ by definition. Applying this formula to $\delta_a(P)$ we deduce (2.5) for $n = 2$. The general case is obtained by a similar induction.

Although \mathbb{S} is only assumed to be a commutative ring, we still have the expected formula

Lemma 2.2. *For $k \geq 1$, for any $P \in \mathbb{S}[X]$ and $a \in \mathbb{S}$, we have the equality $k!\delta_a^k P(a) = \partial_X^k P(a)$ in \mathbb{S} .*

Proof. Notice that $\delta_a P(X)$ is of degree $\deg(P) - 1$ in the variable X . Thus if $d > \deg(P)$, we have $\delta_a^d(P) = 0$. We deduce that for any polynomial $P \in \mathbb{S}[X]$,

$$P(X) = P(a) + (X - a)\delta_a(P)(a) + \cdots + (X - a)^d\delta_a^{d \deg(P)}(P)(a).$$

By the binomial identity, for any $n \in \mathbb{N}$, $X^n = \sum_{i=0}^n \binom{n}{i} a^{n-i} (X - a)^i$. By identification of the coefficients in the basis $(X - a)^k$ of $\mathbb{S}[X]$, we deduce that for all $k \leq n$, we have $\delta_a^k(X^n)(a) = \binom{n}{k} a^{n-k}$. As we have $\partial_X^k(X^n) = \frac{n!}{(n-k)!} X^{n-k}$, we deduce that for any $k, n \in \mathbb{N}$, $k!\delta_a^k(X^n)(a) = \partial_X^k(X^n)(a)$. By linearity, we also have this identity for any polynomial $P \in \mathbb{S}[X]$. \square

As a consequence, if \mathbb{S} is assumed to be a \mathbb{Q} -algebra then (2.5) shows that for all integers $n \geq 1$

$$P(X) = P(a) + (X - a)\partial_X P(a) + \cdots + (X - a)^n \frac{\partial_X^n P(a)}{n!} + (X - a)^{n+1} \delta_a^{n+1}(P).$$

If n is bigger than the degree of P , then we get the Taylor expansion formula

$$P(X) = P(a) + \sum_{n \geq 1} \frac{\partial_X^n P(a)}{n!} (X - a)^n.$$

We now fix a notation that we will use all along the paper. We will mainly manipulate polynomials in the three variables X_1, X_2 and X_3 over a commutative

ring \mathbb{S} . Introducing a new indeterminate X_4 , for any polynomial $P \in \mathbb{S}[X_1, X_2, X_3]$ we set

$$\delta_{3,4}(P) := \frac{P(X_1, X_2, X_3) - P(X_1, X_2, X_4)}{X_3 - X_4} \in \mathbb{S}[X_1, X_2, X_3, X_4]$$

so that we have

$$P(X_1, X_2, X_4) = P(X_1, X_2, X_3) + (X_4 - X_3)\delta_{3,4}(P).$$

Observe that $\delta_{3,4}(P)$ is nothing but $\delta_{X_4}(P) \in (\mathbb{S}[X_1, X_2, X_4])[X_3]$ and therefore that all the above definitions and formulas can be expressed with this notation. Therefore, we have

$$(2.6) \quad \delta_{3,4}P = \partial_3 P(X_3) + (X_4 - X_3)\delta_{3,4}^2 P,$$

$$(2.7) \quad \begin{aligned} 2\delta_{3,4}^2 P &= \partial_3^2 P(X_3) + 2(X_4 - X_3)\delta_{3,4}^3 P \\ &= \frac{2}{(X_4 - X_3)^2} (P(X_1, X_2, X_3) - P(X_1, X_2, X_4) - (X_4 - X_3)\partial_3 P(X_1, X_2, X_3)). \end{aligned}$$

The above definition can be extended to any variable. We denote by $\delta_{i,j}^2$ the corresponding operation, X_i playing the role of X_3 , and X_j the role of X_4 . If the indices i, j are omitted, we implicitly refer to the variables X_3 and X_4 . Notice that $\delta_{3,4}P|_{X_4=X_3} = \partial_3 P(X_3)$. Also, by convention we set $\delta_{i,i}P = \partial_i P$.

We will need the two following properties: for any polynomials L, Q , we have

$$(2.8) \quad \delta_{3,4}(LQ) = \delta_{3,4}(L)Q(X_3) + L(X_4)\delta_{3,4}(Q),$$

$$(2.9) \quad \delta_{3,4}^2(LQ) = \delta_{3,4}^2(L)Q(X_3) + \delta_{3,4}(L)\partial_3(Q)(X_3) + L(X_4)\delta_{3,4}^2(Q).$$

To prove (2.8), we remark that

$$L(X_3)Q(X_3) - L(X_4)Q(X_4) = Q(X_3)(L(X_4) - L(X_3)) + L(X_4)(Q(X_4) - Q(X_3))$$

and divide by $(X_4 - X_3)$ to get the formula. To prove (2.9), we substitute (2.6) in the previous relation and obtain

$$\begin{aligned} &(X_4 - X_3)\partial_3(LQ) + (X_4 - X_3)^2\delta_{3,4}^2(QL) \\ &= (X_4 - X_3)(\partial_3(L)Q(X_3) + L(X_3)\partial_3(Q)) + (X_4 - X_3)^2\delta_{3,4}^2(QL) \\ &= Q(X_3)((X_4 - X_3)\partial_3(L) + (X_4 - X_3)^2\delta_{3,4}^2(L)) + L(X_4)((X_4 - X_3)\partial_3(Q) \\ &\quad + (X_4 - X_3)^2\delta_{3,4}^2(Q)) \end{aligned}$$

so that

$$\begin{aligned} (X_4 - X_3)^2\delta_{3,4}^2(QL) &= (X_4 - X_3)^2\delta_{3,4}^2(L)Q(X_3) \\ &\quad + (X_4 - X_3)(L(X_4) - L(X_3))\partial_3(Q) + (X_4 - X_3)^2L(X_4)\delta_{3,4}^2(Q), \end{aligned}$$

from which we deduce the relation (2.9).

Finally, mention that if \mathbb{S} is assumed to be a \mathbb{Q} -algebra then we have

$$\begin{aligned} P(X_1, X_2, X_4) &= P(X_1, X_2, X_3) + (X_4 - X_3)\partial_3 P(X_3) + \cdots + \\ &\quad \frac{1}{(i-1)!} (X_4 - X_3)^{i-1}\partial_3^{i-1}P(X_3) + (X_4 - X_3)^i\delta_{3,4}^i P \end{aligned}$$

or equivalently $\delta_{3,4}^i P = \sum_{k \geq i} \frac{1}{k!} (X_4 - X_3)^{k-1} \partial_3^k P(X_3)$. Similarly, we have

$$(2.10) \quad \delta_{3,4}^i P|_{X_4=X_3} = \frac{1}{i!} \partial_3^i P.$$

Notice that we will very often omit variables X_1 and X_2 in the sequel, especially in the proofs to avoid to overload the computations and the text.

2.4. A Bertini's lemma. The elimination of the variables X_1, X_2, X_3, X_4 between polynomials constructed from the polynomials

$$P_k(X_1, X_2, X_3) := \sum_{0 \leq i, j; i+j \leq d_k} U_{i,j}^{(k)} X_1^i X_2^j X_3^{d_k-i-j} \in \mathbb{U}[X_1, X_2, X_3],$$

$k = 1, \dots, r$, where \mathbb{U} denotes the universal coefficients ring $\mathbb{U} := \mathbb{Z}[U_{i,j}^{(k)}; 0 \leq i, j; i+j \leq d_k, k = 1, \dots, r]$, yields polynomials in \mathbb{U} for which we are going to give irreducible factorizations. As in geometric applications, this computation is applied with the coefficients $U_{i,j}^{(k)}$ replaced by generic polynomials $\sum_{|\alpha| \leq d_k-i-j} a_{i,j,\alpha}^{(k)} \mathbf{x}^\alpha$ of degree $d_k - i - j \geq 0$, where we consider the coefficients $a_{i,j,\alpha}^{(k)}$ as indeterminates where \mathbf{x} denotes a set of indeterminates (x_1, \dots, x_t) with $t \geq 1$. Therefore, we will need several times the following lemma:

Lemma 2.3. *Suppose that R is an irreducible homogeneous polynomial of \mathbb{U} . Then the polynomial $R(\mathbf{x}, \mathbf{a})$ obtained by substituting the coefficients $U_{i,j}^{(k)}$ by generic polynomials $\sum_{|\alpha| \leq d_k-i-j} a_{i,j,\alpha}^{(k)} \mathbf{x}^\alpha$ is irreducible in $\mathbb{Z}[\mathbf{x}, a_{i,j,\alpha}^{(k)}]$.*

Proof. Assume that R is not irreducible and can be splitted into the product of two factors in $\mathbb{Z}[\mathbf{x}, a_{i,j,\alpha}^{(k)}]$:

$$R(a_{i,j,\alpha}^{(k)}, \mathbf{x}) = R_1(a_{i,j,\alpha}^{(k)}, \mathbf{x}) R_2(a_{i,j,\alpha}^{(k)}, \mathbf{x}).$$

By sending \mathbf{x} to $\mathbf{0}$ we observe that $R(a_{i,j,\alpha}^{(k)}, \mathbf{0})$ is an irreducible polynomial in the $a_{i,j,\alpha}^{(k)}$ since R is irreducible so that we only have to rename each coefficient $a_{i,j,\alpha}^{(k)}$ by $U_{i,j}^{(k)}$. It follows that either $R_1(a_{i,j,\alpha}^{(k)}, \mathbf{0})$ or $R_2(a_{i,j,\alpha}^{(k)}, \mathbf{0})$ must be an invertible element in \mathbb{Z} . But since R_1 and R_2 are homogeneous in the coefficients $a_{i,j,\alpha}^{(k)}$ this implies that either R_1 or R_2 is an invertible element in \mathbb{Z} . \square

3. RESULTANT OF RESULTANTS

In all this section, we suppose given four positive integers d_1, d_2, d_3, d_4 and four homogeneous polynomials

$$P_k(X_1, X_2, X_3) = \sum_{0 \leq i, j; i+j \leq d_k} U_{i,j}^{(k)} X_1^i X_2^j X_3^{d_k-i-j} \in \mathbb{U}[X_1, X_2, X_3], k = 1, \dots, 4,$$

where \mathbb{U} denotes the universal ring of coefficients

$$\mathbb{U} := \mathbb{Z}[U_{i,j}^{(k)}; 0 \leq i, j; i+j \leq d_k, k = 1, \dots, 4].$$

We denote by X_4 a new indeterminate. Our first result is the most general situation of an iterated resultant with four different polynomials.

Theorem 3.1. *Defining*

$$\begin{aligned} R_{12} &:= \text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)) \in \mathbb{U}[X_2], \\ R_{34} &:= \text{Res}_{X_3}(P_3(1, X_2, X_3), P_4(1, X_2, X_3)) \in \mathbb{U}[X_2], \end{aligned}$$

we have the following equality in \mathbb{U} :

$$\text{Res}_{X_2}(R_{12}, R_{34}) =$$

$$\text{Res}_{X_1, \dots, X_4}(P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_3), P_3(X_1, X_2, X_4), P_4(X_1, X_2, X_4)).$$

Moreover, the above quantity is non-zero, irreducible and multi-homogeneous with respect to the set of coefficients $(U_{i,j}^{(1)})_{i,j}$, $(U_{i,j}^{(2)})_{i,j}$, $(U_{i,j}^{(3)})_{i,j}$, $(U_{i,j}^{(4)})_{i,j}$ of multi-degree $(d_2 d_3 d_4, d_1 d_3 d_4, d_1 d_2 d_4, d_1 d_2 d_3)$.

Proof. First of all, we observe that the iterated resultant $\text{Res}_{X_2}(R_{12}, R_{34})$ and the resultant

$$\mathcal{R} := \text{Res}(P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_3), P_3(X_1, X_2, X_4), P_4(X_1, X_2, X_4))$$

are both non-zero polynomials, for they both specialize to the quantity $(-1)^{d_1 d_2 d_3 d_4}$ if the polynomials $P_i(X_1, X_2, X_3)$, $i = 1, \dots, 4$, are specialized to $X_1^{d_1}$, $X_3^{d_2}$, $X_2^{d_3}$ and $X_3^{d_4}$ respectively. Note also that the statement about the multi-degree of \mathcal{R} follows from the homogeneity property of resultants.

To prove the irreducibility of \mathcal{R} , we proceed by induction on the positive integer $d := d_1 + d_2 + d_3 + d_4 \geq 4$. For $d = 4$, \mathcal{R} equals the determinant

$$\begin{vmatrix} U_{1,0}^{(1)} & U_{1,0}^{(2)} & U_{1,0}^{(3)} & U_{1,0}^{(4)} \\ U_{0,1}^{(1)} & U_{0,1}^{(2)} & 0 & 0 \\ U_{0,0}^{(1)} & U_{0,0}^{(2)} & U_{0,0}^{(3)} & U_{0,0}^{(4)} \\ 0 & 0 & U_{0,1}^{(3)} & U_{0,1}^{(4)} \end{vmatrix} \in \mathbb{U}$$

which is checked to be irreducible. Thus, we assume that \mathcal{R} is irreducible up to a given integer $p \geq 4$ and we will prove that \mathcal{R} is irreducible if $d = p + 1$. To do this, first observe that one of the integers d_1, d_2, d_3, d_4 must be greater or equal to 2. We can assume that $d_1 \geq 2$ without loss of generality. Consider the specialization ϕ leaving invariant the polynomials P_2, P_3 and P_4 and sending the polynomial P_1 to the product $L_1 Q_1$ where L_1 and Q_1 are both generic forms of respective degree 1 and $d_1 - 1 \geq 1$. Then, by multiplicativity of resultants we have the equality

$$\phi(\mathcal{R}) = \text{Res}(L_1, P_2, P_3, P_4) \text{Res}(Q_1, P_2, P_3, P_4),$$

whose right hand side is a product of two irreducible polynomials by our induction hypothesis. As the specialization ϕ is homogeneous (in terms of the coefficients of the P_i 's, L_1 and Q_1), the number of irreducible factors of \mathcal{R} can not decrease under the specialization ϕ and we deduce that \mathcal{R} is the product of two irreducible polynomials \mathcal{R}_1 and \mathcal{R}_2 . But then, one of these two factors must depend on the coefficients of P_1 , say \mathcal{R}_1 , and therefore $\phi(\mathcal{R}_1)$ must depend on the coefficients of L_1 and Q_1 . This implies that \mathcal{R}_2 is an invertible element in \mathbb{Z} and consequently that \mathcal{R} is irreducible.

It remains to prove the claimed equality. To do this, we rewrite $P_k(1, X_2, X_3)$, for all $k = 1, \dots, 4$, as

$$P_k(1, X_2, X_3) = \sum_{i=0}^{d_k} \left(\sum_{j=0}^{d_k-i} U_{i,j}^{(k)} X_2^j \right) X_3^i \in \mathbb{U}[X_2, X_3],$$

and we then easily see from well-known properties of the Sylvester resultant that

- R_{12} is bi-homogeneous in the set of coefficients $(U_{i,j}^{(1)})$ and $(U_{i,j}^{(2)})$ of bi-degree (d_2, d_1) ,

- R_{12} is a polynomial in $\mathbb{U}[X_2]$ of degree $d_1 d_2$,
- $R_{12} \in (P_1(1, X_2, X_3), P_2(1, X_2, X_3)) \subset \mathbb{U}[X_2, X_3]$.

Of course, completely analogous results hold for R_{34} , in particular

$$R_{34} \in (P_3(1, X_2, X_4), P_4(1, X_2, X_4)) \subset \mathbb{U}[X_2, X_4].$$

Again, $\text{Res}_{X_2}(R_{12}, R_{34}) \in (R_{12}, R_{34}) \subset \mathbb{U}[X_2]$ and we deduce that

$$\begin{aligned} \text{Res}_{X_2}(R_{12}, R_{34}) &\in \\ &(P_1(1, X_2, X_3), P_2(1, X_2, X_3), P_3(1, X_2, X_4), P_4(1, X_2, X_4)) \subset \mathbb{U}[X_2, X_3, X_4]. \end{aligned}$$

After homogenization with the variable X_1 , it follows that there exists an integer N such that

$$\begin{aligned} X_1^N \text{Res}_{X_2}(R_{12}, R_{34}) &\in \\ &(P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_3), P_3(X_1, X_2, X_4), P_4(X_1, X_2, X_4)) \end{aligned}$$

in $\mathbb{U}[X_1, X_2, X_3, X_4]$ (notice that this does not imply directly that $\text{Res}_{X_2}(R_{12}, R_{34})$ is an inertia form because $P_1(X_1, X_2, X_3)$, $P_2(X_1, X_2, X_3)$, $P_3(X_1, X_2, X_4)$ and $P_4(X_1, X_2, X_4)$ are not generic polynomials). It implies by the divisibility property of resultants, that $\mathcal{R} = \text{Res}(P_1(X_3), P_2(X_3), P_3(X_4), P_4(X_4))$ divides the quantity

$$\begin{aligned} \text{Res}(X_1^N \text{Res}_{X_2}(R_{12}, R_{34}), P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_3), P_3(X_1, X_2, X_4)) = \\ \text{Res}_{X_2}(R_{12}, R_{34})^{d_1 d_2 d_3} \text{Res}(X_1, P_1(X_3), P_2(X_3), P_3(X_4))^N. \end{aligned}$$

Since \mathcal{R} is irreducible and since the second term in the right hand side of the above product does not depend on the coefficients of P_4 , we deduce that \mathcal{R} divides $\text{Res}_{X_2}(R_{12}, R_{34})$. Now, from the degree properties of R_{12} and R_{13} we deduce that $\text{Res}_{X_2}(R_{12}, R_{34})$ is, similarly to \mathcal{R} , multi-homogeneous with respect to the set of coefficients $(U_{i,j}^{(1)})_{i,j}, (U_{i,j}^{(2)})_{i,j}, (U_{i,j}^{(3)})_{i,j}, (U_{i,j}^{(4)})_{i,j}$ of multi-degree

$$(d_2 d_3 d_4, d_1 d_3 d_4, d_1 d_2 d_4, d_1 d_2 d_3).$$

This shows that $\text{Res}_{X_2}(R_{12}, R_{34})$ and \mathcal{R} are equal up to multiplication by an invertible element in \mathbb{Z} . To determine this invertible element, we take again the specialization sending P_1 to $X_1^{d_1}$, P_2 to $X_3^{d_2}$, P_3 to $X_2^{d_3}$, P_4 to $X_3^{d_4}$, and check that \mathcal{R} specializes to $(-1)^{d_1 d_2 d_3 d_4}$, as well as $\text{Res}_{X_2}(R_{12}, R_{34})$. \square

A specialization of this theorem gives the following result.

Corollary 3.2. *Given four polynomials $f_k(\mathbf{x}, y, z), k = 1, \dots, 4$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha,i,j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, then the iterated resultant $\text{Res}_y(\text{Res}_z(f_1, f_2), \text{Res}_z(f_3, f_4)) \in \mathbb{S}[\mathbf{x}]$ is of degree at most $d_1 d_2 d_3 d_4$ in \mathbf{x} and we have

$$\begin{aligned} \text{Res}_y(\text{Res}_z(f_1, f_2), \text{Res}_z(f_3, f_4)) = \\ \text{Res}_{y,z,z'}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), f_3(\mathbf{x}, y, z'), f_4(\mathbf{x}, y, z')) \in \mathbb{S}[\mathbf{x}]. \end{aligned}$$

Moreover, if the polynomials f_1, f_2, f_3 and f_4 are sufficiently generic then this iterated resultant is irreducible and has exactly degree $d_1 d_2 d_3 d_4$ in \mathbf{x} .

Proof. It is a corollary of Theorem 3.1. Indeed, the claimed equality is deduced from the equality given in Theorem 3.1 by substituting X_2 by y , X_3 by z , by seeing X_1 as the homogenization variable of the variables (X_2, X_3) and by specializing each coefficient $U_{i,j}^{(k)}$ to $\sum_{|\alpha| \leq d_k - i - j} a_{i,j,\alpha}^{(k)} \mathbf{x}^\alpha$. Moreover, since each coefficient $U_{i,j}^{(k)}$ is specialized to a polynomial in \mathbf{x} of degree at most d_k we deduce the claimed degree bound as a consequence of the isobarity formula for resultants.

If the polynomials f_1, f_2, f_3 and f_4 are sufficiently generic then it is clear that the degree bound is reached: this and the irreducibility statement is a consequence of Lemma 2.3. \square

The formula proved in Theorem 3.1 can be specialized to get the factorization of an iterated resultant which may occur quite often in practical situations (implicitization of a rational surface, projection of the intersection of two surfaces in projective space, . . . e.g. [24, 6]).

Proposition 3.3. *Assume that $d_1 \geq 2$ and set*

$$\begin{aligned} R_{12} &:= \text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)) \in \mathbb{U}[X_2], \\ R_{13} &:= \text{Res}_{X_3}(P_1(1, X_2, X_3), P_3(1, X_2, X_3)) \in \mathbb{U}[X_2]. \end{aligned}$$

Then, we have the equality

$$\begin{aligned} \text{Res}_{X_2}(R_{13}, R_{12}) &= \text{Res}_{X_1: X_2: X_3}(P_1, P_2, P_3) \times \\ &\quad \text{Res}_{X_1: \dots: X_4}(P_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4} P_1(X_3, X_4)) \end{aligned}$$

where the right hand side is a product of two irreducible polynomials in \mathbb{U} .

Moreover, the iterated resultant $\text{Res}_{X_2}(R_{13}, R_{12})$ is multi-homogeneous with respect to the set of coefficients $(U_{i,j}^{(1)})_{i,j}, (U_{i,j}^{(2)})_{i,j}, (U_{i,j}^{(3)})_{i,j}$ of degree $2d_1d_2d_3, d_1^2d_3$ and $d_1^2d_2$ respectively.

Proof. The claimed equality is easily obtained using formal properties of resultants by specialization of the formula proved in Theorem 3.1:

$$\begin{aligned} &\text{Res}(P_1(X_3), P_3(X_3), P_1(X_4), P_2(X_4)) \\ &= \text{Res}(P_1(X_3), P_2(X_4), P_3(X_3), P_1(X_4)) \\ &= \text{Res}(P_1(X_3), P_2(X_4), P_3(X_3), P_1(X_4) - P_1(X_3)) \\ &= \text{Res}(P_1(X_3), P_2(X_4), P_3(X_3), (X_4 - X_3)\delta_{3,4}(P_1)) \\ &= \text{Res}(P_1, P_2, P_3) \text{Res}(P_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(P_1)). \end{aligned}$$

The multi-degree computation and the irreducibility of $\text{Res}(P_1, P_2, P_3)$ are known properties of resultants. The only point which requires a proof is the irreducibility of the factor (which is easily seen to be non-zero by a straightforward specialization)

$$\mathcal{D} := \text{Res}(P_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(P_1)(X_3, X_4)).$$

To do this, we proceed similarly to what we did in Theorem 3.1, that is, by induction on the integer $d := d_1 + d_2 + d_3 \geq 4$. We can check by hand (or with a computer) that \mathcal{D} is an irreducible polynomial in \mathbb{U} if $(d_1, d_2, d_3) = (2, 1, 1)$. We thus assume that \mathcal{D} is irreducible up to a given integer $p \geq 4$ and we will prove that \mathcal{D} is irreducible if $d = p + 1$. If $d_2 \geq 2$ (resp. $d_3 \geq 2$) then we can specialize P_2 (resp. P_3) as a product of a generic linear form and a generic form of degree $d_2 - 1 \geq 1$ (resp. $d_3 - 1 \geq 1$) and conclude, exactly as we did in Theorem 3.1, that \mathcal{D} is then

irreducible. Otherwise, then $d_1 \geq 3$ and we specialize P_1 to the product of a generic linear form

$$L_1(X_1, X_2, X_3) := aX_1 + bX_2 + cX_3$$

and a generic form Q_1 of degree $d_1 - 1 \geq 2$. We call ϕ the map corresponding to this specialization. By (2.8), we have

$$\delta_{3,4}(L_1Q_1)(X_3, X_4) = cQ_1(X_3) + L_1(X_4)\delta_{3,4}(Q_1)(X_3, X_4)$$

and we deduce after some manipulations on resultants that

$$\begin{aligned} \phi(\mathcal{D}) &= \text{Res}(L_1(X_3)Q_1(X_3), P_2(X_4), P_3(X_3), cQ_1(X_3) + L_1(X_4)\delta_{3,4}(Q_1)) \\ &= \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), L_1(X_4)\delta_{3,4}(Q_1)(X_3, X_4)) \times \\ &\quad \text{Res}(L_1(X_3), P_2(X_4), P_3(X_3), cQ_1(X_3) + L_1(X_4)\delta_{3,4}(Q_1)(X_3, X_4)) \\ &= \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(Q_1)(X_3, X_4)) \times \\ &\quad \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), L_1(X_4)) \times \\ &\quad \text{Res}(L_1(X_3), P_2(X_4), P_3(X_3), (L_1(X_4) - L_1(X_3))\delta_{3,4}(Q_1) + cQ_1(X_3)) \\ &= \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(Q_1)(X_3, X_4)) \times \\ &\quad \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), L_1(X_4)) \times \\ &\quad \text{Res}(L_1(X_3), P_2(X_4), P_3(X_3), cQ_1(X_4)) \\ &= c^{d_2d_3} \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(Q_1)(X_3, X_4)) \times \\ &\quad \text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), L_1(X_4)) \times \\ &\quad \text{Res}(L_1(X_3), P_2(X_4), P_3(X_3), Q_1(X_4)). \end{aligned}$$

Either by our induction hypothesis or by Theorem 3.1, it turns out that the three resultants involved in the right hand side of the above computation are irreducible in \mathbb{U} . So if \mathcal{D} were reducible, say $\mathcal{D} = \mathcal{D}_1\mathcal{D}_2$, then each factor should be homogeneous in the coefficients $(U_{i,j}^{(1)})_{i,j}$ and hence $\phi(\mathcal{D}_1)$ and $\phi(\mathcal{D}_2)$ should be homogeneous in the coefficients of Q_1 and L_1 of the same degree. But

$$\begin{aligned} \deg_{L_1, Q_1}(c) &= (1, 0), \\ \deg_{L_1, Q_1}(\text{Res}(L_1(X_3), P_2(X_4), P_3(X_3), Q_1(X_4))) &= ((d_1 - 1)d_2d_3, d_2d_3), \\ \deg_{L_1, Q_1}(\text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), L_1(X_4))) &= ((d_1 - 1)d_2d_3, d_2d_3), \\ \deg_{L_1, Q_1}(\text{Res}(Q_1(X_3), P_2(X_4), P_3(X_3), \delta_{3,4}(Q_1)(X_3, X_4))) &= (0, 2d_1d_2d_3 - 3d_2d_3), \end{aligned}$$

which implies that either \mathcal{D}_1 or \mathcal{D}_2 is an invertible element in \mathbb{Z} . \square

As a consequence of this proposition, we get the following result which improves Theorem 3.1 and Theorem 3.2 in [26].

Corollary 3.4. *Given three polynomials $f_k(\mathbf{x}, y, z)$, $k = 1, \dots, 3$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha, i, j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, then the iterated resultant $\text{Res}_y(\text{Res}_z(f_1, f_2), \text{Res}_z(f_1, f_3)) \in \mathbb{S}[\mathbf{x}]$

is of degree at most $d_1^2 d_2 d_3$ in \mathbf{x} and we have

$$\begin{aligned} \text{Res}_y(\text{Res}_z(f_1, f_2), \text{Res}_z(f_1, f_3)) &= (-1)^{d_1 d_2 d_3} \times \\ &\quad \text{Res}_{y,z}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), f_3(\mathbf{x}, y, z)) \times \\ &\quad \text{Res}_{y,z,z'}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), f_3(\mathbf{x}, y, z'), \delta_{z,z'}(f_1)). \end{aligned}$$

Moreover, if the polynomials f_1, f_2, f_3 are sufficiently generic then this iterated resultant has exactly degree $d_1^2 d_2 d_3$ in \mathbf{x} and both resultants on the right hand side of the above equality are distinct and irreducible.

This corollary can be interpreted geometrically as follows. For simplicity we assume that \mathbf{x} is a unique variable x and that f_1, f_2 and f_3 are three polynomial in x, y, z . The resultant $R_{12} := \text{Res}_z(f_1, f_2)$ defines the projection of the intersection curve between the two surfaces $\{f_1 = 0\}$ and $\{f_2 = 0\}$. Similarly, $R_{13} := \text{Res}_z(f_1, f_3)$ defines the projection of the intersection curve between the two surfaces $\{f_1 = 0\}$ and $\{f_3 = 0\}$. Then the roots of $\text{Res}_y(R_{12}, R_{13})$ can be decomposed into two distinct sets: the set of roots x_0 such that there exists y_0 and z_0 such that $f_1(x_0, y_0, z_0) = f_2(x_0, y_0, z_0) = f_3(x_0, y_0, z_0)$, and the set of roots x_1 such that there exist two distinct points (x_1, y_1, z_1) and (x_1, y'_1, z'_1) such that $f_1(x_1, y_1, z_1) = f_2(x_1, y_1, z_1)$ and $f_1(x_1, y'_1, z'_1) = f_3(x_1, y'_1, z'_1)$. The first set gives rise to the term $\text{Res}_{x,y,z}(f_1, f_2, f_3)$ in the factorization of the iterated resultant $\text{Res}_y(\text{Res}_{12}, \text{Res}_{13})$, and the second set of roots corresponds to the second factor.

Remark 3.5. Before going further, it is a good point to emphasize the fact that all the formulas presented in this paper are universal in the sens that they remain true for any specialization of the coefficients of the given polynomials. It should be noticed that this is true if and only if we take the univariate resultants in their expecting degrees. For instance, in the above corollary, $\text{Res}_z(f_1, f_2)$ denotes the resultant w.r.t. the variable z of the polynomials f_1, f_2 which are seen as polynomials of degree d_1 and d_2 respectively (even if their degree is actually lower for a given specialization). If one does not take care of this point, then one loses the universal property of these formulas as in [26, §7] where it is observed that if $f_1 := y^2 + z + x$, $f_2 := -y^2 + z$ and $f_3 := y^2 + z$ then $\text{Res}_{y,z}(f_1, f_2, f_3) = 0$ (for there is a base point at infinity) but $\text{Res}_y(\text{Res}_z(f_1, f_2), \text{Res}_z(f_1, f_3)) = 4x^2 \neq 0$, considering f_1, f_2 and f_3 as polynomials in z of degree 1 and not 2. In such a case, formulas similar to the ones we proved above require the use of more sophisticated resultants which can take into account some particular structure of polynomial systems. For instance, in this example one can take into account the presence of the base point defined by the ideal (y^2, w) , where w is the homogenizing variable, by considering the residual resultant [5, 7] (or the multi-homogeneous resultant which is the same here): we have the decomposition

$$(f_1, f_2, f_3) = (y^2, w) \begin{pmatrix} 1 & -1 & 1 \\ z + xw & z & z \end{pmatrix}$$

and we can check that the residual resultant equals $4x^2$ (up to a sign).

4. RESULTANTS OF DISCRIMINANTS

In this section, we will give the factorization of three iterated resultants corresponding to two cases of a resultant of a discriminant and a resultant and to the

case of a resultant of two discriminants. As we will see, the first case we are going to treat yields the two others by suitable specializations.

4.1. Resultant of a discriminant and a resultant.

Proposition 4.1. *Assume that $d_1 \geq 2$ and set*

$$\begin{aligned} D_1 &:= \text{Disc}_{X_3}(P_1(1, X_2, X_3)) \in \mathbb{U}[X_2], \\ R_{23} &:= \text{Res}_{X_3}(P_2(1, X_2, X_3), P_3(1, X_2, X_3)) \in \mathbb{U}[X_2]. \end{aligned}$$

Then the following equality holds in \mathbb{U} :

$$\begin{aligned} &(U_{0,0}^{(1)})^{d_2 d_3} \text{Res}_{X_2}(D_1, R_{23}) = \\ &\text{Res}_{X_1, \dots, X_4}(P_1(X_1, X_2, X_3), \partial_3 P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_4), P_3(X_1, X_2, X_4)). \end{aligned}$$

Moreover, the iterated resultant $\text{Res}_{X_2}(D_1, R_{23}) \in \mathbb{U}$ is irreducible and multi-homogeneous with respect to the coefficients $(U_{i,j}^{(1)})_{i,j}$, $(U_{i,j}^{(2)})_{i,j}$ and $(U_{i,j}^{(3)})_{i,j}$ of multi-degree

$$(2d_2 d_3 (d_1 - 1), d_1 (d_1 - 1) d_3, d_1 (d_1 - 1) d_2).$$

Proof. By (2.3), we easily get that

$$\begin{aligned} \mathcal{D} := (U_{0,0}^{(1)})^{d_2 d_3} \text{Res}_{X_2}(D_1, R_{23}) &= \text{Res}_{X_3}(U_{0,0}^{(1)} D_1, R_{23}) = \\ &\text{Res}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), \partial_3 P_1(1, X_2, X_3)), R_{23}) \end{aligned}$$

and hence, using the formula proved in Theorem 3.1 where we specialize the polynomials P_2, P_3 and P_4 to the polynomials $\partial_3 P_1, P_2$ and P_3 respectively, we deduce that, in \mathbb{U} ,

$$\mathcal{D} = \text{Res}(P_1(X_1, X_2, X_3), \partial_3 P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_4), P_3(X_1, X_2, X_4)).$$

The classical multi-degree formula for resultants gives the claimed result concerning the multi-degree of the iterated resultant $\text{Res}_{X_2}(D_1, R_{23})$. Now, we proceed by induction on the integer $d := d_1 + d_2 + d_3 \geq 4$ (remember $d_1 \geq 2$, $d_2 \geq 1$ and $d_3 \geq 1$) to prove the irreducibility of the iterated resultant $\text{Res}_{X_3}(D_1, R_{23})$.

First, if $d = 4$, that is to say $d_1 = 2$ and $d_2 = d_3 = 1$, then we check by hand that \mathcal{D} equals an irreducible polynomial in \mathbb{U} times $U_{0,0}^{(1)}$.

We now assume that $d \geq 5$. If $d_3 \geq 2$ then we consider the specialization ϕ which sends P_3 to the product of the generic linear form L_3 and the generic homogeneous polynomial Q_3 of degree $d_3 - 1 \geq 1$ and leave P_1 and P_2 invariant. We get, using the multiplicativity of resultants

$$\begin{aligned} \phi(\mathcal{D}) &= \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), L_3(X_4)) \\ &\quad \text{Res}(P_1(X_3), \partial_3 P_1(X_3), L_1(X_4), P_2(X_4), Q_3(X_4)). \end{aligned}$$

Therefore, by the induction hypothesis $\phi(\mathcal{D}) = (U_{0,0}^{(1)})^{d_2 d_3} R_1 R_2$ where R_1 and R_2 are irreducible polynomials such that R_1 depends only on the coefficients of L_3 and R_2 depends only on the coefficients of Q_3 . Since ϕ is a homogeneous specialization, each irreducible factor of \mathcal{D} which depends on P_3 must depend on L_3 and Q_3 , we deduce that \mathcal{D} has only one irreducible factor depending on P_3 . Moreover, the irreducible factors of \mathcal{D} which do not depend on P_3 are left invariant by ϕ so we deduce that \mathcal{D} equals $(U_{0,0}^{(1)})^{d_2 d_3}$ times an irreducible factor and we are done. If $d_2 \geq 2$ then we can argue exactly in the same way.

So it only remains to consider the case where $d_2 = d_3 = 1$. Let ψ be the specialization which sends P_1 to the product of the generic linear form $L_1 := aX_1 + bX_2 + cX_3$ times the generic homogeneous polynomial Q_1 of degree $d_1 - 1$ and leaves P_2 and P_3 invariant. By the basic properties of resultants and our induction hypothesis we get:

$$\begin{aligned}
\psi(\mathcal{D}) &= \text{Res}(L_1(X_3)Q_1(X_3), cQ_1(X_3) + L_1(X_3)\partial_3(Q_1(X_3)), P_2(X_4), P_3(X_4)) \\
&= \text{Res}(L_1(X_3), cQ_1(X_3), P_2(X_4), P_3(X_4)) \times \\
&\quad \text{Res}(Q_1(X_3), L_1(X_3)\partial_3(Q_1(X_3)), P_2(X_4), P_3(X_4)) \\
&= (-1)^{d_1-1} c \text{Res}(L_1(X_3), Q_1(X_3), P_2(X_4), P_3(X_4))^2 \times \\
&\quad \text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), P_2(X_4), P_3(X_4)) \\
&= (-1)^{d_1-1} c (U_{0,0}^{(1)})^{d_1-1} \text{Res}(L_1(X_3), Q_1(X_3), P_2(X_4), P_3(X_4))^2 \times R
\end{aligned}$$

where R is an irreducible polynomial which does not depend on the coefficients of L_1 and does depend on the coefficients of Q_1, P_2 of P_3 ; in particular it has degree $2(d_1 - 2)$ in the coefficients of Q_1 . Observe that $\text{Res}(L_1(X_3), Q_1(X_3), P_2(X_4), P_3(X_4))$ is irreducible by theorem 3.1 and has degree $d_1 - 1$ in the coefficients of L_1 and 1 in the coefficients of Q_1 . Since ψ is an homogeneous specialization, each irreducible factor of \mathcal{D} which depends on P_1 must depend on L_1 and Q_1 and have the same degree with respect to the coefficients of these two polynomials. From this property and the above computation we deduce that \mathcal{D} has only one irreducible factor that depends on P_1 . Moreover, since the others irreducible factors of \mathcal{D} are left invariant by ψ we deduce that \mathcal{D} equals $(U_{0,0}^{(1)})^{d_1}$ times an irreducible polynomial. \square

Corollary 4.2. *Given three polynomials $f_k(\mathbf{x}, y, z), k = 1, 2, 3$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha,i,j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, then the iterated resultant $\text{Res}_y(\text{Disc}_z(f_1), \text{Res}_z(f_2, f_3)) \in \mathbb{S}[\mathbf{x}]$ is of degree at most $d_1(d_1 - 1)d_2d_3$ in \mathbf{x} and we have

$$\begin{aligned}
(a_{0,0,d}^{(1)})^{d_2d_3} \text{Res}_y(\text{Disc}_z(f_1), \text{Res}_z(f_2, f_3)) &= \\
&\text{Res}_{y,z}(f_1(\mathbf{x}, y, z), \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), f_3(\mathbf{x}, y, z)).
\end{aligned}$$

Moreover, if the polynomials f_1, f_2, f_3 are sufficiently generic then this iterated resultant has exactly degree $d_1(d_1 - 1)d_2d_3$ in \mathbf{x} and the iterated resultant $\text{Res}_y(\text{Disc}_z(f_1), \text{Res}_z(f_2, f_3))$ is irreducible.

We can now specialize the formula of Proposition 4.1 to get the factorization of two kinds of iterated resultants: the resultant of two discriminants and the resultant of a discriminant of a polynomial f and a resultant of f and another polynomial. We begin with the simplest one.

4.2. Resultant of two discriminants of distinct polynomials.

Proposition 4.3. *Assume that $d_1 \geq 2$ and $d_2 \geq 2$ and set*

$$\begin{aligned}
D_1 &:= \text{Disc}_{X_3}(P_1(1, X_2, X_3)) \in \mathbb{U}[X_2], \\
D_2 &:= \text{Disc}_{X_3}(P_2(1, X_2, X_3)) \in \mathbb{U}[X_2].
\end{aligned}$$

Then the following equality holds in \mathbb{U} :

$$(U_{0,0}^{(1)})^{d_2(d_2-1)}(U_{0,0}^{(2)})^{d_1(d_1-1)}\text{Res}_{X_2}(D_1, D_2) = \\ \text{Res}_{X_1, \dots, X_4}(P_1(X_1, X_2, X_3), \partial_3 P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_4), \partial_3 P_2(X_1, X_2, X_4)).$$

Moreover, the resultant $\text{Res}_{X_2}(D_1, D_2) \in \mathbb{U}$ is irreducible and bi-homogeneous with respect to the coefficients $(U_{i,j}^{(1)})_{i,j}$ and $(U_{i,j}^{(2)})_{i,j}$ of bi-degree

$$(2d_2(d_1-1)(d_2-1), 2d_1(d_1-1)(d_2-1)).$$

Proof. We set $\mathcal{D} := (U_{0,0}^{(1)})^{d_2(d_2-1)}(U_{0,0}^{(2)})^{d_1(d_1-1)}\text{Res}_{X_2}(D_1, D_2)$. By (2.3), we have

$$(U_{0,0}^{(2)})^{d_1(d_1-1)}\text{Res}_{X_2}(D_1, D_2) = \text{Res}_{X_2}(D_1, U_{0,0}^{(2)}D_2) = \\ \text{Res}_{X_2}(D_1, \text{Res}_{X_3}(P_2(1, X_2, X_3), \partial_3 P_2(1, X_2, X_3)))$$

and hence, using the formula proved in Proposition 4.1, where we specialize the polynomial P_3 to the polynomial $\partial_3 P_2$, we deduce that, in \mathbb{U} , \mathcal{D} is equal to

$$\text{Res}_{X_1, \dots, X_4}(P_1(X_1, X_2, X_3), \partial_3 P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_4), \partial_3 P_2(X_1, X_2, X_4)).$$

The classical multi-degree formula for resultants gives the claimed result concerning the bi-degree of the iterated resultant $\text{Res}_{X_2}(D_1, D_2)$. To prove its irreducibility, we proceed by induction on the integer $d := d_1 + d_2 \geq 4$.

First, if $d = 4$, that is to say $d_1 = d_2 = 2$, then we check by hand (or with a computer) that \mathcal{D} equals an irreducible polynomial in \mathbb{U} times the factor $(U_{0,0}^{(1)})^2(U_{0,0}^{(2)})^2$. We now assume that $d \geq 5$. Without loss of generality we can also assume that $d_1 \geq d_2$ (since the problem is completely symmetric in P_1 and P_2) and hence that $d_1 \geq 3$. Consider the specialization ϕ which sends P_1 to the product of the generic linear form $L_1 := aX_1 + bX_2 + cX_3$ times the generic homogeneous polynomial Q_1 of degree $d_1 - 1$ and leave P_2 invariant. Using properties of resultants we get:

$$\begin{aligned} \phi(\mathcal{D}) &= \text{Res}(L_1(X_3)Q_1(X_3), cQ_1(X_3) + L_1(X_3)\partial_3(Q_1(X_3)), P_2(X_4), \partial_3 P_2(X_4)) \\ &= \text{Res}(L_1(X_3), cQ_1(X_3), P_2(X_4), \partial_3 P_2(X_4)) \times \\ &\quad \text{Res}(Q_1(X_3), L_1(X_3)\partial_3(Q_1(X_3)), P_2(X_4), \partial_3 P_2(X_4)) \\ &= c^{d_2(d_2-1)}\text{Res}(L_1(X_3), Q_1(X_3), P_2(X_4), \partial_3 P_2(X_4))^2 \times \\ &\quad \text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), P_2(X_4), \partial_3 P_2(X_4)). \end{aligned}$$

Using our inductive hypothesis and Proposition 4.1 we deduce that

$$\phi(\mathcal{D}) = (cU_{0,0}^{(1)'})^{d_2(d_2-1)}(U_{0,0}^{(2)})^{d_1(d_1-1)}R_1^2 \times R_2$$

$(U_{0,0}^{(1)'}$ being the coefficient of $X_3^{d_1-1}$ in Q_1) where R_1 is an irreducible polynomial of degree $d_2(d_2-1)(d_1-1)$ in the coefficients of L_1 and $d_2(d_2-1)$ in the coefficients of Q_1 , and R_2 is an irreducible polynomial independent of the coefficients of L_1 and of degree $2d_2(d_2-1)(d_1-2)$ in the coefficients of Q_1 . Note that we already know that $(U_{0,0}^{(1)})^{d_2(d_2-1)}(U_{0,0}^{(2)})^{d_1(d_1-1)}$ is a factor of \mathcal{D} . Moreover, since ϕ is an homogeneous specialization, each remaining irreducible factor of \mathcal{D} which depends on P_1 must depend on L_1 and Q_1 with the same degree, so we deduce that $R_1^2 R_2$ comes from the same irreducible factor of \mathcal{D} and we conclude that \mathcal{D} equals $(U_{0,0}^{(1)})^{d_2(d_2-1)}(U_{0,0}^{(2)})^{d_1(d_1-1)}$ times an irreducible polynomial in \mathbb{U} . \square

Corollary 4.4. *Given two polynomials $f_k(\mathbf{x}, y, z)$, $k = 1, 2$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha,i,j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where $d_1, d_2 \geq 2$, \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is a commutative ring, then the iterated resultant $\text{Res}_y(\text{Disc}_z(f_1), \text{Disc}_z(f_2)) \in \mathbb{S}[\mathbf{x}]$ is of degree at most $d_1(d_1 - 1)d_2(d_2 - 1)$ in \mathbf{x} and we have

$$(a_{0,0,d_1}^{(1)})^{d_2(d_2-1)} (a_{0,0,d_2}^{(2)})^{d_1(d_1-1)} \text{Res}_y(\text{Disc}_z(f_1), \text{Disc}_z(f_2)) = \\ \text{Res}_{y,z}(f_1(\mathbf{x}, y, z), \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), \frac{\partial f_2}{\partial z}(\mathbf{x}, y, z)).$$

Moreover, if the polynomials f_1, f_2, f_3 are sufficiently generic then this iterated resultant has exactly degree $d_1(d_1 - 1)d_2(d_1 - 1)$ in \mathbf{x} and is irreducible.

4.3. Resultant of a discriminant and a resultant sharing one polynomial.

We now turn to the second specialization of Proposition 4.1 which is a little more intricate than the previous one. Note that this iterated resultant has also been studied in [26, Theorem 3.3]. In order to improve the previous analysis, we begin with two technical results. We recall that we sometimes omit the variables X_1, X_2 , i.e. we note $P(X_3)$ instead of $P(X_1, X_2, X_3)$ to not overload the text.

Lemma 4.5. *For $d_1 \geq 2$,*

$$(U_{0,0}^{(1)})^{d_1 d_2} \text{Res}_{X_2}(D_1, R_{12}) = \text{Res}_{X_1:X_2:X_3}(P_1, \partial_3 P_1(X_3), P_2(X_3))^2 \times \\ \text{Res}_{X_1:X_2:X_3:X_4}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), \delta_{3,4}^{(2)} P_1(X_3, X_4)).$$

Proof. By the formula proved in Proposition 4.1 where we specialize the polynomial P_2 to the polynomial P_1 and the polynomial P_3 to the polynomial P_2 , we obtain after some manipulations

$$(U_{0,0}^{(1)})^{d_1 d_2} \text{Res}_{X_2}(D_1, R_{12}) = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_1(X_4), P_2(X_4)) \\ = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), (X_4 - X_3)\delta_{3,4} P_1) \\ (4.1) \quad = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_3)) \times \\ \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), \delta_{3,4}^{(2)}(P_1)).$$

Using formula (2.6) we can push forward this computation and get

$$(U_{0,0}^{(1)})^{d_1 d_2} \text{Res}_{X_2}(D_1, R_{12}) = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_3)) \times \\ \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), (X_4 - X_3)\delta_{3,4}^{(2)} P_1) \\ (4.2) \quad = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_3))^2 \times \\ \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), \delta_{3,4}^{(2)} P_1).$$

□

Lemma 4.6. *Suppose that $d_1 \geq 3$. In \mathbb{U} , we have the equality*

$$\text{Res}_{X_1:\dots:X_4}(P_1, \partial_3 P_1, \delta_{3,4}^{(2)} P_1, P_2(X_4)) = (U_{0,0}^{(1)})^{d_1 d_2} \mathfrak{T}(P_1, P_2)$$

where $\mathfrak{T}(P_1, P_2)$ is an irreducible polynomial in \mathbb{U} of bi-degree

$$((3d_1 - 1)(d_1 - 2)d_2, d_1(d_1 - 1)(d_1 - 2))$$

in the coefficients of (P_1, P_2) .

Proof. We will denote by \mathcal{R} the above resultant. We first use a geometric argument to justify that \mathcal{R} is the product of a certain power of the coefficient $U_{0,0}^{(1)}$ and a certain power of an irreducible polynomial that we will denote $\mathfrak{T}(P_1, P_2) \in \mathbb{U}$.

We rewrite the polynomial P_1 as

$$P_1(X_3) = U_{0,0}^{(1)} X_3^{d_1} + c_1 X_3^{d_1-1} + \cdots + c_{d_1-2} X_3^2 + c_{d_1-1} X_3 + c_{d_1}$$

where the c_i 's are homogeneous polynomials in $\mathbb{U}[X_1, X_2]$ of degree i respectively; we have

$$\partial_3 P = d_1 U_{0,0}^{(1)} X_3^{d_1-1} + (d_1 - 1) c_1 X_3^{d_1-2} + \cdots + 2c_{d_1-2} X_3 + c_{d_1-1}$$

and

$$\delta_{3,4}^{(2)} P_1 = U_{0,0}^{(1)} \left(\frac{(\sum_{i=0}^{d_1-1} X_3^i X_4^{d_1-1-i}) - d_1 X_3^{d_1}}{X_4 - X_3} \right) + \cdots + c_{d_1-3} (X_4 + 2X_3) + c_{d_1-2}.$$

Embedding \mathbb{Z} into the algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} , the variety defined by the equation $\mathcal{R} = 0$ is the projection of the incidence variety

$$\mathcal{W} := \{(x_1 : x_2 : x_3 : x_4) \times (u_{i,j}^{(k)}) \in \mathbb{P}_{\overline{\mathbb{Q}}}^3 \times \mathbb{A}_{\overline{\mathbb{Q}}}^N \text{ such that}$$

$$P_1(x_3) = \partial_3 P_1(x_3) = \delta_{3,4}^{(2)} P_1(x_3, x_4) = P_2(x_4) = 0\}$$

(where $\mathbb{A}_{\overline{\mathbb{Q}}}^N$ denotes the affine space whose coordinates are the $N := (d_1 + 2)(d_1 + 1) + (d_2 + 2)(d_2 + 1)$ indeterminate coefficients, over $\overline{\mathbb{Q}}$) by the canonical projection on the second factor

$$\pi_2 : \mathcal{W} \subset \mathbb{P}^3 \times \mathbb{A}^N \rightarrow \mathbb{A}^N.$$

Consider the canonical projection of \mathcal{W} onto the first factor $\pi_1 : \mathcal{W} \rightarrow \mathbb{P}^3$, which is surjective, and denote by D the line in \mathbb{P}^3 which is defined by the equations $X_1 = 0$ and $X_2 = 0$. On the one hand, we observe that for all $x \in \mathbb{P}^3 \setminus D$ the fiber $\pi_1^{-1}(x)$ is a linear space of codimension 4 in \mathbb{A}^N ; therefore the algebraic closure of $\mathcal{W}|_{\mathbb{P}^3 \setminus D}$ is an irreducible variety of dimension $N - 1$ in $\mathbb{P}^3 \times \mathbb{A}^N$ whose projection by π_2 gives an irreducible hypersurface in \mathbb{A}^N . We denote by $\mathfrak{T}(P_1, P_2)$ a defining equation of this irreducible hypersurface. On the other hand, for all $x \in D$ the fiber $\pi_1^{-1}(x)$ is always included in the linear space of equation $U_{0,0}^{(1)} = 0$ (it is actually exactly this linear space if $x = (0 : 0 : 1 : 0)$ and the linear space $U_{0,0}^{(1)} = U_{0,0}^{(2)} = 0$ otherwise). Consequently, since $\pi_2(\mathcal{W}) = \{U_{0,0}^{(1)} = 0\} \cup \pi_2(\overline{\mathcal{W}|_{\mathbb{P}^3 \setminus D}})$ we deduce that \mathcal{R} is of the form

$$\mathcal{R} = c(d_1, d_2) (U_{0,0}^{(1)})^{a(d_1, d_2)} \mathfrak{T}(P_1, P_2)^{r(d_1, d_2)} \in \mathbb{U}$$

where $c(d_1, d_2)$, $a(d_1, d_2)$, $r(d_1, d_2)$ are constants that we have to determine (note that $\mathfrak{T}(P_1, P_2)$ can be chosen in \mathbb{U} because we know that the variety defined by $\mathcal{R} = 0$ can be defined by an equation in \mathbb{U} since it is a resultant of polynomials in \mathbb{U}).

Now, we will prove by induction on the integer $d := d_1 + d_2 \geq 4$ that

$$(4.3) \quad \mathcal{R} = (U_{0,0}^{(1)})^{d_1 d_2} \mathfrak{T}(P_1, P_2) \in \mathbb{U}$$

that is, for all integers d_1, d_2 such that $d_1 \geq 3$ and $d_2 \geq 1$, we have $c(d_1, d_2) = r(d_1, d_2) = 1$ and $a(d_1, d_2) = d_1 d_2$.

First, by Lemma 4.5 and using the computation (4.2), we note that we always have $c(d_1, d_2) \geq d_1 d_2$.

We check by hand (or with a computer) that the induction hypothesis (4.3) is true for $d = 4$, i.e. $d_1 = 3$ and $d_2 = 1$ and we assume that $d \geq 5$. If $d_2 \geq 2$ we specialize P_2 to the product of two generic homogeneous polynomials, say P'_2 and P''_2 ; then $\mathcal{R}(P_1, P_2)$ specializes, by multiplicativity of the resultants, to $(U_{0,0}^{(1)})^{d_1 d_2} \mathcal{R}(P_1, P'_2) \mathcal{R}(P_1, P''_2)$. Since each irreducible factor of $\mathcal{R}(P_1, P_2)$ must depend on P'_2 and P''_2 and since we already know that $c(d_1, d_2) \geq d_1 d_2$, we deduce that \mathcal{R} satisfies (4.3) for all couple (d_1, d_2) such that $d_2 \geq 2$.

We now turn to the case where $d_1 \geq 4$ and $d_2 = 1$. Consider the homogeneous specialization ϕ which sends P_1 to $X_3 Q_1$ where Q_1 is the generic homogeneous polynomial of degree $d_1 - 1$. Using the properties of resultants we get

$$\begin{aligned} \phi(\mathcal{R}) &= \text{Res}(X_3 Q_1(X_3), Q_1(X_3) + X_3 \partial_3 Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), P_2(X_4)) \\ &= \text{Res}(X_3, Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), P_2(X_4)) \times \\ &\quad \text{Res}(Q_1(X_3), X_3 \partial_3 Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), P_2(X_4)) \\ &= \text{Res}(X_3, Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), P_2(X_4))^2 \times \\ &\quad \text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), P_2(X_4)). \end{aligned}$$

And since, by (2.9), $\delta_{3,4}^{(2)}(X_3 Q_1) = \partial_3 Q_1(X_3) + X_4 \delta_{3,4}^{(2)} Q_1$, we deduce that

$$\begin{aligned} \phi(\mathcal{R}) &= \text{Res}(X_3, Q_1(X_3), \delta_{3,4}^{(2)}(X_3 Q_1), \\ &\quad P_2(X_4))^2 \text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), X_4, P_2(X_4)) \times \mathcal{R}(Q_1, P_2). \end{aligned}$$

Moreover, (2.6) implies that $\delta_{3,4}^{(2)}(X_3 Q_1) = \partial_3 Q_1(X_3) + X_4 \delta_{3,4}^{(2)} Q_1$ evaluated at $X_3 = 0$ is equal to

$$\delta Q_1(0, X_4) = \frac{Q_1(X_4) - Q_1(0)}{X_4}.$$

So finally, we deduce that

$$\begin{aligned} \phi(\mathcal{R}) &= (U_{0,0}^{(1)})^{(d_1-1)} \text{Res}(Q_1(X_1, X_2, 0), \delta Q_1(X_1, X_2, 0, X_4), P_2(X_1, X_2, X_4))^2 \\ &\quad \text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), P_2(X_1, X_2, 0)) \mathfrak{I}(Q_1, P_2). \end{aligned}$$

Since $\mathfrak{I}(Q_1, P_2)$ is irreducible (by our induction hypothesis), it follows that $r(d_1, d_2)$ must equal 1 for each irreducible factor of the above specialization must appear to a power which is a multiple of $r(d_1, d_2)$. Also, since $P_2(X_1, X_2, 0)$ is a generic linear form in X_1 and X_2 it turns out that $\text{Res}(Q_1(X_3), \partial_3 Q_1(X_3), P_2(X_1, X_2, 0))$ is, up to a linear change of coordinates, a discriminant of a univariate polynomial: it equals $U_{0,0}^{(1)}$ times an irreducible polynomial in \mathbb{U} . Moreover, it is easy to see that $U_{0,0}^{(1)}$ does not divide $\text{Res}(Q_1(X_1, X_2, 0), \delta Q_1(X_1, X_2, 0, X_4), P_2(X_1, X_2, X_4))$ (for this resultant does not vanish under this condition). Therefore, we deduce that $a(d_1, d_2) \leq d_1 d_2$ and then that $a(d_1, d_2) = d_1 d_2$. Finally, the three resultants in the above specialization formula are clearly primitive² (either by the induction hypothesis for the last one or either because it stays primitive after the change of coordinate induced by the linear polynomial P_2) and it follows that $c(d_1, d_2) = 1$.

The formula on the degree is obtained as a direct consequence of the known degree formula for the resultants of several homogeneous polynomials. \square

²the gcd of their coefficients is 1

Proposition 4.7. *Suppose that $d_1 \geq 2$. We set*

$$\begin{aligned} D_1 &:= \text{Disc}_{X_3}(P_1(1, X_2, X_3)) \in \mathbb{U}, \\ R_{12} &:= \text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)) \in \mathbb{U}. \end{aligned}$$

Then the following equality holds in \mathbb{U} :

$$\text{Res}_{X_2}(D_1, R_{12}) = \text{Res}_{X_1: X_2: X_3}(P_1, P_2, \partial_3 P_1)^2 \mathfrak{T}(P_1, P_2)$$

where the irreducible polynomial $\mathfrak{T}(P_1, P_2)$ has been defined in Lemma 4.6 for $d_1 \geq 3$; if $d_1 = 2$ we set $\mathfrak{T}(P_1, P_2) := 1 \in \mathbb{U}$.

Moreover, $\text{Res}_{X_1: X_2: X_3}(P_1, P_2, \partial_3 P_1) \in \mathbb{U}$ is irreducible and the iterated resultant $\text{Res}_{X_2}(D_1, R_{12})$ is bi-homogeneous w.r.t. the coefficients $(U_{i,j}^{(1)})_{i,j}$ and $(U_{i,j}^{(2)})_{i,j}$ of bi-degree $(3d_1 d_2 (d_1^2 - 1), d_1^2 (d_1 - 1))$.

Proof. First, the classical multi-degree formula for resultants gives the claimed result for the bi-degree of the iterated resultant $\text{Res}_{X_2}(D_1, R_{12})$. By Lemma 4.5, we have

$$(4.4) \quad (U_{0,0}^{(1)})^{d_1 d_2} \text{Res}_{X_2}(D_1, R_{12}) = \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_3))^2 \times \\ \text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), \delta_{3,4}^{(2)} P_1)$$

and by Lemma 4.6, we know that

$$\text{Res}(P_1(X_3), \partial_3 P_1(X_3), P_2(X_4), \delta_{X_3, X_4}^{(2)} P_1) = (U_{0,0}^{(1)})^{d_1 d_2} \mathfrak{T}(P_1, P_2),$$

where $\mathfrak{T}(P_1, P_2)$ is an irreducible polynomial, which implies the claimed formula.

To conclude the proof, it only remains to prove the irreducibility of the resultant

$$\mathcal{R} := \text{Res}_{X_1: X_2: X_3}(P_1, P_2, \partial_3 P_1).$$

We proceed as we already did several times: by induction on the integer $d := d_1 + d_2 \geq 3$. We check by hand that \mathcal{R} is irreducible if $d_1 = 2$ and $d_2 = 1$ and suppose that $d \geq 4$. If $d_2 \geq 2$ then one specializes P_2 to a product of two generic forms and we conclude using the multiplicativity property of the resultant. Otherwise, we have $d_1 \geq 3$ and one specializes P_1 to $L_1 Q_1$ where Q_1 is the generic homogeneous polynomial of degree $d_1 - 1$ and $L_1 := aX_1 + bX_2 + cX_3$ is the generic linear form; this sends \mathcal{R} to

$$(-1)^{d_1 d_2} c^{d_2} \text{Res}(Q_1, P_2, \partial_3 Q_1) \text{Res}(Q_1, P_2, L_1)^2$$

where $\text{Res}(Q_1, P_2, \partial_3 Q_1)$ is irreducible by our induction hypothesis and also where $\text{Res}(Q_1, P_2, L_1)$ is irreducible for it is the resultant of three generic polynomials. Examining the degrees in the coefficients of Q_1 and L_1 of the above factors and using the fact that each irreducible factor of \mathcal{R} must specialize to a polynomial having the same degree in the coefficients of Q_1 and L_1 , we deduce that \mathcal{R} is irreducible. \square

A specialization of this proposition gives the following result which covers and precises [26, Theorem 3.3].

Corollary 4.8. *Given two polynomials $f_k(\mathbf{x}, y, z), k = 1, 2$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha, i, j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where $d_1, d_2 \geq 2$, \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, then the iterated resultant

$$\text{Res}_y(\text{Disc}_z(f_1), \text{Res}_z(f_1, f_2)) \in \mathbb{S}[\mathbf{x}]$$

is of degree at most $d_1^2 d_2 (d_1 - 1)$ in \mathbf{x} and we have

$$\begin{aligned} \text{Res}_y(\text{Disc}_z(f_1), \text{Res}_z(f_1, f_2)) = \\ \text{Res}_{y,z}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z))^2 \mathfrak{T}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z)) \end{aligned}$$

where we recall that, in $\mathbb{S}[x]$, we have $\mathfrak{T}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z)) = 1$ if $d_1 = 2$ and otherwise

$$\begin{aligned} (a_{0,0,d}^{(1)})^{d_1 d_2} \mathfrak{T}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z)) = \\ \text{Res}_{y,z,z'}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z'), \delta_{z,z'}^{(2)} f_1(\mathbf{x}, y, z, z')) \end{aligned}$$

with

$$\delta_{z,z'}^{(2)} f_1(\mathbf{x}, y, z, z') := \frac{\delta_{z,z'}(f_1)(\mathbf{x}, y, z, z') - \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z)}{z' - z} \in \mathbb{S}[\mathbf{x}][y, z, z'].$$

Moreover, if the polynomials f_1, f_2, f_3 are sufficiently generic then this iterated resultant has exactly degree $d_1^2 d_2 (d_1 - 1)$ in \mathbf{x} and both

$$\text{Res}_{y,z}(f_1(\mathbf{x}, y, z), f_2(\mathbf{x}, y, z), \frac{\partial f_1}{\partial z}(\mathbf{x}, y, z))$$

and $\mathfrak{T}(f_1, f_2)$ are irreducible polynomials.

5. DISCRIMINANT OF A RESULTANT

In this section, we suppose given two positive integers $d_1, d_2 \geq 2$ and two homogeneous polynomials

$$P_k(X_1, X_2, X_3) := \sum_{0 \leq i, j; i+j \leq d_k} U_{i,j}^{(k)} X_1^i X_2^j X_3^{d_k - i - j} \in \mathbb{U}[X_1, X_2, X_3], \quad k = 1, 2,$$

where, as usual, \mathbb{U} denotes the universal ring of coefficients $\mathbb{Z}[U_{i,j}^{(k)}]$. We will hereafter focus on the factorization of a discriminant of a resultant.

Lemma 5.1. *In $\mathbb{U}[X_2]$, setting $X_1 = 1$, we have*

$$\partial_2 \text{Res}_{X_3}(P_1, P_2) = (-1)^{d_1 + d_2} \begin{vmatrix} \partial_2 P_1 & \partial_3 P_1 \\ \partial_2 P_2 & \partial_3 P_2 \end{vmatrix} \text{SRes}_{X_3}^{(1)}(P_1, P_2) \text{ modulo } (P_1, P_2).$$

Proof. Introduce a new indeterminate U and set

$$\begin{aligned} P_1(1, X_2, X_3 + U) &= a_{d_1} X_3^{d_1} + a_{d_1-1} X_3^{d_1-1} + \dots + a_1 X_3 + a_0, \\ P_2(1, X_2, X_3 + U) &= b_{d_2} X_3^{d_2} + b_{d_2-1} X_3^{d_2-1} + \dots + b_1 X_3 + b_0, \end{aligned}$$

where the a_i 's and the b_j 's are polynomials in $\mathbb{U}[X_2, U]$. Expanding the resultant

$$\text{Res}_{X_3}(P_1, P_2) = \begin{vmatrix} a_{d_1} & 0 & \cdots & 0 & b_{d_2} & 0 & 0 \\ a_{d_1-1} & a_{d_1} & & \vdots & b_{d_2-1} & \ddots & 0 \\ \vdots & & \ddots & 0 & \vdots & & b_{d_2} \\ a_0 & & & a_{d_1} & b_1 & & b_{d_2-1} \\ 0 & a_0 & & a_{d_1-1} & b_0 & & \vdots \\ \vdots & & \ddots & \vdots & 0 & \ddots & b_1 \\ 0 & \cdots & 0 & a_0 & 0 & 0 & b_0 \end{vmatrix}$$

with respect to its two last rows, we get

$$\text{Res}_{X_3}(P_1, P_2) = (-1)^{d_1+d_2} \begin{vmatrix} a_0 & b_0 \\ a_1 & b_1 \end{vmatrix} \text{SRes}_{X_3}^{(1)}(P_1, P_2) + a_0^2 \Delta_1 + a_0 b_0 \Delta_2 + b_0^2 \Delta_3,$$

where Δ_i ($i = 1, 2, 3$) are polynomials in the a_i 's and b_j 's. Taking the derivative with respect to the variable X_2 , we deduce that

(5.1)

$$\partial_2 \text{Res}_{X_3}(P_1, P_2) = (-1)^{d_1+d_2} \begin{vmatrix} \partial_2 a_0 & \partial_2 b_0 \\ a_1 & b_1 \end{vmatrix} \text{SRes}_{X_3}^{(1)}(P_1, P_2) \text{ modulo } (a_0, b_0).$$

Now, it is easy to check that we have

$$a_0 = P_1(1, X_2, U), \quad b_0 = P_2(1, X_2, U), \quad a_1 = \partial_3 P_1(1, X_2, U) \quad b_1 = \partial_3 P_2(1, X_2, U).$$

Moreover, by invariance property (change of bases formula)

$$\begin{aligned} \text{Res}_{X_3}(P_1(1, X_2, X_3 + U), P_2(1, X_2, X_3 + U)) &= \\ \text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)) &= \text{Res}_U(P_1(1, X_2, U), P_2(1, X_2, U)) \end{aligned}$$

and the same is true for the subresultant $\text{SRes}_{X_3}^{(1)}(P_1, P_2)$. Therefore we deduce that (5.1) is nothing but the claimed equality by substituting X_3 by U . \square

This lemma implies the following factorization.

Proposition 5.2. *In \mathbb{U} , we have the equality*

$$\begin{aligned} \text{Disc}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))) &= (-1)^{(d_2+1)(d_1 d_2 + d_1 - 1)} \times \\ &\quad \text{Res}_{X_1: X_2: X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) \times \text{Disc}_{X_1: X_2: X_3}(P_1, P_2). \end{aligned}$$

This iterated resultant is bi-homogeneous with respect to the sets of coefficients $U_{i,j}^{(1)}$ and $U_{i,j}^{(2)}$ respectively of bi-degree $(2d_2(d_1 d_2 - 1); 2d_1(d_1 d_2 - 1))$.

Proof. Set $\mathcal{R}_0 := \text{Res}(P_1, P_2, X_1)$, $J_1 := \begin{vmatrix} \partial_2 P_1 & \partial_3 P_1 \\ \partial_2 P_2 & \partial_3 P_2 \end{vmatrix}$ and recall that, by definition, we have $\mathcal{R}_0 \text{Disc}_{X_1: X_2: X_3}(P_1, P_2) = \text{Res}(P_1, P_2, J_1)$. Setting

$$\mathcal{D} := \text{Disc}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))) \in \mathbb{U},$$

it follows from (2.3) that

$$\begin{aligned} \mathcal{R} &:= \mathcal{R}_0 \mathcal{D} = \\ \text{Res}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)), \partial_2 \text{Res}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))) & \end{aligned}$$

Now, using Lemma 5.1 we deduce that \mathcal{R} belongs to the ideal

$$(P_1(1, X_2, X_3), P_2(1, X_2, X_3), J_1(1, X_2, X_3) \text{SRes}_{X_3}^{(1)}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)))$$

which gives, after homogenization by X_1 , the existence of an integer $N \geq 1$ such that

$$X_1^N \mathcal{R} \in (P_1(X_1, X_2, X_3), P_2(X_1, X_2, X_3), J_1(X_1, X_2, X_3) \text{SRes}_{X_3}^{(1)}(P_1(X_1, X_2, X_3))).$$

Therefore, using the divisibility property (and others) of the resultants we deduce that

$$\begin{aligned} \text{Res}(P_1, P_2, J_1 \text{SRes}_{X_3}^{(1)}(P_1, P_2)) &= \text{Res}(P_1, P_2, J_1) \times \text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) \\ &= \mathcal{R}_0 \times \text{Disc}(P_1, P_2) \times \text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) \end{aligned}$$

divides $\text{Res}(P_1, P_2, X_1^N \mathcal{R}) = \mathcal{R}_0^N \mathcal{R}^{d_1 d_2} = \mathcal{R}_0^{N+d_1 d_2} \mathcal{D}^{d_1 d_2}$ in \mathbb{U} . We know that \mathcal{R}_0 and $\text{Disc}(P_1, P_2)$ are irreducible polynomials in \mathbb{U} ; just by comparing their degree we see that \mathcal{R}_0 does not divide $\text{Disc}(P_1, P_2)$. Also, just by looking to the defining matrix of the subresultant, we have

$$\text{SRes}_{X_3}^{(1)}(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2}) = (-X_2)^{(d_1-1)(d_2-1)}$$

so that $\mathcal{R}_0(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2}) = \text{Res}(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2}, X_1) = 0$ in \mathbb{U} and also

$$\begin{aligned} \text{Res}(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2}, \text{SRes}_{X_3}^{(1)}(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2})) &= \\ \text{Res}(X_1^{d_1} + X_2^{d_1-1} X_3, X_3^{d_2}, (-X_2)^{(d_1-1)(d_2-1)}) &= \\ = \text{Res}(X_1, X_3, -X_2)^{d_1 d_2 (d_1-1)(d_2-1)} &= 1. \end{aligned}$$

Since \mathcal{R}_0 is irreducible, it does not divide $\text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$ and therefore

$$\text{Disc}(P_1, P_2) \text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) \text{ divides } \mathcal{D} \text{ in } \mathbb{U}.$$

By known degree properties, \mathcal{D} has degree $d_2(2d_1 d_2 - 2)$ in the coefficients of the polynomial P_1 and that the product

$$\text{Disc}(P_1, P_2) \text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$$

has degree

$$d_2(2(d_1 - 1) + d_2 - 1) + d_2(d_1 - 1)(d_2 - 1) + d_1 d_2 (d_2 - 1) = d_2(2d_1 d_2 - 2)$$

(note that $\text{SRes}_{X_3}^{(1)}(P_1, P_2)$ is an homogeneous polynomial in X_1, X_2 of degree $(d_1 - 1)(d_2 - 1)$ which is also homogeneous in the coefficients of P_1 , resp. P_2 , of degree $d_2 - 1$, resp. $d_1 - 1$) With a similar computation for the degree with respect to the coefficients of P_2 , we deduce that \mathcal{D} and the product

$$\text{Disc}(P_1, P_2) \text{Res}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$$

are equal in \mathbb{U} up to a non-zero element in \mathbb{Z} , that we denote $c(d_1, d_2)$.

To finish the proof, it remains to determine $c(d_1, d_2) \in \mathbb{U}$ using, as usual, a suitable specialization. We choose the specialization ϕ such that

$$\phi(P_1) = X_3^{d_1} + AX_1^{d_1-1} X_2 + BX_1^{d_1} \text{ and } \phi(P_2) = X_1^{d_2-1} X_3 - X_2^{d_2}.$$

It is easy to compute

$$\text{Res}_{X_3}(X_3^{d_1} + AX_2 + B, X_3 - X_2^{d_2}) = (-1)^{d_1} (X_2^{d_1 d_2} + AX_2 + B)$$

and hence to deduce that

$$\begin{aligned} \text{Disc}_{X_2}(\text{Res}_{X_3}(\phi(P_1)(1, X_2, X_3), \phi(P_2)(1, X_2, X_3))) &= \\ \text{Disc}_{X_2}((-1)^{d_1}(X_2^{d_1 d_2} + AX_2 + B)) &= \\ \text{Res}_{X_2}(X_2^{d_1 d_2} + AX_2 + B, d_1 d_2 X_2^{d_1 d_2 - 1} + A). \end{aligned}$$

Denoting \mathcal{D} the above quantity, we have

$$\begin{aligned} (d_1 d_2)^{d_1 d_2 - 1} \mathcal{D} &= \text{Res}_{X_2}(d_1 d_2 X_2^{d_1 d_2} + d_1 d_2 AX_2 + d_1 d_2 B, d_1 d_2 X_2^{d_1 d_2 - 1} + A) \\ &= \text{Res}_{X_2}((d_1 d_2 - 1)AX_2 + d_1 d_2 B, d_1 d_2 X_2^{d_1 d_2 - 1} + A) \end{aligned}$$

and finally

$$(d_1 d_2)^{d_1 d_2 - 1} \mathcal{D} = (-1)^{d_1 d_2 - 1} (d_1 d_2)^{d_1 d_2 - 1} \begin{vmatrix} (d_1 d_2 - 1)A & 0 & \cdots & 0 & d_1 d_2 \\ d_1 d_2 B & (d_1 d_2 - 1)A & \cdots & \cdots & 0 \\ 0 & d_1 d_2 B & \vdots & 0 & \cdots \\ \vdots & \cdots & \vdots & (d_1 d_2 - 1)A & 0 \\ 0 & \cdots & 0 & d_1 d_2 B & A \end{vmatrix},$$

that is to say,

$$\mathcal{D} = (-1)^{d_1 d_2 - 1} (d_1 d_2 - 1)^{d_1 d_2 - 1} U^{d_1 d_2} + (d_1 d_2)^{d_1 d_2} B^{d_1 d_2 - 1}.$$

Now, since two consecutive integers are always relatively prime, we deduce that $c(d_1, d_2) = \pm 1 \in \mathbb{Z}$. To determine exactly this integer, we compute the other side of the claimed equality. For simplicity, we will consider the specialization ψ which is similar to the specialization ϕ with in addition $A = 0$. We have

$$\begin{aligned} &\text{Disc}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3 - X_2^{d_2}) \\ &= \frac{\text{Res}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3 - X_2^{d_2}, d_1 d_2 X_2^{d_2 - 1} X_3^{d_1 - 1})}{\text{Res}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3 - X_2^{d_2}, X_1)} \\ &= (-1)^{d_1(d_2 + 1)} (d_1 d_2)^{d_1 d_2} \text{Res}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3)^{d_2 - 1} \text{Res}(BX_1^{d_1}, -X_2^{d_2})^{d_1 - 1} \\ &= (-1)^{d_1(d_2 + 1)} (d_1 d_2)^{d_1 d_2} (-1)^{d_1(d_2 - 1)^2} B^{d_2 - 1} B^{d_2(d_1 - 1)} \\ &= (-1)^{d_1(d_2 + 1) + d_1(d_2 - 1)^2} (d_1 d_2)^{d_1 d_2} B^{d_1 d_2 - 1}. \end{aligned}$$

It is not hard to see from the definition of the subresultant that

$$\text{SRes}_{X_3}^{(1)}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3 - X_2^{d_2}) = X_1^{(d_1 - 1)(d_2 - 1)}$$

and also to compute

$$\text{Res}_{X_1: X_2: X_3}(X_3^{d_1} + BX_1^{d_1}, X_1^{d_2 - 1} X_3 - X_2^{d_2}, X_1^{(d_1 - 1)(d_2 - 1)}) = (-1)^{(d_1 - 1)(d_2 - 1)}.$$

Gathering all these specializations, we obtain $c(d_1, d_2) = (-1)^{(d_2 + 1)(d_1 d_2 + d_1 - 1)}$, as claimed. \square

At this point, the factorization given in the above proposition is not complete since we only know that one factor is irreducible. The following result shows that the second factor is not irreducible, but is the square of an irreducible polynomial and moreover that it can be interpreted as a particular iterated resultant itself.

Lemma 5.3. *Introducing a new indeterminate X_4 , we have the following equalities in \mathbb{U} :*

$$\begin{aligned} & \text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) = \\ & \text{Res}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)), \text{SRes}_{X_3}^{(1)}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))) \\ & = \text{Res}_{X_1:X_2:X_3:X_4}(P_1, \delta_{3,4}P_1, P_2, \delta_{3,4}P_2). \end{aligned}$$

Proof. First, for simplicity we set

$$\begin{aligned} \mathcal{R}_1 := & \text{Res}_{X_2, X_3}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)), \\ & \text{SRes}_{X_3}^{(1)}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))). \end{aligned}$$

We know that $\text{Res}_{X_3}(P_1, P_2)$ has degree d_1d_2 in X_2 and bi-degree (d_2, d_1) in \mathbb{U} , and that $\text{SRes}_{X_3}^{(1)}(P_1, P_2)$ has degree $(d_1 - 1)(d_2 - 1)$ in X_2 and bi-degree $(d_2 - 1, d_1 - 1)$ in \mathbb{U} . It follows that \mathcal{R}_1 has bi-degree $((2d_1 - 1)d_2(d_2 - 1), (2d_2 - 1)d_1(d_1 - 1))$ which is exactly the same than the bi-degree of $\text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$ (by a straightforward computation). Moreover, we have

$$\mathcal{R}_1 \in (P_1(1, X_2, X_3), P_2(1, X_2, X_3), \text{SRes}_{X_3}^{(1)}(P_1(1, X_2, X_3), P_2(1, X_2, X_3)))$$

which implies, after homogenization by X_1 and a suitable use of the divisibility property of the resultants that

$$\text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) \text{ divides } \text{Res}_{X_1:X_2:X_3}(P_1, P_2, X_1)^N \mathcal{R}_1^{d_1d_2}$$

in \mathbb{U} , where N denotes a positive integer. We have already seen in the proof of Proposition 5.2, that $\mathcal{R}_0 := \text{Res}(P_1, P_2, X_1)$ is irreducible and does not divide $\text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$. It follows that the later divides \mathcal{R} and since they have the same bi-degree in \mathbb{U} we deduce that they equal up to multiplication by a constant. To determine this constant, we consider the specialization $P_1 = X_1^{d_1} + X_2^{d_1-1}X_3$ and $P_2 = X_3^{d_2}$ for which we have already seen that $\text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)) = 1$ in the proof of Proposition 5.2. Since we also have

$$\mathcal{R}_1(P_1, P_2) = \text{Res}_{X_2}((-X_1)^{d_1d_2}, (-X_2)^{(d_1-1)(d_2-1)}) = 1.$$

we deduce that $\mathcal{R}_1 = \text{Res}_{X_1:X_2:X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$ in \mathbb{U} .

We now turn to the proof of the third claimed equality. Introduce a new indeterminate U and define

$$\begin{aligned} P_1(X_1, X_2, X_3 + U) &= a_{d_1}X_3^{d_1} + a_{d_1-1}X_3^{d_1-1} + \cdots + a_1X_3 + a_0, \\ P_2(X_1, X_2, X_3 + U) &= b_{d_2}X_3^{d_2} + b_{d_2-1}X_3^{d_2-1} + \cdots + b_1X_3 + b_0, \end{aligned}$$

where the a_i 's and the b_j 's are polynomials in $\mathbb{U}[X_1, X_2, U]$. The subresultant $\text{SRes}_{X_3}^{(1)}(P_1, P_2)$ is defined as the determinant of the matrix

$$\begin{pmatrix} a_{d_1} & 0 & \cdots & 0 & 0 & b_{d_2} & 0 & 0 & 0 \\ a_{d_1-1} & a_{d_1} & & \vdots & \vdots & b_{d_2-1} & \ddots & 0 & 0 \\ \vdots & a_{d_1-1} & \ddots & 0 & 0 & \vdots & & b_{d_2} & 0 \\ a_2 & & \ddots & a_{d_1} & 0 & \vdots & & b_{d_2-1} & b_{d_2} \\ a_1 & a_2 & & a_{d_1-1} & a_{d_1} & b_2 & & \vdots & b_{d_2-1} \\ a_0 & a_1 & \ddots & \vdots & a_{d_1-1} & b_1 & & \vdots & \vdots \\ 0 & \ddots & \ddots & a_2 & \vdots & b_0 & \ddots & b_2 & \vdots \\ \vdots & & \ddots & a_1 & a_2 & 0 & \ddots & b_1 & b_2 \\ 0 & \cdots & 0 & a_0 & a_1 & 0 & 0 & b_0 & b_1 \end{pmatrix},$$

determinant which remains unchanged if we add, for all $i = 1, \dots, d_1 + d_2 - 3$, the line number i times

$$\frac{X_3^{d_1+d_2-1-i} - X_4^{d_1+d_2-1-i}}{X_3 - X_4} = \sum_{0 \leq i, j; i+j=d_1+d_2-2-i} X_3^i X_4^j$$

to the last line which then becomes of the form

$$(0 \quad \cdots \quad 0 \quad a_0 R_1 \quad \delta_{3,4}(P_1) \quad 0 \quad \cdots \quad 0 \quad b_0 R_2 \quad \delta_{3,4}(P_2))$$

where R_1 and R_2 are polynomials in $\mathbb{U}[X_1, X_2, X_3, X_4, U]$. It follows that, by developing this determinant with respect to the last line,

$$\begin{aligned} \text{SRes}_{X_3}^{(1)}(P_1(X_1, X_2, X_3 + U), P_2(X_1, X_2, X_3 + U)) \\ \in (a_0, \delta_{3,4}(P_1), b_0, \delta_{3,4}(P_2)) \subset \mathbb{U}[X_1, X_2, X_3, X_4]. \end{aligned}$$

Moreover, $a_0 = P_1(X_1, X_2, U)$, $b_0 = P_2(X_1, X_2, U)$ and by invariance of the subresultant under the change of coordinates $U \leftarrow X_3 + U$ we deduce that

$$\begin{aligned} \text{SRes}_{X_3}^{(1)}(P_1(X_1, X_2, U), P_2(X_1, X_2, U)) \\ \in (P_1(U), \delta_{3,4}P_1(U, X_4), P_2(U), \delta_{3,4}P_2(U, X_4)) \end{aligned}$$

which implies, after a substitution of U by X_3 and a suitable use of the divisibility property of the resultants, that

$$\begin{aligned} \text{Res}_{X_1: \dots : X_4}(P_1(X_3), \delta_{3,4}P_1(X_3, X_4), P_2(X_3), \delta_{3,4}P_2(X_3, X_4)) \\ \text{divides } \text{Res}_{X_1: X_2: X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2)). \end{aligned}$$

An easy computation shows that these two resultants have the same bi-degree w.r.t. the coefficients of P_1 and P_2 ; therefore they are equal up to sign in \mathbb{U} (we have already seen that

$$\text{Res}_{X_1: X_2: X_3}(P_1, P_2, \text{SRes}_{X_3}^{(1)}(P_1, P_2))$$

is a primitive polynomial in \mathbb{U} through a particular specialization). To determine the sign we consider again the specialization $P_1 = X_1^{d_1} + X_2^{d_1-1}X_3$ and $P_2 = X_3^{d_2}$ for which it is easy to see that both resultants then specialize to 1. \square

The following result can be seen as the main explanation of [26, theorem 3.4].

Proposition 5.4. *Introducing a new indeterminate X_4 , there exists a non-zero irreducible polynomial $\mathfrak{D}(P_1, P_2)$ in \mathbb{U} such that*

$$\text{Res}_{X_1, \dots, X_4}(P_1, \delta_{3,4}(P_1), P_2, \delta_{3,4}(P_2)) = \mathfrak{D}(P_1, P_2)^2.$$

It is bi-homogeneous with respect to the set of coefficients $(U_{i,j}^{(1)})_{i,j}$ and $(U_{i,j}^{(2)})_{i,j}$ of bi-degree

$$\left(\frac{(2d_1 - 1)d_2(d_2 - 1)}{2}, \frac{(2d_2 - 1)d_1(d_1 - 1)}{2} \right).$$

Proof. Let us denote by \mathfrak{R} the above resultant. Embedding \mathbb{Z} into the algebraic closure of \mathbb{Q} , the variety defined by the equation $\mathfrak{R} = 0$ is the projection of the incidence variety

$$\mathcal{W} := \{(x_1 : x_2 : x_3 : x_4) \times (u_{i,j}^{(k)}) \in \mathbb{P}^3 \times \mathbb{A}^N \text{ such that } P_1 = \delta_{3,4}(P_1) = P_2 = \delta_{3,4}(P_2) = 0\}$$

(where N is the number of indeterminate coefficients) by the canonical projection on the second factor $\pi_2 : \mathcal{W} \subset \mathbb{P}^3 \times \mathbb{A}^N \rightarrow \mathbb{A}^N$. But for a generic point $(a_{i,j}^{(k)}) \in \mathbb{A}^N$ such that $\mathfrak{R} = 0$, we have at least two pre-image in \mathcal{W} since if $(x_1 : x_2 : x_3 : x_4)$ is such pre-image then $(x_1 : x_2 : x_4 : x_3)$ is also a pre-image (which is generically different). It follows that the co-restriction of π_2 to the variety $\mathfrak{R} = 0$ has degree at least 2 and hence that, in \mathbb{U} ,

$$\mathfrak{R} = c(\mathbf{d}) \times \mathfrak{R}_1^{r_1(\mathbf{d})} \times \mathfrak{R}_2^{r_2(\mathbf{d})} \times \dots \times \mathfrak{R}_p^{r_p(\mathbf{d})}$$

where p is a positive integer, $c(\mathbf{d})$ is also a positive integer but may depend on $\mathbf{d} := (d_1, d_2)$ and $r_1(\mathbf{d}), \dots, r_p(\mathbf{d})$ are positive integers greater or equal to 2 and may also depend on \mathbf{d} . Note that we know by Lemma 5.3 that $c(\mathbf{d}) = 1$. To determine the other quantities we will proceed by induction on $d_1 + d_2 \geq 4$ (remember that $d_1 \geq 2$ and $d_2 \geq 2$). First, we can check by hand (or with a computer) that the claim is true if $d_1 = d_2 = 2$: \mathfrak{R} is the square of an irreducible polynomial in \mathbb{U} of bi-degree (3,3). Now, assume that $d_1 + d_2 > 4$; without loss of generality one may assume that $d_1 \geq 3$. Consider the homogeneous specialization ϕ which sends P_1 to the product $L_1 Q_1$ where Q_1 is generic homogeneous polynomials of degree $d_1 - 1 \geq 2$ and $L_1 := aX_1 + bX_2 + cX_3$ is a generic linear form. We have, using

obvious notations,

$$\begin{aligned}
 (5.2) \quad \phi(\mathfrak{R}) &= \text{Res}(L_1(X_3)Q_1(X_3), Q_1(X_3)\delta_{3,4}(L_1) + L_1(X_4)\delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= \text{Res}(L_1(X_3), cQ_1(X_3) + L_1(X_4)\delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), L_1(X_4)\delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= \text{Res}(L_1(X_3), cQ_1(X_3) + (L_1(X_4) - L_1(X_3))\delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), L_1(X_4), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), \delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= c^{d_2(d_2-1)} \text{Res}(L_1(X_3), Q_1(X_3) + (X_4 - X_3)\delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), L_1(X_4), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), \delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= c^{d_2(d_2-1)} \text{Res}(L_1(X_3), Q_1(X_4), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), L_1(X_4), P_2(X_3), \delta_{3,4}(P_2)) \times \\
 &\quad \text{Res}(Q_1(X_3), \delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= c^{d_2(d_2-1)} \text{Res}(L_1(X_3), Q_1(X_4), P_2(X_3), \delta_{3,4}(P_2))^2 \times \\
 &\quad \text{Res}(Q_1(X_3), \delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2)),
 \end{aligned}$$

since exchanging the role of X_3 and X_4 , we have

$$\begin{aligned}
 &\text{Res}(L_1(X_3), Q_1(X_4), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= \text{Res}(L_1(X_4), Q_1(X_3), P_2(X_4), \delta_{3,4}(P_2)) \\
 &= \text{Res}(L_1(X_4), Q_1(X_3), P_2(X_3), \delta_{3,4}(P_2)) \\
 &= \text{Res}(Q_1(X_3), L_1(X_4), P_2(X_3), \delta_{3,4}(P_2)).
 \end{aligned}$$

Observe that this resultant is irreducible by Proposition 3.3 and that the last one is the square of an irreducible polynomial by our induction hypothesis. Moreover,

- c has bi-degree $(1, 0)$ in terms of the coefficients of L_1 and Q_1 respectively,
- $\text{Res}(L_1(X_4), Q_1(X_3), P_2(X_3), \delta_{3,4}(P_2))$ has also bi-degree $(d_2(d_2 - 1)(d_1 - 1), d_2(d_2 - 1))$ and
- $\text{Res}(Q_1(X_3), \delta_{3,4}(Q_1), P_2(X_3), \delta_{3,4}(P_2))$ is the square of an irreducible polynomial which has bi-degree $(0, d_2(d_2 - 1)(d_1 - 1) - \frac{d_2(d_2-1)}{2})$.

Since the specialization ϕ is homogeneous, each irreducible factor \mathfrak{R}_i of \mathfrak{R} must give through this specialization irreducible polynomial(s) having the same degree with respect to the coefficients of L_1 and Q_1 . With the bi-degree given above, we deduce that \mathfrak{R} can at most have two irreducible factors (and moreover that they specialize to the same polynomial via ϕ). It follows that we have $p = 1$ and $r_1(\mathbf{d}) = 2$. \square

Remark 5.5. Some technical limitations of the theory of *anisotropic resultants* as exposed in [18, 19] prevent the explicit description of the “squareroot” of the above resultant. More precisely, suppose given two sequences of integers (m_1, \dots, m_n) and (d_1, \dots, d_n) such that $m_j | d_i$ for all couple $(i, j) \in \{1, \dots, n\}^2$. If g_1, \dots, g_n are n homogeneous polynomials of degree m_1, \dots, m_n respectively in the graded ring $\mathbb{S}[X_1, \dots, X_n]$ (with $\deg(X_i) = 1$ for all $i = 1, \dots, n$) and f_1, \dots, f_n are n isobaric polynomials of weight d_1, \dots, d_n respectively in the graded ring ${}^a\mathbb{S}[X_1, \dots, X_n]$ (now with $\deg(X_i) = m_i$ for all $i = 1, \dots, n$) then an easy adaptation of the proof

of the base change formula [18, 5.12] shows that

$$\text{Res}(f_1(g_1, \dots, g_n), \dots, f_n(g_1, \dots, g_n)) = {}^a\text{Res}(f_1, \dots, f_n)^\Delta \text{Res}(g_1, \dots, g_n)^{\frac{d_1 \dots d_n}{m_1 \dots m_n}}$$

in \mathbb{S} where $\Delta := \frac{m_1 \dots m_n}{\gcd(m_1, \dots, m_n)}$.

In our case, denoting $\delta^0(P) := P(X_1, X_2, X_3) - P(X_1, X_2, X_4)$, it is easy to see that

$$\text{Res}(\delta^0(P_1), \delta^1(P_1), \delta^0(P_2), \delta^1(P_2)) = 2^{(d_1-1)(d_2-1)(d_1+d_2)} \times \\ \text{Res}(P_1(X_1, X_2, X_3), \delta(P_1)(X_1, X_2, X_3, X_4), P_2(X_1, X_2, X_3), \delta(P_2)(X_1, X_2, X_3, X_4)).$$

And since, for all $i = 1, 2$, $\delta^0(P_i)$ and $\delta^1(P_i)$ are symmetric polynomials with respect to the variables X_3 and X_4 we deduce that there exists four *quasi*-homogeneous polynomials $Q_i(X_1, \dots, X_4)$, $i = 1, \dots, 4$, with $\deg(X_1) = \deg(X_2) = \deg(X_3) = 1$, $\deg(X_4) = 2$ and such that, for instance,

$$Q_1(X_1, X_2, X_3 + X_4, X_3 X_4) = \delta^0(P_1)(X_1, X_2, X_3, X_4).$$

By using the above adapted base change formula, we should obtain

$$\text{Res}(\delta^0(P_1), \delta^1(P_1), \delta^0(P_2), \delta^1(P_2)) = \pm^a \text{Res}(Q_1, Q_2, Q_3, Q_4)^2.$$

It turns out that Q_1, Q_2, Q_3, Q_4 are isobaric of weights $(d_1, d_1 - 1, d_2, d_2 - 2)$ respectively and hence the condition of the existence of their anisotropic resultant as in [19, §2] are not fulfilled. However, in a personal communication Jouanolou informed us that it is possible to extend the theory of anisotropic resultant to our particular setting and conclude to the existence of such an anisotropic resultant.

Gathering the results of this section, we obtain the full factorization of the discriminant of a resultant.

Theorem 5.6. *Introducing a new indeterminate X_4 , we have in \mathbb{U} :*

$$\text{Disc}_{X_2}(\text{Res}_{X_3}(P_1(1, X_2, X_3), P_2(1, X_2, X_3))) = (-1)^{(d_2+1)(d_1 d_2 + d_1 - 1)} \times \\ \text{Disc}_{X_1: X_2: X_3}(P_1, P_2) \mathfrak{D}(P_1, P_2)^2$$

where $\text{Disc}_{X_1: X_2: X_3}(P_1, P_2)$ and $\mathfrak{D}(P_1, P_2)$ are irreducible polynomials in \mathbb{U} . This iterated resultant is bi-homogeneous with respect to the sets of coefficients $U_{i,j}^{(1)}$ and $U_{i,j}^{(2)}$ respectively of bi-degree

$$(2d_2(d_1 d_2 - 1); 2d_1(d_1 d_2 - 1)).$$

As usual, we can specialize this result to obtain the following:

Corollary 5.7. *Given two polynomials $f_k(\mathbf{x}, y, z)$, $k = 1, 2$, of the form*

$$f_k(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d_k} a_{\alpha, i, j}^{(k)} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, the iterated resultant $\text{Disc}_y(\text{Res}_z(f_1, f_2)) \in \mathbb{S}[\mathbf{x}]$ is of degree at most $d_1 d_2 (d_1 d_2 - 1)$ in \mathbf{x} and can be factorized, up to sign, as

$$\text{Disc}_y(\text{Res}_z(f_1, f_2)) = (-1)^{(d_2+1)(d_1 d_2 + d_1 - 1)} \text{Disc}_{y,z}(f_1, f_2) \mathfrak{D}(f_1, f_2)^2.$$

Moreover, if the polynomials f_1 and f_2 are sufficiently generic, then this iterated resultant has exactly degree $d_1 d_2 (d_1 d_2 - 1)$ in \mathbf{x} and the two terms in the right hand




FIGURE 1. A quartic surface and the projection of a polar curve.

side of the above equality are respectively an irreducible polynomial and the square of an irreducible polynomial.

6. DISCRIMINANT OF A DISCRIMINANT

In this section, we are interested in analyzing two iterated discriminants. Before going into the algebraic study, let us consider the problem from a geometric point of view. Suppose we are given an implicit surface $f(x, y, z) = 0$. Computing the discriminant of f in z consists in projecting the apparent contour (or polar) curve in the z direction, which is defined by $f(x, y, z) = 0, \partial_z f(x, y, z) = 0$. Computing the discriminant in y of this discriminant in z of f consists in computing the position of lines parallel to the y axis, which are tangent to the projected curve.

We illustrate it by some explicit computations, with the polynomial

$$f(x, y, z) = z^4 - y^3z + 2z^3 - yz^2 - y^2 - xz + 1.$$

Its discriminant in z is a polynomial of degree 12 and the discriminant in y of this discriminant can be factorized as:

$$\begin{aligned} &5540271966595842048 (14348907 x^{10} - 93002175 x^9 + 273574017 x^8 - \\ &909290448 x^7 + 2868603336 x^6 - 5353192260 x^5 + 9038030571 x^4 - 17693165669 x^3 + \\ &17648229264 x^2 - 4081683588 x + 218938829)(x - 1)^2(125 x - 173)^2(47832147 x^4 + \\ &147495688 x^3 - 245928792 x^2 - 212731008 x + 230501936)^3. \end{aligned}$$

Figures³ 1 and 2 illustrate the situation, where we represent the surface $f = 0$ and the projection of its apparent contour (the x -direction is pointing to the top of the image and z -direction to the left). The first factor of degree 10 has 4 real roots corresponding to the smooth points of the surface with a tangent plane orthogonal to the x -direction. The second factor of multiplicity 2 corresponds to points of the polar variety which project in the (x, y) -plane onto the same point.

³The topology computation and visualization have been performed by the softwares AXEL (<http://axel.inria.fr/>) and SYNAPS (<http://synaps.inria.fr/>).




FIGURE 2. The polar curve of the quartic surface projected on a plane.

Geometrically speaking, we have a double folding of the surface in the z -direction above these values. There are two such real points in our example. The last factor of multiplicity 3 corresponds to cusp points on the discriminant curve, which are the projection of a “fronce” or a pleat of the surface. There are 4 such real points. Notice that the branches of the discriminant curve between two of these cusp points and one of the double folding points form a very tiny loop, which is difficult to observe at this scale.

These phenomena can be explained from an algebraic point of view, as we will see hereafter. They have also been analyzed from a singularity theory point of view. A well-know result in singularity theory, due to H. Whitney (see [31, 25, 2]) asserts that the singularities of the projection of a generic surface onto a plane are of 3 types:

- a regular point on the contour curve corresponding to a fold of the surface,
- a cusp corresponding to a pleat,
- a double point corresponding to the projection of two transversal folds.

These are stable singularities, which remain by a small perturbation of the surface or of the direction of projection. For a more complete analysis of the singularities of the apparent contour, see also [29].

We are now going to analyze the algebraic side of these geometric properties for generic polynomials of a given degree. Suppose given a positive integer $d \geq 3$ and a homogeneous polynomial

$$P(X_1, X_2, X_3) := \sum_{0 \leq i, j; i+j \leq d} U_{i,j} X_1^i X_2^j X_3^{d-i-j} \in \mathbb{U}[X_1, X_2, X_3],$$

where \mathbb{U} is the universal coefficients ring $\mathbb{Z}[U_{i,j}]$. In this section we will study the discriminant of the discriminant of P . Here is the first factorization we can get as a specialization of the iterated resultant we studied in the previous section.

Proposition 6.1. *We have the equality in \mathbb{U} :*

$$U_{0,0}^{2d^2-2d-1} \text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3))) = \text{Disc}_{X_1:X_2:X_3}(P) \times \\ \text{Res}_{X_1:X_2:X_3}(P, \partial_3 P, \partial_3^2 P) \times \text{Res}_{X_1:\dots:X_4}(P, \delta_{3,4} P, \partial_3 P, \delta_{3,4} \partial_3 P).$$

The iterated discriminant $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3)))$ is homogeneous with respect to the set of coefficients $U_{i,j}$ of degree $4(d-1)(d^2-d-1)$.

Proof. This is essentially a specialization of the formula given in Proposition 5.6, which yields in our case the equality

$$\text{Disc}_{X_2}(\text{Res}_{X_3}(P(1, X_2, X_3), \partial_3 P(1, X_2, X_3))) = \text{Disc}(P, \partial_3 P) \times \\ \text{Res}(P, \delta_{3,4} P, \partial_3 P, \delta_{3,4} \partial_3 P).$$

On the one hand, we know by definition of the discriminant and properties of resultants, that

$$(6.1) \quad \begin{aligned} \text{Res}(P, \partial_3 P, X_1) \text{Disc}(P, \partial_3 P) \\ &= \text{Res}(P, \partial_3 P, \partial_2 P \partial_3^2 P - \partial_3 P \partial_2 \partial_3 P) \\ &= \text{Res}(P, \partial_3 P, \partial_2 P \partial_3^2 P) \end{aligned}$$

$$(6.2) \quad = \text{Res}(P, \partial_3 P, \partial_2 P) \times \text{Res}(P, \partial_3 P, \partial_3^2 P).$$

Moreover, by Euler identity we have $dP = X_1 \partial_1 P + X_2 \partial_2 P + X_3 \partial_3 P$ from we deduce that

$$\begin{aligned} d^{d-1} \text{Res}(P, \partial_3 P, X_1) &= \text{Res}(X_2 \partial_2 P, \partial_3 P, X_1) \\ &= \text{Res}(X_2, \partial_3 P, X_1) \times \text{Res}(\partial_2 P, \partial_3 P, X_1) \\ &= d U_{0,0} \text{Res}(\partial_2 P, \partial_3 P, X_1). \end{aligned}$$

Again by Euler identity, the relation (2.4) yields

$$\begin{aligned} d^{(d-1)^2} \text{Res}(P, \partial_3 P, \partial_2 P) &= \text{Res}(X_1 \partial_1 P, \partial_3 P, \partial_2 P) \\ &= \text{Res}(X_1, \partial_3 P, \partial_2 P) \text{Res}(\partial_1 P, \partial_3 P, \partial_2 P) \\ &= d^{d^2-3d+3} \text{Res}(\partial_2 P, \partial_3 P, X_1) \text{Disc}(P) \end{aligned}$$

and by substitution in (6.2) and simplification by $\text{Res}(\partial_2 P, \partial_3 P, X_1)$ and d^{d-2} , we get that

$$U_{0,0} \text{Disc}(P, \partial_3 P) = \text{Disc}(P) \text{Res}(P, \partial_3 P, \partial_3^2 P)$$

so that

$$U_{0,0} \text{Disc}_{X_2}(\text{Res}_{X_3}(P(1, X_2, X_3), \partial_3 P(1, X_2, X_3))) = \text{Disc}(P) \times \\ \text{Res}(P, \partial_3 P, \partial_3^2 P) \times \text{Res}_{X_1:\dots:X_4}(P(X_3), \delta_{X_3, X_4}(P), \partial_3 P(X_3), \delta_{X_3, X_4}(\partial_3 P)).$$

On the other hand,

$$\begin{aligned} \text{Disc}_{X_2}(\text{Res}_{X_3}(P(1, X_2, X_3), \partial_3 P(1, X_2, X_3))) \\ &= \text{Disc}_{X_2}(U_{0,0} \text{Disc}_{X_3}(P(1, X_2, X_3))) \\ &= U_{0,0}^{2(d(d-1)-1)} \text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3))) \\ &= U_{0,0}^{2d^2-2d-2} \text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3))) \end{aligned}$$

since $\text{Disc}_{X_3}(P(1, X_2, X_3))$ is of degree $d(d-1)$ in X_2 and the discriminant of a polynomial of degree D is homogeneous of degree $2(D-1)$ in its coefficients. The claimed formula then follows immediately.

Concerning the degree, observe that $\text{Disc}_{X_3}(P(1, X_2, X_3))$ has coefficients of degree $2(d-1)$ in the coefficients of P . The discriminant of a polynomial of degree D being homogeneous of degree $2(D-1)$ in its coefficients, we obtain that $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3)))$ is of degree

$$2(d(d-1)-1) \times 2(d-1) = 4(d-1)(d^2-d-1)$$

since the degree of $\text{Disc}_{X_3}(P(1, X_2, X_3))$ in X_2 is $D = d(d-1)$. \square

In the factorization of $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P))$ given in this proposition, we only know that the factor $\text{Disc}(P)$ is known to be irreducible in \mathbb{U} . The remaining of this section is devoted to the study of the full factorization of the two other factors. We begin with the study of the factor appearing in Proposition 6.1 which corresponds to the resultant of P , its first and second derivatives with respect to X_3 .

Lemma 6.2. *Let \mathbb{S} be a commutative ring and suppose given a linear form $L = aX_1 + bX_2 + cX_3 \in \mathbb{S}[X_1, X_2, X_3]$ and a homogeneous polynomial $Q \in \mathbb{S}[X_1, X_2, X_3]$ of degree $d-1 \geq 2$. Then*

$$\text{Res}(LQ, \partial_3(LQ), \partial_3^2(LQ)) = 2^{2(d-1)} c^{3d-4} \text{Res}(L, Q, \partial_3(Q))^3 \times \text{Res}(Q, \partial_3(Q), \partial_3^2(Q)).$$

Proof. It is a straightforward computation using the properties of the resultants:

$$\begin{aligned} \text{Res}(LQ, \partial_3(LQ), \partial_3^2(LQ)) &= \text{Res}(LQ, cQ + L\partial_3(Q), 2c\partial_3(Q) + L\partial_3^2(Q)) \\ &= \text{Res}(L, cQ, 2c\partial_3(Q)) \times \\ &\quad \text{Res}(Q, L\partial_3(Q), 2c\partial_3(Q) + L\partial_3^2(Q)) \\ &= 2^{d-1} c^{2d-3} \text{Res}(L, Q, \partial_3(Q)) \text{Res}(Q, L, 2c\partial_3(Q)) \times \\ &\quad \text{Res}(Q, \partial_3(Q), L\partial_3^2(Q)) \\ &= 2^{2(d-1)} c^{2d-3+d-1} \text{Res}(L, Q, \partial_3(Q))^2 \times \\ &\quad \text{Res}(Q, \partial_3(Q), L) \text{Res}(Q, \partial_3(Q), \partial_3^2(Q)) \\ &= 2^{2(d-1)} c^{3d-4} \text{Res}(L, Q, \partial_3(Q))^3 \times \\ &\quad \text{Res}(Q, \partial_3(Q), \partial_3^2(Q)). \end{aligned}$$

\square

Proposition 6.3. *The following equality holds in \mathbb{U} :*

$$\text{Res}_{X_1:X_2:X_3}(P, \partial_3 P, \partial_3^2 P) = 2^{d(d-1)} U_{0,0}^2 \mathfrak{F}(P)$$

where $\mathfrak{F}(P)$ is an irreducible homogeneous polynomial in \mathbb{U} of degree $3d(d-2)$.

Proof. We rewrite P as

$$P = U_{0,0} X_3^d + c_1 X_3^{d-1} + \cdots + c_{d-2} X_3^2 + c_{d-1} X_3 + c_d$$

where the c_i 's are homogeneous polynomials in $\mathbb{U}[X_1, X_2]$ of degree i respectively; we have

$$\begin{aligned} \partial_3 P &= dA_{0,0} X_3^{d-1} + (d-1)c_1 X_3^{d-2} + \cdots + 2c_{d-2} X_3 + c_{d-1}, \\ \partial_3^2 P &= d(d-1)U_{0,0} X_3^{d-2} + (d-1)(d-2)c_1 X_3^{d-3} + \cdots + 2c_{d-2}. \end{aligned}$$

Consider the incidence variety

$$\mathcal{W} := \{(x_1 : x_2 : x_3) \times (a_{i,j}) \in \mathbb{P}^2 \times \mathbb{A}^{d(d+1)/2} \text{ such that } P = \partial_3 P = \partial_3^2 P = 0\}$$

whose canonical projection onto the second factor, i.e. by $\pi_2 : \mathcal{W} \rightarrow \mathbb{A}^{d(d+1)/2}$, is the variety of pure codimension one defined by the equation $\text{Res}(P, \partial_3 P, \partial_3^2 P) = 0$. Considering the canonical projection onto its first factor $\pi_1 : \mathcal{W} \rightarrow \mathbb{P}^2$, which is surjective, we observe that $\pi_1^{-1}(0 : 0 : 1)$ is the hyperplane $\{U_{0,0} = 0\}$ in $\mathbb{A}^{d(d+1)/2}$ and that $\pi_1^{-1}(x_1 : x_2 : x_3)$ is a linear space of codimension 3 in $\mathbb{A}^{d(d+1)/2}$ if $(x_1 : x_2 : x_3) \neq (0 : 0 : 1) \in \mathbb{P}^2$ (just observe that in this case the three conditions $c_d = 0, c_{d-1} = 0, c_{d-2} = 0$ are non trivial and that $\partial_3 P$ does not depend on c_d and that $\partial_3^2 P$ does not depend on c_d and c_{d-1}). It follows that $\mathcal{W} = \mathcal{W}_0 \cup \mathcal{W}_3$ where \mathcal{W}_0 is the irreducible variety defined by $\{U_{0,0} = 0\}$ and \mathcal{W}_3 , which is the closure of the fiber bundle $\pi_1^{-1}(\mathbb{P}^2 \setminus (0 : 0 : 1)) \rightarrow \mathbb{P}^2 \setminus (0 : 0 : 1)$ in $\mathbb{P}^2 \times \mathbb{A}^{d(d+1)/2}$, is an irreducible variety of codimension 3. We deduce that $\pi_2(\mathcal{W})$ is the union of two irreducible varieties: $\pi_2(\mathcal{W}_0)$ whose defining equation is $\{U_{0,0} = 0\}$ and $\pi_2(\mathcal{W}_3)$. Therefore, we obtain that, in \mathbb{U} ,

$$\text{Res}(P, \partial_3 P, \partial_3^2 P) = c(d)U_{0,0}^{a(d)}R_d^{r(d)}$$

where $c(d), a(d), r(d)$ are all positive integers (in particular, they are nonzero) which may only depend on d , the degree of P , and R_d is an irreducible polynomial in \mathbb{U} . In order to determine $c(d), a(d), r(d)$ we will use a particular specialization and some properties of the resultants.

First, it is easy to check by hand (or with a computer), that the case $d = 3$ gives, in \mathbb{U} ,

$$\text{Res}(P, \partial_3 P, \partial_3^2 P) = 2^6 U_{0,0}^2 \mathfrak{F}(P)$$

where $\mathfrak{F}(P)$ is an irreducible polynomial. Moreover, if ϕ denotes the homogeneous specialization which sends all the coefficients of c_d to 0, we have $\phi(P) = X_3 Q$, where Q is a generic homogeneous polynomial of degree $d - 1$. By Lemma 6.2, we get

$$\phi(\text{Res}(P, \partial_3 P, \partial_3^2 P)) = 2^{2(d-1)} \text{Res}(Q, \partial_3 Q, X_3)^3 \text{Res}(Q, \partial_3 Q, \partial_3^2 Q).$$

So, if we proceed by induction on the integer d , we obtain

$$\begin{aligned} \phi(\text{Res}(P, \partial_3 P, \partial_3^2 P)) &= 2^{2(d-1)} \text{Res}(Q, \partial_3 Q, X_3)^3 \times 2^{(d-1)(d-2)} U_{0,0}^{a(d-1)} \mathfrak{F}(Q) \\ &= 2^{d(d-1)} U_{0,0}^2 \text{Res}(Q, \partial_3 Q, X_3)^3 \mathfrak{F}(Q) \end{aligned}$$

where we notice that $\text{Res}(Q, \partial_3 Q, X_3) = \text{Res}(c_{d-1}, c_{d-2})$ is an irreducible polynomial in \mathbb{U} (it is the resultant of two generic homogeneous polynomials) which does not depend on $U_{0,0}$. Since we must also have

$$\phi(\text{Res}(P, \partial_3 P, \partial_3^2 P)) = c(d)U_{0,0}^{a(d)} \phi(\mathfrak{F}(P))^{r(d)}$$

we deduce by comparison that, for all $d \geq 3$,

- $c(d)$ divides $2^{d(d-1)}$,
- $r(d) = 1$ (since the specialization of R_d by ϕ produces an irreducible and reduced factor),
- $a(d) \leq 2$.

It is easy to see that 2 divides $\partial_3^2 P$ and hence that $2^{d(d-1)}$ divides $\text{Res}(P, \partial_3 P, \partial_3^2 P)$. This implies that $2^{d(d-1)}$ divides $c(d)$ and hence, since we already noted that $c(d)$ divides $2^{d(d-1)}$, $c(d) = 2^{d(d-1)}$. To conclude the proof, it remains to show that

$a(d) \geq 2$ for all $d \geq 3$, that is to say that $U_{0,0}^2$ divides $\text{Res}(P, \partial_3 P, \partial_3^2 P)$ in \mathbb{U} . Moreover, it is sufficient to show that $U_{0,0}^2$ divides

$$\mathfrak{R} := \text{Res}(d(d-1)P, (d-1)\partial_3 P, \partial_3^2 P)$$

since

$$\text{Res}(d(d-1)P, (d-1)\partial_3 P, \partial_3^2 P) = d(d-1)^{(d-1)(d-2)} \times (d-1)^{d(d-2)} \times \text{Res}(P, \partial_3 P, \partial_3^2 P).$$

Using Euler identity several times, we get

$$\begin{aligned} \mathfrak{R} &= \text{Res}((d-1)X_1\partial_1 P + (d-1)X_2\partial_2 P, (d-1)\partial_3 P, \partial_3^2 P) \\ &= \text{Res}(X_1^2\partial_1^2 P + 2X_1X_2\partial_1\partial_2 P + X_2^2\partial_2^2 P, X_1\partial_1\partial_3 P + X_2\partial_2\partial_3 P, \partial_3^2 P). \end{aligned}$$

It is clear that $\partial_3^2 P \in (X_1, X_2, U_{0,0}X_3)$ and that

$$X_1\partial_1\partial_3 P + X_2\partial_2\partial_3 P \in (X_1, X_2) \subset (X_1, X_2, U_{0,0}X_3).$$

Moreover, $X_1^2\partial_1^2 P + 2X_1X_2\partial_1\partial_2 P + X_2^2\partial_2^2 P \in (X_1, X_2)^2 \subset (X_1, X_2, U_{0,0}X_3)^2$. Therefore, the divisibility property of resultants implies that

$$\text{Res}(X_1, X_2, U_{0,0}X_3)^{2 \times 1 \times 1} \text{ divides } \mathfrak{R},$$

and since $\text{Res}(X_1, X_2, U_{0,0}X_3) = U_{0,0}$, we are done.

Finally, the degree of $\mathfrak{F}(P)$ in the coefficients of P is given by the formula

$$d(d-1) + d(d-2) + (d-1)(d-2) - 2 = 3d(d-2).$$

□

We now turn to the study of the third factor appearing in the factorization of $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P))$ in Proposition 6.1.

Proposition 6.4. *Assuming that $d \geq 4$, we have the following equality between homogeneous polynomials in \mathbb{U} of degree $2(d-1)(2d^2 - 4d + 1)$:*

$$\begin{aligned} 2^{d(d-1)} \text{Res}_{X_1, \dots, X_4}(P, \delta_{3,4}P, \partial_3 P, (\delta_{3,4}\partial_3)P) = \\ \text{Res}_{X_1, X_2, X_3}(P, \partial_3 P, \partial_3^2 P)^2 \times \text{Res}_{X_1, \dots, X_4}(P, \partial_3 P, \delta_{3,4}^2 P, (\delta_{3,4}^2\partial_3 - 2\delta_{3,4}^3)P). \end{aligned}$$

Proof. Using known properties of resultants and equalities (2.6) and (2.7), we get the equalities

$$\begin{aligned} \text{Res}(P, \delta_{3,4}P, \partial_3 P, \delta_{3,4}\partial_3 P) &= \text{Res}(P, \partial_3 P + (X_4 - X_3)\delta_{3,4}^2 P, \partial_3 P, \delta_{3,4}\partial_3 P) \\ &= \text{Res}(P, (X_4 - X_3)\delta_{3,4}^2 P, \partial_3 P, \delta_{3,4}\partial_3 P) \\ &= \text{Res}(P, (X_4 - X_3), \partial_3 P, \delta_{3,4}\partial_3 P) \times \\ &\quad \text{Res}(P, \delta_{3,4}^2 P, \partial_3 P, \partial_3^2 P + (X_4 - X_3)\delta_{3,4}^2 \partial_3 P) \\ &= \text{Res}(P, \partial_3 P, \partial_3^2 P) \times \\ &\quad \text{Res}(P, \delta_{3,4}^2 P, \partial_3 P, \partial_3^2 P + (X_4 - X_3)\delta_{3,4}^2 \partial_3 P). \end{aligned}$$

Let us denote

$$\mathcal{S}(P) := \text{Res}(P, \delta_{3,4}^2 P, \partial_3 P, \partial_3^2 P + (X_4 - X_3)\delta_{3,4}^2 \partial_3 P).$$

Since $\partial_3^2 P = 2\delta_{3,4}^2 P - 2(X_4 - X_3)\delta_{3,4}^3 P$ in \mathbb{U} , we deduce that, in \mathbb{U} ,

$$\begin{aligned}
 2^{d(d-1)(d-2)}\mathcal{S}(P) &= \text{Res}(P, 2\delta_{3,4}^2 P, \partial_3 P, \partial_3^2 P + (X_4 - X_3)\delta_{3,4}^2 \partial_3 P) \\
 &= \text{Res}(P, 2\delta_{3,4}^2 P, \partial_3 P, (X_4 - X_3)(\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P) \\
 &= \text{Res}(P, 2\delta_{3,4}^2 P, \partial_3 P, (X_4 - X_3)) \times \\
 &\quad \text{Res}(P, 2\delta_{3,4}^2 P, \partial_3 P, (\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P) \\
 &= \text{Res}(P, \partial_3^2 P, \partial_3 P, (X_4 - X_3)) \times \\
 &\quad \text{Res}(P, 2\delta_{3,4}^2 P, \partial_3 P, (\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P) \\
 &= 2^{d(d-1)(d-3)}\text{Res}(P, \partial_3^2 P, \partial_3 P) \times \\
 &\quad \text{Res}(P, \delta_{3,4}^2 P, \partial_3 P, (\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P).
 \end{aligned}$$

Therefore, we have

$$\begin{aligned}
 2^{d(d-1)}\text{Res}(P, \delta_{3,4} P, \partial_3 P, (\delta_{3,4} \partial_3)P) &= \text{Res}(P, \partial_3 P, \partial_3^2 P)^2 \times \\
 &\quad \text{Res}((P, \partial_3 P, \delta_{3,4}^2 P, (\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P)).
 \end{aligned}$$

The claimed formula for the degree of $\text{Res}(P, \delta_{3,4} P, \partial_3 P, (\delta_{3,4} \partial_3)P)$ follows immediately by specialization of the formula given in Lemma 5.4. \square

As a consequence, to get the full factorization of $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P))$ it only remains to study the factorization of the term

$$\mathcal{R}(P) := \text{Res}_{X_1:\dots:X_4}(P, \partial_3 P, \delta_{3,4}^2 P, (\delta_{3,4}^2 \partial_3 - 2\delta_{3,4}^3)P)$$

whose degree in the coefficients of P is

$$2(d-1)(2d^2 - 4d + 1) - 2(d(d-1) + d(d-2) + (d-1)(d-2)) = 2(2d-3)(d^2 - 3d + 1).$$

This is the aim of the next proposition. We begin with two technical lemmas.

Lemma 6.5. *Let \mathbb{S} be a commutative ring and suppose given a linear form $L = aX_1 + bX_2 + cX_3 \in \mathbb{S}[X_1, X_2, X_3]$ and a homogeneous polynomial $Q \in \mathbb{S}[X_1, X_2, X_3]$ of degree $d-1 \geq 2$. Then*

$$\begin{aligned}
 \text{Res}_{X_1:\dots:X_4}(LQ, \partial_3(LQ), \delta_{3,4}(LQ), \delta_{3,4}\partial_3(LQ)) &= 2^{2(d-1)}c^{2(d-1)(3d-5)} \times \\
 \text{Res}_{X_1:X_2:X_3}(L, Q, \partial_3 Q)^6 \times \text{Res}_{X_1:\dots:X_4}(L(X_4), Q, \partial_3 Q, \delta_{3,4}^2 Q)^4 \times \\
 &\quad \text{Res}_{X_1:\dots:X_4}(Q, \delta_{3,4} Q, \partial_3 Q, \delta_{3,4}\partial_3 Q).
 \end{aligned}$$

Proof. Set $P := LQ$. Applying (5.2) with $P_2 = \partial_3(P)$ and

$$\partial_3 P(X_4) = \partial_3 P(X_3) + (X_4 - X_3)\delta_{3,4}\partial_3 P,$$

we obtain the decomposition

$$\begin{aligned}
 (6.3) \quad &\text{Res}(P, \partial_3(P), \delta_{3,4}(P), \delta_{3,4}\partial_3(P)) \\
 &= c^{(d-1)(d-2)} \times \text{Res}(L(X_4), Q(X_3), \partial_3(P), \delta_{3,4}\partial_3(P))^2 \times \\
 &\quad \text{Res}(Q, \delta_{3,4}(Q), \partial_3(P), \delta_{3,4}\partial_3(P)) \\
 &= c^{(d-1)(d-2)} R_1^2 R_2.
 \end{aligned}$$

Using the relations

$$\begin{aligned}
 \partial_3(P) &= \partial_3(L)Q + L\partial_3(Q), \\
 \delta_{3,4}\partial_3(P) &= \partial_3(L)\delta_{3,4}(Q) + \delta_{3,4}(L)\partial_3(Q) + L(X_4)\delta_{3,4}\partial_3(Q),
 \end{aligned}$$

we can decompose further the previous expressions:

$$\begin{aligned}
R_1 &:= \text{Res}(L(X_4), Q, \partial_3(P), \delta_{3,4}\partial_3(P)) \\
&= \text{Res}(L(X_4), Q, L\partial_3(Q), \partial_3(L)\delta_{3,4}(Q) + \delta_{3,4}(L)\partial_3(Q)) \\
&= \text{Res}(L(X_4), Q, L, c\delta_{3,4}(Q) + c\partial_3(Q)) \times \text{Res}(L(X_4), Q, \partial_3(Q), c\delta_{3,4}(Q)) \\
&= \text{Res}(L(X_4), Q, c(X_3 - X_4), c(\delta_{3,4}(Q) + \partial_3(Q))) \times \\
&\quad c^{(d-1)(d-2)}\text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}(Q)) \\
&= c^{(d-1)+(d-1)(d-2)}\text{Res}(L, Q, 2\partial_3(Q)) \times \\
&\quad c^{(d-1)(d-2)}\text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}(Q)) \\
&= 2^{d-1}c^{(d-1)(2d-3)}\text{Res}(L, Q, \partial_3(Q)) \times \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}(Q)).
\end{aligned}$$

Now, using the relation $\delta_{3,4}(Q) = \partial_3(Q) + (X_4 - X_3)\delta_{3,4}^2(Q)$, we have

$$\begin{aligned}
(6.4) \quad &\text{Res}(Q, \partial_3(Q), \delta_{3,4}(Q), L(X_4)) \\
&= \text{Res}(Q, \partial_3(Q), (X_4 - X_3)\delta_{3,4}^2(Q), L(X_4)) \\
&= \text{Res}(L, Q, \partial_3(Q)) \times \text{Res}(Q, \partial_3(Q), \delta_{3,4}^2(Q), L(X_4)).
\end{aligned}$$

We deduce that

$$R_1 = 2^{d-1}c^{(d-1)(2d-3)}\text{Res}(L, Q, \partial_3(Q))^2 \times \text{Res}(Q, \partial_3(Q), \delta_{3,4}^2(Q), L(X_4)).$$

Similarly, we have

$$\begin{aligned}
R_2 &= \text{Res}(Q, \delta_{3,4}(Q), \partial_3(P), \delta_{3,4}\partial_3(P)) \\
&= \text{Res}(Q, \delta_{3,4}(Q), L\partial_3(Q), \delta_{3,4}(L)\partial_3(Q) + L(X_4)\delta_{3,4}\partial_3(Q)) \\
&= \text{Res}(Q, \delta_{3,4}(Q), L, \delta_{3,4}(L)\partial_3(Q) + L(X_4)\delta_{3,4}\partial_3(Q)) \times \\
&\quad \text{Res}(Q, \delta_{3,4}(Q), \partial_3(Q), L(X_4)\delta_{3,4}\partial_3(Q)) \\
&= \text{Res}(Q, \delta_{3,4}(Q), L, c\partial_3(Q)(X_4)) \times \text{Res}(Q, \delta_{3,4}(Q), \partial_3(Q), L(X_4)) \\
&\quad \times \text{Res}(Q, \delta_{3,4}(Q), \partial_3(Q), \delta_{3,4}\partial_3(Q)) \\
&= c^{(d-1)(d-2)}\text{Res}(Q, \partial_3(Q), \delta_{3,4}(Q), L(X_4))^2 \times \\
&\quad \text{Res}(Q, \delta_{3,4}(Q), \partial_3(Q), \delta_{3,4}\partial_3(Q)),
\end{aligned}$$

since

$$c\partial_3(Q) + L(X_4)\delta_{3,4}\partial_3(Q) - L(X_3)\delta_{3,4}\partial_3(Q) = c\partial_3(Q)(X_4),$$

and

$$\text{Res}(Q, \delta_{3,4}(Q), L, \partial_3(Q)(X_4)) = \text{Res}(Q, \partial_3(Q), \delta_{3,4}(Q), L(X_4)).$$

using the relation (6.4), we deduce that

$$\begin{aligned}
R_2 &= c^{(d-1)(d-2)}\text{Res}(L, Q, \partial_3(Q))^2 \times \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q))^2 \times \\
&\quad \text{Res}(Q, \delta_{3,4}(Q), \partial_3(Q), \delta_{3,4}\partial_3(Q)).
\end{aligned}$$

By (6.3), squaring R_1 and taking the product with R_2 and $c^{(d-1)(d-2)}$, we obtain the expected decomposition. \square

Lemma 6.6. *Let \mathbb{S} be a commutative ring and suppose given a linear form $L = aX_1 + bX_2 + cX_3 \in \mathbb{S}[X_1, X_2, X_3]$ and a homogeneous polynomial $Q \in \mathbb{S}[X_1, X_2, X_3]$ of degree $d - 1 \geq 2$. Then*

$$\mathcal{R}(LQ) = c^{6d^2 - 22d + 18} \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q))^4 \mathcal{R}(Q).$$

Proof. Set $P := LQ$. By Proposition 6.4 and Lemma 6.2, we have

$$\begin{aligned} & 2^{d(d-1)} \text{Res}(P, \delta_{3,4}(P), \partial_3(P), \delta_{3,4}\partial_3(P)) \\ &= \text{Res}(P, \partial_3(P), \partial_3^2(P))^2 \mathcal{R}(P) \\ &= \left(2^{2(d-1)} c^{3d-4} \text{Res}(L, Q, \partial_3(Q))^3 \text{Res}(Q, \partial_3(Q), \partial_3^2(Q)) \right)^2 \mathcal{R}(P) \\ &= 2^{4(d-1)} c^{6d-8} \text{Res}(L, Q, \partial_3(Q))^6 \text{Res}(Q, \partial_3(Q), \partial_3^2(Q))^2 \mathcal{R}(P). \end{aligned}$$

Moreover, by Lemma 6.5 and Proposition 6.4 we also have

$$\begin{aligned} & \text{Res}(P, \delta_{3,4}(P), \partial_3(P), \delta_{3,4}\partial_3(P)) \\ &= 2^{2(d-1)} c^{2(d-1)(3d-5)} \text{Res}(L, Q, \partial_3 Q)^6 \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q))^4 \times \\ & \quad \text{Res}(Q, \delta_{3,4}Q, \partial_3 Q, \delta_{3,4}\partial_3 Q) \\ &= 2^{-d(d-1)} c^{2(d-1)(3d-5)} \text{Res}(L, Q, \partial_3 Q)^6 \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q))^4 \times \\ & \quad \text{Res}(Q, \partial_3 Q, \partial_3^2 Q)^2 \mathcal{R}(Q) \end{aligned}$$

and the claimed formula follows by comparison. \square

Proposition 6.7. *Assuming that $d \geq 4$, we have*

$$\mathcal{R}(P) = U_{0,0}^{2d(d-1)-6} \mathfrak{U}(P)^2,$$

where $\mathfrak{U}(P)$ is irreducible in \mathbb{U} of degree $2d(d-2)(d-3)$.

Proof. We first prove that $U_{0,0}^{2d(d-1)-6}$ divides $\mathcal{R}(P)$ in \mathbb{U} . Gathering the results of the propositions 6.1, 6.3, 6.4 and 6.7 we obtain the following equality in \mathbb{U} :

$$(6.5) \quad U_{0,0}^{2d(d-1)-6} \text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3))) = U_{0,0} \text{Disc}_{X_1:X_2:X_3}(P) \mathfrak{F}(P)^3 \mathcal{R}(P).$$

In order to prove that $U_{0,0}^{2d(d-1)-6}$ divides $\mathcal{R}(P)$, it is thus sufficient to prove that $U_{0,0}$ divides $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3)))$. Rewrite the polynomial P as

$$P := U_{0,0} X_3^d + a_{d-1} X_3^{d-1} + \cdots + a_1 X_3 + a_0$$

where the a_i 's are homogeneous polynomials in X_1, X_2 . If one specializes $U_{0,0}$ to 0 then P specializes to a polynomial of degree $d-1$ in X_3 and hence, by a well-known property of discriminants we have

$$\text{Disc}_{X_3}(P(1, X_2, X_3)) = (-1)^d a_{d-1} (1, X_2)^2 \text{Disc}'_{X_3}(P(1, X_2, X_3))$$

where $\text{Disc}'_{X_3}(P(1, X_2, X_3))$ denotes the discriminant of P as a polynomial of degree $d-1$ (and not d). But then, the discriminant with respect to X_2 of the above quantity equals 0 since it contains a square factor. This implies that $U_{0,0}$ divides $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3)))$.

Observe now that by Proposition 5.4 and Proposition 6.4, $\mathcal{R}(P)$ is a square and hence can be decomposed as

$$\mathcal{R}(P) = c(d)^2 U_{0,0}^{2d(d-1)-6} \mathfrak{R}_1^{2r_1(d)}(P) \mathfrak{R}_2^{2r_2(d)}(P) \cdots \mathfrak{R}_s(P)^{2r_s(d)},$$

where $\mathfrak{R}_i(P)$ are irreducible polynomials and $c(d) \in \mathbb{Z}$. As we did several times, we will prove the claimed factorization by induction on the degree d of P . For $d = 4$, we find by explicit computation that

$$\mathcal{R}(P) := 2^8 U_{0,0}^{18} \mathfrak{U}(P)^2$$

where $\mathfrak{U}(P)$ is an irreducible polynomial in \mathbb{U} of the expected degree. Assume that for a generic polynomial Q such that $\deg(Q) < d$, we have

$$\mathcal{R}(Q) = U_{0,0}(Q)^{2(d-1)(d-2)-6}\mathfrak{U}(Q)^2$$

where $\mathfrak{U}(Q)$ is an irreducible polynomial of bi-degree $(0, 2(d-1)(d-3)(d-4))$ in (L, Q) .

Consider the specialization $\phi(P) = LQ$ where L and Q are generic polynomials of degree 1 and $(d-1)$ respectively. We denote by c , resp. $U'_{0,0}$, the coefficient of X_3 , resp. X_3^{d-1} , of L , resp. Q . Each factor $\phi(\mathfrak{R}_i)$ must decompose into a product of irreducible factors such that the degree in the coefficients of L is equal to the degree in the coefficients of Q . By Lemma 6.6, we have

$$\mathcal{R}(LQ) = c^{6d^2-22d+18} \text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q))^4 \mathcal{R}(Q)$$

and by Proposition 4.6, we have

$$\text{Res}(L(X_4), Q, \partial_3(Q), \delta_{3,4}^2(Q)) = U_{0,0}'^{d-1} \mathfrak{T}(Q, L)$$

where $\mathfrak{T}(Q, L)$ is irreducible of degree $((d-1)(d-2)(d-3), (3d-4)(d-3))$ in (L, Q) . Using now the induction hypothesis, we deduce that

$$\begin{aligned} \mathcal{R}(LQ) &= c^{6d^2-22d+18} U_{0,0}'^{4(d-1)+2(d-1)(d-2)-6} \mathfrak{T}(Q, L)^4 \mathfrak{U}(Q)^2 \\ &= c^{6d^2-22d+18} U_{0,0}'^{2d(d-1)-6} \mathfrak{T}(Q, L)^4 \mathfrak{U}(Q)^2 \\ &= (cU_{0,0}')^{2d(d-1)-6} c^{4(d-2)(d-3)} \mathfrak{T}(Q, L)^4 \mathfrak{U}(Q)^2. \end{aligned}$$

An explicit analysis, as the ones we did several times before in this paper, shows that the product $c^{2(d-2)(d-3)} \mathfrak{T}(Q, L)^2 \mathfrak{U}(Q)$ must come from the same factor $\mathfrak{R}_i(P)$. Indeed, counting 1 for the degree of a coefficient of L and -1 for the degree of a coefficient of Q , the degree of a term $c^l \mathfrak{T}(Q, L)^k \mathfrak{U}(Q)$ with $l \leq 2(d-2)(d-3)$ is at most

$$\begin{aligned} 2(d-2)(d-3) + k((d-1)(d-2)(d-3) - (3d-4)(d-3)) - 2(d-1)(d-3)(d-4) \\ = 2(d-3)(d^2 - 6d + 6)(k-2). \end{aligned}$$

which is < 0 for $k = 0, 1$ with $d \geq 5$ and 0 for $k = 2$. This proves that for $d \geq 5$, any sub-product $c^{2(d-2)(d-3)} \mathfrak{T}(Q, L)^2 \mathfrak{U}(Q)$ has not the same degree in the coefficients of L and Q , except when $k = 2$ for the product of all terms. We deduce that $\mathfrak{U}(P)$ is irreducible, which concludes the proof of the proposition. \square

Gathering all our results, we get the following theorem:

Theorem 6.8. *Assuming that $d \geq 4$, we have the equality in \mathbb{U}*

$$\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3))) = U_{0,0} \text{Disc}_{X_1: X_2: X_3}(P) \mathfrak{F}(P)^3 \mathfrak{U}(P)^2.$$

The iterated discriminant $\text{Disc}_{X_2}(\text{Disc}_{X_3}(P(1, X_2, X_3)))$ is homogeneous with respect to the set of coefficients $U_{i,j}$ of degree $4(d-1)(d^2-d-1)$.

Proof. It is a direct consequence of (6.5) and Proposition 6.7. \square

Corollary 6.9. *Given a polynomial $f(\mathbf{x}, y, z)$ of the form*

$$f(\mathbf{x}, y, z) = \sum_{|\alpha|+i+j \leq d} a_{\alpha, i, j} \mathbf{x}^\alpha y^i z^j \in \mathbb{S}[\mathbf{x}][y, z],$$

where \mathbf{x} denotes a set of variables (x_1, \dots, x_n) for some integer $n \geq 1$ and \mathbb{S} is any commutative ring, the iterated discriminant $\text{Disc}_y(\text{Disc}_z(f)) \in \mathbb{S}[\mathbf{x}]$ has degree at most $d(d-1)(d^2-d-1)$ in \mathbf{x} . If the polynomial f is sufficiently generic and \mathbb{S} is an infinite field, we have

$$\text{Disc}_y(\text{Disc}_z(f)) = a_{0,0,d} \text{Disc}_{y,z}(f) \mathfrak{F}(f)^3 \mathfrak{U}(f)^2.$$

where

- $\text{Disc}_{y,z}(f)$ is an irreducible polynomial in \mathbf{x} of degree $d(d-1)^2$,
- $\mathfrak{F}(f)$ is irreducible in \mathbf{x} of degree $d(d-1)(d-2)$ and we have the equality $\text{Res}_{y,z}(f, \partial_z f, \partial_z^2 f) = 2^{d(d-1)} a_{0,0,d}^2 \mathfrak{F}(f)$.
- $\mathfrak{U}(f)$ is an irreducible polynomial in \mathbf{x} of degree $\frac{1}{2}d(d-1)(d-2)(d-3)$, such that

$$\text{Res}_{y,z,z'}(f, \delta_{z,z'}(f), \partial_z f, \delta_{z,z'}(\partial_z f)) = a_{0,0,d}^4 \mathfrak{F}(f)^2 \mathfrak{U}(f)^2$$

and also

$$\text{Res}_{y,z,z'}(f, \partial_z f, \delta_{z,z'}^2(f), (\delta_{z,z'}^2 \partial_z - 2\delta_{z,z'}^3)f) = a_{0,0,d}^{2d(d-1)-6} \mathfrak{U}(f)^2.$$

Remark 6.10. In the decomposition formula given in the above corollary, we can replace $\text{Disc}_{y,z}(f)$ by a resultant up to the constant factor $a_{0,0,d}^2$.

Acknowledgments: This work was partially supported by the french ANR GECKO.

REFERENCES

- [1] F. Apéry and J.-P. Jouanolou, *Élimination: le cas d'une variable*, Hermann, Collection Méthodes, 2006.
- [2] V. Arnold, A. Varchenko, and S. M Gusein-Zade, *Singularités des applications différentiables*, Edition Mir, Moscou, 1986.
- [3] S. Basu, R. Pollack, and M.-F. Roy, *Algorithms in real algebraic geometry*, Springer-Verlag, Berlin, 2003, ISBN 3-540-00973-6.
- [4] E. Bézout, *Théorie Générale des Équations Algébriques*, Paris : Ph.-D. Pierres, 1779.
- [5] L. Busé, M. Elkadi, and B. Mourrain, *Resultant over the residual of a complete intersection*, J. Pure Appl. Algebra **164** (2001), no. 1-2, 35–57, Effective methods in algebraic geometry (Bath, 2000).
- [6] L. Busé, M. Elkadi, and B. Mourrain, *Using projection operators in computer aided geometric design*, Topics in Algebraic Geometry and Geometric Modeling., Contemporary Mathematics, 2003, pp. 321–342.
- [7] L. Busé, *Étude du résultant sur une variété algébrique*, Ph.D. thesis, Université de Nice Sophia Antipolis, 2001.
- [8] ———, *Resultants of determinantal varieties*, J. Pure Appl. Algebra **193** (2004), no. 1-3, 71–97.
- [9] G. E. Collins, *Quantifier elimination for real closed fields by cylindrical algebraic decomposition*, Automata theory and formal languages (Second GI Conf., Kaiserslautern, 1975), Springer, Berlin, 1975, pp. 134–183. Lecture Notes in Comput. Sci., Vol. 33.
- [10] M. Coste., *An introduction to semi-algebraic geometry*, RAAG network school, 2002.
- [11] D. Cox, John Little, and Donal O'Shea, *Using algebraic geometry*, Graduate Texts in Mathematics, vol. 185, Springer-Verlag, New York, 1998.
- [12] C. D'Andrea, *Macaulay style formulas for sparse resultants*, Trans. Amer. Math. Soc. **354** (2002), no. 7, 2595–2629 (electronic).
- [13] I.Z. Emiris and J.F. Canny, *Efficient incremental algorithms for the sparse resultant and the mixed volume*, J. Symbolic Computation **20** (1995), no. 2, 117–149.
- [14] I. M. Gel'fand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1994.

- [15] R. M. Hardt, *Triangulation of subanalytic sets and proper light subanalytic maps*, Invent. Math. **38** (1976/77), no. 3, 207–217.
- [16] O. Henrici, *On certain formulæ concerning the theory of discriminants*, Proc. of London Math. Soc. (1868), 104–116.
- [17] ———, *On the singularities of curves envelopes*, Proc. of London Math. Soc. (1869), 177–195.
- [18] J.-P. Jouanolou, *Le formalisme du résultant*, Adv. Math. **90** (1991), no. 2, 117–263.
- [19] ———, *Résultant anisotrope, compléments et applications*, Electron. J. Combin. **3** (1996), no. 2, Research Paper 2, approx. 91 pp. (electronic), The Foata Festschrift.
- [20] ———, *Formes d’inertie et résultant: un formulaire*, Adv. Math. **126** (1997), no. 2, 119–250.
- [21] W. Krull, *Funktionaldeterminanten und Diskriminanten bei Polynomen in mehreren Unbestimmten*, Monatsh. Math. Phys. **48** (1939), 353–368.
- [22] S. Lang, *Algebra*, third ed., Graduate Texts in Mathematics, vol. 211, Springer-Verlag, New York, 2002.
- [23] F.S. Macaulay, *Some formulae in elimination*, Proc. London Math. Soc. **1** (1902), no. 33, 3–27.
- [24] D. Manocha and J. F. Canny, *Implicit representation of rational parametric surfaces*, J. Symbolic Comput. **13** (1992), no. 5, 485–510.
- [25] J.N. Mather, *Generic projections*, Annals of Mathematics **98** (1973), 226–245.
- [26] S. McCallum, *Factors of iterated resultants and discriminants*, J. Symbolic Comput. **27** (1999), no. 4, 367–385.
- [27] ———, *Repeated discriminants*, Preprint of Macquarie University, December 10th 2001.
- [28] B. Mourrain and J.P. Tércourt, *Isotopic meshing of a real algebraic surface*, Technical Report 5508, INRIA Sophia Antipolis, 2005.
- [29] O.A. Platonova, *Projection of smooth surfaces*, J. of Mathematical Sciences **35** (1986), no. 6, 2796–808.
- [30] B. L. Van der Waerden, *Modern algebra, vol. ii*, New-York, Frederick Ungar Publishing Co, 1948.
- [31] H. Whitney, *On singularities of mappings of euclidean spaces. i. mapping of the plane into the plane*, Annals of Mathematics **62** (1955), no. 3, 374–410.

GALAAD, INRIA, B.P. 93, 06902 SOPHIA ANTIPOLIS, FRANCE
E-mail address: `Laurent.Buse@inria.fr`

GALAAD, INRIA, B.P. 93, 06902 SOPHIA ANTIPOLIS, FRANCE
E-mail address: `mourrain@sophia.inria.fr`