

Convergence of a non-monotone scheme for Hamilton-Jacobi-Bellman equations with discontinuous data

Olivier Bokanowski, Nadia Megdich, Hasnaa Zidani

► **To cite this version:**

Olivier Bokanowski, Nadia Megdich, Hasnaa Zidani. Convergence of a non-monotone scheme for Hamilton-Jacobi-Bellman equations with discontinuous data. *Numerische Mathematik*, Springer Verlag, 2010, 115 (1), pp.1–44. 10.1007/s00211-009-0271-1 . inria-00193157

HAL Id: inria-00193157

<https://hal.inria.fr/inria-00193157>

Submitted on 30 Nov 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Convergence of a non-monotone scheme for
Hamilton-Jacobi-Bellman equations with
discontinuous data***

Olivier Bokanowski — Nadia Megdich — Hasnaa Zidani

N° ????

Novembre 2007

Thème NUM



*Rapport
de recherche*

Convergence of a non-monotone scheme for Hamilton-Jacobi-Bellman equations with discontinuous data

Olivier Bokanowski ^{*}, Nadia Megdich [†], Hasnaa Zidani [‡]

Thème NUM — Systèmes numériques
Projets Commands

Rapport de recherche n° 7000 — Novembre 2007 — 44 pages

Abstract: We prove the convergence of a non-monotonous scheme for a one-dimensional first order Hamilton-Jacobi-Bellman equation of the form $v_t + \max_{\alpha} (f(x, \alpha)v_x) = 0$, $v(0, x) = v_0(x)$. The scheme is related to the HJB-UltraBee scheme suggested in [7]. We show for general discontinuous initial data a first-order convergence of the scheme, in L^1 -norm, towards the viscosity solution. We also illustrate the non-diffusive behavior of the scheme on several numerical examples.

Key-words: HJB equation, L^1 - error estimate, antidiffusive scheme, non-monotone scheme, discontinuous initial data

^{*} Lab. Jacques-Louis Lions, Université Paris 6 & 7, CP 7012, 175 rue du Chevaleret, 75013 Paris, boka@math.jussieu.fr

[†] UMA - ENSTA, 32 Bd Victor, 75015 Paris. Also at projet Commands, CMAP - INRIA Futurs, Ecole Polytechnique, 91128 Palaiseau, Nadia.Megdich@ensta.fr

[‡] UMA - ENSTA, 32 Bd Victor, 75015 Paris. Also at projet Commands, CMAP - INRIA Futurs, Ecole Polytechnique, 91128 Palaiseau, Hasnaa.Zidani@ensta.fr

Convergence d'un schéma non monotone pour les équations HJB avec donnée initiale discontinue

Résumé : On étudie un schéma non monotone pour l'équation Hamilton Jacobi Bellman du premier ordre, en dimension 1. Le schéma qu'on considère est lié au schéma anti-diffusif, appelé UltraBee, proposé dans [7]. Dans ce papier, on prouve la convergence, en norme L^1 , à l'ordre 1, pour une condition initiale discontinue. Le caractère anti-diffusif du schéma est illustré par quelques exemples numériques.

Mots-clés : équation HJB, schéma non monotone, estimation d'erreur en norme L^1 , schéma anti-diffusif, condition initiale discontinue

1 Introduction

We consider the following first order Hamilton-Jacobi-Bellman equation:

$$\vartheta_t(t, x) + \max_{\alpha \in \mathcal{A}} (f(x, \alpha) - \vartheta_x(t, x)) = 0, \quad t > 0, x \in \mathbb{R}, \quad (1a)$$

$$\vartheta(0, x) = v_0(x), \quad x \in \mathbb{R}, \quad (1b)$$

with discontinuous initial data v_0 . In optimal control theory, the solution ϑ of equation (1) corresponds to the value function of the optimized problem [3, 2]. It is usual that this function, as well as the “final” cost v_0 , is discontinuous (for instance for target or Rendez-Vous problems).

In the continuous case (v_0 is continuous), there are several contributions dealing with numerical schemes for the discretization of HJB equations [8, 10, 1, 13]. The work of Barles and Souganidis [4] gives a general framework for the convergence of approximated solutions towards the viscosity solution, under generic monotonicity stability and consistency assumptions. In that case, an error bound in $\Delta x^{\frac{\gamma}{2}}$ is exhibited, Δx being the mesh size, whenever the function v_0 is bounded γ -Hölder.

Nevertheless, when we deal with discontinuous initial data v_0 , classical monotone schemes are no more adapted. In fact, if we attempt to use these schemes, we observe an increasing numerical diffusion around discontinuities, and this is due to the fact that monotone schemes use at some level finite differences and/or interpolation technics.

In this paper, we analyse an explicit scheme for the numerical resolution of (1), closely related to the HJB-UltraBee scheme proposed in [7]. We give a convergence proof, show anti-dissipative properties of the scheme, and give a L^1 -error estimate.

The UltraBee scheme has been developed to study compressible gas dynamics [9], and more precisely to solve the transport equation. A generalization to HJB equations and many academic tests have been done to evaluate the behaviour of the scheme when dealing with discontinuities [7]. Its comparison with the viability algorithm [5] [15] was encouraging to study more deeply convergence results.

Let us stress on that this scheme is explicit and non-monotonous (neither ϵ -monotone in the sense of R. Abgrall [1]). As far as we know, there are few non-monotone schemes that have been proved to converge for HJ equations. In [12], Lions and Souganidis show the convergence of a TVD second order scheme, but which is implicit.

For a large class of discontinuous initial data v_0 , and under some assumptions on the dynamics $f(x, \alpha)$ (see assumption (H3) in Section 2), we obtain a first-order error bound in L^1 norm, of the following form:

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq C(L, t_n, v_0) \Delta x \quad \forall t_n \geq 0, \quad (2)$$

where ϑ is the viscosity solution of (1), V is the numerical approximation and $C(L, t_n, v_0)$ is a positive constant which depends only on t_n , on L (Lipschitz constant of $x \rightarrow f(x, \alpha)$, see Section 2.1), t_n , and on the total variation of v_0 (see Definition 2.2).

This is the first result of this kind to our knowledge, in the case of discontinuous viscosity solutions. Furthermore, in some particular cases, such as the eikonal equation $(\vartheta_t + |\vartheta_x| = 0)$, corresponding to $\mathcal{A} := \{\pm 1\}$ and $f(x, \alpha) = \alpha$, the constant C does not depend of t_n . This shows a "non-diffusive" behavior of the scheme, as identified in [9], for the advection case with constant sign velocity.

The paper is organized as follows: Sections 2 and 3 are devoted to the proof of (2) in the case of piece-wise constant initial data. In Section 4, we prove (2) for more general discontinuous initial data. In Section 5, we weaken some assumptions made before on the velocities and prove a similar estimate for a modified scheme. We conclude in Section 6 by some numerical illustrations, in particular showing the non-diffusive behavior of the proposed scheme. The appendix contains some useful technical results.

2 Preliminaries

2.1 Notations and preliminary results

We denote by $v_0 : \mathbb{R} \rightarrow \mathbb{R}$ a bounded lower semicontinuous (l.s.c.) function, \mathcal{A} a compact set, and $f : \mathbb{R} \times \mathcal{A} \rightarrow \mathbb{R}$ a continuous function satisfying:

$$\exists L \geq 0, \forall \alpha \in \mathcal{A}, \forall x, y \in \mathbb{R}, |f(y, \alpha) - f(x, \alpha)| \leq L|y - x|. \quad (3)$$

It is known that, under these assumptions, equation (1) admits a unique bounded l.s.c. *bilateral viscosity* solution [3, 2]. For convenience of the reader, we recall in the following definition the viscosity notion we use.

Definition 2.1. *A bounded l.s.c. function ϑ is a bilateral viscosity solution of (1) if,*

i) for any $\phi \in C^1((0, +\infty) \times \mathbb{R})$ and at any local minimum $(t, x) \in]0, +\infty[\times \mathbb{R}$ of $\vartheta - \phi$,

$$\phi_t(t, x) + \max_{\alpha \in \mathcal{A}}(f(x, \alpha) \phi_x(t, x)) = 0.$$

ii) $\liminf_{\substack{t \rightarrow 0^+ \\ y \rightarrow x}} \vartheta(t, y) = v_0(x), \quad \forall x \in \mathbb{R}.$

If we set for all $x \in \mathbb{R}$

$$f_m(x) := \min_{\alpha \in \mathcal{A}} f(x, \alpha) \quad \text{and} \quad f_M(x) := \max_{\alpha \in \mathcal{A}} f(x, \alpha),$$

then equation (1) can be rewritten in the equivalent form:

$$\vartheta_t(t, x) + \max(f_m(x) \vartheta_x(t, x), f_M(x) \vartheta_x(t, x)) = 0, \quad t > 0, x \in \mathbb{R}, \quad (4a)$$

$$\vartheta(0, x) = v_0(x), \quad x \in \mathbb{R}. \quad (4b)$$

Notice that by (3) and the definitions of f_m and f_M , we have $f_m(x) \leq f_M(x)$, $\forall x \in \mathbb{R}$, and also

(H1) f_m and f_M are L -Lipschitz continuous.

For simplicity of presentation, we first suppose the simplifying additional assumptions:

(H2) f_m and f_M are of constant sign,

(H3) f_m and f_M are increasing functions of x .

These assumptions will be weakened in Section 5.

Remark 2.1. Assumptions (H1)-(H3) are satisfied for the particular case of the Eikonal equation: $\vartheta_t + c|\vartheta_x| = 0$, where c is a given constant and $c \geq 0$ (taking $f_M(x) = -f_m(x) = c$).

We now define exact and approximated characteristics that will be very useful throughout the paper. Let $x_j := j \Delta x$ be a uniform mesh with $\Delta x > 0$ and $j \in \mathbb{Z}$, and denote:

$$x_{j+\frac{1}{2}} := (j + \frac{1}{2}) \Delta x, \quad \text{and} \quad I_j :=]x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}[.$$

As the dynamics f_m and f_M are lipschitz continuous, then for any $x \in \mathbb{R}$ we can define characteristics X_x^M and X_x^m as the solutions of the Cauchy problems:

$$\begin{cases} \dot{X}_x^M(t) = f_M(X_x^M(t)), \\ X_x^M(0) = x, \end{cases} \quad \text{and} \quad \begin{cases} \dot{X}_x^m(t) = f_m(X_x^m(t)), \\ X_x^m(0) = x. \end{cases} \quad (5)$$

We also define approximated piece-wise constant velocity functions f_M^S and f_m^S , such that, $\forall j \in \mathbb{Z}$:

$$f_M^S(x) = f_M(x_j), \quad \forall x \in I_j, \\ f_M^S(x_{j+\frac{1}{2}}) = \begin{cases} 0 & \text{if } f_M(x_j) f_M(x_{j+1}) \leq 0, \\ f_M(x_j) & \text{otherwise,} \end{cases}$$

and

$$f_m^S(x) = f_m(x_j), \quad \forall x \in I_j, \\ f_m^S(x_{j+\frac{1}{2}}) = \begin{cases} 0 & \text{if } f_m(x_j) f_m(x_{j+1}) \leq 0, \\ f_m(x_j) & \text{otherwise.} \end{cases}$$

In general, the differential equation

$$\dot{\chi}_x(t) = f_M^S(\chi_x(t)), \quad a.e. \ t \geq 0, \quad \chi_x(0) = x, \quad (6)$$

may have more than one absolutely continuous solution. The non-uniqueness comes from the behavior on boundary points ($x_{j+\frac{1}{2}}$) in the case when the velocity vanishes (or changes sign). Throughout this paper, we shall denote by $X_x^{M,S}$ the function defined by:

$$X_x^{M,S} \text{ solution of (6), and} \quad (7a)$$

$$\text{if } \exists t^* \geq 0 \text{ s.t. } \begin{cases} X_x^{M,S}(t^*) = x_{j+\frac{1}{2}}, \\ f_M(x_j) f_M(x_{j+1}) \leq 0 \end{cases} \text{ then } X_x^{M,S}(t) = x_{j+\frac{1}{2}} \quad \forall t \geq t^*. \quad (7b)$$

We have uniqueness of such solution (see appendix A). We construct $X_x^{m,S}$ in a similar way.

Lemma 2.1. *Assume that (H1) holds. Let a, b be in \mathbb{R} . The following assertions are satisfied:*

(i) *For every $t \geq 0$, we have:*

$$\max(|X_a^{M,S}(t) - X_a^M(t)|, |X_b^{m,S}(t) - X_b^m(t)|) \leq \frac{1}{2}Lt e^{Lt} \Delta x,$$

where L is the Lipschitz constant of f_M and f_m (see (H1)).

(ii) *Let $s \leq t$ and assume that $X_a^{m,S}(\theta) \geq X_b^{M,S}(\theta)$, for every $\theta \in [s, t]$. Then*

$$|X_a^{m,S}(t) - X_b^{M,S}(t)| + \Delta x \leq e^{L(t-s)}(|X_a^{m,S}(s) - X_b^{M,S}(s)| + \Delta x).$$

On the other hand, if $X_a^m(\theta) \geq X_b^M(\theta)$, for every $\theta \in [s, t]$, then

$$|X_a^m(t) - X_b^M(t)| \leq e^{L(t-s)}(|X_a^m(s) - X_b^M(s)|).$$

(iii) *Assume (H1) and (H3). If $a > b$, then the functions $t \mapsto X_a^{M,S}(t) - X_b^{m,S}(t)$ and $t \mapsto X_a^M(t) - X_b^m(t)$ are increasing for $t \geq 0$.*

(iv) *Assume (H1) – (H3). If $\frac{\Delta x}{2} < a - b$, then the function $t \mapsto X_a^{M,S}(t) - X_b^m(t)$ is increasing for $t \geq 0$.*

(v) *Assume (H1) – (H3). If $X_b^{m,S}(\theta) \geq X_a^{m,S}(\theta)$, for every $\theta \in [s, t]$. Then we have*

$$|X_b^{m,S}(t) - X_a^{m,S}(t)| + \Delta x \leq e^{L(t-s)}(|X_b^{m,S}(s) - X_a^{m,S}(s)| + \Delta x).$$

On the other hand, if $X_a^{M,S}(\theta) \leq X_b^{M,S}(\theta)$, for every $\theta \in [s, t]$, then

$$|X_b^{M,S}(t) - X_a^{M,S}(t)| + \Delta x \leq e^{L(t-s)}(|X_b^{M,S}(s) - X_a^{M,S}(s)| + \Delta x).$$

Proof. (i) Let $x \in \mathbb{R}$ and $j \in \mathbb{Z}$ such that $x \in I_j$. For $y \in \mathbb{R}$, the following inequality holds

$$\begin{aligned} |f_M^S(x) - f_M(y)| &= |f_M(x_j) - f_M(x) + f_M(x) - f_M(y)| \\ &\leq \frac{1}{2}L\Delta x + L|x - y|. \end{aligned}$$

Therefore, for all $t \geq 0$, we get:

$$\begin{aligned} |X_a^{M,S}(t) - X_a^M(t)| &= \left| \int_0^t (f_M^S(X_a^{M,S}(s)) - f_M(X_a^M(s))) ds \right| \\ &\leq \frac{1}{2}Lt\Delta x + L \int_0^t |X_a^{M,S}(s) - X_a^M(s)| ds. \end{aligned}$$

Hence by using Gronwall's Lemma we obtain the desired estimate for $|X_a^{M,S}(t) - X_a^M(t)|$. The bound for $|X_b^{m,S}(t) - X_b^m(t)|$ is obtained in the same way.

(ii) Let $\delta(\theta) := X_a^{m,S}(\theta) - X_b^{M,S}(\theta) + \Delta x$. We have

$$\begin{aligned} \frac{d}{d\theta}\delta(\theta) &= f_m^S(X_a^{m,S}(\theta)) - f_M^S(X_b^{M,S}(\theta)) \\ &\leq f_M^S(X_a^{m,S}(\theta)) - f_M^S(X_b^{M,S}(\theta)) \\ &\leq L\left(X_a^{m,S}(\theta) - X_b^{M,S}(\theta) + \Delta x\right) = L\delta(\theta). \end{aligned}$$

The result follows by using a Gronwall estimate. The proof for the other estimate is similar.

(iii) Define $\delta(t) := X_a^{M,S}(t) - X_b^{m,S}(t)$, and $t^* := \inf\{t > 0, \delta(t) < 0\}$. As $\delta(0) > 0$, then $t^* > 0$. Then for all $t \in [0, t^*[$, $\delta(t) \geq 0$ and we have:

$$\frac{d}{dt}\delta(t) = f_M^S(X_a^{M,S}(t)) - f_m^S(X_b^{m,S}(t)) \geq f_m^S(X_a^{M,S}(t)) - f_m^S(X_b^{m,S}(t)),$$

which is positive, for $t \in [0, t^*[$, by (H3). We deduce that δ is increasing on $[0, t^*[$. Suppose that t^* is finite, then we get by continuity of $X_a^{M,S}$ and $X_b^{m,S}$ that $\delta(t^*) \geq \delta(0) > 0$. This contradiction shows that $t^* = +\infty$ and δ is increasing for all $t \geq 0$. (The proof is similar for $t \mapsto X_a^M(t) - X_b^m(t)$.)

(iv) Similar arguments as in (iii)

(v) The proof is obtained as in (ii). \square

Lemma 2.2. *Let v_0 be a bounded l.s.c. function on \mathbb{R} , and assume that (H1) holds. Then, the unique viscosity solution of (4) is given by:*

$$\vartheta(t, x) = \min_{y \in [X_x^M(-t), X_x^m(-t)]} v_0(y), \quad \forall t > 0, x \in \mathbb{R}. \quad (8)$$

Proof. Notice that equation (4a) can be rewritten as follows

$$\vartheta_t(t, x) + \max_{\alpha \in [0,1]} \{((1-\alpha)f_m(x) + \alpha f_M(x)) \cdot \vartheta_x(t, x)\} = 0, \quad t > 0, x \in \mathbb{R}. \quad (9)$$

The unique viscosity solution of equation (9) satisfying the initial condition (4b) (see [2]) is given by

$$\vartheta(t, x) = \min_{\alpha \in L^\infty(\mathbb{R}^+, [0,1])} v_0(X_x^\alpha(-t)) = \min_{y \in [X_x^M(-t), X_x^m(-t)]} v_0(y),$$

where X_x^α is the solution of $X_x^\alpha(0) = x$ and $\dot{X}_x^\alpha(t) = (1 - \alpha(t))f_m(X_x^\alpha(t)) + \alpha(t)f_M(X_x^\alpha(t))$ for $t \geq 0$ with $\alpha \in L^\infty(\mathbb{R}^+, [0, 1])$. \square

We also consider the function ϑ^S which is defined in an analogous way as in (8), but with the approximated characteristics $X_x^{M,S}, X_x^{m,S}$ instead of X_x^M and X_x^m :

$$\vartheta^S(t, x) := \min_{y \in [X_x^{M,S}(-t), X_x^{m,S}(-t)]} v_0(y), \quad \forall t > 0, x \in \mathbb{R}. \quad (10)$$

This approximate function will play an important role throughout the paper.

Proposition 2.1. *Under assumption (H1), we have¹:*

$$\|\vartheta(t, \cdot) - \vartheta^S(t, \cdot)\|_{L^1(\mathbb{R})} \leq Lte^{Lt} TV(v_0) \Delta x. \quad (11)$$

Proof. By using Lemma B.1 (taking $a_x^1 = X_x^M(-t)$, $b_x^1 = X_x^m(-t)$, and $a_x^2 = X_x^{M,S}(-t)$, $b_x^2 = X_x^{m,S}(-t)$) (whose inverse functions are $X_x^M(t)$, $X_x^m(t)$, and $X_x^{M,S}(t)$, $X_x^{m,S}(t)$ respectively) together with Lemma 2.1 (i), we obtain the L^1 -norm estimate. \square

Using Lemma B.3, we also obtain

Proposition 2.2. *Under assumption (H1),*

$$TV(\vartheta^S(t, \cdot)) \leq TV(v_0), \quad \forall t \geq 0.$$

For $E \subset \mathbb{R}$ a given set, we shall use in all the sequel the notation $\mathbf{1}_E$ for the function defined by:

$$\mathbf{1}_E(x) := \begin{cases} 1 & \text{if } x \in E, \\ 0 & \text{otherwise.} \end{cases}$$

2.2 The HJB-UltraBee scheme

Let $\Delta t > 0$ be a constant time step, and $t_n := n\Delta t$ for $n \geq 0$. We set the following notation for local "CFL" numbers:

$$\nu_j^m := \frac{\Delta t}{\Delta x} f_m(x_j) \quad \text{and} \quad \nu_j^M := \frac{\Delta t}{\Delta x} f_M(x_j),$$

and $\nu^m = \{\nu_j^m, j \in \mathbb{Z}\}$, $\nu^M = \{\nu_j^M, j \in \mathbb{Z}\}$. An adaptation of the UltraBee scheme has been proposed and numerically tested for the HJB equation [7, 5, 6]. Let us recall this formulation. We first introduce the notation for the average values of initial data:

$$V_j^0 := \frac{1}{\Delta x} \int_{x_{j-\frac{1}{2}}}^{x_{j+\frac{1}{2}}} v_0(x) dx, \quad j \in \mathbb{Z}. \quad (12)$$

As we deal with an explicit scheme, we will assume in all the paper that the mesh size satisfies the CFL condition:

$$\max_{x \in \mathbb{R}} \max(|f_m(x)|, |f_M(x)|) \frac{\Delta t}{\Delta x} \leq 1. \quad (13)$$

¹Throughout this paper, we will use the following definition for the variation of a real-valued function w .

Definition 2.2. *Let w be a real-valued function. the total variation of w is defined by:*

$$TV(w) := \sup \left\{ \sum_{j=1, \dots, k} |w(y_{j+1}) - w(y_j)|; k \in \mathbb{N}^*, \text{ and } (y_j)_{1 \leq j \leq k+1} \text{ increasing sequence} \right\}.$$

Algorithm 1 :

Initialization: We compute the initial averages $V^0 = (V_j^0)_{j \in \mathbb{Z}}$ as defined in (12).

Loop: For $n \geq 0$, We compute $V^{n+1} = (V_j^{n+1})_{j \in \mathbb{Z}}$ by:

- Define “fluxes” $V_{j+\frac{1}{2}}^n(\nu)$ for $\nu \in \{\nu^m, \nu^M\}$ as follows:
If $\nu_j \geq 0$ for every $j \in \mathbb{Z}$, define:

$$V_{j+1/2}^n(\nu) := \begin{cases} \min(\max(V_{j+1}^n, b_j^+(\nu)), B_j^+(\nu)) & \text{if } \nu_j > 0 \\ V_{j+1}^n & \text{if } \nu_j = 0 \text{ and } V_j^n \neq V_{j-1}^n \\ V_j^n & \text{if } \nu_j = 0 \text{ and } V_j^n = V_{j-1}^n, \end{cases}$$

where

$$\begin{cases} b_j^+(\nu) := \max(V_j^n, V_{j-1}^n) + \frac{1}{\nu_j}(V_j^n - \max(V_j^n, V_{j-1}^n)), \\ B_j^+(\nu) := \min(V_j^n, V_{j-1}^n) + \frac{1}{\nu_j}(V_j^n - \min(V_j^n, V_{j-1}^n)), \end{cases} \quad (14)$$

If $\nu_j \leq 0$ for every $j \in \mathbb{Z}$, define:

$$V_{j-1/2}^n(\nu) := \begin{cases} \min(\max(V_{j-1}^n, b_j^-(\nu)), B_j^-(\nu)) & \text{if } \nu_j < 0 \\ V_{j-1}^n & \text{if } \nu_j = 0 \text{ and } V_j^n \neq V_{j+1}^n \\ V_j^n & \text{if } \nu_j = 0 \text{ and } V_j^n = V_{j+1}^n, \end{cases}$$

where

$$\begin{cases} b_j^-(\nu) := \max(V_j^n, V_{j+1}^n) + \frac{1}{|\nu_j|}(V_j^n - \max(V_j^n, V_{j+1}^n)), \\ B_j^-(\nu) := \min(V_j^n, V_{j+1}^n) + \frac{1}{|\nu_j|}(V_j^n - \min(V_j^n, V_{j+1}^n)). \end{cases} \quad (15)$$

- For $\nu \in \{\nu^m, \nu^M\}$, let

$$V_j^{n+1}(\nu) := V_j^n - \nu_j \left(V_{j+\frac{1}{2}}^n(\nu) - V_{j-\frac{1}{2}}^n(\nu) \right).$$

- Finally, set $V_j^{n+1} := \min(V_j^{n+1}(\nu^m), V_j^{n+1}(\nu^M))$ for every $j \in \mathbb{Z}$.

In all the sequel, we shall use the notation:

$$\mathcal{S}_{UB}(V^n) := \left(\min_{j \in \mathbb{Z}} (V_j^{n+1}(\nu^m), V_j^{n+1}(\nu^M)) \right).$$

Under assumption (H2), we notice that the resulting scheme is well defined. We associate to the scheme values $(V_j^n)_j$, the l.s.c. step function $V(t_n, \cdot)$ defined for every $t_n \geq 0$, $x \in \mathbb{R}$ by

$$V(t_n, x) := \begin{cases} V_j^n & \text{if } x \in]x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}[, \\ \min(V_j^n, V_{j+1}^n) & \text{if } x = x_{j+\frac{1}{2}}. \end{cases} \quad (16)$$

2.3 A first case where fronts do not meet

We consider here the case of an initial data of the form

$$v_0(x) := \mathbf{1}_{]-\infty, b[}(x) + \mathbf{1}_{]a, +\infty[}(x), \quad (17)$$

where $a, b \in \mathbb{R} \cup \{\pm\infty\}$ and $a > b$. Our aim in this section is to study in this simple case the relationship between the viscosity solution at time t_n , $\vartheta(t_n, \cdot)$ and the values V^n computed by the HJB-UB algorithm. We show, under the CFL condition, an L^1 -error estimate in Δx stated in the following theorem:

Theorem 2.1. *We assume that (H1) – (H3) and the CFL condition (13) are satisfied. Let v_0 be defined by (17), Δx be such that $a \geq b + 3\Delta x$, and ϑ be the viscosity solution of (4). Then*

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (1 + Lt_n e^{Lt_n}) TV(v_0) \Delta x, \quad \forall n \geq 0. \quad (18)$$

Before dealing with the proof of Theorem 2.1, it will be useful to have the analytic expression of the viscosity solution ϑ (known in this case, since v_0 has a simple form):

Remark 2.2. *Assume assumption (H1) is satisfied, $a > b$, and v_0 as in (17). Then using Lemma 2.2 and by a direct calculation we obtain that the l.s.c. viscosity solution of (4) is given by :*

$$\vartheta(t, x) := \mathbf{1}_{]-\infty, X_b^m(t)[}(x) + \mathbf{1}_{]X_a^M(t), +\infty[}(x). \quad \forall t \geq 0, x \in \mathbb{R}.$$

Also, the function ϑ^S defined by (10) satisfies:

$$\vartheta^S(t, x) = \mathbf{1}_{]-\infty, X_b^{m,S}(t)[}(x) + \mathbf{1}_{]X_a^{M,S}(t), +\infty[}(x), \quad \forall t \geq 0, x \in \mathbb{R}.$$

In the following we denote by $\bar{\vartheta}^{S,n}$ the cell averages of $\vartheta^S(t_n, \cdot)$, defined by

$$\bar{\vartheta}_j^{S,n} := \frac{1}{\Delta x} \int_{I_j} \vartheta^S(t_n, x) dx \quad \text{for } j \in \mathbb{Z}, n \in \mathbb{N}. \quad (19)$$

Lemma 2.3. *Assume that (H1) – (H3) hold, v_0 is defined by (17), and $a \geq b + 3\Delta x$. Then the values V_j^n computed by algorithm 1 satisfy:*

$$V_j^n = \bar{\vartheta}_j^{S,n}, \quad \forall n \geq 0, \forall j \in \mathbb{Z}.$$

Proof. Case 1. We consider the case of $v_0(x) := \mathbf{1}_{]a, +\infty[}(x)$, and proceed by recursion on $n \geq 0$ (the case $v_0(x) = \mathbf{1}_{]-\infty, b[}(x)$ may be treated in a similar way). Let $j \in \mathbb{Z}$ be such that the discontinuity position $x_n := X_a^{M,S}(t_n)$ lies in $]x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}[$.

Then we have $V_k^n = 0$ for $k < j$, $V_j^n \in [0, 1[$, and $V_k^n = 1$ for $k > j$. By straightforward calculations, if $\nu^M \geq 0$, we can verify that

$$\begin{aligned} \bar{\vartheta}_{j-1}^{S,n+1} = V_{j-1}^{n+1} &= 0, \\ \bar{\vartheta}_j^{S,n+1} = V_j^{n+1} &= \max(0, V_j^n - \nu_j^M), \\ \bar{\vartheta}_{j+1}^{S,n+1} = V_{j+1}^{n+1} &= \begin{cases} 1 - \frac{\nu_{j+1}^M}{\nu_j^M} \max(0, \nu_j^M - V_j^n) & \text{if } \nu_j^M > 0 \\ 1 & \text{if } \nu_j^M = 0 \end{cases} \end{aligned}$$

and, if $\nu^M \leq 0$,

$$\begin{aligned} \bar{\vartheta}_{j-1}^{S,n+1} = V_{j-1}^{n+1} &= \begin{cases} \frac{|\nu_{j-1}^M|}{|\nu_j^M|} (\max(1, V_j^n + |\nu_j^M|) - 1) & \text{if } \nu_j^M < 0, \\ 0 & \text{if } \nu_j^M = 0 \end{cases} \\ \bar{\vartheta}_j^{S,n+1} = V_j^{n+1} &= \min(1, V_j^n + |\nu_j^M|), \\ \bar{\vartheta}_{j+1}^{S,n+1} = V_{j+1}^{n+1} &= 1, \end{aligned}$$

and in all cases, $\bar{\vartheta}_k^{S,n+1} = V_k^{n+1} = 0$, $\forall k \leq j - 2$, and $\bar{\vartheta}_k^{S,n+1} = V_k^{n+1} = 1$, $\forall k \geq j + 2$. This means that the HJB-UltraBee scheme computes exactly $V^{n+1}(\nu^M)$ from V^n for the advection with velocity f_M^S . In the same way we obtain that $V^{n+1}(\nu^m)$ computes exactly the average values from V^n for the advection with velocity f_m^S . Hence we have, for every $k \in \mathbb{Z}$,

$$V_k^{n+1}(\nu^M) = \frac{1}{\Delta x} \int_{I_k} \mathbf{1}_{]X_{x_n}^{M,S}(\Delta t), \infty[}(x) dx, \quad (20)$$

$$V_k^{n+1}(\nu^m) = \frac{1}{\Delta x} \int_{I_k} \mathbf{1}_{]X_{x_n}^{m,S}(\Delta t), \infty[}(x) dx. \quad (21)$$

Since $X_{x_n}^{m,S}(\Delta t) \leq X_{x_n}^{M,S}(\Delta t)$ we deduce that $V_k^{n+1}(\nu^M) \leq V_k^{n+1}(\nu^m)$, and, for all $k \in \mathbb{Z}$,

$$V_k^{n+1} = V_k^{n+1}(\nu^M).$$

This concludes the proof of $\bar{\vartheta}^{S,n+1} = V^{n+1}$.

Case 2. Consider the case of $v_0(x) := \mathbf{1}_{]-\infty, b[}(x) + \mathbf{1}_{]a, \infty[}(x)$. As $a - b \geq 3\Delta x$, by Lemma 2.1 (iii), we get $X_a^{M,S}(t) \geq 3\Delta x + X_b^{m,S}(t)$ for $t \geq 0$. This means that there are at least two successive cells with value $V_j^n = V_{j+1}^n = 0$ separating $X_b^{m,S}(t_n)$ and $X_a^{M,S}(t_n)$. Then as in Case 1 we obtain for $k \geq j + 1$ that (20) and (21) are also valid, and thus

$$\text{for } k \geq j + 1, \quad V_k^{n+1} = V_k^{n+1}(\nu^M) = \bar{\vartheta}_k^{S,n+1}$$

(i.e., an exact evolution following the discontinuity position $X_a^{M,S}(t_{n+1})$). Also, in the same way as in Case 1, we obtain

$$\text{for } k \leq j, \quad V_k^{n+1} = V_k^{n+1}(\nu^m) = \bar{\vartheta}_k^{S,n+1}$$

(i.e, an exact evolution following the discontinuity position $X_b^{m,S}(t_{n+1})$). This concludes to $V_k^{n+1} = \bar{v}_k^{S,n+1}$ for all $k \in \mathbb{Z}$. \square

Proof of Theorem 2.1: Since $V_j^n = \bar{v}_j^{S,n}$, for all $j \in \mathbb{Z}$ and $n \geq 0$, we obtain:

$$\|\vartheta^S(t_n, \cdot) - V(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq \Delta x TV(\vartheta^S(t_n, \cdot)) = 2\Delta x = TV(v_0) \Delta x. \quad (22)$$

Inequalities (22) and (11) lead to the desired result (18). \square

Remark 2.3. Lemma 2.3 shows the behaviour of the HJB-UltraBee scheme: when the two discontinuities are far from each other, the algorithm is able to recover, from the average values, their exact positions $X_a^{M,S}(t)$ and $X_b^{m,S}(t)$. Then the scheme makes these discontinuities evolve with the piece-wise constant velocities f_M^S and f_m^S . This is due to the fact that, as long as the discontinuities are separated by more than $3\Delta x$ from each other, the extrema of ϑ^S can be identified by the scheme.

This interpretation of the scheme extends some results of [11] (for the advection case) to HJB equations. We will see in the next section that this exact computation of the averages of ϑ^S is no more possible when two discontinuities lie in two successive cells, or in the same cell.

3 Case of piece-wise constant initial data

In this section we assume that there exists an increasing sequence of real values $(y_i)_{i=0, \dots, p+1}$ with $y_0 = -\infty$, $y_{p+1} = +\infty$, and $(\gamma_i)_{i=0, \dots, p}$ such that

$$v_0(x) = \begin{cases} \sum_{i=0, \dots, p} \gamma_i \mathbf{1}_{]y_i, y_{i+1}[}(x) & \text{for } x \in \mathbb{R} \setminus \{y_1, \dots, y_p\}, \\ \min(\gamma_{i-1}, \gamma_i) & \text{for } x = y_j \text{ and for } i = 1, \dots, p. \end{cases} \quad (23)$$

With this definition, v_0 is a l.s.c. piece-wise constant function. We also assume that the mesh size Δx satisfies:

$$\Delta x \leq \frac{1}{3} \min_{1 \leq i \leq p-1} (y_{i+1} - y_i). \quad (24)$$

We have derived in the previous section an error estimate when the discontinuities keep far enough from each other (more precisely, when they are separated by at least two entire cell intervals). In general, two discontinuities may become very close. Two critical cases may happen, see Figure 1.

In these cases, one time step of the UltraBee scheme given in algorithm 1 may not compute the average values exactly. In fact, it would do a false interpretation of the maximum value and of the discontinuity localization. The idea is to anticipate this critical situation. Hence when two discontinuities are too close, a truncation is done such that just one discontinuity remains in its right location (see Fig. 2). Here we modify the HJB-UltraBee scheme around maxima when one of the two critical cases of Fig. 1 occur.




Figure 1: Critical cases of truncation

Algorithm 2

Initialization: We compute the averages $(V_j^0)_{j \in \mathbb{Z}}$ by (12)

Loop: For $n \geq 0$:

A) Compute $W := \mathcal{S}_{UB}(V^n)$ (HJB-UltraBee step).

B) (Truncation step)

• For all indexes j such that

$$\left\{ \begin{array}{l} W_j > \max(W_{j-1}, W_{j+1}), \text{ and } W_j = V_j^n \\ \text{or} \\ W_j > W_{j-1}, W_{j+1} > W_{j+2}, \text{ and } W_j < V_j^n \end{array} \right\} \text{ and } V_j^n - W_{j-2} < V_j^n - W_{j+2},$$

set

$$\begin{aligned} V_{j-1}^{n+1} &:= W_{j-2}, & V_j^{n+1} &:= W_{j-2}, \\ \text{and } V_{j+1}^{n+1} &:= W_{j+2} + \frac{W_{j-2} - W_{j+2}}{V_j^n - W_{j+2}}(W_{j+1} - W_{j+2}), \end{aligned}$$

• For all indexes j such that

$$\left\{ \begin{array}{l} W_j > \max(W_{j-1}, W_{j+1}), \text{ and } W_j = V_j^n \\ \text{or} \\ W_{j-1} > W_{j-2}, W_j > W_{j+1}, \text{ and } W_j < V_j^n \end{array} \right\} \text{ and } V_j^n - W_{j-2} \geq V_j^n - W_{j+2},$$

set

$$\begin{aligned} V_{j+1}^{n+1} &:= W_{j+2}, & V_j^{n+1} &:= W_{j+2}, \\ \text{and } V_{j-1}^{n+1} &:= W_{j-2} + \frac{W_{j+2} - W_{j-2}}{V_j^n - W_{j-2}}(W_{j-1} - W_{j-2}). \end{aligned}$$

• Otherwise set $V_j^{n+1} := W_j$.




Figure 2: Truncation step for critical case 2

Hereafter the truncation step will be denoted by

$$V^{n+1} := T_{V^n}(W).$$

Remark 3.1. *The truncation step modifies values only near local strict maxima of $(V_j^n)_{j \in \mathbb{Z}}$.*

The test $V_j^n - W_{j-2} < V_j^n - W_{j+2}$ (resp. $V_j^n - W_{j-2} \geq V_j^n - W_{j+2}$) allows to check if the left jump is strictly smaller (resp. equal or greater) than the right jump near a local maxima, see Figure 2. In this case, the truncation corresponds to a recomputation of the average values such that the right (resp. left) discontinuity position is correctly coded. Hence the truncation allows to get rid of the left discontinuity and to keep the right one in its correct location. This truncation step aims to improve the treatment by the UltraBee scheme and to prevent the presence of two discontinuities in the same cell or in adjacent² cells.

The main result of the section is the following.

²We mean by adjacent here the neighboring cells from the left and from the right

Theorem 3.1. *Assume that (H1) – (H3) are satisfied. Let v_0 be a piece-wise constant function as in (23). We assume that the mesh size satisfies (13) and (24). Let ϑ be the viscosity solution of (4), and let V be defined by (16) and algorithm 2. We have*

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (Lt_n + 4)e^{Lt_n} TV(v_0) \Delta x, \quad \forall n \geq 0.$$

Notice that the total variation of v_0 given by (23) is $TV(v_0) := \sum_{i=1, \dots, p-1} |\gamma_{i+1} - \gamma_i|$.

Remark 3.2. *We shall also see in the proof, for instance in the case of the eikonal equation, that $\bar{\vartheta}_j^{S,n} = V_j^n$ as long as the discontinuities are far enough one from each other.*

3.1 A first simple case when two fronts may meet

Let a, b be in \mathbb{R} , with $b \geq a + 3\Delta x$. Consider the following initial data:

$$v_0(x) = \mathbf{1}_{]a, b[}(x), \quad a.e. x \in \mathbb{R}. \quad (25)$$

Remark 3.3. *Thanks to (8), under assumption (H1), the unique l.s.c. viscosity solution of (4) is given by (for $t \geq 0, x \in \mathbb{R}$):*

$$\vartheta(t, x) := \begin{cases} \mathbf{1}_{]X_a^M(t), X_b^m(t)[}(x), & \text{if } X_a^M(t) < X_b^m(t), \\ 0 & \text{otherwise.} \end{cases} \quad (26)$$

Also, by definition of ϑ^S (see (10)), for $t \geq 0$ and $x \in \mathbb{R}$ we get

$$\vartheta^S(t, x) := \begin{cases} \mathbf{1}_{]X_a^{M,S}(t), X_b^{m,S}(t)[}(x), & \text{if } X_a^{M,S}(t) < X_b^{m,S}(t), \\ 0 & \text{otherwise} \end{cases} \quad (27)$$

(the approximated characteristics $X_a^{M,S}$ and $X_b^{m,S}$ are defined as in (7)).

As in (19) we denote by $\bar{\vartheta}^{S,n}$ the average cell values of $\vartheta^S(t_n, \cdot)$.

Lemma 3.1. *Assume that (H1) – (H3) are satisfied. Let v_0 be as in (25), and the mesh size satisfy (13) and $\Delta x \leq \frac{b-a}{3}$. For every $n \geq 0$, we have*

$$\|V(t_n, \cdot) - \bar{\vartheta}^{S,n}(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq 4\Delta x e^{Lt_n}. \quad (28)$$

Proof. For $n \in \mathbb{N}$, let j_n and ℓ_n be two integers such that $X_a^{M,S}(t_n) \in]x_{j_n - \frac{1}{2}}, x_{j_n + \frac{1}{2}}]$ and $X_b^{m,S}(t_n) \in]x_{\ell_n - \frac{1}{2}}, x_{\ell_n + \frac{1}{2}}]$.

Two cases may occur:

- (i) $\ell_n \geq j_n + 3 \forall n \geq 0$.
- (ii) There exists a first index $n \geq 1$ such that $\ell_n < j_n + 3$.

Assume (i). By the results of section 2.3 we know that the scheme computes the exact averages of ϑ^S as long as the fronts are separated by at least two cells, that is, in the case for all $k \leq n$, $\ell_k \geq j_k + 3$. In particular, no truncation step has occurred in this case. We then have $V_j^n = \overline{\vartheta_j^{S,n}}$ and can conclude as in (22) to

$$\|\vartheta^S(t_n, \cdot) - V(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq \Delta x TV(\vartheta^S(t_n, \cdot)) = 2\Delta x = TV(v_0) \Delta x. \quad (29)$$

Assume now (ii). For $k < n$, the estimate (29) holds (no truncation done yet). For $k \geq n$, a truncation has occurred at step n and thus $V^k = 0$. For $k \geq n$, we then have (using Lemma 2.1(ii)) $\|V(t_k, \cdot) - \vartheta^S(t_k, \cdot)\|_{L^1} = \|\vartheta^S(t_k, \cdot)\|_{L^1} = \max(0, X_b^{m,S}(t_k) - X_a^{M,S}(t_k)) \leq e^{L(t_k - t_n)}(X_b^{m,S}(t_n) - X_a^{M,S}(t_n) + \Delta x) \leq e^{Lt_k} 4\Delta x$. This gives the desired bound. \square

Proof of theorem 3.1 in the case v_0 is given by (25): It is now a simple consequence of (28) and of Proposition 2.1.

3.2 Proof of Theorem 3.1 in the general case

We notice that since v_0 is a step function, $\vartheta^S(t, \cdot)$ is also a step function. We need to define a truncation ϑ^\sharp of ϑ^S that will be connected to the scheme values.

For a given piece-wise constant l.s.c. function w (of the form (23)), we define the truncation function $Trunc(w)$ as follows. For $x \in \mathbb{R}$, set

$$\begin{aligned} z_1^x &:= \sup\{z, z \leq x, w(z) \neq w(x)\} \in [-\infty, \infty[, \\ z_2^x &:= \inf\{z, z \geq x, w(z) \neq w(x)\} \in]-\infty, \infty] \end{aligned}$$

(i.e. the closest left and right discontinuities of w to x). Let j_1 be such that $z_1^x \in]x_{j_1 - \frac{1}{2}}, x_{j_1 + \frac{1}{2}}]$ and j_2 be such that $z_2^x \in [x_{j_2 - \frac{1}{2}}, x_{j_2 + \frac{1}{2}}[$. Then set (see Fig. 2(b)):

$$Trunc(w)(x) := \begin{cases} \max(w(z_1^x), w(z_2^x)) & \text{if } j_2 \in \{j_1 + 1, j_1 + 2\} \\ & \text{and } w(x) > \max(w(z_1^x), w(z_2^x)), \\ w(x) & \text{otherwise.} \end{cases}$$

We now define the function ϑ^\sharp by:

- $\vartheta^\sharp(0, \cdot) := v_0$, and
- $\forall n \geq 0$, $\vartheta^\sharp(t_{n+1}, \cdot) = Trunc(w_{n+1})$ where

$$w_{n+1}(x) := \min_{y \in [X_x^{M,S}(-\Delta t), X_x^{m,S}(-\Delta t)]} \vartheta^\sharp(t_n, y). \quad (30)$$

In the next result we derive an L^1 -error bound for $\vartheta^\sharp - \vartheta^S$. We also prove that the cell averages of ϑ^\sharp are exactly the values given by algorithm 2.

Lemma 3.2. *Assume (H1)-(H3).*

(i) $\forall n \geq 0$, $\|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (4e^{Lt_n} - 1)TV(v_0) \Delta x$.

(ii) $\forall j \in \mathbb{Z}$, $\forall n \geq 0$, we have

$$\frac{1}{\Delta x} \int_{I_j} \vartheta^\sharp(t_n, x) dx = V_j^n.$$

Proof. (i) First we notice that for $t \geq 0$, $\vartheta^\sharp(t, \cdot)$ is a piece-wise constant l.s.c function (this can be proved, as for ϑ^S , by using a recursion argument).

At a given time t_n , let $(] \alpha_i, \beta_i[)_{i=1, \dots, p}$ be the local maxima intervals of $\vartheta^S(t_n, \cdot)$ (i.e. $\vartheta^S(t_n, x) \equiv \text{const} = \mu_i$ on $] \alpha_i, \beta_i[$ and $\mu_i > \max(\vartheta^S(t_n, \alpha_i), \vartheta^S(t_n, \beta_i))$).

Then we obtain

- (a) $\vartheta^\sharp(t_n, \cdot)$ and $\vartheta^S(t_n, \cdot)$ can only differ on the intervals $\cup_{i=1, \dots, p}] \alpha_i, \beta_i[$,
- (b) $\forall i$, if $\vartheta^\sharp(t_n, \cdot)$ and $\vartheta^S(t_n, \cdot)$ differ on $] \alpha_i, \beta_i[$, then $\forall x \in] \alpha_i, \beta_i[$, $\vartheta^\sharp(t_n, x) = \max(\vartheta^S(t_n, \alpha_i), \vartheta^S(t_n, \beta_i))$.

Indeed, by Lemma 2.1(iii), the length of the minima regions of ϑ^S can only increase, so they will not disappear and create new local extrema; Also by (H3) the fronts of an increasing region of ϑ^S cannot get closer, as well as the fronts of a decreasing region of ϑ^S .

Now if $\vartheta^S(t_n, \cdot)$ and $\vartheta^\sharp(t_n, \cdot)$ differ on some interval $] \alpha_i, \beta_i[$, we have for $x \in] \alpha_i, \beta_i[$:

$$\begin{aligned} \|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(] \alpha_i, \beta_i[)} &= \left(\vartheta^S(t_n, x) - \max(\vartheta^S(t_n, \alpha_i), \vartheta^S(t_n, \beta_i)) \right) |\beta_i - \alpha_i| \\ &\leq TV(\vartheta^S(t_n, \cdot);] \alpha_i, \beta_i[) |\beta_i - \alpha_i|. \end{aligned}$$

Also we know that a truncation has occurred at some time $t_k < t_n$. At time t_k , the discontinuity positions corresponding to α_i and β_i are located in $\alpha_i^0 := X_{\alpha_i}^{M,S}(-t_n - t_k)$ and $\beta_i^0 := X_{\beta_i}^{m,S}(-t_n - t_k)$ respectively. Since the truncation has occurred, α_i^0 and β_i^0 are separated by less than two cell intervals, and $|\beta_i^0 - \alpha_i^0| \leq 3\Delta x$. By Lemma 2.1(ii), we thus have

$$\begin{aligned} |\beta_i - \alpha_i| &\leq e^{L(t_n - t_k)} (|\beta_i^0 - \alpha_i^0| + \Delta x) - \Delta x \\ &\leq (4e^{Lt_n} - 1)\Delta x. \end{aligned}$$

Summing these bounds for all local maxima intervals $] \alpha_i, \beta_i[$ we obtain $\|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (4e^{Lt_n} - 1)\Delta x TV(\vartheta^S(t_n, \cdot))$. Then we conclude the proof of (i) using Proposition. 2.2.

(ii) Now, we prove recursively that for all $n \geq 0$:

$$(P_n) \quad V_j^n = \frac{1}{\Delta x} \int_{I_j} \vartheta^\sharp(t_n, x) dx, \quad \forall j \in \mathbb{Z}$$

and

$$(Q_n) \quad \left\{ \begin{array}{l} \text{Any two successive discontinuity positions in } \vartheta^\sharp(t_n, \cdot) \\ \text{are separated by at least two cell intervals} \end{array} \right.$$

((Q_n) amounts to have $\vartheta^\sharp(t_n, \cdot) = \sum_i \mu_i^n \mathbf{1}_{]y_i, y_{i+1}[}$ a.e. , with $y_i \in]x_{j_1^i - \frac{1}{2}}, x_{j_1^i + \frac{1}{2}}]$, $y_{i+1} \in]x_{j_2^i - \frac{1}{2}}, x_{j_2^i + \frac{1}{2}}[$ and $j_2^i \geq j_1^i + 3$.)

For $n = 0$, $\vartheta^\sharp(0, \cdot) = v_0 = \vartheta(0, \cdot)$ and (P_0) follows, also (Q_0) is true by the definition of v_0 and of the mesh step Δx .

Now let us suppose (P_n) and (Q_n). Let $W := \mathcal{S}_{UB}(V^n)$, and define w^{n+1} as in (30). Since V^n codes the average values of $\vartheta^\sharp(t_n, \cdot)$ where the discontinuity positions are separated by at least two cells, the Ultra-Bee scheme computes the correct averages for one time-step evolution, and we will have

$$W_j = \frac{1}{\Delta x} \int_{I_j} w^{n+1}(x) dx$$

(this uses the same arguments as in the proof of Lemma 2.3).

As long as the discontinuity positions in w^{n+1} are separated by at least two entire cells, $\vartheta^\sharp(t_{n+1}, \cdot) = \text{Trunc}(w^{n+1}) = w^{n+1}$, and we have (Q_{n+1}). Also V_j^{n+1} values in step B) of algorithm 2 are unchanged. Hence $V_j^{n+1} = W_j$ which proves (P_{n+1}).

Because of assumption (H3), the only case the discontinuity positions of w^{n+1} may be separated by less than two entire cells is around the local maxima. (Recall that by (H3), discontinuity positions in a monotonous part of $\vartheta^\sharp(t_n, \cdot)$ can only get far from each other. This is also true for discontinuity positions around a local minimum, by Lemma 2.1(iii)).

Now let us assume that $\vartheta^\sharp(t_{n+1}, x) := \text{Trunc}(w^{n+1})(x) \neq w^{n+1}(x)$ for some $x \in \mathbb{R}$. This means that x is surrounded by two discontinuities of w^{n+1} , denoted z_1^x and z_2^x , which are separated by one cell (critical case 1), or which lie in two successive cells (critical case 2), see Fig. 1. Let us consider for instance that we have a critical case 2 (critical case 1 being similar). We may also assume that $V_j^n - W_{j-2} < V_j^n - W_{j+2}$ as illustrated in Fig. 2.

In this case we see that the average values in cells I_{j-1} and I_j can be set to W_{j-2} (hence the redefinition $V_{j-1}^{n+1} = V_j^{n+1} := W_{j-2}$ in algorithm 2). Then the remaining discontinuity position $\bar{x} \in I_{j+1}$ can be computed by two different ways: first, by

$$\frac{\bar{x} - x_{j+\frac{1}{2}}}{\Delta x} = \frac{W_{j+1} - W_{j+2}}{V_j^n - W_{j+2}},$$

and second, if V_{j+1}^{n+1} codes the correct average value on I_{j+1} after truncation,

$$\frac{\bar{x} - x_{j+\frac{1}{2}}}{\Delta x} = \frac{V_{j+1}^{n+1} - W_{j+2}}{W_{j-2} - W_{j+2}}$$

(recall that the scheme values should code the average of an exact piece-wise function). Thus we obtain the desired definition of V_j^{n+1} in terms of the (W_k) as in algorithm 2 B), which proves (P_{n+1}).

On the other hand, after the truncation step, there are no more critical cases in $\vartheta^\sharp(t_{n+1}, \cdot)$ and thus we have (Q_{n+1}). \square

Proof of theorem 3.1 in the general piece-wise constant case: Combining Proposition 2.1 and Lemma 3.2(i) we obtain

$$\begin{aligned} \|\vartheta(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} &\leq \|\vartheta(t_n, \cdot) - \vartheta^S(t_n, \cdot)\|_{L^1(\mathbb{R})} + \|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} \\ &\leq (Lt_n + 4)e^{Lt_n} - 1)TV(v_0) \Delta x. \end{aligned} \quad (31)$$

Then, using Lemma 3.2(ii), we obtain

$$\|\vartheta^\sharp(t_n, \cdot) - V(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq TV(\vartheta^\sharp(t_n, \cdot))\Delta x. \quad (32)$$

By construction of ϑ^\sharp (and using Lemma B.3), we also have $TV(\vartheta^\sharp(t, \cdot)) \leq TV(\vartheta^S(t, \cdot)) \leq TV(v_0)$. Together with (31), (32) we obtain the desired bound of Theorem 3.1. \square

4 Case of a general discontinuous initial data

In this section we generalize Theorem 3.1 (where we supposed that v_0 was a piece-wise constant function) to more general discontinuous initial data v_0 .

We assume that $v_0 : \mathbb{R} \rightarrow \mathbb{R}$ is a l.s.c. function, with $TV(v_0) < \infty$, and that v_0 has a finite number of extrema. More precisely we consider the following assumption :

$$(H4) \left\{ \begin{array}{l} \text{There exist } A_1, \dots, A_{q+1} \text{ and } B_1, \dots, B_q \text{ real numbers with} \\ A_1 = -\infty \leq B_1 < A_2 < \dots < B_q \leq A_{q+1} = +\infty, \\ \text{(with possibly } B_1 = -\infty \text{ or } B_q = +\infty), \text{ such that } v_0 \nearrow \text{ on each } [A_i, B_i[, \\ v_0 \searrow \text{ on each }]B_i, A_{i+1}], \text{ and } v_0(B_i) = \min(v_0(B_i^-), v_0(B_i^+)). \end{array} \right.$$

In particular A_i are local minima of v_0 , and B_i are local maxima of v_0 .

We also consider Δx small enough such that the minima and maxima of v_0 are separated at least by $3\Delta x$:

$$\Delta x < \frac{1}{3} \min_{i=1, \dots, q} \min(B_i - A_i, A_{i+1} - B_i). \quad (33)$$

Let a regular grid with mesh size Δx , and let $\Delta t > 0$.

We denote by v_0^P the l.s.c. function associated to v_0 as follows. We set $U_j :=]x_{3j-\frac{1}{2}}, x_{3j+\frac{5}{2}}[$, for all $j \in \mathbb{Z}$,

- If $\overline{U_j} \cap \{(A_k)_{k=2, \dots, q}, (B_k)_{k=1, \dots, q}\} = \emptyset$, set

$$v_0^P(x) := \frac{1}{3\Delta x} \int_{U_j} v_0(y) dy, \quad \forall x \in U_j. \quad (34a)$$

- otherwise if $A_k \in \overline{U_j}$ (resp. $B_k \in \overline{U_j}$) set

$$v_0^P(x) := v_0(A_k) \text{ (resp. } v_0(B_k)) \quad \forall x \in U_j. \quad (34b)$$

- Extend v_0^P by lower semi-continuity :

$$v_0^P(x_{3j-\frac{1}{2}}) = \min\left(v_0^P(x_{3j-\frac{1}{2}}^+), v_0^P(x_{3j-\frac{1}{2}}^-)\right). \quad (34c)$$

Then the scheme values are initialized as usual but starting from v_0^P , i.e.

$$V_j^0 := \frac{1}{\Delta x} \int_{[x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}]} v_0^P(y) dy = v_0^P(x_j) \quad (35)$$

This initialization ensures that the initial step function v_0^P will satisfy (23)-(24), and that the coded discontinuities are separated by at least $3\Delta x$. The aim of step (34) is also to keep the correct (local) extremal value of v_0 . This is motivated by the fact that the exact minima and maxima values should propagate. It allows to obtain better long-time approximations.

The general convergence result is the following.

Theorem 4.1. *We assume (H1)-(H3), and consider a l.s.c. function v_0 satisfying (H4), and such and $TV(v_0) < \infty$. We also assume that mesh steps Δx and Δt satisfy the CFL condition (13) and (33). Let ϑ be the unique viscosity solution of (4a)-(4b). We consider (V_j^0) defined by (34)-(35) and $(V^n)_{n \geq 1}$ given by algorithm 2. Let V be defined by (16). Then*

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (Lt_n + 7)e^{Lt_n} TV(v_0) \Delta x, \quad \forall n \geq 0.$$

Remark 4.1. *We can treat a discontinuous initial data of the form $v_0 := \mathbf{1}_{\mathbb{R} \setminus \{0\}}$. In this case we have $v_0^P = \mathbf{1}_{\mathbb{R} \setminus [-\frac{\Delta x}{2}, \frac{\Delta x}{2}]}$.*

We shall need preliminary estimates

Lemma 4.1. *We have*

$$\|v_0 - v_0^P\|_{L^1(\mathbb{R})} \leq 3\Delta x TV(v_0).$$

Proof. The result is immediate by the definition of v_0^P . □

Now let ϑ^P be the l.s.c. viscosity solution of (4a) with initial data v_0^P , i.e.,

$$\vartheta^P(0, x) := v_0^P(x), \quad \forall x \in \mathbb{R}.$$

The following estimate is essential in the analysis (the proof is postponed to the end of the section).

Proposition 4.1. *We assume (H1). Let u_0 and v_0 be two l.s.c. functions, such that $v_0 - u_0 \in L^1(\mathbb{R})$. We suppose furthermore that*

- (i) u_0 satisfies assumption (H4) (with $(A_i)_{i=2, \dots, q}$ local minima);
- (ii) for all interval $I \subset \mathbb{R}$,

$$\begin{cases} u_0 \nearrow \text{ on } I \Rightarrow v_0 \nearrow \text{ on } I, \\ u_0 \searrow \text{ on } I \Rightarrow v_0 \searrow \text{ on } I; \end{cases} \quad (36)$$

- (iii) for any local minima A_i of u_0 ($i = 2, \dots, q$), $u_0(A_i) = v_0(A_i)$.

Let u and v be defined by

$$u(t, x) := \min_{y \in [X_x^M(-t), X_x^m(-t)]} u_0(y), \quad \text{and} \quad v(t, x) := \min_{y \in [X_x^M(-t), X_x^m(-t)]} v_0(y).$$

We have

$$\|v(t, \cdot) - u(t, \cdot)\|_{L^1(\mathbb{R})} \leq e^{Lt} \|v_0 - u_0\|_{L^1(\mathbb{R})} \quad \forall t \geq 0. \quad (37)$$

Proof of Theorem 4.1. By construction of v_0^P , and under assumption (H4), the function v_0^P is increasing on intervals where v_0 is increasing, and decreasing on intervals where v_0 is decreasing, and the local minima of v_0^P are the same as the ones of v_0 . Hence we can apply Proposition 4.1 to compare ϑ and ϑ^P and obtain together with Lemma 4.1:

$$\|\vartheta(t, \cdot) - \vartheta^P(t, \cdot)\|_{L^1(\mathbb{R})} \leq e^{Lt} (3\Delta x TV(v_0)), \quad \forall t \geq 0. \quad (38)$$

Now by Theorem 3.1 we also have

$$\|V(t_n, \cdot) - \vartheta^P(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (Lt_n + 4)e^{Lt_n} TV(v_0^P) \Delta x, \quad \forall n \geq 0.$$

Furthermore, it is easy to see that $TV(v_0^P) \leq TV(v_0)$. Together with (38) this concludes the proof of Theorem 4.1. \square

We now conclude the section with the proof of Proposition 4.1.

Proof of Proposition 4.1: *Step 1.* We first study the case when u_0 is a monotonous increasing function. In this case v_0 is also increasing. $u(t, x) = u_0(X_x^M(-t))$ and $v(t, x) = v_0(X_x^M(-t))$. Hence, using the change of variable $y = X_x^M(t)$, we get

$$\begin{aligned} \|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} &= \int_{\mathbb{R}} |u(t, y) - v(t, y)| dy \\ &= \int_{\mathbb{R}} |u(t, X_x^M(t)) - v(t, X_x^M(t))| \left| \frac{dy}{dx}(t) \right| dx. \end{aligned}$$

Here, as f_M is Lipschitz then, by the Rademacher theorem, it is almost everywhere differentiable and we get: $\frac{dy}{dx}(t) = \exp(\int_0^t f'_M(X_x^M(s)) ds)$. In particular, $|\frac{dy}{dx}(t)| \leq e^{Lt}$ for all $t \geq 0$, and we obtain the bound

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} \leq e^{Lt} \|u_0 - v_0\|_{L^1(\mathbb{R})}.$$

The proof is similar when u_0 is a decreasing function.

Step 2. Now we study the case when u_0 has essentially only one local maximum located in $\overline{B_1}$. More precisely, we suppose that $u_0 \nearrow$ on $(-\infty, B_1)$ and $u_0 \searrow$ on (B_1, ∞) . The representation Lemma C.2 allows to write u_0 as:

$$u_0 = \min(u_{01}, u_{02}), \quad \text{with } u_{01} \nearrow \text{ and } u_{02} \searrow$$

where u_{01} and u_{02} are defined as in Lemma C.2. Then the viscosity solution is given by:

$$u(t, x) = \min(u_{01}(X_x^M(-t)), u_{02}(X_x^M(-t))), \quad t \geq 0, x \in \mathbb{R}.$$

By assumption (ii), v_0 also satisfies $v_0 \nearrow$ on $(-\infty, B_1)$ and $v_0 \searrow$ on (B_1, ∞) . We can also write $v_0 = \min(v_{01}, v_{02})$ where v_{01} and v_{02} are also defined as in Lemma C.2, and we have

$$v(t, x) = \min(v_{01}(X_x^M(-t)), v_{02}(X_x^m(-t))) \quad t \geq 0, x \in \mathbb{R}. \quad (39)$$

Next, we define b as a meeting point for the following two curves at a given time t :

$$x \rightarrow v_{01}(X_x^M(-t)) \quad \text{and} \quad x \rightarrow v_{02}(X_x^m(-t)).$$

Then

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} = \int_{-\infty}^b |u(t, x) - v(t, x)| dx + \int_b^{+\infty} |u(t, x) - v(t, x)| dx.$$

We notice that for $x \leq b$, $u(t, x) = u_{01}(X_x^M(-t))$ and $v(t, x) = v_{01}(X_x^M(-t))$. We thus get

$$\begin{aligned} \int_{-\infty}^b |u(t, x) - v(t, x)| dx &= \int_{-\infty}^b |u_{01}(X_x^M(-t)) - v_{01}(X_x^M(-t))| dx \\ &= \int_{-\infty}^{X_b^M(-t)} |u_{01}(x) - v_{01}(x)| \exp\left(\int_0^t f'_M(X_x^M(s)) ds\right) dx \\ &\leq e^{LT} \|u_{01} - v_{01}\|_{L^1(-\infty, X_b^M(-t))}, \end{aligned} \quad (40)$$

In the same way:

$$\int_{(b, +\infty)} |u(t, x) - v(t, x)| dx \leq e^{LT} \|u_{02} - v_{02}\|_{L^1(X_b^m(-t), +\infty)} \quad (41)$$

Since $f_M \geq f_m$, then $X_b^M(-t) \leq X_b^m(-t)$. Hence combining (40) and (41) we obtain (37).

Notice that for Step 1 and Step 2 there is no need of assumption (iii).

Step 3. We turn now to the proof in the general case (u_0 as in (H4)). Using assumptions (23)-(24) we can decompose u_0 into monotonous parts: there exist an integer $q \geq 1$ and real numbers A_1, \dots, A_{q+1} , B_1, \dots, B_q as in (H4) (with possibly $B_1 = -\infty$ or $B_q = +\infty$), and such that $u_0 \nearrow$ on each $[A_i, B_i[$ and \searrow on each $]B_i, A_{i+1}]$.

We first consider the time interval $[0, \tau_1[$ such that the number of local maxima q of u keeps constant.³ We note that around a local minima A_i , the solutions u and v will stay constant in the following sense: if we set $I_i^t := [X_{A_i}^m(t), X_{A_i}^M(t)]$ for $t \in [0, \tau_1]$, we have

$$u(t, x) = u_0(A_i) \quad \text{and} \quad v(t, x) = u_0(A_i), \quad \forall x \in I_i^t.$$

Hence, in order to bound $\|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})}$, we have to estimate the difference on the remaining intervals $J_i^t := [X_{A_i}^M(t), X_{A_{i+1}}^m(t)]$:

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} = \sum_{i=1}^q \|u(t, \cdot) - v(t, \cdot)\|_{L^1(J_i^t)},$$

³ For instance we consider $\tau_1 > 0$ to be the first time such that $\min_{i=1, \dots, q} X_{A_{i+1}}^m(t) - X_{A_i}^M(t)$ vanishes, with $a_i := \inf\{x, u_0(y) = u_0(A_i) \forall x \leq y \leq A_i\}$ and $b_i := \sup\{x, u_0(y) = u_0(A_i) \forall A_i \leq y \leq x\}$.

(the J_i^t are disjoint as long as $t < \tau_1$). Notice that $\vartheta(t, \cdot)$ and $\vartheta^P(t, \cdot)$ admit only one maximum on J_i^t . Then as in Step 2 we can show that:

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1(J_i^t)} \leq e^{Lt} \|u_0 - v_0\|_{L^1(J_i^0)} = e^{Lt} \|u_0 - v_0\|_{L^1([A_i, A_{i+1}])}.$$

Summing the previous bounds we obtain for $t \in [0, \tau_1]$:

$$\begin{aligned} \|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} &\leq e^{Lt} \sum_{i=1}^q \|u_0 - v_0\|_{L^1([A_i, A_{i+1}])} \\ &\leq e^{Lt} \|u_0 - v_0\|_{L^1(\mathbb{R})}. \end{aligned} \quad (42)$$

Now we consider the case when the number of maxima q may lower, and proceed recursively on q . We obtain on a time interval $[\tau_1, \tau_2]$, where the number of maxima is constant and equal to $q - 1$, the similar bound:

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1(\mathbb{R})} \leq e^{L(t-\tau_1)} \|u(\tau_1, \cdot) - v(\tau_1, \cdot)\|_{L^1(\mathbb{R})}, \quad t \in [\tau_1, \tau_2].$$

Then with (42) we obtain for all $t \leq \tau_2$ the desired bound, and so on. This concludes the proof of Proposition 4.1. \square

5 Case of changing sign velocities

We explain in this section how to get a general error estimate using mainly assumption (H1). Instead of assumptions (H2) and (H3) we shall use less restrictive assumptions that will be detailed in Theorem 5.1 below. Let $v_0 : \mathbb{R} \rightarrow \mathbb{R}$ be a l.s.c. function such that $TV(v_0) < \infty$. We consider a regular grid (x_j) as in Section 2, and define f_M^S and f_m^S as in Section 2.1.

The following algorithm introduces two modifications to algorithm 2: *left and right fluxes* (denoted by $V^{n,L}$ and $V^{n,R}$), for computing the UltraBee estimates, and a *prediction step*. We will explain later on the relevance of these modifications.

Algorithm 3

Initialization: We compute the initial averages (V_j^0) as in (34)-(35).

Loop: For $n \geq 0$, we compute V^{n+1} from V^n in three steps:

A) *Evolution by a modified HJB-UltraBee scheme:*

- Define “fluxes” $V_{j+\frac{1}{2}}^{n,L}(\nu)$, $V_{j+\frac{1}{2}}^{n,R}(\nu)$ for $\nu \in \{\nu^m, \nu^M\}$ as follows:

If $\nu_j \geq 0$, set

$$V_{j+\frac{1}{2}}^{n,L}(\nu) := \begin{cases} \min(\max(V_{j+1}^n, b_j^+(\nu)), B_j^+(\nu)) & \text{if } \nu_j > 0 \\ V_{j+1}^n & \text{if } \nu_j = 0 \text{ and } V_j^n \neq V_{j-1}^n \\ V_j^n & \text{if } \nu_j = 0 \text{ and } V_j^n = V_{j-1}^n, \end{cases}$$

If $\nu_j \leq 0$, set

$$V_{j-1/2}^{n,R}(\nu) := \begin{cases} \min(\max(V_{j-1}^n, b_j^-(\nu)), B_j^-(\nu)) & \text{if } \nu_j < 0 \\ V_{j-1}^n & \text{if } \nu_j = 0 \text{ and } V_j^n \neq V_{j+1}^n \\ V_j^n & \text{if } \nu_j = 0 \text{ and } V_j^n = V_{j+1}^n, \end{cases}$$

(where b_j^+ , b_j^- , B_j^+ and B_j^- are defined by (14)-(15)).

If $\nu_j \geq 0$ and $\nu_{j+1} > 0$, set $V_{j+\frac{1}{2}}^{n,R}(\nu) := V_{j+\frac{1}{2}}^{n,L}(\nu)$.

If $\nu_{j+1} \leq 0$ and $\nu_j < 0$, set $V_{j+\frac{1}{2}}^{n,L}(\nu) := V_{j+\frac{1}{2}}^{n,R}(\nu)$.

If $\nu_j < 0$ and $\nu_{j+1} > 0$, then set

$$V_{j+\frac{1}{2}}^{n,R}(\nu) := \begin{cases} V_{j+1}^n & \text{if } V_{j+1}^n = V_{j+2}^n \\ V_j^n & \text{otherwise} \end{cases} \quad \text{and} \quad V_{j+\frac{1}{2}}^{n,L}(\nu) := \begin{cases} V_j^n & \text{if } V_j^n = V_{j-1}^n \\ V_{j+1}^n & \text{otherwise.} \end{cases} \quad (43)$$

- For $\nu \in \{\nu^m, \nu^M\}$, let $V_j^{n+1}(\nu) := V_j^n - \nu_j \left(V_{j+\frac{1}{2}}^{n,L}(\nu) - V_{j-\frac{1}{2}}^{n,R}(\nu) \right)$.
- Set $V^{n+1,1} := \min \left(V^{n+1}(\nu^m), V^{n+1}(\nu^M) \right)$.

B) *Truncation*: Set $V^{n+1,2} := T_{V^n}(V^{n+1,1})$ as in algorithm 2.

C) *Prediction*: Set $W := V^{n+1,2}$.

- (decreasing critical cases)

$$\text{Let } J^- := \left\{ \begin{array}{l} j \in \mathbb{Z}, \left[W_{j-1} > W_j > W_{j+1} > W_{j+2} \text{ and } V_j^n < W_j \right] \\ \text{or } \left[W_{j-2} > W_{j-1} > W_j > W_{j+1} > W_{j+2} \text{ and } V_j^n = W_j \right] \end{array} \right\}$$

and $J^{-,*} := J^- \setminus \{j \in J^-, \text{ s.t. } j+2 \in J^-\}$.

For $j \in J^{-,*}$, set

$$\begin{aligned} V_{j-1}^{n+1} &:= W_{j-2}, \quad V_j^{n+1} := W_{j-2}, \text{ and} \\ V_{j+1}^{n+1} &:= W_{j+2} + \frac{W_{j+1} - W_{j+2}}{V_j^n - W_{j+2}} (W_{j-2} - W_{j+2}). \end{aligned} \quad (44)$$

- (increasing critical cases)

$$\text{Let } J^+ := \left\{ \begin{array}{l} j \in \mathbb{Z}, \left[W_{j+1} > W_j > W_{j-1} > W_{j-2} \text{ and } V_j^n < W_j \right] \\ \text{or } \left[W_{j+2} > W_{j+1} > W_j > W_{j-1} > W_{j-2} \text{ and } V_j^n = W_j \right] \end{array} \right\}$$

and $J^{+,*} := J^+ \setminus \{j \in J^+, \text{ s.t. } j-2 \in J^+\}$.

For $j \in J^{+,*}$, set

$$\begin{aligned} V_{j+1}^{n+1} &:= W_{j+2}, \quad V_j^{n+1} := W_{j+2}, \\ \text{and } V_{j-1}^{n+1} &:= W_{j-2} + \frac{W_{j-1} - W_{j-2}}{V_j^n - W_{j-2}} (W_{j+2} - W_{j-2}). \end{aligned} \quad (45)$$

- otherwise set $V_j^{n+1} = W_j$ (i.e. for j such that $\{j-1, j, j+1\} \cap (J^{-,*} \cup J^{+,*}) = \emptyset$).

Then we have the following general error estimate:

Theorem 5.1. *We suppose (H1). Let v_0 be an l.s.c. function satisfying (H4) and such that $TV(v_0) < \infty$ (in particular v_0 has a finite number of extrema).*

Let $\Delta x > 0$ and $\Delta t > 0$ satisfy the CFL condition (13) and (33). Let V_j^n be defined by algorithm 3, V as in (16), and ϑ be the viscosity solution of (4).

(i) If v_0 is piece-wise constant as in (23), with p discontinuities, and if

$$(H5a) \quad \exists \varepsilon > 0, \forall x \in \mathbb{R}, f_m(x) + \varepsilon \leq f_M(x) \quad \text{and} \quad \Delta x < \frac{\varepsilon}{2L},$$

or

$$(H5b) \quad \forall x \in \mathbb{R}, f_m(x) \leq 0, f_M(x) \geq 0,$$

then

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (Lt_n + 4p)e^{Lt_n} TV(v_0) \Delta x, \quad \forall n \geq 0. \quad (46)$$

(ii) If

$$(H5c) \quad f_m = f_M \quad \text{and is an increasing function,}$$

then

$$\|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq (1 + Lt_n e^{Lt_n}) TV(v_0) \Delta x, \quad \forall n \geq 0. \quad (47)$$

Remark 5.1. *Assumption (H5b) is satisfied by the eikonal equation $\vartheta_t + c(x)|\vartheta_x| = 0$, where c is a L -lipschitz positive function.*

Remark 5.2. *As in [7], when f_M (resp. f_m) changes sign, it is important to use two fluxes $V_{j+\frac{1}{2}}^{n,L}$ and $V_{j+\frac{1}{2}}^{n,R}$, which may be different on the cell's interface containing the zero of f_M (resp. f_m). The choice made for these fluxes insure the stability, consistency and TVD⁴ properties, see [7, Remark 2.1].*

Remark 5.3. *When assumptions (H1) – (H3) hold, the discontinuities around a minimum could only get far from each other. However, if we assume only (H1), the discontinuities may become closer. A truncation in this feature would produce an error which is not always controlled by the mesh size Δx . Hence we assume one of the (H5) assumptions in order to avoid this truncation. Indeed, with (H5) two discontinuities around a local minimum cannot become closer than $2\Delta x$ (see Lemma 5.3).*

⁴Total variation diminishing

Remark 5.4. When removing assumption (H3), two successive discontinuities in a monotone zone of v_0 can become very close. This motivates the prediction step: if the discontinuities are too close, we keep only one of them as shown in Figure 3. The algorithm then codes correctly the exact localization of the remaining discontinuity. This prediction step is handled only in the two critical cases explicated in figure 4, i.e. when the discontinuities are separated by less than two entire cells.




Figure 3: Prediction step

5.1 Preliminaries

We first prove in the following Lemma that the scheme computes the exact average values of ϑ^S in some elementary cases, where truncation and prediction steps are not used.

Lemma 5.1. We assume (H1) and (13). Let a, b, α, β be real numbers, with $\beta \geq 0$.

Let $v_0(x) := \alpha + \beta \mathbf{1}_{]a, +\infty[}(x)$ (resp. $v_0(x) := \alpha + \beta \mathbf{1}_{]-\infty, b[}(x)$).

(i) We have $\vartheta^S(t, x) := \alpha + \beta \mathbf{1}_{]X_a^{M,S}(t), +\infty[}(x)$ (resp. $\vartheta^S(t, x) := \alpha + \beta \mathbf{1}_{]-\infty, X_b^{m,S}(t)[}(x)$),




Figure 4: Critical cases of prediction

(ii) $\forall n \geq 0, \forall j \in \mathbb{Z}$:

$$V_j^n = \frac{1}{\Delta x} \int_{I_j} \vartheta^S(t_n, x) dx \quad (48)$$

Proof. Part (i) is obtained by direct verifications using (10). For (ii), we shall treat the case of $v_0(x) = \mathbf{1}_{]a, +\infty[}(x)$ (the other cases being similar). We proceed as in Lemma 2.3. We prove the statement by recursion. Let us denote by $x_n := X^{M,S}(t_n)$ the discontinuity position at time t_n . Note that it suffices to prove that

$$V_j^{n+1}(\nu^M) = \frac{1}{\Delta x} \int_{I_j} \mathbf{1}_{]X_{x_n}^{M,S}(\Delta t), +\infty[}(x) dx \quad \forall j \in \mathbb{Z}. \quad (49)$$

Indeed we have in the same way:

$$V_j^{n+1}(\nu^m) = \frac{1}{\Delta x} \int_{I_j} \mathbf{1}_{]X_{x_n}^{m,S}(\Delta t), +\infty[}(x) dx \quad \forall j \in \mathbb{Z}.$$

And this will prove that $V_j^{n+1} = V_j^{n+1}(\nu^M)$ as desired. In the case when for all $j \in \mathbb{Z}$, $\nu_j^M \geq 0$, (or if for all $j \in \mathbb{Z}$, $\nu_j^M \leq 0$), the result comes from Lemma 2.3. It remains to treat the case when (ν_j^M) changes signs. We denote $\nu := \nu^M$ for simplicity.

Assume that $x_n \in]x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}]$ (i.e., $V_k^n = 0$ for $k < j$, $V_j^n \in [0, 1[$, and $V_k^n = 1$ for $k > j$). We furthermore assume that $x_n \neq x_{j+\frac{1}{2}}$ (the case of $x_n = x_{j+\frac{1}{2}}$ can be treated in a similar way). In particular we have $V_{j-1}^n = V_{j-2}^n$ and $V_{j+1}^n = V_{j+2}^n$. Let us show that the values $(V_k^{n+1}(\nu))_{k=j-1, j, j+1}$ are correctly computed by the modified UltraBee scheme. We first state the following Lemma:

Lemma 5.2. (i) $V_{j+\frac{1}{2}}^{n,L/R} \in [\min(V_j^n, V_{j+1}^n), \max(V_j^n, V_{j+1}^n)]$ (Consistency).

(ii) (Stability) $V_j^n = V_{j-1}^n$ and $\nu_j \geq 0 \Rightarrow V_j^{n+1} = V_j^n$.

(iii) (Stability) $V_j^n = V_{j+1}^n$ and $\nu_j \leq 0 \Rightarrow V_j^{n+1} = V_j^n$.

Proof of Lemma 5.2 Assertion (i) can be obtained as in [7].

(ii) If $\nu_j = 0$, is immediate. If $\nu_j > 0$, we notice that $V_{j+\frac{1}{2}}^{n,L} = V_j^n$ because $b_j^+ = B_j^+ = V_j^n$. On the other hand, if $\nu_{j-1} > 0$, then $V_{j+\frac{1}{2}}^{n,R} = V_{j-\frac{1}{2}}^{n,L} = V_j^n$ (using $V_{j-1}^n = V_j^n$ and the consistency property). If otherwise $\nu_{j-1} \leq 0$, then $V_{j+\frac{1}{2}}^{n,R} \in \{V_j^n, V_{j-1}^n\}$ (see definition), hence $V_{j+\frac{1}{2}}^{n,R} = V_j^n$.

(iii) the proof of this assertion is similar to (ii). \square

We come back to the proof of Lemma 5.1.

In the case $\nu_j = 0$, we have $V_j^{n+1}(\nu) = V_j^n$, which is the correct value (the characteristics does not evolve in I_j). If $\nu_{j+1} > 0$, we already know from Section 2 that $V_{j+1}^{n+1}(\nu)$ is correctly computed (we are in a subcase of $\nu_j \geq 0, \nu_{j+1} \geq 0$). The case when $\nu_{j+1} < 0$ is similar. We can use the same arguments to see that $V_{j-1}^{n+1}(\nu)$ is also correctly computed.

Now we assume that $\nu_j > 0$ (the case when $\nu_j < 0$ can be treated in a similar way), and study the different remaining cases.

1) Case $\nu_{j-1} > 0$. We can also suppose that $\nu_{j+1} \leq 0$ (otherwise, $(\nu_k)_{k=j-1, j, j+1} \geq 0$ and this has already been treated).

We first have $V_{j-1}^{n+1} = V_{j-1}^n$ (using Lemma 5.2(i)), the correct expected value.

Also $V_j^{n+1} = V_j^n - \nu_j(V_{j+\frac{1}{2}}^{n,L} - V_{j-\frac{1}{2}}^{n,R})$, where $V_{j+\frac{1}{2}}^{n,L}$ is computed as in Algorithm 1 (Section 2) for positive velocities, and where $V_{j-\frac{1}{2}}^{n,R} = V_{j-\frac{1}{2}}^{n,L} = V_{j-1}^n$. Hence the computation of V_j^{n+1} is as in the case of positive velocities, and gives the correct expected value.

Finally, $V_{j+1}^{n+1} = V_{j+1}^n$ using Lemma (iii).

2) Case $\nu_{j-1} < 0$. First, $V_{j-1}^{n+1} = V_{j-1}^n - \nu_{j-1}(V_{j-\frac{1}{2}}^{n,L} - V_{j-\frac{3}{2}}^{n,R})$, where $V_{j-\frac{1}{2}}^{n,L} = V_{j-1}^n$ (by definition in Algorithm 3), $V_{j-\frac{3}{2}}^{n,R} \in [V_{j-1}^n; V_{j-2}^n] = \{V_{j-1}^n\}$ (by consistency), hence $V_{j-1}^{n+1} = V_{j-1}^n$. Then, $V_j^{n+1} = V_j^n - \nu_j(V_{j+\frac{1}{2}}^{n,L} - V_{j-\frac{1}{2}}^{n,R})$ where $V_{j+\frac{1}{2}}^{n,L}$ has the flux definition with $\nu_j > 0$, and $V_{j-\frac{1}{2}}^{n,R}$ takes the value V_{j-1}^n (since $V_j^n \neq V_{j+1}^n$ using the fact that $x_n \neq x_{j-\frac{1}{2}}$). Hence in the interval I_j , the estimate of the fluxes are the same as in the case of positive velocities, and are thus correct (exact evolution of the average values locating the discontinuity position in I_j or I_{j+1}). Finally, we have only to check the value of V_{j+1}^{n+1} in the case $\nu_{j+1} < 0$. In this case, $V_{j+\frac{1}{2}}^{n,R} = V_{j+1}^n$ (by Lemma 5.2(ii)), and $V_{j+\frac{3}{2}}^{n,L} = V_{j+1}^n$ (by Lemma 5.2(i)), hence $V_{j+1}^{n+1} = V_{j+1}^n$.

3) Case $\nu_{j-1} = 0$. We first obtain $V_{j-1}^{n+1} = V_{j-1}^n$. Then V_j^{n+1} and V_{j+1}^{n+1} are computed as in the case $\nu_{j-1} < 0$ (proof is left to the reader). This concludes the proof of Lemma 5.1 \square

Remark 5.5. *Indeed we have proved a more precise result: if at some time t_n we have that $\vartheta^S(t_n, \cdot)$ is a step-wise constant function with all successive discontinuities separated by at least two entire intervals, and if (48) holds for $j \in \mathbb{Z}$, then also we have*

$$V_j^{n+1} = \frac{1}{\Delta x} \int_{I_j} \vartheta^S(t_{n+1}, x) dx \quad \forall j \in \mathbb{Z}.$$

Remark 5.6. Several estimates of Section 2 can be extended here. In particular, under assumption (H1) only, the estimates of Lemma 2.1(i),(ii) and (v) also hold (in particular for changing sign velocities). Proof is left to the reader.

Lemma 5.3. We assume (H1) and one of the (H5) assumptions. Let $a, b \in \mathbb{R}$ be such that $b - a \geq 2\Delta x$. Then for all $t \geq 0$, $X_b^{M,S}(t) - X_a^{m,S}(t) \geq 2\Delta x$.

Proof. If we assume (H5b) or (H5c) it is easy to see that $\frac{d}{dt}(X_b^{M,S}(t) - X_a^{m,S}(t)) \geq 0$, for a.e. $t \geq 0$, hence the result. Now assume (H5a). Suppose there exists $\theta \geq 0$ such that $X_b^{M,S}(\theta) - X_a^{m,S}(\theta) < 2\Delta x$. By continuity there exists $\tau \geq 0$ such that $X_b^{M,S}(\tau) - X_a^{m,S}(\tau) = 2\Delta x$ and

$$X_b^{M,S}(t) - X_a^{m,S}(t) < 2\Delta x, \quad \text{for } t \text{ in a neighborhood } \mathcal{V}(\tau^+) \text{ of } \tau^+.$$

Case 1. We first suppose that $X_a^{m,S}(\tau) \in I_{j-1}$ for some $j \in \mathbb{Z}$ (i.e., $X_a^{m,S}(\tau)$ belongs to the interior of a mesh interval) and thus also $X_b^{M,S}(\tau) \in I_{j+1}$. In particular $\delta : t \rightarrow X_b^{M,S}(t) - X_a^{m,S}(t)$ is differentiable at $t = \tau$ and necessarily we have $\dot{\delta}(\tau) \leq 0$. Hence $f_m(x_{j-1}) \geq f_M(x_{j+1})$. Then

$$f_M(x_{j-1}) - 2L\Delta x \leq f_M(x_{j+1}) \leq f_m(x_{j-1}) \leq f_M(x_{j-1}) - \varepsilon,$$

and we get $2L\Delta x \geq \varepsilon$ which contradicts (H5a). Thus this case cannot occur.

Case 2. Now we suppose that $X_a^{m,S}(\tau) = x_{j-\frac{1}{2}}$ for some $j \in \mathbb{Z}$ (and thus $X_b^{M,S}(\tau) = x_{j+\frac{1}{2}}$). If the two characteristics $X_a^{m,S}(\theta)$ and $X_b^{M,S}(\theta)$ move such that $X_a^{m,S}(\theta) \in I_j$ and $X_b^{M,S}(\theta) \in I_{j+1}$ for $\theta \in \mathcal{V}(\tau^+)$, then using (H5a) we get

$$f_M(x_{j+1}) \leq f_m(x_j) \leq f_M(x_j) - \varepsilon.$$

Since we also have $f_M(x_{j+1}) \geq f_M(x_j) - L\Delta x$, we obtain

$$f_M(x_j) - L\Delta x \leq f_M(x_j) - \varepsilon \leq f_M(x_j) - 2L\Delta x$$

which leads to $2L \leq L$, a contradiction. Otherwise, if the two characteristics move in the same direction, we obtain a contradiction in the same way as in Case 1. \square

5.2 Proof of Theorem 5.1

We first define a *prediction operator* as follows. Let w be a real piece-wise constant and l.s.c. function. For $x \in \mathbb{R}$, set

$$\begin{aligned} z_1^x &:= \sup\{z, z \leq x, w(z) \neq w(x)\} \in [-\infty, \infty[, \\ z_2^x &:= \inf\{z, z \geq x, w(z) \neq w(x)\} \in]-\infty, \infty] \end{aligned}$$

i.e. the closest left and right discontinuities of w to a given x . Let j and k be such that

$$z_1^x \in]x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}] \quad \text{and} \quad z_2^x \in [x_{k-\frac{1}{2}}, x_{k+\frac{1}{2}}[.$$

Let us denote $w(x^-) := \lim_{y \rightarrow x, y < x} w(y)$ and $w(x^+) := \lim_{y \rightarrow x, y > x} w(y)$. Then set

$$\text{Pred}(w)(x) := \begin{cases} w((z_1^x)^-) & \text{if } w((z_1^x)^-) > w(x) > w((z_2^x)^+), k \in \{j+1, j+2\}, \\ & \text{and } z_2^{z_2^x} \notin]x_{k+2-\frac{1}{2}}, x_{k+2+\frac{1}{2}}[\\ w((z_2^x)^+) & \text{if } w((z_1^x)^-) < w(x) < w((z_2^x)^+), k \in \{j+1, j+2\}, \\ & \text{and } z_1^{z_1^x} \notin]x_{j-2-\frac{1}{2}}, x_{j-2+\frac{1}{2}}[\\ w(x) & \text{otherwise.} \end{cases} \quad (50)$$

Remark 5.7. *The above definition means that we set $\text{Pred}(w)(x) = w((z_1^x)^-)$ in the case x belongs to a decreasing zone of w (for instance), surrounded by two discontinuities of w that are separated at most by one cell, but also such that the next right discontinuity $z_2^{z_2^x}$ is at least separated by two cells from the right discontinuity z_2^x (see Fig. 3).*

Then we define $\vartheta^\sharp(t_n, \cdot)$ for all $n \geq 0$ by :

- $\vartheta^\sharp(0, \cdot) := v_0^P$,
- $\forall n \geq 0, \vartheta^\sharp(t_{n+1}, \cdot) = \text{Pred}(\text{Trunc}(w_{n+1}))$ where w_{n+1} is as in (30).

We also define ϑ^S as in (10) but starting from v_0^P , i.e.:

$$\vartheta^S(t, x) := \min_{y \in [X_x^{M,S}(-t), X_x^{m,S}(-t)]} v_0^P(y), \quad \forall t > 0, x \in \mathbb{R}. \quad (51)$$

Now, we focus on the proof of Theorem 5.1 (i) (in particular we have $v_0^P \equiv v_0$ here). We follow the proofs of Theorem 3.1 and 4.1. For a given piece-wise constant initial data v_0 (with discontinuities separated by at least $3\Delta x$), it remains to prove that the following still holds - in a similar way as in Lemma 3.2.

Lemma 5.4. *Under the assumptions of Theorem 5.1(i), we have*

(i)

$$\forall n \geq 0, \quad \|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} \leq p(4e^{Lt_n} - 1)TV(v_0) \Delta x \quad (52)$$

(ii)

$$\forall j \in \mathbb{Z}, \forall n \geq 0, \quad V_j^n = \frac{1}{\Delta x} \int_{I_j} \vartheta^\sharp(t_n, x) dx. \quad (53)$$

Proof of Theorem 5.1. The proof of (i) follows by using (52) and previous estimates :

$$\begin{aligned} & \|V(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \\ & \leq \|V(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(\mathbb{R})} + \|\vartheta^\sharp(t_n, \cdot) - \vartheta^S(t_n, \cdot)\|_{L^1(\mathbb{R})} + \|\vartheta^S(t_n, \cdot) - \vartheta(t_n, \cdot)\|_{L^1(\mathbb{R})} \\ & \leq TV(v_0)\Delta x + p(4e^{Lt_n} - 1)TV(v_0)\Delta x + Lt_n e^{Lt_n} TV(v_0)\Delta x \\ & \leq (Lt_n + 4p)e^{Lt_n} TV(v_0)\Delta x. \end{aligned}$$

Now to prove (ii), notice that under assumption (H5c), equation (4a) is reduced to an advection equation. In particular, as f_M is increasing then the discontinuities never meet: we do not need any truncation or prediction step in the scheme. The proof of Theorem 5.1(ii) is similar to the proof of Theorem 2.1. \square

In order to prove Lemma 5.4, we first establish the following.

Lemma 5.5. *Assume that v_0 satisfies assumption (H4) (with $(A_i)_{i=2,\dots,q}$ local minima). (i) for all interval $I \subset \mathbb{R}$,*

$$\begin{cases} \vartheta^S \nearrow \text{ on } I \Rightarrow \vartheta^\sharp \nearrow \text{ on } I, \\ \vartheta^S \searrow \text{ on } I \Rightarrow \vartheta^\sharp \searrow \text{ on } I, \end{cases}$$

(ii) *If $X_{A_i}^{M,S}(t_n)$ is a local minima of $\vartheta^S(t_n, \cdot)$ ($i = 2, \dots, q$), then it is also a local minima of $\vartheta^\sharp(t_n, \cdot)$, and we have $\vartheta^\sharp(t_n, y) = \vartheta^S(t_n, y) (= \vartheta^S(t_n, X_{A_i}^{M,S}(t_n)))$ for all $y \in [X_{A_i}^{m,S}(t_n), X_{A_i}^{M,S}(t_n)]$.*

Proof. By Lemma 5.3, two discontinuities positions a_i and b_i limiting a given minimum A_i in v_0 , initially separated by at least $3\Delta x$, stay at more than $3\Delta x$ in $\vartheta^S(t_n, \cdot)$ and thus stay separated at least by two cell intervals. In particular they cannot meet with our assumption.

Also since the operators *Pred* and *Trunc* do not modify the monotonicity and keep the minima values, we obtain that $\vartheta^\sharp(t_n, \cdot)$ will keep the monotonicity regions of $\vartheta^S(t_n, \cdot)$ (as well as the minima regions of $\vartheta^S(t_n, \cdot)$). This proves both (i) and (ii). \square

Remark 5.8. *Note that a minima zone of $\vartheta^S(t_n, \cdot)$ can disappear. This happens only in the case the neighboring left or right maxima zones disappear. By the previous Lemma $\vartheta^\sharp(t_n, \cdot)$ will have a similar property.*

Proof of Lemma 5.4(i). Let $J_i := [X_{A_i}^{M,S}(t_n), X_{A_{i+1}}^{m,S}(t_n)]$. To show (52), by using Lemma 5.5(ii), it is sufficient to obtain the following bound

$$\|\vartheta^S(t_n, \cdot) - \vartheta^\sharp(t_n, \cdot)\|_{L^1(J_i)} \leq p(4e^{Lt_n} - 1)TV(v_0, [A_i, A_{i+1}]) \Delta x \quad (54)$$

(the result will then follow by summation on i). It means that we need to show (52) in the particular case when $v_0 \nearrow$ on $[A_1, B_1]$ and $v_0 \searrow$ on $[B_1, A_2]$. We can assume $A_1 = -\infty$ and $A_2 = \infty$ to simplify.

Let $\vartheta^{\sharp,0}(t_n, \cdot) := \vartheta^S(t_n, \cdot)$. We define recursively the function $\vartheta^{\sharp,k}$ for $k \geq 0$ as follows. Let t_{n_k} be the first time where a prediction should be performed in the function $\vartheta^{\sharp,k-1}(t_{n_k}, \cdot)$ (i.e., two successive discontinuities in a decreasing or in an increasing region of $\vartheta^{\sharp,k-1}(t_{n_k}, \cdot)$ are separated by less than two entire cell intervals). Then set

$$\begin{aligned} \vartheta^{\sharp,k}(t, \cdot) &:= \vartheta^{\sharp,k-1}(t, \cdot) \quad \text{for } t < t_{n_k}, \\ \vartheta^{\sharp,k}(t_{n_k}, \cdot) &:= \text{Pred}(\vartheta^{\sharp,k-1}(t_{n_k}^-, \cdot)), \\ \vartheta^{\sharp,k}(t, x) &:= \min_{y \in [X_x^{M,S}(t-t_{n_k}), X_x^{m,S}(t-t_{n_k})]} \vartheta^{\sharp,k}(t_{n_k}, y) \quad \text{for } t \geq t_{n_k} \text{ and } x \in \mathbb{R}. \end{aligned}$$

This means that $\vartheta^{\#,1}(t_n, \cdot)$ is the function where only the first occurring prediction in $\vartheta^{\#,0} = \vartheta^S$ is taken into account, $\vartheta^{\#,2}(t_n, \cdot)$ is the function where only the first prediction in $\vartheta^{\#,1}$ is taken into account, etc.

We obtain for $k = 1, \dots, p$,

$$\|\vartheta^{\#,k}(t_n, \cdot) - \vartheta^{\#,k-1}(t_n, \cdot)\|_{L^1} \leq (4e^{Lt_n} - 1)TV(v_0) \Delta x \quad (55)$$

Indeed, if $t_n < t_{n_k}$, $\|\vartheta^{\#,k}(t_n, \cdot) - \vartheta^{\#,k-1}(t_n, \cdot)\|_{L^1} = 0$, and if $t_n \geq t_{n_k}$, we obtain the bound (55) by using similar arguments as in the proof of Lemma 3.2(i).

Also we note that $\vartheta^{\#,p}(t_n, \cdot) \equiv \vartheta^{\#}(t_n, \cdot)$, because there are at most p prediction steps that can be done in $\vartheta^{\#}(t_n, \cdot)$. Summing these bounds for $k = 1, \dots, p$ and by a triangular inequality this proves (54). \square

Proof of Lemma 5.4(ii). The proof is obtained by a recursion argument on $n \geq 0$.

Lemma 5.1 shows that an isolated discontinuity of $\vartheta^S(t_n, \cdot)$, as long as it keeps separated from other discontinuity positions by at least two cell intervals, has its evolution correctly coded by algorithm 3 for one time step. Also Lemma 5.3 implies that two discontinuities positions a_i and b_i limiting a given minimum A_i in v_0 (and evolving as $X_{a_i}^{m,S}(t)$ and $X_{b_i}^{M,S}(t)$) stay at more than $3\Delta x$, and thus are separated at least by two cell intervals. Hence the problem of having discontinuity positions in ϑ^S no more separated by two cell intervals can only come from maxima regions or monotonous regions of ϑ^S .

Then as in Lemma 3.2(ii) we can show that the cell averages of $Trunc(w_{n+1})$ are well coded by the scheme values $V_j^{n+1,2}$ after Step A) and Step B).

Finally it remains to prove that the prediction step C) corresponds to the application of prediction operator $Pred$ on the function $Trunc(w_{n+1})$. The proof is very similar to the proof of Lemma 3.2(ii) for the truncation step (i.e. a discontinuity position vanishes and the remaining discontinuity leads to a recomputation of average values). \square

6 Numerical tests

In this section, we apply algorithm 2 and 3 to some examples. These tests show the numerical relevance of the method especially for the truncation step. The L^1 -error is computed by the formula:

$$\text{error} \equiv \sum_j \Delta x \left| V_j^n - \frac{1}{\Delta x} \int_{I_j} \vartheta(t^n, x) dx \right|,$$

where (V_j^n) are the numerical values and ϑ is the exact viscosity solution (4).

Example 1: piece-wise constant initial data. We consider the following Eikonal equation:

$$\vartheta_t(t, x) + |\vartheta_x(t, x)| = 0, \quad t \geq 0, \quad (56)$$

for $x \in (-2, 2)$ and with periodic boundary conditions. Notice that we can take $f_M = 1$ and $f_m = -1$. The CFL condition is $\frac{\Delta t}{\Delta x} \leq 1$ (here we have chosen the CFL number to be $\frac{\Delta t}{\Delta x} = 0.9$). The initial condition is defined as follows:

$$\vartheta(0, x) = \begin{cases} 2 & \text{if } x \in]-1.6, -1[, \\ 1.7 & \text{if } x \in [-1, 0.1[, \\ 0.2 & \text{if } x \in [0.1, 0.6], \\ 1.2 & \text{if } x \in]0.6, 1.2[, \\ 0.7 & \text{if } x \in [1.2, 1.6[, \\ 0 & \text{otherwise.} \end{cases}$$

In Figure 5, we show the exact solution (black line) and the numerical values of algorithm 2 (cross) at different times.

We follow the evolution of two local maxima until they disappear, in particular in the last two time steps: just before they disappear (c) and then just after (d). Notice that the scheme computes exactly the mean value of the solution far from critical situations. In fact the dynamics f_m and f_M are constant here, and the approximated characteristics $X_x^{M,S}$ and $X_x^{m,S}$ coincide with X_x^M and X_x^m respectively. This leads to an exact computation of the average values of $\vartheta \equiv \vartheta^S$ by the scheme (except when discontinuities are too close)

Example 2: piece-wise continuous initial data.

The initial condition is given by

$$\vartheta(0, x) = \begin{cases} -x^2 + 1.5 & \text{if } x \in [-1, 1], \\ 0 & \text{otherwise.} \end{cases}$$

The results are shown in Figure 6.

Example 3: two local maxima.

Here we consider a piece-wise continuous initial data with two local maxima :

$$\vartheta(0, x) = \begin{cases} x^2 + 0.7 & \text{if } x \in [-1, 1], \\ 0 & \text{otherwise.} \end{cases}$$

The results are shown in Figure 7.

Remark 6.1. *In practice, numerical tests show that when the initialization of the algorithm is done with a projection with mesh size Δx (instead of $3\Delta x$), the error gets smaller (up to 10 times smaller).*

Example 4: We consider the eikonal equation (56) on $(-1, 1)$ but now with an initial condition composed of two discontinuities, as follows:

$$\vartheta(0, x) = \begin{cases} 1 & \text{if } x \in]-0.7, 0.9[, \\ 0 & \text{otherwise.} \end{cases}$$







(a) error=0, $t = 0$ (b) error=0, $t = 0.20$ (c) error=0.06, $t = 0.26$ (d) error=0, $t = 0.3$


Figure 5: (Example 1) Piece-wise constant function, #cells=70.




(a) error=0.076, $t = 0$



(b) error=0.047, $t = 0.55$



(c) error=0.020, $t = 0.85$



(d) error=0.055, $t = 0.95$

Figure 6: (Example 2) Evolution of a piece wise continuous function with one maximum, # cells=72, initialization of the algorithm using v_0^P .




Figure 7: (Example 3) Piece wise continuous function with two maxima, # cells=70.

Δx	error	Truncation Time	# cells
0.1	0.16	0.72	20
0.05	0.07	0.76	40
0.025	0.025	0.79	80
0.0125	0.0025	0.8	160

Table 1: (Example 4) Evolution of the error with the mesh size Δx , $CFL = 0.9$

We see in Table 1 that the error is bounded by $2\Delta x$.

Example 5: Eikonal equation with varying velocity


In this example we deal with algorithm 3 in order to illustrate the prediction step. We consider the following Eikonal equation:

$$\vartheta_t(t, x) + |x| \cdot |\vartheta_x(t, x)| = 0, \quad x \in (-1, 1),$$


with periodic boundary conditions. Here we take $f_M(x) = |x|$ and $f_m(x) = -|x|$. The initial condition is given by

$$\vartheta(0, x) = \begin{cases} 0.4 & \text{if } x \in]-0.8, -0.5], \\ 1 & \text{if } x \in [-0.5, 0.6], \\ 0.7 & \text{if } x \in [0.6, 0.7[, \\ 0.2 & \text{if } x \in [0.7, 0.8[, \\ 0 & \text{otherwise.} \end{cases}$$


Since f_m and f_M are not constant, ϑ and ϑ^S will differ. Results are shown in Figure 8. In Fig.8(a), two critical cases of prediction appear simultaneously in the decreasing zone. The algorithm handles only a prediction for the right discontinuity (Fig.8(b)). In Fig.8(c), a prediction step is again needed, and so on.




(a) error=0.095, $t = 0.045$
(before prediction step)




(b) error=0.0487, $t = 0.045$
(after prediction step)




(c) error=0.035, $t = 0.4$
(before prediction step)



(d) error=0.074, $t = 0.4$
(after prediction step)



(e) error=0.12, $t = 0.855$



(f) error=0.027, $t = 2.52$

Figure 8: (Example 5) Prediction steps in Algorithm 3, # cells=40

A Definition of the approximated characteristics

We prove existence and uniqueness of the absolutely continuous solution $X_a^{M,S}(t)$ of the differential equation:

$$\begin{aligned} \dot{X}_a^{M,S}(t) &= f_M^S(X_a^{M,S}(t)) \text{ a.e } t \geq 0, & X_a^{M,S}(0) &= a, & (57a) \\ \text{if } \exists t^* \geq 0 \text{ s.t. } & \begin{cases} X_a^{M,S}(t^*) = x_{j+\frac{1}{2}}, \\ \text{and } f^M(x_j)f^M(x_{j+1}) \leq 0 \end{cases} & \text{then } X_a^{M,S}(t) &= x_{j+\frac{1}{2}} \quad \forall t \geq t^* & (57b) \end{aligned}$$

We define recursively the characteristic as follows. Assume that $a \in]x_{j_0-\frac{1}{2}}, x_{j_0+\frac{1}{2}}]$ for some index j_0 . We consider the case $f_M(x_{j_0}) \geq 0$ (the case $f_M(x_{j_0}) < 0$ being similar). We define $\tau_0 := 0$,

$$\tau_1 := \tau_0 + \frac{x_{j_0+\frac{1}{2}} - a}{f_M(x_{j_0})}, \quad \text{if } f_M(x_{j_0}) > 0,$$

and for $k \geq 1$,

$$\tau_{k+1} := \tau_k + \frac{\Delta x}{f_M(x_{j_0+k})}, \quad \text{if } f_M(x_{j_0+k}) > 0$$

(i.e., $\frac{\Delta x}{f_M(x_{j_0+k})}$ is the time needed for a characteristic to cross the interval I_{j_0+k}). Otherwise, if there exists a first index k^* such that $f_M(x_{j_0+k^*}) \leq 0$, then we define $\tau_{k^*+1} := +\infty$ and stop the iterations. Note that since f_M is Lipschitz, we have either $\lim_{k \rightarrow \infty} \tau_k = +\infty$ (in this case set $k^* = +\infty$), or there exists $k^* < +\infty$ such that $\tau_{k^*+1} := +\infty$.

Now, $t \in [\tau_k, \tau_{k+1}[$ and $k < k^*$, we set

$$\chi(t) := \chi(\tau_k) + (t - \tau_k)f_M(x_{j_0+k})$$

(where $\chi(\tau_0) = a$ and $\chi(\tau_k) = x_{j_0+k-\frac{1}{2}}$ for $k \geq 1$), and if $t \geq \tau_{k^*}$, we set

$$\chi(t) = \chi(\tau_{k^*}).$$

Then $\chi(t)$ is a solution of (57).

In order to show the unicity of the solution of (57), we first notice that the first time t^* when two solutions may differ must be such that $\chi(t^*)$ be on an interface: $\exists j \in \mathbb{Z}$, $\chi(t^*) = x_{j+\frac{1}{2}}$. In the case $f_M(x_j)f_M(x_{j+1}) \leq 0$, by definition we have $\chi(t) = x_{j+\frac{1}{2}}$ for all $t \geq t^*$, and unicity. Otherwise, in the case $f_M(x_j)f_M(x_{j+1}) > 0$, we have necessarily $f_M(x_j) > 0$ (we assume here that $f_M(x_{j_0}) > 0$). Then the only solution for $t > t^*$ in a neighborhood of t^* , is given by

$$\chi(t) = \chi(t^*) + (t - t^*)f_M(x_{j+1}).$$

This shows unicity.

B TV bounds

Lemma B.1. *Let v_0 be an l.s.c. function such that $TV(v_0) < \infty$. For $j = 1, 2$, let $x \rightarrow a_x^j$ and $x \rightarrow b_x^j$ be non-decreasing, one-to-one functions from \mathbb{R} onto \mathbb{R} . We assume that*

- (i) $x \rightarrow a_x^j$ and $x \rightarrow b_x^j$ are non-decreasing, one-to-one functions from \mathbb{R} onto \mathbb{R} (for $j = 1, 2$).
- (ii) $a_x^j \leq b_x^j$ for $j = 1, 2$ and $x \in \mathbb{R}$.
- (iii) there exists $\delta \geq 0$, such that,

$$\forall x \in \mathbb{R}, \quad \max(|(a^2)^{-1}(x) - (a^1)^{-1}(x)|, |(b^2)^{-1}(x) - (b^1)^{-1}(x)|) \leq \delta.$$

(where $(a^j)^{-1}$ and $(b^j)^{-1}$ denotes the reciprocal functions of a^j and b^j resp.).
Let also

$$v_j(x) := \min_{y \in [a_x^j, b_x^j]} v_0(y).$$

Then

$$\|v_1 - v_2\|_{L^1(\mathbb{R})} \leq 2\delta TV(v_0). \quad (58)$$

Proof. We first assume that $a_x^2 = a_x^1, \forall x$. Then we notice that:

$$|v_2(x) - v_1(x)| \leq TV(v_0; [b_x^1; b_x^2]) \quad (59)$$

where $[\alpha; \beta]$ denotes the interval $[\min(\alpha, \beta), \max(\alpha, \beta)]$. (To prove (59), assume for instance the case $b_x^1 \leq b_x^2$. Consider $z_1 \in [a_x^1, b_x^1]$ such that $v_1(x) = v_0(z_1)$ and $z_2 \in [a_x^1, b_x^2]$ such that $v_2(x) = v_0(z_2)$. If the minimum for v_2 can be reached in $[a_x^1, b_x^1]$ then $v_1(x) = v_2(x)$. Otherwise we have $z_2 \in [b_x^1, b_x^2]$ and $v_2(x) = v_0(z_2) < v_0(z_1) = v_1(x)$. Hence

$$\begin{aligned} |v_2(x) - v_1(x)| &= v_0(z_1) - v_0(z_2) \\ &= v_0(z_1) - v_0(b_x^1) + v_0(b_x^1) - v_0(z_2) \\ &\leq v_0(b_x^1) - v_0(z_2) \leq TV(v_0; [b_x^1, b_x^2]). \end{aligned}$$

The case when $b_x^2 \leq b_x^1$ is similar.)

Now we establish the following.

Lemma B.2. *There exists a positive real measure μ such that $\mu(\mathbb{R}) = TV(v_0)$ and $TV(v_0;]-\infty, \beta]) = \int_{y \in]-\infty, \beta[} d\mu(y)$, for a.e $\beta \in \mathbb{R}$.*

Proof. Let \tilde{v}_0 be defined by $\tilde{v}_0(x) := \lim_{y \rightarrow x, y < x} v_0(y)$ for every $x \in \mathbb{R}$ (i.e, $\tilde{v}_0(x) = v_0(x^-)$). Then \tilde{v}_0 is left-continuous, and there exists a positive measure $\tilde{\mu}$ such that $TV(\tilde{v}_0,]-\infty, x]) = \tilde{\mu}(]-\infty, x])$ for every $x \in \mathbb{R}$. (This can be deduced for instance from [14]). We have also $\tilde{\mu}(\{x\}) = |v_0(x^-) - v_0(x^+)|$, for every $x \in \mathbb{R}$.

Also, since $TV(v_0) < \infty$, v_0 admits at most a countable set of discontinuity points denoted $(a_n)_{n \geq 0}$. Let

$$q_n := |v_0(a_n^-) - v_0(a_n)| + |v_0(a_n) - v_0(a_n^+)| - |v_0(a_n^-) - v_0(a_n^+)|$$

($q_n \geq 0$) and

$$\mu := \tilde{\mu} + \sum_{n \in \mathbb{N}} q_n \delta_{x=a_n}$$

where $\delta_{x=a_n}$ is the dirac measure centered in a_n . Then for $x \notin \{a_n, n \in \mathbb{N}\}$, we have $\mu(]-\infty, x]) = \tilde{\mu}(]-\infty, x]) + \sum_{a_n < x} q_n = TV(v_0,]-\infty, x])$. Passing to the limit $x \rightarrow \infty$ we obtain $\mu(\mathbb{R}) = TV(v_0)$. \square

We know come back to the proof of Lemma B.1 In particular we obtain from (59)

$$|v_2(x) - v_1(x)| \leq \mu([b_x^1; b_x^2]), \quad \text{a.e. } x \in \mathbb{R}.$$

Then we have, using the Fubini Theorem :

$$\begin{aligned} \int_{\mathbb{R}} |v_2(x) - v_1(x)| dx &\leq \int_{x \in \mathbb{R}} \left(\int_{y \in \mathbb{R}} 1_{y \in [b_x^1; b_x^2]} d\mu(y) \right) dx \\ &= \int_{y \in \mathbb{R}} \left(\int_{x \in \mathbb{R}} 1_{y \in [b_x^1; b_x^2]} dx \right) d\mu(y) \\ &= \int_{y \in \mathbb{R}} |(b^2)^{-1}(y) - (b^1)^{-1}(y)| d\mu(y) \\ &\leq \int_{y \in \mathbb{R}} \delta d\mu(y) \leq \delta TV(v_0). \end{aligned}$$

Now in the general case when $a_x^2 \neq a_x^1$, we have

$$|v_2(x) - v_1(x)| \leq TV(v_0; [a_x^1; a_x^2]) + TV(v_0; [b_x^1; b_x^2]) \quad (60)$$

(the proof is analogous to (59)). Then both parts of the R.H.S. of (60) can be handled as before and we deduce (58). \square

Lemma B.3. *Let v_0 be an l.s.c. function with $TV(v_0) < \infty$. We assume that for all $x \in \mathbb{R}$, $a_x \leq b_x$, and $x \rightarrow a_x$, $x \rightarrow b_x$ are non-decreasing functions, and consider*

$$w(x) := \min_{y \in [a_x, b_x]} v_0(y).$$

We have

$$TV(w) \leq TV(v_0).$$

Proof. Let $x_0 < x_1, \dots < x_p$ be real numbers. We want to estimate

$$\delta := \sum_{j=1, \dots, p} |w(x_j) - w(x_{j-1})|$$

For all j , we define y_j as the smallest real number of $[a_{x_j}, b_{x_j}]$ such that $w(x_j) := \min_{[a_{x_j}, b_{x_j}]} v_0(y) = v_0(y_j)$. Let us prove that (y_j) is a non-decreasing sequence. It suffices to check that $x_0 \leq x_1 \Rightarrow y_0 \leq y_1$.

a) In the case $b_{x_0} \leq a_{x_1}$, we obtain $y_0 \leq y_1$ trivially.

b) Otherwise, if $\min_{[a_{x_0}, b_{x_0}]} v_0 = \min_{[a_{x_0}, a_{x_1}]} v_0$, then $y_0 \in [a_{x_0}, a_{x_1}]$ and thus $y_0 \leq y_1$.

c) Otherwise, we have $\min_{[a_{x_0}, b_{x_0}]} v_0 = \min_{[a_{x_1}, b_{x_0}]} v_0$. If the minimum $w(y_1) = \min_{[a_{x_1}, b_{x_1}]} v_0$ is reached on $[b_{x_0}, b_{x_1}]$, then $y_0 \leq b_{x_0} \leq y_1$. If $w(y_1)$ is reached on $[a_{x_1}, b_{x_0}]$, then $y_0 = y_1$. Hence we have proved that $y_0 \leq y_1$ in all cases.

Then, by definition of $TV(v_0)$,

$$\delta = \sum_{j=1, \dots, p} |v_0(y_j) - v_0(y_{j-1})| \leq TV(v_0).$$

Taking the supremum over all non-decreasing sequences (x_j) , we obtain the desired result. \square

Lemma B.4. Let v_0 be a real valued function such that $TV(v_0) < \infty$, then

$$TV(v_0^P) \leq TV(v_0).$$

Proof. : The number of discontinuity points of v_0 is of zero measure as $TV(v_0) < \infty$. It is clear that $|v_0(x + \Delta x) - v_0(x)| \leq TV(v_0; [x, x + \Delta x])$ for almost every $x \in \mathbb{R}$. Then we can write

$$\begin{aligned} TV(v_0^P) &= \sum_{j \in \mathbb{Z}} |V_{j+1}^0 - V_j^0| \\ &\leq \frac{1}{\Delta x} \sum_{j \in \mathbb{Z}} \int_{I_j} |v_0(x + \Delta x) - v_0(x)| dx \\ &\leq \frac{1}{\Delta x} \sum_{j \in \mathbb{Z}} \int_{I_j} TV(v_0, [x, x + \Delta x]) dx \\ &= \frac{1}{\Delta x} \sum_{j \in \mathbb{Z}} \int_{[-\frac{1}{2}, \frac{1}{2}[} TV(v_0, [y\Delta x + x_j, y\Delta x + x_{j+1}]) \Delta x dy \\ &= \int_{[-\frac{1}{2}, \frac{1}{2}[} TV(v_0) dy = TV(v_0) \end{aligned}$$

(we recall that $x_j = j\Delta x$ here). To invert the sum and the integral, we have used that a.e $x \in \mathbb{R}$, v_0 is continuous at the points $(x + x_j)_{j \in \mathbb{Z}}$. \square

C Representation Lemma

In this section we denote by ϑ^* the upper semi continuous (u.s.c.) envelope of a real valued function ϑ . We also denote

$$\vartheta(B^-) := \lim_{y \rightarrow B, y < B} \vartheta(y), \quad \vartheta(B^+) := \lim_{y \rightarrow B, y > B} \vartheta(y).$$

We start with an elementary result related to the minimum of two viscosity solutions.

Lemma C.1. *Let v_0, v_{01} and $v_{02} : \mathbb{R} \rightarrow \mathbb{R}$ be l.s.c. functions. Let ϑ, ϑ_1 and ϑ_2 be the l.s.c solutions of (4a) with initial data v_0, v_{01} and v_{02} respectively. If $v_0 = \min(v_{01}, v_{02})$ then for all $t \geq 0, x \in \mathbb{R}$, $\vartheta(t, x) = \min(\vartheta_1(t, x), \vartheta_2(t, x))$.*

Proof. Let $w = \min(\vartheta_1, \vartheta_2)$. It is easy to check that the function w is l.s.c, and satisfies the initial condition $w(0, x) = v_0(x)$ in the sense of Definition 2.1 ii). Let ϕ be C^1 -regular and $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$ a minimum of $w - \phi$ with $w(t, x) = \vartheta_1(t, x)$ (the case $w(t, x) = \vartheta_2(t, x)$ being similar). Then $\vartheta_1 - \phi$ has also a minimum at (t, x) . Since ϑ_1 is a viscosity solution, we obtain that ϕ satisfies (4a). Hence w is a l.s.c. viscosity solution of (4a). By uniqueness of ϑ , we obtain $w = \vartheta$. \square

Notice that the same arguments would not work for the maximum of two viscosity solutions instead of the minimum.

We now give a representation formula for a viscosity solution in a particular case.

Lemma C.2. *Let $v_0 : \mathbb{R} \rightarrow \mathbb{R}$ be a l.s.c function. We assume that there exists B_1 such that $v_0 \nearrow$ for $x < B_1$, and $v_0 \searrow$ for $x > B_1$, and $v_0(B_1) = \min(v_0(B_1^-), v_0(B_1^+))$. We define the functions:*

$$v_{01}(x) := \begin{cases} v_0(x) & \text{if } x < B_1, \\ v_0(B_1^-) & \text{if } x = B_1, \\ v_0^*(B_1) & \text{if } x > B_1, \end{cases} \quad \text{and} \quad v_{02}(x) := \begin{cases} v_0^*(B_1) & \text{if } x < B_1, \\ v_0(B_1^+) & \text{if } x = B_1, \\ v_0(x) & \text{if } x > B_1. \end{cases}$$

Let ϑ_1 (respectively ϑ_2) be the l.s.c. viscosity solution of (4a) with initial condition v_{01} (respectively v_{02}). Then $v_0 = \min(v_{01}, v_{02})$ and

$$\vartheta(t, x) := \min(\vartheta_1(t, x), \vartheta_2(t, x)), \quad t \geq 0, x \in \mathbb{R}$$

is the l.s.c. viscosity solution of (4a)-(4b). In particular, we have

$$\vartheta(t, x) = \min \left(v_{01}(X_x^M(-t)), v_{02}(X_x^m(-t)) \right).$$

Proof. : It is easy to check that $v_0 = \min(v_{01}, v_{02})$. Then we apply Lemma C.1 to obtain $\vartheta = \min(\vartheta_1, \vartheta_2)$. Let $I_x^t := [X_x^M(-t), X_x^m(-t)]$. We also have $\vartheta_1(t, x) = \inf_{y \in I_x^t} v_{01}(y) = v_{01}(X_x^M(-t))$ as v_{01} is an increasing function, and similarly $\vartheta_2(t, x) = v_{02}(X_x^m(-t))$ as v_{02} is decreasing. \square

References

- [1] R. Agbrall and S. Augoula. High order numerical discretization for hamilton-jacobi equations on triangular meshes. *J. Scientific Computing*, 15(2):197–229, 2000.
- [2] M. Bardi and I. Capuzzo-Dolcetta. *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*. Systems and Control: Foundations and Applications. Birkhäuser, Boston, 1997.
- [3] G. Barles. *Solutions de viscosité des équations de Hamilton-Jacobi*, volume 17 of *Mathématiques & Applications (Berlin)*. Springer-Verlag, Paris, 1994.
- [4] G. Barles and P.E. Souganidis. Convergence of approximation schemes for fully non-linear second order equations. *Asymptotic Analysis*, 4:271–283, 1991.
- [5] O. Bokanowski, S. Martin, R. Munos, and H. Zidani. An anti-diffusive scheme for viability problems. *Applied Numerical Mathematics*, 56(Issue 9, in Numerical Methods for Viscosity Solutions and Applications):1135–1254, 2006.
- [6] O. Bokanowski, N. Megdich, and H. Zidani. An adaptative antidissipative method for optimal control problems. *Arima*, 5:256–271, 2006.
- [7] O. Bokanowski and H. Zidani. Anti-diffusive schemes for linear advection and application to Hamilton-Jacobi-Bellman equations. *J. Sci. Computing*, 30(1):1–33, 2007.
- [8] M.G. Crandall and P.-L. Lions. Two approximations of solutions of Hamilton Jacobi equations. *Math. Comp.*, 43:1–19, 1984.
- [9] B. Després and F. Lagoutière. Contact discontinuity capturing schemes for linear advection and compressible gas dynamics. *J. Sci. Comput.*, 16:479–524, 2001.
- [10] M. Falcone; R. Ferretti. Semi-lagrangian schemes for Hamilton-Jacobi equations, discrete representation formulae and godunov methods. *J. Computational Physics*, 175:559–575, 2002.
- [11] F. Lagoutiere. A non-dissipative entropic scheme for convex scalar equations via discontinuous cell reconstruction. *C. R. Acad. Sci.*, 338(7):549–554, 2004.
- [12] P.L. Lions and P.E. Souganidis. Convergence of muscl and filtered schemes for scalar conservation laws and hamilton jacobi equations. *Numerische Mathematik*, (69):441–470, 1995.
- [13] S. Osher and C-W. Shu. High essentially nonoscillatory schemes for Hamilton-Jacobi equations. *SIAM J. Numer. Anal.*, 28(4):907–922, 1991.
- [14] W. Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.

- [15] P. Saint-Pierre. Approximation of viability kernel. *Appl. Math. Optim.*, 29:187–209, 1994.



Unité de recherche INRIA Futurs
Parc Club Orsay Université - ZAC des Vignes
4, rue Jacques Monod - 91893 ORSAY Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399