

Markov processes and parabolic partial differential equations

Mireille Bossy, Nicolas Champagnat

► **To cite this version:**

Mireille Bossy, Nicolas Champagnat. Markov processes and parabolic partial differential equations. Cont Rama. Encyclopedia of Quantitative Finance, John Wiley & Sons Ltd. Chichester, UK, pp.1142-1159, 2010. <inria-00193883v2>

HAL Id: inria-00193883

<https://hal.inria.fr/inria-00193883v2>

Submitted on 25 Jan 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Markov processes and parabolic partial differential equations

Mireille Bossy*, Nicolas Champagnat*

January 25, 2008

Abstract

In the first part of this article, we present the main tools and definitions of Markov processes' theory: transition semigroups, Feller processes, infinitesimal generator, Kolmogorov's backward and forward equations and Feller diffusion. We also give several classical examples including stochastic differential equations (SDEs) and backward SDEs (BSDEs). The second part of this article is devoted to the links between Markov processes and parabolic partial differential equations (PDEs). In particular, we give Feynman-Kac formula for linear PDEs, we present Feynman-Kac formula for BSDEs, and we give some examples of the correspondence between stochastic control problems and Hamilton-Jacobi-Bellman (HJB) equations and between optimal stopping problems and variational inequalities. Several examples of finan-

*TOSCA project-team

INRIA Sophia Antipolis – Méditerranée Research Center

2004 route des Lucioles, BP-93

06902 Sophia Antipolis Cedex, France

Mireille.Bossy@sophia.inria.fr and Nicolas.Champagnat@sophia.inria.fr

cial applications are given to illustrate each of these results, including European options, Asian options and American put options.

Keywords: Markov processes; parabolic partial differential equations; semigroup; Feller processes; strong Markov property; infinitesimal generator; Kolmogorov's backward and forward equations; Fokker-Planck equation; stochastic differential equations; backward stochastic differential equations; heat equation; Feynman-Kac formula; Hamilton-Jacobi-Bellman equation; variational inequality.

1 Introduction

We consider a stochastic process $(X_t, t \geq 0)$ adapted to a filtration $(\mathcal{F}_t, t \geq 0)$. The σ -field \mathcal{F}_t represents the information available at time t : any event that is determined by the path of X_s for $s \in [0, t]$ belongs to \mathcal{F}_t .

A Markov process evolves in a memoryless way: its future law depends on the past only through the present position of the process. In terms of conditional expectations, this can be written as follows: a process $(X_t, t \geq 0)$ adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ is a Markov process if

$$\mathbb{E}(f(X_{t+s}) \mid \mathcal{F}_t) = \mathbb{E}(f(X_{t+s}) \mid X_t) \quad (1)$$

for all $s, t \geq 0$ and f bounded and measurable.

The interest of such a process in financial models becomes clear when one observes that the price of an option, or more generally the value at time t of any future claim with maturity T , is given by the general formula

$$\text{value at time } t = \mathbb{E}(\text{payoff at time } T \mid \mathcal{F}_t),$$

for a convenient probability law. The Markov property is a frequent assumption in financial models because it provides powerful tools (semigroup, theory of partial differential equations, or *PDEs*,...) for the quantitative analysis of such problems.

This is illustrated by the classical example of the Black-Scholes PDE. Assume that the price of a stock follows a Black-Scholes model $(S_t, t \geq 0)$ with constant drift r and volatility σ . The price V of a call option on this stock can be characterized by $V_t = u(t, S_t)$ where the function $u(t, x)$ is differentiable w.r.t. t and twice differentiable w.r.t. x and solves the Black-Scholes PDE

$$\begin{cases} \frac{\partial u}{\partial t} + rx \frac{\partial u}{\partial x} + \frac{\sigma^2}{2} x^2 \frac{\partial^2 u}{\partial x^2} - ru = 0, & \forall (t, x) \in [0, T) \times (0, +\infty) \\ u(T, x) = (x - K)^+, & \forall x \in (0, +\infty). \end{cases} \quad (2)$$

The price as defined above can also be characterized in terms of the Markov process $(S_t, t \geq 0)$:

$$V_t = \mathbb{E}(e^{-r(T-t)} f(S_T) \mid \mathcal{F}_t)$$

with $f(x) = (x - K)^+$ and, because of the Markov property (1), $u(t, S_t) = \mathbb{E}(e^{-r(T-t)} f(S_T) \mid S_t)$. The equality $u(t, x) = \mathbb{E}(e^{-r(T-t)} f(S_T) \mid S_t = x)$ is a consequence of the Markov property of $(S_t, t \geq 0)$. The goal of this article is to clarify and generalize this relation between PDEs and Markov processes and to illustrate the role of Markovian models in various financial problems..

The natural approach is opposite to the previous example: we start from the description of the stochastic process and then obtain the PDEs from quantities related to this process. We give a general overview of the links between Markov processes and PDEs without coming into all details and we focus on the case of Markov processes solution to stochastic differential equations (SDEs).

This article is organized as follows. In Section 2, we precisely define Markov processes and introduce the fundamental tools: their semigroup and infinitesimal generator. The section ends with classical examples of Markov processes. In Section 3, we explore in details the links between Markov processes and PDEs: we give the Feynman-Kac formula for linear PDEs, we explore the links between backward stochastic differential equations (BSDEs) and quasi-linear and semi-linear PDEs, and we give the probabilistic interpretation of some Hamilton-Jacobi-Bellman (HJB) equations and variational inequalities. Finally, Section 4 gives

concluding remarks on the use and limits of the approach and on simulations.

Main references: [10, 11, 15, 16, 18, 19, 23]

2 Markov processes

2.1 Definition and first properties

We will restrict ourselves to \mathbb{R}^d -valued Markov processes. The set of Borel subsets of \mathbb{R}^d is denoted by \mathcal{B} . In the following, we will denote Markov processes by $(X_t, t \geq 0)$, or simply X when no confusion is possible.

Markov property and transition semigroup

A Markov process retains no memory of where it has been in the past. Only the current state of the process influences its future dynamics. The following definition makes this formal.

Definition 2.1 *Let $(X_t, t \geq 0)$ be a stochastic process defined on a probability filtered space $(\Omega, \mathcal{F}_t, \mathbb{P})$ with values in \mathbb{R}^d . X is a Markov process if*

$$\mathbb{P}(X_{t+s} \in \Gamma \mid \mathcal{F}_t) = \mathbb{P}(X_{t+s} \in \Gamma \mid X_t) \quad \mathbb{P} - \text{a.s.} \quad (3)$$

for all $s, t \geq 0$ and $\Gamma \in \mathcal{B}$. Equation (3) is called the Markov property of the process X . The Markov process is called time-homogeneous if the law of X_{t+s} conditionally on $X_t = x$ is independent of t .

Observe that (3) is equivalent to (1) and that X is a time-homogeneous Markov process if there exists a positive function P defined on $\mathbb{R}_+ \times \mathbb{R}^d \times \mathcal{B}$ such that

$$P(s, X_t, \Gamma) = \mathbb{P}(X_{t+s} \in \Gamma \mid \mathcal{F}_t)$$

holds \mathbb{P} -a.s. for all $t, s \geq 0$ and $\Gamma \in \mathcal{B}$. P is called the *transition function* of the time homogeneous Markov process X .

For the moment, we restrict ourselves to the time-homogeneous case.

Proposition 2.2 *The transition function P of a time-homogeneous Markov process X satisfies*

1. $P(t, x, \cdot)$ is a probability measure on \mathbb{R}^d for any $t \geq 0$ and $x \in \mathbb{R}^d$,
2. $P(0, x, \cdot) = \delta_x$ (unit mass at x) for any $x \in \mathbb{R}^d$,
3. $P(\cdot, \cdot, \Gamma)$ is measurable for any $\Gamma \in \mathcal{B}$

and for any $s, t \geq 0$, $x \in \mathbb{R}^d$, $\Gamma \in \mathcal{B}$, P satisfies the Chapman-Kolmogorov property

$$P(t + s, x, \Gamma) = \int_{\mathbb{R}^d} P(s, y, \Gamma) P(t, x, dy).$$

From an analytical viewpoint, we can think of the transition function as a Markov semigroup¹ $(P_t, t \geq 0)$, defined by

$$P_t f(x) := \int_{\mathbb{R}^d} P(t, x, dy) f(dy) = \mathbb{E}(f(X_t) \mid X_0 = x), \quad (4)$$

in which case the Chapman-Kolmogorov equation becomes the *semigroup property*

$$P_s P_t = P_{t+s}, \quad s, t \geq 0. \quad (5)$$

Conversely, given a Markov semigroup $(P_t, t \geq 0)$ and a probability measure ν on \mathbb{R}^d , it is always possible to construct a Markov process X with initial law ν that satisfies (4) (see [10, Th.4.1.1]). The links between PDEs and Markov processes are based on this equivalence between semigroups and Markov processes. This can be expressed through a single object: the infinitesimal generator.

Strong Markov property, Feller processes and infinitesimal generator

Recall that a random time τ is called a \mathcal{F}_t -stopping time if $\{\tau \leq t\} \in \mathcal{F}_t$ for any $t \geq 0$.

¹A Markov semigroup family $(P_t, t \geq 0)$ on \mathbb{R}^d is a family of bounded linear operators of norm 1 on the set of bounded measurable functions on \mathbb{R}^d equipped with the L^∞ norm, that satisfy (5).

Definition 2.3 A Markov process $(X_t, t \geq 0)$ with transition function $P(t, x, \Gamma)$ is strong Markov if, for any \mathcal{F}_t -stopping time τ ,

$$\mathbb{P}(X_{\tau+t} \in \Gamma \mid \mathcal{F}_\tau) = P(t, X_\tau, \Gamma)$$

for all $t \geq 0$ and $\Gamma \in \mathcal{B}$.

Let $C_0(\mathbb{R}^d)$ denote the space of bounded continuous functions on \mathbb{R}^d which vanish at infinity, equipped with the L^∞ norm denoted by $\|\cdot\|$.

Definition 2.4 A Feller semigroup² is a strongly continuous³, positive, Markov semigroup $(P_t, t \geq 0)$ such that $P_t : C_0(\mathbb{R}^d) \rightarrow C_0(\mathbb{R}^d)$ and

$$\begin{aligned} \forall f \in C_0(\mathbb{R}^d), \quad 0 \leq f \Rightarrow 0 \leq P_t f \\ \forall f \in C_0(\mathbb{R}^d), \quad \forall x \in \mathbb{R}^d, \quad P_t f(x) \rightarrow f(x) \quad \text{as } t \rightarrow 0 \end{aligned} \quad (6)$$

For a Feller semigroup, the corresponding Markov process can be constructed as a strong Markov process.

Theorem 2.5 ([10] Th.4.2.7) Given a Feller semigroup $(P_t, t \geq 0)$ and any probability measure ν on \mathbb{R}^d , there exists a filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{P})$ and a strong Markov process $(X_t, t \geq 0)$ on this space with values in \mathbb{R}^d with initial law ν and with transition function P_t . A strong Markov process whose semigroup is Feller is called a Feller process.

We are now in a position to introduce the key notion of *infinitesimal generator* of a Feller process.

²This is not the most general definition of Feller semigroups (see [23, Def.III.6.5]). In our context, because we only introduce analytical objects from stochastic processes, the semigroup (P_t) is naturally defined on the set of bounded measurable functions.

³The strong continuity of a semigroup is usually defined as $\|P_t f - f\| \rightarrow 0$ as $t \rightarrow 0$ for all $f \in C_0(\mathbb{R}^d)$. However, in the case of Feller semigroups, this is equivalent to the weaker formulation (6) (see [23, Lemma III.6.7]).

Definition 2.6 For a Feller process $(X_t, t \geq 0)$, the infinitesimal generator of X is the (generally unbounded) linear operator $L : \mathcal{D}(L) \rightarrow C_0(\mathbb{R}^d)$ defined as follows. We write $f \in \mathcal{D}(L)$ if, for some $g \in C_0(\mathbb{R}^d)$, we have

$$\frac{\mathbb{E}(f(X_t) \mid X_0 = x) - f(x)}{t} \rightarrow g(x)$$

when $t \rightarrow 0$ for the norm $\|\cdot\|$, and we then define $Lf = g$.

By Theorem 2.5, an equivalent definition can be obtained by replacing X by its Feller semigroup $(P_t, t \geq 0)$. In particular, for all $f \in \mathcal{D}(L)$,

$$Lf(x) = \lim_{t \rightarrow 0} \frac{P_t f(x) - f(x)}{t} = \lim_{t \rightarrow 0} \frac{\mathbb{E}(f(X_t) \mid X_0 = x) - f(x)}{t}. \quad (7)$$

An important property of the infinitesimal generator is that it allows one to construct fundamental martingales associated with a Feller process.

Theorem 2.7 ([23], III.10) Let X be a Feller process on $(\Omega, \mathcal{F}_t, \mathbb{P})$ with infinitesimal generator L such that $X_0 = x \in \mathbb{R}^d$. For all $f \in \mathcal{D}(L)$,

$$f(X_t) - f(x) - \int_0^t Lf(X_s) ds$$

defines a \mathcal{F}_t -martingale. In particular,

$$\mathbb{E}(f(X_t)) = f(x) + \mathbb{E}\left(\int_0^t Lf(X_s) ds\right). \quad (8)$$

As explained above, the law of a Markov process is characterized by its semigroup. In most cases, a Feller semigroup can be itself characterized by its infinitesimal generator (the precise conditions for this to hold are given by the Hille-Yosida Theorem, see [23, Th.III.5.1]). For almost all Markov financial models, these conditions are well established and always satisfied (see the examples of Section 2.2). As illustrated by (8), when $\mathcal{D}(L)$ is large enough, the infinitesimal generator captures the law of the whole dynamics of a Markov process and provides an analytical tool to study the Markov process. The other major mathematical tool used in finance is the stochastic calculus, which applies to semi-martingales (see [20]). It is

therefore crucial for applications to characterize under which conditions a Markov process is a semi-martingale. This question is answered for very general processes in [6]. We mention that this is always the case for Feller diffusions, defined below.

Feller diffusions

Let us consider the particular case of continuous Markov processes, which include the solutions of stochastic differential equations (SDEs).

Definition 2.8 *A Feller diffusion on \mathbb{R}^d is a Feller process X on \mathbb{R}^d that has continuous paths, and such that the domain $\mathcal{D}(L)$ of the generator L of X contains the space $C_K^\infty(\mathbb{R}^d)$ of infinitely differentiable functions of compact support.*

Feller diffusions are Markov processes admitting a second-order differential operator as infinitesimal generator.

Theorem 2.9 *For any $f \in C_K^\infty(\mathbb{R}^d)$, the infinitesimal generator L of a Feller diffusion has the form*

$$Lf(x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2 f}{\partial x_i \partial x_j}(x) + \sum_{i=1}^d b_i(x) \frac{\partial f}{\partial x_i}(x), \quad (9)$$

where the functions $a_{ij}(\cdot)$ and $b_i(\cdot)$, $1 \leq i, j \leq d$ are continuous and the matrix $a = (a_{ij}(x))_{1 \leq i, j \leq d}$ is non-negative definite symmetric for all $x \in \mathbb{R}^d$.

Kolmogorov backward and forward equations

Observe by (7) that the semigroup P_t of a Feller process X satisfies the following differential equation: for all $f \in \mathcal{D}(L)$,

$$\frac{d}{dt} P_t f = L P_t f. \quad (10)$$

This equation is called Kolmogorov's backward equation. In particular, if L is a differential operator (for example if X is a Feller diffusion), the function $u(t, x) =$

$P_t f(x)$ is solution of the PDE

$$\begin{cases} \frac{\partial u}{\partial t} = Lu \\ u(0, x) = f(x). \end{cases} \quad (11)$$

Conversely, if this PDE admits a unique solution, then its solution is given by

$$u(t, x) = \mathbb{E}(f(X_t) \mid X_0 = x). \quad (12)$$

This is the simplest example of a probabilistic interpretation of the solution of a PDE in terms of a Markov process.

Moreover, since Feller semigroups are strongly continuous, it is easy to check that the operators P_t and L commute. Therefore, (10) may be rewritten as

$$\frac{d}{dt} P_t f = P_t L f.$$

This equation is known as Kolmogorov's forward equation. It is the weak formulation of the equation

$$\frac{d}{dt} \mu_t^x = L^* \mu_t^x,$$

where the probability measure μ_t^x on \mathbb{R}^d denotes the law of X_t conditioned on $X_0 = x$ and where L^* is the adjoint operator of L . In particular, with the notation of Theorem 2.9, if X is a Feller diffusion and if $\mu_t^x(dy)$ admits a density $q(x; t, y)$ with respect to Lebesgue's measure on \mathbb{R}^d (which holds for example if the functions $b_i(x)$ and $a_{ij}(x)$ are bounded and locally Lipschitz, if the functions $a_{ij}(x)$ are globally Hölder and if the matrix $a(x)$ is uniformly positive definite [11, Th.6.5.2]), the forward Kolmogorov equation is the weak form (in the sense of the distribution theory) of the PDE

$$\frac{\partial}{\partial t} q(x; t, y) = - \sum_{i=1}^d \frac{\partial}{\partial y_i} (b_i(y) q(x; t, y)) + \sum_{i,j=1}^d \frac{\partial^2}{\partial y_i \partial y_j} (a_{ij}(y) q(x; t, y)).$$

This equation is known as Fokker-Planck equation and gives another family of PDEs which have a probabilistic interpretations. Fokker-Planck equation has applications in finance for quantiles, value at risk or risk measure computations [24],

whereas Kolmogorov's backward equation (10) or (11) is more suited to financial problems related to the hedging of derivatives products or portfolio allocation (see Section 3 for examples and for the probabilistic interpretation of more general PDEs).

Some comments on time-inhomogeneous Markov processes

The law of a time-inhomogeneous Markov process is described by the doubly-indexed family of operators $(P_{s,t}, 0 \leq s \leq t)$ where, for any bounded measurable f and any $x \in \mathbb{R}^d$,

$$P_{s,t}f(x) = \mathbb{E}(f(X_t) | X_s = x).$$

Then, the semigroup property becomes, for $s \leq t \leq r$,

$$P_{s,t}P_{t,r} = P_{s,r}.$$

The Definition 2.4 of Feller semigroups can be generalized to time-inhomogeneous processes as follows. The time-inhomogeneous Markov process X is called a *Feller time-inhomogeneous process* if $(P_{s,t}, 0 \leq s \leq t)$ is a family of positive, Markov linear operators on $C_0(\mathbb{R}^d)$ which is strongly continuous in the sense

$$\forall s \geq 0, x \in \mathbb{R}^d, f \in C_0(\mathbb{R}^d), \quad \|P_{s,t}f - f\| \rightarrow 0 \quad \text{as } t \rightarrow s.$$

In this case, it is possible to generalize the notion of infinitesimal generator. For any t , let

$$L_t f(x) = \lim_{s \rightarrow 0} \frac{P_{t,t+s}f(x) - f(x)}{s} = \lim_{s \rightarrow 0} \frac{\mathbb{E}[f(X_{t+s}) | X_t = x] - f(x)}{s}$$

for any $f \in C_0(\mathbb{R}^d)$ such that $L_t f \in C_0(\mathbb{R}^d)$ and the limit above holds in the sense of the norm $\|\cdot\|$. The set of such $f \in C_0(\mathbb{R}^d)$ is called the domain $\mathcal{D}(L_t)$ of the operator L_t . $(L_t, t \geq 0)$ is called the family of *time-inhomogeneous infinitesimal generators* of the process X .

All the results on Feller processes stated above can be easily transposed to the time-inhomogeneous case, observing that, if $(X_t, t \geq 0)$ is a time-inhomogeneous

Markov process on \mathbb{R}^d , then $(\tilde{X}_t, t \geq 0)$ where $\tilde{X}_t = (t, X_t)$ is a time-homogeneous Markov process on $\mathbb{R}_+ \times \mathbb{R}^d$. Moreover, if X is time-inhomogeneous Feller, it is elementary to check that the process \tilde{X} is time-homogeneous Feller as defined in Definition 2.4. Its semigroup $(\tilde{P}_t, t \geq 0)$ is linked to the time-inhomogeneous semigroup by the relation

$$\tilde{P}_t f(s, x) = \mathbb{E}[f(s+t, X_{s+t}) \mid X_s = x] = (P_{s, s+t} f(s+t, \cdot))(x)$$

for all bounded and measurable $f : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$. If \tilde{L} denotes the infinitesimal generator of the process \tilde{X} , it is elementary to check that, for any $f(t, x) \in \mathcal{D}(\tilde{L})$ which is differentiable w.r.t. t , with derivative uniformly continuous in (t, x) , $x \mapsto f(t, x)$ belongs to $\mathcal{D}(L_t)$ for any $t \geq 0$ and

$$\tilde{L}f(t, x) = \frac{\partial f}{\partial t}(t, x) + (L_t f(t, \cdot))(x).$$

On this observation, it is possible to apply Theorem 2.9 to time-inhomogeneous Feller diffusions, defined as continuous time-inhomogeneous Feller processes with infinitesimal generators $(L_t, t \geq 0)$ such that $C_K^\infty(\mathbb{R}^d) \subset \mathcal{D}(L_t)$ for any $t \geq 0$. For such processes, there exist continuous functions b_i and a_{ij} , $1 \leq i, j \leq d$ from $\mathbb{R}_+ \times \mathbb{R}^d$ to \mathbb{R} such that the matrix $a(t, x) = (a_{i,j}(t, x))_{1 \leq i, j \leq d}$ is symmetric non-negative definite and

$$L_t f(x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(t, x) \frac{\partial^2 f}{\partial x_i \partial x_j}(x) + \sum_{i=1}^d b_i(t, x) \frac{\partial f}{\partial x_i}(x) \quad (13)$$

for all $t \geq 0$, $x \in \mathbb{R}^d$ and $f \in C_K^\infty(\mathbb{R}^d)$.

For more details on time-inhomogeneous Markov processes, we refer to [11].

2.2 Examples

In financial models, the following Markov processes are the most used.

The Brownian motion

The standard one-dimensional Brownian motion $(B_t, t \geq 0)$ is a Feller diffusion in \mathbb{R} ($d = 1$) such that $B_0 = 0$ and for which the parameters of Theorem 2.9

are $b = 0$ and $a = 1$. The Brownian motion is the fundamental prototype of Feller diffusions. Other diffusions are inherited from this process since they can be expressed as solutions to SDEs driven by independent Brownian motions (see below). Similarly, the standard d -dimensional Brownian motion is a vector of d independent standard one-dimensional Brownian motions and corresponds to the case $b_i = 0$ and $a_{ij} = \delta_{ij}$ for $1 \leq i, j \leq d$, where δ_{ij} is the Kronecker delta function ($\delta_{ij} = 1$ if $i = j$ and 0 otherwise).

The Black-Scholes model

In the Black-Scholes model, the underlying asset price S_t follows a geometric Brownian motion with constant drift μ and volatility σ .

$$S_t = S_0 \exp((\mu - \sigma^2/2)t + \sigma B_t),$$

where B is a standard Brownian motion. With Itô's formula, it is easily checked that S is a Feller diffusion with infinitesimal generator

$$Lf(x) = \mu x f'(x) + \frac{1}{2} \sigma^2 x^2 f''(x).$$

Itô's formula also yields

$$S_t = S_0 + \mu \int_0^t S_s ds + \sigma \int_0^t S_s dB_s$$

which can be written as the stochastic differential equation (SDE)

$$dS_t = \mu S_t dt + \sigma S_t dB_t. \tag{14}$$

The correspondance between the SDE and the second-order differential operator L appears below as a general fact.

Stochastic differential equations

SDEs are probably the most used Markov models in finance. Solutions of SDEs are particular examples of Feller diffusions. When the parameters b_i and a_{ij} of

Theorem 2.9 are sufficiently regular, a Feller process X with generator (9) can be constructed as the solution of the SDE

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t, \quad (15)$$

where $b(x) \in \mathbb{R}^d$ is $(b_1(x), \dots, b_d(x))$, where the $d \times r$ matrix $\sigma(x)$ satisfies $a_{ij}(x) = \sum_{k=1}^r \sigma_{ik}(x)\sigma_{jk}(x)$ (i.e. $a = \sigma\sigma'$) and where B_t is a r -dimensional standard Brownian motion. For example, when $d = r$, one can take for $\sigma(x)$ the symmetric square root matrix of the matrix $a(x)$.

The construction of a Markov solutions to the SDE (15) with generator (9) is possible if the matrix σ can be chosen such that b and σ are globally Lipschitz with linear growth [15, Th.5.2.9], or if b and a are bounded and continuous functions [15, Th.5.4.22]. In the second case, the SDE has a solution in a weaker sense. Uniqueness (at least in law) and the strong Markov property hold if b and σ are locally Lipschitz [15, Th.5.2.5], or if b and a are Hölder-continuous and the matrix a is uniformly positive definite [15, Rmk.5.4.30, Th.5.4.20]. In the one-dimensional case, existence and uniqueness for the SDE (15) can be proved under weaker assumptions [15, Sec.5.5].

In all these cases, the Markov property allows one to identify the SDE (15) with its generator (9). This will allow us to make the link between parabolic PDEs and the corresponding SDE in the Section 3.

Similarly, one can associate to the time-inhomogeneous SDE

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t \quad (16)$$

the time-inhomogeneous generators (13). Existence for this SDE holds if b_i and σ_{ij} are globally Lipschitz in x and and locally bounded (uniqueness holds if b_i and σ_{ij} are only locally Lipschitz in x). As above, in this case, a solution to (16) is strong Markov. We refer the reader to [18] for more details.

Backward stochastic differential equations

Backward stochastic differential equations (BSDEs) are solutions of SDEs with fixed random terminal conditions. Let us motivate the definition of a BSDE by continuing the study of the elementary example of the introduction of this article.

Consider an asset S_t modeled by the Black-Scholes SDE (14) and assume that it is possible to borrow and lend cash at a constant risk-free interest rate r . A self-financed trading strategy is determined by an initial portfolio value and the amount π_t of the portfolio value placed in the risky asset at time t . Given the stochastic process $(\pi_t, t \geq 0)$, the portfolio value V_t at time t solves the SDE

$$dV_t = rV_t dt + \pi_t(\mu - r)dt + \sigma\pi_t dB_t, \quad (17)$$

where B is the Brownian motion driving the dynamics (14) of the risky asset S .

Assume that this portfolio serves to hedge a call option with strike K and maturity T . This problem can be expressed as finding a couple of processes (V_t, π_t) adapted to the Brownian filtration $\mathcal{F}_t = \sigma(B_s, s \leq t)$ such that

$$V_t = (S_T - K)^+ - \int_t^T (rV_s + \pi_s(\mu - r))ds - \int_t^T \sigma\pi_s dB_s. \quad (18)$$

Such SDEs with terminal condition and with unknown process driving the Brownian integral are called BSDEs. This particular BSDE admits a unique solution (see Section 3.3) and can be explicitly solved. Since V_0 is \mathcal{F}_0 -adapted, it is non-random and therefore V_0 is the usual free arbitrage price of the option. In particular, choosing $\mu = r$, we recover the usual formula for the free arbitrage price $V_0 = \mathbb{E}[e^{-rT}(S_T - K)^+]$, and the quantity of risky asset π_t/S_t in the portfolio is given by the Black-Scholes Δ -hedge $\frac{\partial u}{\partial x}(t, S_t)$, where $u(t, x)$ is the solution of the PDE (2). Applying Itô formula to $u(t, S_t)$, an elementary computation shows that $u(t, S_t)$ solves the same SDE (17) as V_t , with the same terminal condition. Therefore, by uniqueness, $V_t = u(t, S_t)$.

Usually, for more general BSDEs, $(\pi_t, t \geq 0)$ is an implicit process given by the martingale representation theorem. We give in Section 3.3 results on the existence and uniqueness of solutions of BSDEs, and on their links with non-linear PDEs.

Non continuous Markov processes

In financial models, it is sometimes natural to consider non-continuous Markov processes, for examples when one wants to take into account external shocks, such as earning announcements, changes of interest rates, political assassinations or currency collapses. This can sometimes be done by adding Poisson processes to the dynamics, modelling discrete jumps. This is relevant for example when one wants to study incomplete markets or credit risk for intensity models (e.g. default intensity or Cox processes).

In other situations, Markov processes related to Lévy processes must be considered. In particular, it is possible to define the analogue of SDEs where the Brownian motion is replaced by a Lévy process. We refer to [14] for an introduction on the topic. Financial applications include the CGMY model, the NIG model or the generalized hyperbolic model. In a financial context, statistical tests have been recently developed and experimented on historical price records [1].

About Markov processes and the dimension of the state space

In many pricing/hedging problems, the dimension of the pricing PDE is greater than the state space of the undelyings. In such cases, the financial problem is apparently related to non-Markov stochastic processes. However, it can usually be expressed in terms of Markov processes if one increases the dimension of the process considered. For example, in the context of Markov short rates $(r_t, t \geq 0)$, the pricing of a zero-coupon bond is expressed in terms of the process $R_t = \int_0^t r_s ds$ which is not Markovian, whereas the couple (r_t, R_t) is. For Asian options on a Markov asset, the couple formed by the asset and its integral is Markovian. If the asset involves a stochastic volatility solution to a SDE, then the couple formed by the asset value and its volatility is Markov. As mentionned above, another important example is given by time-inhomogeneous Markov processes which become time-homogeneous when one considers the couple formed by the current time and

the original process.

In some cases, the dimension of the system can be reduced while preserving the Markovian nature of the problem. In the case of the portfolio management of multidimensional Black-Scholes prices with deterministic volatility matrix, mean return vector and interest rate, the dimension of the problem is actually reduced to one (this is the case, e.g. in Merton's problem). When the volatility matrix, the mean return vector and the interest rate are Markov processes of dimension d' , the dimension of the problem is reduced to $d' + 1$.

3 Parabolic PDEs associated to Markov processes

Computing the value of any future claim with fixed maturity (for example, the price of a European option on an asset solution to a SDE), or solving an optimal portfolio management problem, amounts to solve a parabolic second order PDE, that is a PDE of the form

$$\frac{\partial u}{\partial t}(t, x) + L_t u(t, x) = f(t, x, u(t, x), \nabla u(t, x)), \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$$

where $\nabla u(t, x)$ is the gradient of $u(t, x)$ w.r.t. x and the linear differential operators L_t has the form (13).

The goal of this section is to explain the links between these PDEs and the original diffusion process, or some intermediate Markov process. We will distinguish between *linear* parabolic PDEs, where the function $f(t, x, y, z)$ does not depend on z and is linear in y , *semi-linear* parabolic PDEs, where the function $f(t, x, y, z)$ does not depend on z but is nonlinear in y , and *quasi-linear* parabolic PDEs, where the function $f(t, x, y, z)$ is nonlinear in (y, z) . We will also discuss the links between diffusion processes and some *fully non-linear* PDEs (Hamilton-

Jacobi-Bellman equations or variational inequalities) of the form

$$F\left(t, \frac{\partial u}{\partial t}(t, x), u(t, x), \nabla u(t, x), Hu(t, x)\right) = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$$

for some non-linear function F , where Hu denotes the Hessian matrix of u w.r.t. the space variable x .

Such problems involve several notions of solutions discussed in the literature. In Sections 3.1 and 3.2, we consider *classical solutions*, i.e. solutions that are continuously differentiable w.r.t. the time variable, and twice continuously differentiable w.r.t. the space variables. In Sections 3.3 and 3.4, because of the nonlinearity of the problem, classical solutions may not exist, and one must consider the weaker notion of viscosity solutions.

In Section 3.1, we consider heat-like equations where the solution can be explicitly computed. Section 3.2 deals with linear PDEs, Section 3.3 with quasi- and semilinear PDEs and their links with BSDEs and Section 3.4 deals with optimal control problems.

3.1 Brownian motion, Black-Scholes model, Ornstein-Uhlenbeck process and the heat equation

The *heat equation* is the first example of parabolic PDE with basic probabilistic interpretation (for which there is no need of stochastic calculus).

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = \frac{1}{2}\Delta u(t, x), & (t, x) \in (0, +\infty) \times \mathbb{R}^d \\ u(0, x) = f(x), & x \in \mathbb{R}^d, \end{cases} \quad (19)$$

where Δ denotes the Laplacian operator of \mathbb{R}^d . When f is a bounded measurable function, it is well-known that the solution of this problem is given by the formula

$$u(t, x) = \int_{\mathbb{R}^d} f(y)g(x; t, y)dy \quad (20)$$

where

$$g(x; t, y) = \frac{1}{(2\pi t)^{d/2}} \exp\left(-\frac{|x-y|^2}{2t}\right).$$

$|\cdot|$ denotes the Euclidean norm on \mathbb{R}^d . g is often called the *fundamental solution* of the heat equation. We recognize that $g(x; t, y)dy$ is the law of $x + B_t$ where B is a standard d -dimensional Brownian motion. Therefore, (20) may be rewritten as

$$u(t, x) = \mathbb{E}[f(x + B_t)],$$

which provides a simple probabilistic interpretation of the solution of the heat equation in \mathbb{R}^d as a particular case of (12). Note that (19) involves the infinitesimal generator of the Brownian motion $(1/2)\Delta$.

Let us mention two other cases where the link between PDEs and stochastic processes can be done without stochastic calculus. The first one is the Black-sholes model, solution to the SDE

$$dS_t = S_t(\mu dt + \sigma dB_t).$$

When $d = 1$, its infinitesimal generator is $Lf(x) = \mu x f'(x) + (\sigma^2/2)x^2 f''(x)$ and its law at time t when $S_0 = x$ is $l(x; t, y)dy$ where

$$l(x; t, y) = \frac{1}{\sigma y \sqrt{2\pi t}} \exp \left[-\frac{1}{2\sigma^2 t} \left(\log \frac{y}{x} - (\mu - \sigma^2/2)t \right)^2 \right].$$

Then, for any bounded and measurable f , elementary computations show that

$$u(t, x) = \int_0^\infty f(y)l(x; t, y)dy$$

satisfies

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = Lu(t, x), & (t, x) \in (0, +\infty)^2 \\ u(0, x) = f(x), & x \in (0, +\infty). \end{cases}$$

Here again, this formula gives immediately the probabilistic interpretation

$$u(t, x) = \mathbb{E}[f(S_t) | S_0 = x].$$

The last example is the Ornstein-Uhlenbeck process in \mathbb{R}

$$dX_t = \beta X_t dt + \sigma dB_t,$$

with $\beta \in \mathbb{R}$, $\sigma > 0$ and $X_0 = x$. The infinitesimal generator of this process is $Af(x) = \beta x f'(x) + (\sigma^2/2)f''(x)$. It can be easily checked that X_t is a Gaussian random variable with mean $x \exp(\beta t)$ and variance $\sigma^2(\exp(2\beta t) - 1)/2\beta$ with the convention that $(\exp(2\beta t) - 1)/2\beta = t$ if $\beta = 0$. Therefore, its probability density function is given by

$$h(x; t, y) = \sqrt{\frac{\beta}{\sigma^2 \pi (\exp(2\beta t) - 1)}} \exp \left[-\frac{2\beta(y - x \exp(\beta t))^2}{\sigma^2 (\exp(2\beta t) - 1)} \right].$$

Then, for any bounded and measurable f ,

$$u(t, x) = \int_{\mathbb{R}} f(y) h(x; t, y) dy = \mathbb{E}[f(X_t) \mid X_0 = x]$$

is solution of

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = Au(t, x), & (t, x) \in (0, +\infty) \times \mathbb{R} \\ u(0, x) = f(x), & x \in \mathbb{R}. \end{cases}$$

3.2 Generic linear case: Feynman-Kac formula

The probabilistic interpretations of the previous PDEs can be generalized to a large class of linear parabolic PDEs with arbitrary second order differential operator, interpreted as the infinitesimal generator of a Markov process. Assume that the vector $b(t, x) \in \mathbb{R}^d$ and the $d \times r$ matrix $\sigma(t, x)$ are uniformly bounded and locally Lipschitz functions on $[0, T] \times \mathbb{R}^d$ and consider the SDE in \mathbb{R}^d

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t \tag{21}$$

where B is a standard r -dimensional Brownian motion. Set $a = \sigma\sigma'$ and assume also that the $d \times d$ matrix $a(t, x)$ satisfies the uniform ellipticity condition: there exists $\gamma > 0$ such that for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $\xi \in \mathbb{R}^d$,

$$\sum_{i,j=1}^d a_{ij}(t, x)\xi_i\xi_j \geq \gamma|\xi|^2. \tag{22}$$

Let $(L_t)_{t \geq 0}$ be the family of time-inhomogeneous infinitesimal generators of the Feller diffusion X_t solution to the SDE (21), given by (13). Consider the Cauchy

problem

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + L_t u(t, x) + c(t, x)u(t, x) = f(t, x), & (t, x) \in [0, T] \times \mathbb{R}^d \\ u(T, x) = g(x), & x \in \mathbb{R}^d \end{cases} \quad (23)$$

where $c(t, x)$ is uniformly bounded and locally Hölder on $[0, T] \times \mathbb{R}^d$, $f(t, x)$ is locally Hölder on $[0, T] \times \mathbb{R}^d$, $g(x)$ is continuous on \mathbb{R}^d and

$$|f(t, x)| \leq A \exp(a|x|), \quad |g(x)| \leq A \exp(a|x|), \quad \forall (t, x) \in [0, T] \times \mathbb{R}^d$$

for some constants $A, a > 0$. Under these conditions, it follows easily from Theorems 6.4.5 and 6.4.6 of [11] that (23) admits a unique classical solution u such that

$$|u(t, x)| \leq A' \exp(a|x|), \quad \forall (t, x) \in [0, T] \times \mathbb{R}^d \quad (24)$$

for some constant $A' > 0$.

The following result is known as *Feynman-Kac formula* and can be deduced from (24) using exactly the same method as for [11, Th.6.5.3] and using the fact that, under our assumptions, X_t has finite exponential moments [11, Th.6.4.5].

Theorem 3.1 *Under the previous assumptions, the solution of the Cauchy problem (23) is given by*

$$\begin{aligned} u(t, x) = & \mathbb{E} \left[g(X_T) \exp \left(\int_t^T c(s, X_s) ds \right) \mid X_t = x \right] \\ & - \mathbb{E} \left[\int_t^T f(s, X_s) \exp \left(\int_t^s c(\alpha, X_\alpha) d\alpha \right) ds \mid X_t = x \right]. \end{aligned} \quad (25)$$

Let us mention that this result can be extended to parabolic linear PDEs on bounded domains [11, Th.6.5.2] and to elliptic linear PDEs on bounded domains [11, Th.6.5.1].

Example 1: European options

The Feynman-Kac formula has many applications in finance. Let us consider the case of a European option on a one-dimensional Markov asset $(S_t, t \geq 0)$ with

payoff $g(S_u, 0 \leq u \leq T)$. The free arbitrage value at time t of this option is

$$V_t = \mathbb{E}[e^{-r(T-t)}g(S_u, t \leq u \leq T) \mid \mathcal{F}_t].$$

By the Markov property (1), this quantity only depends on S_t and t [11, Th.2.1.2]. The Feynman-Kac formula (25) allows one to characterize V in the case where g depends only on S_T and S is a Feller diffusion.

Most often, the asset SDE

$$dS_t = S_t(\mu(t, S_t)dt + \sigma(t, S_t)dB_t)$$

cannot satisfy the uniform ellipticity assumption (22) in the neighborhood of 0. Therefore, Theorem 3.1 does not apply directly. This is a general difficulty for financial models. However, in most cases (and in all the examples below), it can be overcome by taking the logarithm of the asset price. In our case, we assume that the process $(\log S_t, 0 \leq t \leq T)$ is a Feller diffusion on \mathbb{R} with time inhomogeneous generator

$$L_t\phi(y) = \frac{1}{2}a(t, y)\phi''(y) + b(t, y)\phi'(y)$$

that satisfy the assumptions of Theorem 3.1. This holds for example for the Black-Scholes model (14). This assumption implies in particular that S is a Feller diffusion on $(0, +\infty)$ whose generator takes the form

$$\tilde{L}_t\phi(x) = \frac{1}{2}\tilde{a}(t, x)x^2\phi''(x) + \tilde{b}(t, x)x\phi'(x)$$

where $\tilde{a}(t, x) = a(t, \log x)$ and $\tilde{b}(t, x) = b(t, \log x) + a(t, \log x)/2$.

Assume also that $g(x)$ is continuous on \mathbb{R}_+ with polynomial growth when $x \rightarrow +\infty$. Then, by Theorem 3.1, the function

$$v(t, y) = \mathbb{E} \left[e^{-r(T-t)}g(S_T) \mid \log S_t = y \right]$$

is solution to the Cauchy problem

$$\begin{cases} \frac{\partial v}{\partial t}(t, y) + L_tv(t, y) - rv(t, y) = 0, & (t, y) \in [0, T) \times \mathbb{R} \\ v(T, y) = g(\exp(y)), & y \in \mathbb{R}. \end{cases}$$

Making the change of variable $x = \exp(y)$, $u(t, x) = v(t, \log x)$ is solution to

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + \tilde{b}(t, x)x\frac{\partial u}{\partial x}(t, x) + \frac{1}{2}\tilde{a}(t, x)x^2\frac{\partial^2 u}{\partial x^2}(t, x) - rv(t, x) = 0, & (t, x) \in [0, T] \times (0, +\infty) \\ u(T, x) = g(x), & x \in (0, +\infty) \end{cases} \quad (26)$$

and $V_t = u(t, S_t)$. The Black-Scholes PDE (2) is a particular case of this result.

Example 2: an Asian option

We give an example of a path-dependent option for which the uniform ellipticity condition of the matrix a does not hold. An Asian option is an option where the payoff is determined by the average of the underlying price over the period considered. Consider the Asian call option

$$\left(\frac{1}{T} \int_0^T S_u du - K \right)^+$$

on a Black-Scholes asset $(S_t, t \geq 0)$ following

$$dS_t = rS_t dt + \sigma S_t dB_t$$

where B is a standard one-dimensional Brownian motion. The free arbitrage price at time t is

$$\mathbb{E} \left[e^{-r(T-t)} \left(\frac{1}{T} \int_0^T S_u du - K \right)^+ \mid S_t \right].$$

In order to apply the Feynman-Kac formula, one must express this quantity as the (conditional) expectation of the value at time T of some Markov quantity. This can be done by introducing the process

$$A_t = \int_0^t S_u du, \quad 0 \leq t \leq T.$$

It is straightforward to check that (S, A) is a Feller diffusion on $(0, +\infty)^2$ with infinitesimal generator

$$Lf(x, y) = rx \frac{\partial f}{\partial x}(x, y) + \frac{\sigma^2}{2} x^2 \frac{\partial^2 f}{\partial x^2}(x, y) + \frac{1}{T} x \frac{\partial f}{\partial y}(x, y).$$

Although considering the change of variable $(\log S, A)$, Theorem 3.1 does not apply to this process because the infinitesimal generator is degenerate (without second-order derivative in y). Formally, the Feynman-Kac formula would give that

$$u(t, x, y) := \mathbb{E}[e^{-r(T-t)}(A_T/T - K)^+ \mid S_t = x, A_t = y]$$

is solution to the PDE

$$\begin{cases} \frac{\partial u}{\partial t} + \frac{\sigma^2}{2} x^2 \frac{\partial^2 u}{\partial x^2} + rx \frac{\partial u}{\partial x} + \frac{1}{T} x \frac{\partial u}{\partial y} - ru = 0, & (t, x, y) \in [0, T) \times (0, +\infty) \times \mathbb{R} \\ u(T, x, y) = (y/T - K)^+, & (x, y) \in (0, +\infty) \times \mathbb{R}. \end{cases} \quad (27)$$

Actually, it is possible to justify the previous statement in the specific case of a one-dimensional Black-Scholes asset: u can be written as

$$u(t, x, y) = e^{-r(T-t)} x \varphi\left(t, \frac{KT - y}{x}\right)$$

(see [22]) where $\varphi(t, z)$ is the solution of the one-dimensional parabolic PDE

$$\begin{cases} \frac{\partial \varphi}{\partial t}(t, z) + \frac{\sigma^2}{2} z^2 \frac{\partial^2 \varphi}{\partial z^2}(t, z) - \left(\frac{1}{T} + rz\right) \frac{\partial \varphi}{\partial z}(t, z) + r\varphi(t, z) = 0, & (t, z) \in [0, T) \times \mathbb{R} \\ \varphi(T, z) = -(z)^+/T, & z \in \mathbb{R}. \end{cases}$$

From this, it is easy to check that u solves (27).

Note that this relies heavily on the fact the underlying asset follows the Black-Scholes model. As far as we know no rigorous justification of Feynman-Kac formula is available for Asian options on more general assets.

3.3 Quasi- and semi-linear PDEs and BSDEs

The link between quasi- and semi-linear PDEs and BSDEs is motivated by the following formal argument. Consider the semi-linear PDE

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + L_t u(t, x) = f(u(t, x)) & (t, x) \in (0, T) \times \mathbb{R} \\ u(T, x) = g(x) & x \in \mathbb{R} \end{cases} \quad (28)$$

where (L_t) is the family of infinitesimal generators of a time-inhomogeneous Feller diffusion $(X_t, t \geq 0)$. Assume that this PDE admits a classical solution $u(t, x)$. Assume also that we can find a unique adapted process $(Y_t, 0 \leq t \leq T)$ such that

$$Y_t = \mathbb{E}[g(X_T) - \int_t^T f(Y_s)ds \mid \mathcal{F}_t], \quad \forall t \in [0, T]. \quad (29)$$

Now, by Itô's formula applied to $u(t, X_t)$,

$$u(t, X_t) = \mathbb{E}[g(X_T) - \int_t^T f(u(s, X_s))ds \mid \mathcal{F}_t].$$

Therefore, $Y_t = u(t, X_t)$ and the stochastic process Y provides a probabilistic interpretation of the solution of the PDE (28). Now, by the martingale decomposition theorem, if Y satisfies (29), there exists an adapted process $(Z_t, 0 \leq t \leq T)$ such that

$$Y_t = g(X_T) - \int_t^T f(Y_s)ds - \int_t^T Z_s dB_s, \quad \forall t \in [0, T]$$

where B is the same Brownian motion as the one driving the Feller diffusion X . In other words, Y is solution of the SDE $dY_t = f(Y_t)dt + Z_t dB_t$ with *terminal* condition $Y_T = g(X_T)$.

The following definition of a BSDE generalizes the previous situation. Given functions $b_i(t, x)$ and $\sigma_{ij}(t, x)$ that are globally Lipschitz in x and locally bounded ($1 \leq i, j \leq d$) and a standard d -dimensional Brownian motion B , consider the unique solution X of the time-inhomogeneous SDE

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t \quad (30)$$

with initial condition $X_0 = x$. Consider also two functions $f : [0, T] \times \mathbb{R}^d \times \mathbb{R}^k \times \mathbb{R}^{k \times d} \rightarrow \mathbb{R}^k$ and $g : \mathbb{R}^d \rightarrow \mathbb{R}^k$. We say that $((Y_t, Z_t), t \geq 0)$ solve the BSDE

$$dY_t = f(t, X_t, Y_t, Z_t)dt + Z_t dB_t$$

with terminal condition $g(X_T)$ if Y and Z are progressively measurable processes with respect to the Brownian filtration $\mathcal{F}_t = \sigma(B_s, s \leq t)$ such that, for any $0 \leq t \leq T$,

$$Y_t = g(X_T) - \int_t^T f(s, X_s, Y_s, Z_s)ds - \int_t^T Z_s dB_s. \quad (31)$$

The example of Section 2.2 corresponds to $g(x) = (x - K)^+$, $f(t, x, y, z) = -ry + z(\mu - r)/\sigma$ and $Z_t = \sigma\pi_t$. Note that the role of the implicit unknown process Z is to make Y adapted.

The existence and uniqueness of (Y, Z) solving (31) hold under the assumptions that $g(x)$ is continuous with (at most) polynomial growth in x , $f(t, x, y, z)$ is continuous with polynomial growth in x and linear growth in y and z , and f is uniformly Lipschitz in y and z . Let us denote by (A) all these assumptions. We refer to [19] for the proof of this result and the general theory of BSDEs.

Consider the quasi-linear parabolic PDE

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + L_t u(t, x) = f(t, x, u(t, x), \nabla_x u(t, x)\sigma(t, x)), & (t, x) \in (0, T) \times \mathbb{R}^d \\ u(T, x) = g(x), & x \in \mathbb{R}^d. \end{cases} \quad (32)$$

The following results give the links between the BSDE (31) and the PDE (32).

Theorem 3.2 ([17], Th.4.1) *Assume that $b(t, x)$, $\sigma(t, x)$, $f(t, x, y, z)$ and $g(x)$ are continuous and differentiable w.r.t. to the space variables x, y, z with uniformly bounded derivatives. Assume also that b , σ and f are uniformly bounded and that $a = \sigma\sigma'$ is uniformly elliptic. Then (32) admits a unique classical solution u and*

$$Y_t = u(t, X_t) \quad \text{and} \quad Z_t = \nabla_x u(t, X_t)\sigma(t, X_t).$$

Theorem 3.3 ([19], Th.2.4) *Assume (A) and that $b(t, x)$ and $\sigma(t, x)$ are globally Lipschitz in x and locally bounded. Define the function $u(t, x) = Y_t^{t,x}$, where $Y^{t,x}$ is the solution to the BSDE (31) on the time interval $[t, T]$ where X is solution to the SDE (30) with initial condition $X_t = x$. Then u is a viscosity solution of (32).*

Theorem 3.2 gives an interpretation of the solution of a BSDE in terms of the solution of a quasi-linear PDE. In particular, in the example of BSDE of Section 2.2, it gives the usual interpretation of the hedging strategy $\pi_t = Z_t/\sigma$ as the Δ -hedge of the option price. Note also that Theorem 3.2 implies that the process (X, Y, Z) is Markov—a fact which is not obvious from the definition.

Conversely, Theorem 3.3 shows how to construct a viscosity solution of a quasi-linear PDE from BSDEs.

BSDEs provide an indirect tool to compute quantities related to a solution X of a SDE (such as the hedging price and strategy of an option based on the process X). BSDEs also have links with general stochastic control problems, that we will not mention. Here we give an example of application to the pricing of an American put option.

Example: pricing of an American put option

Consider a Black-Scholes underlying asset S and assume for simplicity that the risk-free interest rate r is zero. The price of an American put option on S with strike K and maximal exercise policy T is given by

$$\sup_{0 \leq \tau \leq T} \mathbb{E}^*[(K - S_\tau)^+]$$

where τ is a stopping time and where \mathbb{P}^* is the risk-neutral probability measure, under which the process S is simply a Black-Scholes asset with zero drift.

In the case of a European put option, the price is given by the solution of the BSDE

$$Y_t = (K - S_T)^+ - \int_t^T Z_s dB_s, \quad (33)$$

by a similar argument of Section 2.2. In the case of an American put option, the price at time t is necessarily bigger than $(K - S_t)^+$. It is therefore natural to include this condition by considering the BSDE (33) reflected on the obstacle $(K - S_t)^+$. Mathematically, this corresponds to the problem of finding adapted processes Y , Z and R such that

$$\begin{cases} Y_t = (K - S_T)^+ - \int_t^T Z_s dB_s + R_T - R_t \\ Y_t \geq (K - S_t)^+ \\ R \text{ is continuous, increasing, } R_0 = 0 \text{ and } \int_0^T [Y_t - (K - S_t)^+] dR_t = 0. \end{cases} \quad (34)$$

The process R increases only when $Y_t = (K - S_t)^+$ in such a way that Y cannot cross this obstacle. The existence of a solution of this problem is a particular case of general results (see [9]). As a consequence of the following theorem, this reflected BSDE gives a way to compute the price of the American put option.

Theorem 3.4 ([9], Th.7.2) *The American put option has the price Y_0 , where (Y, Z, R) solves the reflected BSDE (34).*

The essential argument of the proof is the following. Fix $t \in [0, T]$ and a stopping time $\tau \in [t, T]$. Since

$$Y_\tau - Y_t = R_t - R_\tau + \int_t^\tau Z_s dB_s$$

and since R is increasing, $Y_t = \mathbb{E}^*[Y_\tau + R_\tau - R_t \mid \mathcal{F}_t] \geq \mathbb{E}^*[(K - S_\tau)^+ \mid \mathcal{F}_t]$. Conversely, if $\tau_t^* = \inf\{u \in [t, T] : Y_u = (K - S_u)^+\}$ (with the convention $\inf \emptyset = T$), since $Y > (K - S)^+$ on $[t, \tau_t^*)$, R is constant on this interval and

$$Y_t = \mathbb{E}^*[Y_{\tau_t^*} + R_{\tau_t^*} - R_t \mid \mathcal{F}_t] = \mathbb{E}^*[(K - S_{\tau_t^*})^+].$$

Therefore,

$$Y_t = \operatorname{ess\,sup}_{t \leq \tau \leq T} \mathbb{E}^*[(K - S_\tau)^+ \mid \mathcal{F}_t] \tag{35}$$

which gives another interpretation for the solution Y of the reflected BSDE. Applying this for $t = 0$ yields $Y_0 = \sup_{\tau \leq T} \mathbb{E}^*[(K - S_\tau)^+]$ as stated.

Moreover, as shown by the previous computation, the process Y provides an interpretation of the optimal exercise policy as the first time where Y hits the obstacle $(K - S)^+$. This fact is actually natural from (35): the optimal exercise policy is the first time where the current payoff equals the maximal future expected payoff.

As it will appear in the next section, as the solution of an optimal stopping problem, if $S_0 = x$, the price of this American put option is $u(0, x)$ where u is the

solution of the nonlinear PDE

$$\begin{cases} \min \left\{ u(t, x) - (K - x)^+; -\frac{\partial u}{\partial t}(t, x) - \frac{\sigma^2}{2} x^2 \frac{\partial^2 u}{\partial x^2}(t, x) \right\} = 0, & (t, x) \in (0, T) \times (0, +\infty) \\ u(T, x) = (K - x)^+, & x \in (0, +\infty). \end{cases} \quad (36)$$

Therefore, similarly as in Theorem 3.3, the reflected BSDE (34) provides a probabilistic interpretation of the solution of this PDE.

The (formal) essential argument of the proof of this result can be summarized as follows (for details, see [16, Sec.V.3.1]). Consider the solution u of (36) and apply Itô's formula to $u(t, S_t)$. Then, for any stopping time $\tau \in [0, T]$,

$$u(0, x) = \mathbb{E}[u(\tau, S_\tau)] - \mathbb{E} \left[\int_0^\tau \left(\frac{\partial u}{\partial t}(t, S_t) + \frac{\sigma^2}{2} S_t^2 \frac{\partial^2 u}{\partial x^2}(t, S_t) \right) ds \right].$$

Since u is solution of (36), $u(0, x) \geq \mathbb{E}[u(\tau, S_\tau)] \geq \mathbb{E}[(K - S_\tau)^+]$. Hence, $u(0, x) \geq \sup_{0 \leq \tau \leq T} \mathbb{E}[(K - S_\tau)^+]$.

Conversely, if $\tau^* = \inf\{0 \leq t \leq T : u(t, S_t) = (K - S_t)^+\}$, then

$$\frac{\partial u}{\partial t}(t, S_t) + \frac{\sigma^2}{2} S_t^2 \frac{\partial^2 u}{\partial x^2}(t, S_t) = 0 \quad \forall t \in [0, \tau^*]$$

and $u(\tau^*, S_{\tau^*}) = (K - S_{\tau^*})^+$. Therefore, for $\tau = \tau^*$ all the inequalities in the previous computation are equalities and $u(0, x) = \sup_{0 \leq \tau \leq T} \mathbb{E}[(K - S_\tau)^+]$.

3.4 Optimal control, Hamilton-Jacobi-Bellman equations and variational inequalities

We discuss only two main families of optimal control problems: the problems with finite horizon and the optimal stopping problems. Other classes of optimal problems appearing in finance are mentioned in the end of this section.

Finite horizon problems

The study of optimal control problems with finite horizon is motivated, for example, by the questions of portfolio management, quadratic hedging of options or superhedging cost for uncertain volatility models.

Let us consider a controlled diffusion X^α in \mathbb{R}^d solution to the SDE

$$dX_t^\alpha = b(X_t^\alpha, \alpha_t)dt + \sigma(X_t^\alpha)dB_t, \quad (37)$$

where B is a standard r -dimensional Brownian motion and the control α is a given progressively measurable process taking values in some compact metric space A . Such a control is called *admissible*. For simplicity, we consider the time-homogeneous case and we assume that the control does not act on the diffusion coefficient σ of the SDE. Assume that $b(x, a)$ is bounded, continuous, and Lipschitz in the variable x , uniformly in $a \in A$. Assume also that σ is Lipschitz and bounded. For any $a \in A$, we introduce the linear differential operator

$$L^a \varphi = \frac{1}{2} \sum_{i,j=1}^d \left(\sum_{k=1}^d \sigma_{ik}(x) \sigma_{jk}(x) \right) \frac{\partial^2 \varphi}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(x, a) \frac{\partial \varphi}{\partial x_i},$$

which is the infinitesimal generator of X^α when α is constant equal to $a \in A$.

A typical form of finite horizon optimal control problems in finance consists in computing

$$u(t, x) = \inf_{\alpha \text{ admissible}} \mathbb{E} \left[e^{-rT} g(X_T^\alpha) + \int_t^T e^{-rt} f(X_t^\alpha, \alpha_t) dt \mid X_t^\alpha = x \right], \quad (38)$$

where f and g are continuous and bounded functions, and to find an optimal control α^* that realizes the minimum. Moreover, it is desirable to find a *Markov* optimal control, i.e. an optimal control having the form $\alpha_t^* = \psi(t, X_t)$. Indeed, in this case, the controlled diffusion X^{α^*} is a Markov process.

In the case of non-degenerate diffusion coefficient, we have the following link between the optimal control problems and a semilinear PDEs.

Theorem 3.5 *Under the additional assumption that σ is uniformly elliptic, u is the unique bounded classical solution of the Hamilton-Jacobi-Bellman (HJB) equation*

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + \inf_{a \in A} \{L^a u(t, x) + f(x, a)\} - ru(t, x) = 0, & (t, x) \in (0, T) \times \mathbb{R}^d \\ u(T, x) = g(x), & x \in \mathbb{R}^d. \end{cases} \quad (39)$$

Furthermore, a Markov control $\alpha_t^* = \psi(t, X_t)$ is optimal for a fixed initial condition x and initial time $t = 0$ if and only if

$$L^{\psi(t,x)}u(t, x) + f(x, \psi(t, x)) = \inf_{a \in A} \{L^a u(t, x) + f(x, a)\}$$

for almost every $(t, x) \in [0, T] \times \mathbb{R}^d$.

This is Theorem III.2.3 of [4] restricted to the case of precise controls (see below).

Here again, the essential argument of the proof can be easily (at least formally) written: Consider any admissible control α and the corresponding controlled diffusion X^α with initial condition $X_0 = x$. By Itô's formula applied to $e^{-rt}v(t, X_t^\alpha)$ where v is the solution of (39),

$$\mathbb{E}[e^{-rT}v(T, X_T^\alpha)] = v(0, x) + \int_0^T e^{-rt} \mathbb{E} \left[\frac{\partial v}{\partial t}(t, X_t^\alpha) + L^{\alpha_t}v(t, X_t^\alpha) + rv(t, X_t^\alpha) \right] ds.$$

Therefore, by (39),

$$v(0, x) \leq \mathbb{E} \left[e^{-rT}g(X_T^\alpha) + \int_t^T e^{-rt}f(X_t^\alpha, \alpha_t)dt \mid X_t^\alpha = x \right]$$

for any admissible control α . Now, for the Markov control α^* defined in Theorem 3.5, all the inequalities in the previous computation are equalities. Hence $v = u$.

The cases where σ is not uniformly elliptic or where σ also depends of the current control α_t are much more difficult. In both cases, it is necessary to enlarge the set of admissible control by considering *relaxed controls*, i.e. controls that belong to the set $\mathcal{P}(A)$ of probability measures on A . For such a control α , the terms $b(x, \alpha_t)$ and $f(x, \alpha_t)$ in (37) and (38) are replaced by $\int b(x, a)\alpha_t(da)$ and $\int f(x, a)\alpha_t(da)$, respectively. The admissible controls of the original problem correspond to relaxed controls that are Dirac masses at each time. These are called *precise controls*.

The value \tilde{u} of this new problem is defined as in (38), but the infimum is taken over all progressively measurable processes α taking values in $\mathcal{P}(A)$. It is possible to prove under general assumptions that both problems give the same value: $\tilde{u} = u$ (cf. [4, Cor.I.2.1] or [8, Th.2.3]).

In these cases, one usually cannot prove the existence of a classical solution of (39). The weaker notion of viscosity solution is generally the correct one. In all cases treated in the literature, $u = \tilde{u}$ solves the same HJB equation as in Theorem 3.5, except that the infimum is taken over $\mathcal{P}(A)$ instead of A (cf. [4, Th.IV.2.2] for the case without control on σ). Contrary to the case of Theorem 3.5, it is not trivial at all in the general case to obtain a result on precise controls from the result on relaxed controls. This is due to the fact that usually no result is available on the existence and the characterization of a Markov relaxed optimal control. The only examples where it has been done require restrictive assumptions (cf. [8, Cor.6.8]). However, in most of the financial applications, the value function u is the most useful information. In practice, one usually only needs to compute a control that give an expected value arbitrarily close to the optimal one.

Optimal stopping problems

Optimal stopping problems arise in finance for example for the American options pricing (when to sell a claim, an asset?) or in production models (when to extract or product a good? when to stop production?).

Let us consider a Feller diffusion X in \mathbb{R}^d solution to the SDE

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t, \quad (40)$$

where B is a standard d -dimensional Brownian motion. As in (13), let $(L_t)_{t \geq 0}$ denote its family of time-inhomogeneous infinitesimal generators. Denote by $\Pi(t, T)$ the set of stopping times valued in $[t, T]$.

A typical form of optimal stopping problems consists in computing

$$u(t, x) = \inf_{\tau \in \Pi(t, T)} \mathbb{E} \left[e^{-r(\tau-t)} g(\tau, X_\tau) + \int_t^\tau e^{-r(s-t)} f(s, X_s) ds \mid X_t = x \right], \quad (41)$$

and to characterize an optimal stopping time.

Assume that $b(t, x)$ is bounded and continuously differentiable with bounded derivatives and that $\sigma(t, x)$ is bounded, continuously differentiable w.r.t. t and

twice continuously differentiable w.r.t. x with bounded derivatives. Assume also that σ is uniformly elliptic. Finally, assume that $g(t, x)$ is differentiable w.r.t. t and twice differentiable w.r.t. x and that

$$|f(t, x)| + \left| \frac{\partial g}{\partial t}(t, x) \right| + \sum_{i=1}^d \left| \frac{\partial g}{\partial x_i}(t, x) \right| \leq C e^{\mu|x|}$$

for positive constants C and μ .

Theorem 3.6 ([3], Sec.III.4.9) *Under the previous assumptions, $u(t, x)$ admits first-order derivatives w.r.t. t and second-order derivatives w.r.t. x that are L^p for all $1 \leq p < \infty$. Moreover, u is solution of the variational inequality*

$$\begin{cases} \max \{u(t, x) - g(t, x); -\frac{\partial u}{\partial t}(t, x) - L_t u(t, x) + ru(t, x) - f(t, x)\} = 0, & (t, x) \in (0, T) \times \mathbb{R}^d \\ u(T, x) = g(T, x) & x \in \mathbb{R}^d. \end{cases}$$

The proof of this result is based on a similar (formal) justification as the one we gave for (36). We refer to [13] for a similar result under weaker assumptions more suited to financial models when $f = 0$ (this is in particular the case for American options).

In some cases (typically with $f = 0$, see [12]), it can be shown that the infimum in (41) is attained for the stopping time

$$\tau^* = \inf\{t \leq s \leq T : u(s, X_s^{t,x}) = g(s, X_s^{t,x})\}$$

where $X^{t,x}$ is the solution of the SDE (40) with initial condition $X_t^{t,x} = x$.

Generalizations and extensions

An optimal control problem can also be solved through the optimisation of a family of BSDEs related to the laws of the controlled diffusions. On this question, we refer to [21].

In the present section, we only considered very specific optimal control problems. Other important families of optimal control problems are given by impulse

control problems, where the control may induce a jump of the underlying stochastic process, or ergodic control problems, where the goal is to optimize a quantity related to the stationary behavior of the controlled diffusion. Impulse control has applications for example in stock or resource management problems. In the finite horizon case, when the underlying asset follow a model with stochastic or elastic volatility, or when the market is incomplete, other optimal control problems can be considered, such as determining the superhedging cost, or minimizing some risk measure. Various constraints can be included in the optimal control problem, such as maximizing the expectation of an utility with the constraint that this utility has a fixed volatility, or minimizing the volatility for a fixed expected utility. One can also impose Gamma constraints on the control. Another important extension of optimal control problems arises when one wants to solve numerically an HJB equation. Usual discretization methods require to restrict to a bounded domain and to fix artificial boundary conditions. The numerical solution can be interpreted as the solution of an optimal control problem in a bounded domain. In this situation, a crucial question is to quantify the impact on the discretized solution of an error on the artificial boundary condition (which usually cannot be computed exactly).

4 Conclusion

Rewriting a stochastic financial problem in terms of a PDE is a very useful tool for the analysis and for numerical resolution of the problem. Let us conclude with some comments on the limitations of this approach and on the use of probabilistic methods for the numerical resolution of PDEs.

Limitations of the approach

As explained in this article, the stochastic approach to PDEs usually allows one to prove the existence and give an interpretation of a solution. However, for

uniqueness, one must generally use analytic approaches.

The theory of PDEs and the stochastic calculus approach to PDEs have been mostly developed for the equations of physics. However, the equations of finance usually have some specificity, so that they usually do not enter in many of the pre-existing developments of the theory. In particular, in physical models, problems are often posed in bounded domains where the diffusion parameter σ is constant (or piecewise constant) and the difficulties are mainly in nonlinearities or discontinuities. Conversely, in financial models, the problems are *a priori* posed in unbounded domains, or with non-homogeneous boundary conditions, and there are often degeneracy problems for the σ .

We can give some examples where the existing PDE theory basically does not apply at all. This is the case of models with elastic volatility ($\sigma(x) = x^\alpha$, $\alpha < 1$), such as SABR model, Heston model or interest rate models with a CIR model for the volatility. In these situations, the stochastic approach is the only existing one.

About numerical methods

The Feynman-Kac formula for linear PDEs allows one to use Monte Carlo methods to compute the solution of the PDE. They are especially useful when the solution of the PDE has to be computed at a small number of points, or when dimension is large (typically larger or equal to 4), since they provide a rate of convergence which grows linearly with the dimension.

Concerning quasi- or semi-linear PDEs and some optimal control problems (see the example of American put options in Section 3.3), interpretations in terms of BSDEs provide indirect Monte Carlo methods of numerical computation (see [2] for Monte Carlo methods for American options or [5, 7] for general BSDEs schemes). These methods have the advantage that they do not require to consider artificial boundary conditions. However, their speed of convergence to the exact solution is still largely unknown, and could depend on the dimension of the problem.

For high dimensional HJB equations, the analytical discretization methods

lead to important numerical problems. First, these methods need to solve an optimization problem at each node of the discretization grid, which can be very costly in high dimension or difficult depending on the particular constraints imposed on the control. Moreover, these methods require to localize the problem, i.e. to solve the problem in a bounded domain with artificial boundary conditions, which are usually difficult to compute precisely. This localization problem can be solved by computing the artificial boundary condition with a Monte Carlo method based on BSDEs. However, the error analysis of this method is based on the probabilistic interpretation of HJB equations in bounded domains, which is a difficult problem in general.

References

- [1] Y. Aït-Sahalia and J. Jacob. Volatility estimators for discretely sampled Lévy processes. *Ann. Statist.*, 35(1):355–392, 2007.
- [2] V. Bally and G. Pagès. Error analysis of the optimal quantization algorithm for obstacle problems. *Stochastic Process. Appl.*, 106(1):1–40, 2003.
- [3] A. Bensoussan and J.-L. Lions. *Applications of variational inequalities in stochastic control*, volume 12 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, 1982. Translated from the French.
- [4] V. S. Borkar. *Optimal control of diffusion processes*, volume 203 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow, 1989.
- [5] B. Bouchard and N. Touzi. Discrete-time approximation and Monte-Carlo simulation of backward stochastic differential equations. *Stochastic Process. Appl.*, 111(2):175–206, 2004.

- [6] E. Çinlar and J. Jacod. Representation of semimartingale Markov processes in terms of Wiener processes and Poisson random measures. In *Seminar on Stochastic Processes, 1981 (Evanston, Ill., 1981)*, volume 1 of *Progr. Prob. Statist.*, pages 159–242. Birkhäuser Boston, Mass., 1981.
- [7] F. Delarue and S. Menozzi. A forward-backward stochastic algorithm for quasi-linear PDEs. *Ann. Appl. Probab.*, 16(1):140–184, 2006.
- [8] N. El Karoui, D. Hüü Nguyen, and M. Jeanblanc-Picqué. Compactification methods in the control of degenerate diffusions: existence of an optimal control. *Stochastics*, 20(3):169–219, 1987.
- [9] N. El Karoui, C. Kapoudjian, E. Pardoux, S. Peng, and M. C. Quenez. Reflected solutions of backward SDE’s, and related obstacle problems for PDE’s. *Ann. Probab.*, 25(2):702–737, 1997.
- [10] S. N. Ethier and T. G. Kurtz. *Markov processes*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1986. Characterization and convergence.
- [11] A. Friedman. *Stochastic differential equations and applications. Vol. 1*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975. Probability and Mathematical Statistics, Vol. 28.
- [12] S. D. Jacka. Local times, optimal stopping and semimartingales. *Ann. Probab.*, 21(1):329–339, 1993.
- [13] P. Jaillet, D. Lamberton, and B. Lapeyre. Variational inequalities and the pricing of American options. *Acta Appl. Math.*, 21(3):263–289, 1990.
- [14] A. Janicki. *Numerical and statistical approximation of stochastic differential equations with non-gaussian measures*, volume 1 of *A Series of Monographs*. H. Steinhaus Center for Stochastic Methods, Technical University, Wrocław, Poland, 1996.

- [15] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1988.
- [16] D. Lamberton and B. Lapeyre. *Introduction to stochastic calculus applied to finance*. Chapman & Hall, London, 1996. Translated from the 1991 French original by Nicolas Rabeau and François Mantion.
- [17] J. Ma, P. Protter, and J. M. Yong. Solving forward-backward stochastic differential equations explicitly—a four step scheme. *Probab. Theory Related Fields*, 98(3):339–359, 1994.
- [18] B. Øksendal. *Stochastic differential equations*. Universitext. Springer-Verlag, Berlin, sixth edition, 2003. An introduction with applications.
- [19] E. Pardoux. Backward stochastic differential equations and viscosity solutions of systems of semilinear parabolic and elliptic pdes of second order. In B. Ø. L. Decreusefond, J. Gjerde and A. Üstünel, editors, *Stochastic Analysis and Related Topics : The Geilo Workshop*, pages 79–127. Birkhäuser, 1998.
- [20] P. Protter. A partial introduction to financial asset pricing theory. *Stochastic Process. Appl.*, 91(2):169–203, 2001.
- [21] M. C. Quenez. Stochastic control and BSDEs. In *Backward stochastic differential equations (Paris, 1995–1996)*, volume 364 of *Pitman Res. Notes Math. Ser.*, pages 83–99. Longman, Harlow, 1997.
- [22] L. C. G. Rogers and Z. Shi. The value of an Asian option. *J. Appl. Probab.*, 32(4):1077–1088, 1995.
- [23] L. C. G. Rogers and D. Williams. *Diffusions, Markov processes, and martingales. Vol. 1*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Ltd., Chichester, second edition, 1994. Foundations.

- [24] D. Talay and Z. Zheng. Quantiles of the Euler scheme for diffusion processes and financial applications. *Math. Finance*, 13(1):187–199, 2003. Conference on Applications of Malliavin Calculus in Finance (Rocquencourt, 2001).