

# Interacting Multi-Class Transmissions in Large Stochastic Networks

Carl Graham, Philippe Robert

► **To cite this version:**

Carl Graham, Philippe Robert. Interacting Multi-Class Transmissions in Large Stochastic Networks. 2008. <inria-00326156>

**HAL Id: inria-00326156**

**<https://hal.inria.fr/inria-00326156>**

Submitted on 2 Oct 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INTERACTING MULTI-CLASS TRANSMISSIONS IN LARGE STOCHASTIC NETWORKS

CARL GRAHAM AND PHILIPPE ROBERT

ABSTRACT. The mean-field limit of a Markovian model describing the interaction of several classes of permanent connections in a network is analyzed. In the same way as for the TCP algorithm, each of the connections has a self-adaptive behavior in that its transmission rate along its route depends on the level of congestion of the nodes of the route. Since several classes of connections going through the nodes of the network are considered, an original mean-field result in a multi-class context is established. It is shown that, as the number of connections goes to infinity, the behavior of the different classes of connections can be represented by the solution of an unusual non-linear stochastic differential equation depending not only on the sample paths of the process, but also on its *distribution*. Existence and uniqueness results for the solutions of these equations are derived. Properties of their invariant distributions are investigated and it is shown that, under some natural assumptions, they are determined by the solutions of a fixed point equation in a finite dimensional space.

---

## CONTENTS

|   |    |
|---|----|
| 1. Introduction                                     | 1  |
| 2. A Mathematical Model of a Single Connection      | 4  |
| 3. The Stochastic Model of the Network              | 6  |
| 4. Analysis of the Non-Linear Limiting Process      | 8  |
| 5. The Mean-Field Limit for Converging Initial Data | 14 |
| 6. Fixed Points and Invariant Distributions         | 19 |
| References  | 22 |

---

## 1. INTRODUCTION

Mathematical modeling of data transmission in communication networks has been the subject of an intense activity for some time now. For data transmission, the Internet network can be described as a very large distributed system with self-adaptive capabilities to the different congestion events that regularly occur at its numerous nodes. Various approaches have been used in this respect: control theory, ordinary differential equations, Markov processes, optimization techniques, . . . For data transmission one can distinguish two levels of mathematical abstraction:

(1) *Packet Level*.

For this approach, the self-adaptive behavior of the transmission control protocol (TCP) is analyzed. Packets are sent as long as no loss is detected

---

*Date:* October 2, 2008.

*Key words and phrases.* Multi-Class Systems; Mean-Field Interaction; Coupling; Chaoticity; Non-Linear Stochastic Differential Equations; TCP Algorithm; Equilibrium Distributions.

and throughput grows linearly during that time. On the contrary when a loss occurs, the throughput is sharply reduced by a multiplication by some factor  $r < 1$  (the constant  $1/2$  in general). This scheme is known as an Additive Increase and Multiplicative Decrease algorithm (AIMD).

Ott *et al.* [22] is one of the early mathematical studies. In this setting, after a scaling with respect to the loss rate, Dumas *et al.* [8] proves various limit theorems for the resulting processes. The asymptotic evolution of the throughput of a connection can then be described as a Markov process whose infinitesimal generator is given by

$$\Omega(f)(x) = af'(x) + bx(f(rx) - f(x))$$

for  $f$  a  $C^1$ -function from  $\mathbb{R}_+$  to  $\mathbb{R}$ , where  $a, b > 0$  and  $r \in (0, 1)$ . The constant  $a$  is the rate of linear increase and  $r$  is the multiplicative factor. A precise formulation of this procedure is detailed in Section 2. See Dumas *et al.* [8] for a survey of the literature in this domain.

(2) *Flow Level.*

It consists in looking at the state of the network on a quite long time scale so that the self-adaptive procedures of data transmission in the Internet can be described as a variation of a processor-sharing discipline at the level of the network. A connection is seen as a fluid characterized by the throughput it achieves given the other competing flows. A possible mathematical formulation is via an optimization problem: given  $J$  classes of connections, when there  $x_j$  connections of class  $j \in \{1, \dots, J\}$ , the total throughput achieved by the connections of class  $j$  is given by  $\lambda_j$  so that the vector  $(\lambda_j)$  is a solution of

$$\max_{\lambda \in \Lambda} \sum_{j=1}^J x_j U_j(\lambda/x_j),$$

where  $\Lambda$  is the set of admissible throughputs which takes into account the capacity constraints of the network. The functions  $(U_j)$  are defined as *utility* functions, various expressions have been proposed for them. See Kelly *et al.* [17], Massoulié [19] and Massoulié and Roberts [20]. Kelly and Williams [18] gives a heavy-traffic analysis of a network in such a context.

In this paper, the packet level approach will be used to analyze the coexistence of numerous connections in a network with a general number of nodes. Up to now the stochastic models analyzed rigorously have considered a *single* node carrying *similar* flows. In this context, some approximated mean field analyses have been carried out for a single buffer receiving packets of a large number of connections, see Adjih *et al.* [1] and Baccelli *et al.* [3] for example. In particular, few rigorous, explicit, results have been obtained for models with several types of connections (relating different couples of nodes for example) and describing the evolution of the state of the whole network. In the general context of stochastic networks, mean-field approaches have been used in various situations notably by Dobrushin and his co-authors, see Karpelevitch *et al.* [16] for example.

The simultaneous transmission of several classes of permanent connections is analyzed here. A class of connections is characterized, in particular, by the set of nodes it uses and by the rate of increase of its throughput when no loss occurs. Because of the additive part of the AIMD scheme, this last quantity is directly related to the transmission time between the source and the destination of the

connection (the round trip time, RTT). The connections interact as follows: At a fixed node, the loss rate of a given connection depends on the number of packets of different classes of connections going through it. If a loss occurs at some node of the route of a connection, its throughput is multiplied by some factor less than 1 so that the transmission rate of a connection depends on the state of *all* the nodes it is using and therefore on the other connections traversing any of these nodes.

**Mean-Field Analysis of Multi-Class Systems.** The evolution of a *large* number of connections divided into homogeneous classes will be considered. As it will be seen, a Markovian multi-class mean-field analysis has to be developed to investigate the properties of these networks. In this setting, chaoticity (i.e. convergence in law to i.i.d. random variables) is a classical notion. For statistically indistinguishable objects, modeled by systems of exchangeable r.v., chaoticity is equivalent to the fact that the empirical measures satisfy a weak law of large numbers. Sznitman [27] develops compactness-uniqueness methods, as well as coupling methods between exchangeable systems and an i.i.d. system which give stronger results. See also Graham and Méléard [21] and Graham [10, 11].

Many phenomena involve dissimilar objects, called particles, classified in a finite number of types, particles of a class being similar and *numerous*, see *e.g.* Cercignani *et al.* [6] and the review papers Bellomo and Stöcker [4], Graham [11], Grunfeld [14], Struckmaier [26]. Few convergence proofs for such multi-class systems exist, and most references directly use extensions of the limit equations for systems of identical particles without further justification. Some coupling techniques, such as those in Graham and Méléard [13] and Graham [11] using representations of the past history by random graphs and trees, may apply to some multi-class systems, but the models considered here do not have the specific structure required. Graham [12] studies natural notions of multi-exchangeability and chaoticity, and one of its scopes is to enable to adapt Sznitman's compactness-uniqueness methods to multi-class systems, but these methods will not be used here.

**Outline of the Paper.** The mean-field limit of a Markovian model describing the interaction of  $K$  classes of permanent connections in a network with  $J$  nodes is analyzed. Each of the connections has a self-adaptive behavior in that its transmission rate along its route depends on the level of congestion of the nodes of the route. More formally, the state of a class  $k$  connection is given by its throughput, i.e. some non negative number  $w_k \in \mathbb{R}_+$ , and the *utilization*  $u_j$  of a node  $j \in \{1, \dots, J\}$  is given as a weighted sum of all throughputs of the connections using this node. If  $w$  is the state of a class  $k$  connection, it increases linearly with slope  $a_k(w, u)$  and jumps to  $r_k w$  at rate  $b_k(w, u)$  where  $u = (u_j)$  is the utilization vector and  $r \in (0, 1)$  is the multiplicative factor. A sketch of the stochastic evolution of the  $n$ th class  $k$  connection,  $1 \leq n \leq N_k$ , can be given by a fluid description through the ordinary differential equation

$$\frac{d}{dt} w_{k,n}(t) = a_k(w_{k,n}, u(t)) - (1 - r_k) w_{k,n}(t) b_k(w_{k,n}(t), u(t)),$$

where  $u(t) = (u_j(t))$  and, for  $1 \leq j \leq J$ ,

$$u_j(t) = \sum_{k=1}^K A_{jk} \sum_{p=1}^{N_k} w_{k,p}(t),$$

see Section 3 for a more formal description. Related models are considered by Kelly *et al.* [17].

Since several classes of connections going through the nodes of the network are considered, an original mean-field result in a multi-class context is established in the following context: for any  $1 \leq k \leq K$ , the number  $N_k$  of class  $k$  connections goes to infinity such that  $N_k/(N_1 + \dots + N_K)$  goes to  $p_k$ . Due to the lack of symmetry, an extension to multi-class systems of the notion of chaoticity and of Sznitman's coupling methods will be developed. It is shown that, in the mean-field limit, the asymptotic behavior of a "typical" class  $k$  of connection can be represented by the solution of an unusual non-linear stochastic differential equation depending not only on the sample paths of the process, but also on its *distribution*. Existence and uniqueness results for the solutions of these equations are derived. Properties of their invariant distributions are investigated. It is shown in particular that, for a quite large class of models, the invariant distributions have an explicit expression given in terms of the solution of a fixed point equation in  $\mathbb{R}_+^J$ . Several examples of networks are discussed.

Section 2 recalls the main scaling results for the case of one class of connections in a one node network and Section 3 defines the model of the networks analyzed in this paper. The properties of the limiting process associated to the behavior of a given transmission are investigated in Section 4. In Section 5, convenient notions of exchangeability and chaoticity in a multi-class system are introduced and mean-field results are obtained in this context. Section 6 is devoted to the analysis of possible invariant distributions of the limiting process.

## 2. A MATHEMATICAL MODEL OF A SINGLE CONNECTION

This section introduces the mathematical framework used to describe the evolution of the transmission rate of a single connection. In the following section, interactions of several connections within a network are presented and analyzed.

A connection between a source and a destination progressively increases its transmission rate until it receives some indication that the capacity along its path in the network is almost fully utilized. On the other hand, when congestion occurs, the throughput of the connection is drastically reduced. A given connection has a variable  $W$  which gives the maximum number of packets that can be transmitted without receiving any acknowledgment from the destination. The variable  $W$  is called the *congestion window size*. If all the  $W$  packets are successfully transmitted, then  $W$  is increased by 1 (Additive part of the AIMD), so that  $W + 1$  packets can be sent for the next round. Otherwise  $W$  is divided by some factor (Multiplicative part of the AIMD). An AIMD algorithm can thus be described as follows:

$$(2.1) \quad W \rightarrow \begin{cases} W + 1 & \text{if no loss occurs among the } W \text{ packets} \\ \lfloor rW \rfloor & \text{otherwise,} \end{cases}$$

where  $\lfloor x \rfloor$  is the integer part of  $x \in \mathbb{R}$ . The randomness for this model is given by the sequence of events when losses occur. One of the simplest and reasonable model is to consider that the loss probability of a packet is given by  $\varepsilon$  and that losses occur independently for each packet. This is the model analyzed in Ott *et al.* [22] at the fluid level and by Dumas *et al.* [8]. The questionable independence property of losses is discussed thoroughly in Guillemin [15], it is shown that this

assumption is in fact pessimistic in that it underestimates the real efficiency of the algorithm.

Under these assumptions, there is a natural Markov chain  $(W_n^\varepsilon)$  whose transitions are given by Equation (2.1). The variable  $W_n^\varepsilon$  can be interpreted as the “throughput” at the beginning of the  $(n+1)$ th cycle. For a given  $\varepsilon > 0$  little is known on the behavior of this Markov chain, on its invariant probability for example. A scaling procedure has been developed in Dumas *et al.* [8] consisting in looking at the continuous time Markov process

$$\left(\overline{W}_0^\varepsilon(t)\right) = \left(\sqrt{\varepsilon}W_{\lfloor t/\varepsilon \rfloor}^\varepsilon\right).$$

It is shown that, as  $\varepsilon$  goes to 0, the family of Markov processes  $(\overline{W}_0^\varepsilon(t), t \geq 0)$  converges in distribution to a Markov process  $(\overline{W}_0(t), t \geq 0)$  whose infinitesimal generator  $\Omega_0$  is given by

$$(2.2) \quad \Omega_0(f)(x) = f'(x) + x(f(rx) - f(x)),$$

for  $f$  a  $C^1$ -function from  $\mathbb{R}_+$  to  $\mathbb{R}$ . It is also proved that, as  $\varepsilon$  goes to 0, the invariant distribution of  $(\overline{W}_0^\varepsilon(t))$  converges to the invariant distribution of  $(\overline{W}_0(t))$ .

It must be noted that this description does not take into account the delay after which the source node decides to decrease its throughput after a loss: Because of transmission times this event is discovered by the source after some time. This is nevertheless a standard assumption in the mathematical literature of this domain, just because the corresponding mathematical models (stochastic differential equations with delays) are, up to now, quite intractable. The decision to increase is, for the same reasons, also delayed because destination node informs the source node of the successful transmission of a packet by sending an acknowledgment packet. This aspect is nevertheless taken into account in the stochastic model by the linear growth rate  $a$  of the throughput of the connection (the coefficient of  $f'(x)$  which is 1 in Equation (2.2)). The quantity  $a$  will be thought as proportional to the inverse of the RTT (Round Trip Time), the duration of transmission between the two nodes. See Section 3.

In the following, a connection will be described as a Markov process similar to  $(\overline{W}_0(t))$ . Since different classes of connections will be considered, several parameters have to be introduced, a connection will be represented as a Markov process  $(W(t))$  whose infinitesimal generator is given by

$$(2.3) \quad \Omega(f)(x) = af'(x) + bx(f(rx) - f(x))$$

for  $f$  a  $C^1$ -function from  $\mathbb{R}_+$  to  $\mathbb{R}$ . As noted previously, the rate of increase  $a$  is related to the round trip time between the source and the destination. Remember one of the basic (simplified) principle of the TCP Algorithm: The variable  $W$  is increased by 1 only when it is known that all the  $W$  packets have been successfully transmitted and in particular received by the destination. The variable  $b$  driving the loss rate will depend on the state of the node considered. The quantities  $a$ ,  $b$  and  $r$  will also depend on the class of the connection.

The Markov process  $(W(t))$  increases linearly at rate  $b$  and, given  $W(t) = x$ , jumps at rate  $x$  from  $x$  to  $rx$ . As a stochastic process starting from  $x \geq 0$ , it can be represented as the solution  $(W(t))$  of the following stochastic differential equation:

$W(0) = x$  and

$$(2.4) \quad \begin{aligned} dW(t) &= adt - (1-r)W_0(t-)\mathcal{N}([0, bW(t-)], dt) \\ &= adt - (1-r)W(t-) \int \mathbb{1}_{\{0 \leq x \leq bW(t-)\}} \mathcal{N}(dx, dt) \end{aligned}$$

where  $\mathcal{N}$  is a Poisson process on  $\mathbb{R}_+^2$  whose intensity measure is the Lebesgue measure on  $\mathbb{R}_+^2$  and  $W(t-)$  is the left limit of  $W$  at  $t$ . See Davis [7] and Robert [24].

If  $\rho = a/b$  and  $(V(t)) = ((W(t)/\sqrt{\rho}))$ , then the infinitesimal generator of  $(V(t))$  is given by

$$\Omega_V(f)(x) = \frac{a}{\sqrt{\rho}}f'(x) + b\sqrt{\rho}x(f(rx) - f(x)) = \sqrt{ab}(f'(x) + x(f(rx) - f(x)))$$

which is proportional to the infinitesimal generator of Equation (2.2). In particular, the invariant measure of  $(V(t))$  and  $(\bar{W}_0(t))$  are therefore the same. The density of the invariant distribution is given in the following proposition. See Dumas *et al.* [8] and Guillemin *et al.* [15].

**Proposition 2.1.** *If  $W_0$  is a random variable with density*

$$(2.5) \quad H_{r,\rho}(x) = \frac{\sqrt{2\rho/\pi}}{\prod_{n=0}^{+\infty}(1-r^{2n+1})} \sum_{n=0}^{+\infty} \frac{r^{-2n}}{\prod_{k=1}^n(1-r^{-2k})} e^{-\rho r^{-2n}x^2/2}, \quad x \geq 0,$$

where  $\rho = a/b$ , then the solution of Equation (2.4) with initial condition  $W_0$  having density  $H_{r,\rho}$  is a stationary process and its expected value is given by

$$(2.6) \quad \mathbb{E}(W_0) = \sqrt{\frac{2\rho}{\pi}} \prod_{n=1}^{+\infty} \frac{1-r^{2n}}{1-r^{2n-1}}.$$

At equilibrium, the quantity  $\mathbb{E}(W_0/\sqrt{\varepsilon})$  can be seen as a first order expansion (in  $\varepsilon$ ) of the asymptotic mean throughput of the connection.

### 3. THE STOCHASTIC MODEL OF THE NETWORK

The network has  $J \geq 1$  nodes and accommodates  $K \geq 1$  classes of permanent connections. For  $1 \leq k \leq K$ , the number of class  $k$  connections is  $N_k \geq 1$ , one sets

$$N = (N_1, \dots, N_K), \quad \text{and} \quad |N| = N_1 + \dots + N_K.$$

An *allocation matrix*  $A = (A_{jk}, 1 \leq j \leq J, 1 \leq k \leq K)$  with positive coefficients describes the utilization of nodes by the connections. If  $w_{n,k} \geq 0$  is the state of the  $n$ th class  $k$  connection,  $1 \leq n \leq N_k$ , the quantity  $A_{jk}w_{n,k}$  is the utilization of node  $j$  by this connection. A simple example would be of taking  $A_{jk} = 1$  or 0 if a class  $k$  connection uses node  $j$  or not. The total utilization  $u_j$  of node  $j$  by the various connections is given by

$$u_j = \sum_{k=1}^K \sum_{n=1}^{N_k} A_{jk}w_{n,k}.$$

The quantity  $u_j$  represents the level of congestion of node  $j$ , in particular the loss rate of a connection going through it will depend on this variable. A closely related model has been proposed in Kelly *et al.* [17], see also Raghunathan and Kumar [23] in the context of wireless network.

More precisely, for  $1 \leq k \leq K$ , the corresponding parameters  $a$  and  $b$  of Equation (2.3) for a class  $k$  connection are given by functions  $a_k : \mathbb{R}_+ \times \mathbb{R}_+^J \rightarrow \mathbb{R}_+$  and

$b_k : \mathbb{R}_+ \times \mathbb{R}_+^J \rightarrow \mathbb{R}_+$ , so that when the resource utilization vector of the network is  $u = (u_j, 1 \leq j \leq J)$  and if the state of a class  $k$  connection is  $w_k$ :

- Its state increases linearly at rate  $a_k(w_k, u)$ ;
- A loss for this connection occurs at rate  $b_k(w_k, u)$  and in this case its state jumps to  $r_k w_k$ .

In view of Equation (2.3), a natural form for these functions (with slight abuse of notation) is

$$(3.7) \quad a_k(w_k, u) = a_k(u), \text{ and } b_k(w_k, u) = w_k \beta_k(u).$$

For example, one can take

$$(3.8) \quad \begin{cases} a_k(u) = \left( \tau_k + \sum_{j=1}^J t_{jk}(u_j) \right)^{-1}, \\ \beta_k(u) = \delta_k + \sum_{j=1}^J d_{jk}(u_j) \end{cases}$$

where  $\tau_k > 0$  is the round trip time between source and destination and  $\delta_k$  is the loss rate of class  $k$  connections in non-congested network. For  $1 \leq j \leq J$ ,  $t_{jk}(u_j)$  is their delay at node  $j$  when the total utilization is  $u_j$ . In particular  $t_{jk}(0) = 0$  and  $t_{jk}(u) = 0$  when  $A_{jk} = 0$ . Similarly,  $d_{jk}(u_j)$  is the loss rate at node  $j$ , it is 0 when  $A_{jk} = 0$ .

**A Markovian Representation.** The Markov process describing the state of the ongoing connections is given by, for  $t \geq 0$ ,

$$W^N(t) = (W_{n,k}^N(t), 1 \leq n \leq N_k, 1 \leq k \leq K)$$

where  $W_{n,k}^N(t)$  is the state of the  $n$ th flow of class  $k$  at time  $t$ .

As for Equation (2.4), it can be represented by the solution of the following stochastic differential equation: For  $1 \leq k \leq K$  and  $1 \leq n \leq N_k$ ,

$$(3.9) \quad dW_{n,k}^N(t) = a_k(W_{n,k}^N(t-), U^N(t-)) dt - (1 - r_k) W_{n,k}^N(t-) \int \mathbb{1}_{\{0 \leq z \leq b_k(W_{n,k}^N(t-), U^N(t-))\}} \mathcal{N}_{n,k}(dz, dt),$$

with  $U^N(t) = (U_j^N(t), 1 \leq j \leq J)$  and

$$U_j^N(t) = \sum_{k=1}^K A_{jk} N_k W_k^N(t),$$

where  $(\mathcal{N}_{n,k}, 1 \leq k \leq K, 1 \leq n \leq N_k)$  are independent Poisson processes on  $\mathbb{R}_+^2$  whose intensity measure is Lebesgue measure on  $\mathbb{R}_+^2$ . The following proposition is proved with standard arguments.

**Proposition 3.1.** *If the functions  $a_k$  and  $b_k$ ,  $1 \leq k \leq K$ , are locally bounded then there is pathwise existence and uniqueness for a solution of stochastic differential equation (3.9).*

**Mean Field Scaling.** Due to its high dimensional state space, the system of coupled stochastic differential equations (3.9) does not seem to be mathematically tractable as such. A scaling is used to investigate the qualitative properties of the corresponding Markov processes. It is assumed that for any  $1 \leq k \leq K$  the number of class  $k$  connections  $N_k$  goes to infinity so that

$$(3.10) \quad \frac{N_k}{|N|} = \frac{N_k}{N_1 + \dots + N_K} \rightarrow p_k,$$



where  $p_k \geq 0$  and  $p_1 + \dots + p_K = 1$ .

The variable  $U^N$  is accordingly scaled as  $\bar{U}^N = U^N/|N|$  in the functions  $a_k$  and  $b_k$  so that Equation (3.9) becomes: For  $1 \leq k \leq K$  and  $1 \leq n \leq N_k$ ,

$$(3.11) \quad dW_{n,k}^N(t) = a_k \left( W_{n,k}^N(t-), \bar{U}^N(t-) \right) dt \\ - (1 - r_k) W_{n,k}^N(t-) \int \mathbb{1}_{\{0 \leq z \leq b_k(W_{n,k}^N(t-), \bar{U}^N(t-))\}} \mathcal{N}_{n,k}(dz, dt).$$

with  $U^N(t) = (U_j^N(t), 1 \leq j \leq J)$  and

$$\bar{U}_j^N(t) = \sum_{k=1}^K \frac{N_k}{|N|} A_{jk} \bar{W}_k^N(t) \quad \text{with} \quad \bar{W}_k^N(t) = \frac{1}{N_k} \sum_{n=1}^{N_k} W_{n,k}^N(t).$$

This is a multi-class mean-field system with an interaction through the *scaled utilization* vector  $(\bar{U}^N(t))$ .

For  $1 \leq k \leq K$ , it is natural to introduce the *empirical measure* for the stochastic processes associated to class  $k$  connections,

$$\Lambda_k^N = \frac{1}{N_k} \sum_{n=1}^{N_k} \delta_{(W_{n,k}^N(t), t \geq 0)},$$

where  $\delta_{(x(t), t \geq 0)}$  is the Dirac mass at  $(x(t), t \geq 0)$ .

**Notations and Conventions.** For  $x = (x_m) \in \mathbb{R}^M$  for some  $M \in \mathbb{N}$ ,  $\|x\|$  denotes  $\max(|x_m| : 1 \leq m \leq M)$  and if  $z : \mathbb{R}_+ \rightarrow \mathbb{R}^M$  and  $T > 0$ , one defines

$$\|z\|_T \stackrel{\text{def.}}{=} \sup_{0 \leq s \leq T} \|z(s)\|.$$

Depending on the context the expressions

$$(z_m(t), 1 \leq m \leq M, t \geq 0) \text{ or } (z_m(t), 1 \leq m \leq M)$$

are used for the function  $(z(t))$ .

If  $H$  is a complete metric space,  $\mathcal{D}(\mathbb{R}_+, H)$  denotes the Skorohod space of functions with values in  $H$ , continuous on the right and with left limits at any point of  $\mathbb{R}_+$ , see Billingsley [5]. The variable  $\Lambda_k^N$  has values in the set  $\mathcal{P}(\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+))$  of probability distributions on  $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$ .

#### 4. ANALYSIS OF THE NON-LINEAR LIMITING PROCESS

In view of Equation (3.11), because of the symmetry properties within each class of connection, it is natural to expect a mean-field convergence phenomenon to hold when  $N$  gets large: For  $1 \leq k \leq K$ , the evolution of a *typical* class  $k$  connection should converge in law. More formally, the empirical measure  $\Lambda_k^N$  *should* converge in law to the distribution of a stochastic process  $(W(t)) = ([W_k(t), 1 \leq k \leq K], t \geq 0)$  solution of the non-linear stochastic differential equation

$$(4.1) \quad dW_k(t) = a_k(W_k(t-), u_W(t)) dt \\ - (1 - r_k) W_k(t-) \int \mathbb{1}_{\{0 \leq z \leq b_k(W_k(t-), u_W(t))\}} \mathcal{N}_k(dz, dt),$$

with  $u_W(t) = (u_{W,j}(t), 1 \leq j \leq J)$  and, for  $1 \leq j \leq J$ ,

$$u_{W,j}(t) = \sum_{k=1}^K A_{jk} p_k \mathbb{E}(W_k(t)),$$

where  $(\mathcal{N}_k, 1 \leq k \leq K)$  are i.i.d. Poisson point processes on  $\mathbb{R}_+^2$  with Lebesgue characteristic measure.

**A non-linear stochastic differential equation.** In these equations, the interaction between coordinates of  $(W(t))$  depends in a non-linear way of the mean utilization vector  $(u(t))$  which is a linear functional of mean values  $(\mathbb{E}(W_k(t), 1 \leq k \leq K))$ . In particular, the infinitesimal generator of the process  $(W(t))$  depends *on the law* of  $(W(t))$  and not only on its sample path up to time  $t$  as it is usually the case.

The properties of solution of equations (4.1) are analyzed in this section. The convergence results of the mean-field scaling are the subject of Section 5.

**Non-linear martingale problems.** Throughout this section, the natural filtration  $(\mathcal{F}_t)$  used to investigate the solutions  $(W(t))$  of Equations (4.1) is defined as follows: For  $t \geq 0$ ,  $\mathcal{F}_t$  is the  $\sigma$ -field defined by

$$\mathcal{F}_t = \sigma \langle \mathcal{N}_k(A \times B) : 1 \leq k \leq K, A \in \mathcal{B}(\mathbb{R}_+), B \in \mathcal{B}([0, t]) \rangle,$$

where  $\mathcal{B}(H)$  is the space of Borel sets of  $H$  and  $\mathcal{N}(A \times B)$  is the number of points of the point process  $\mathcal{N}$  in  $A \times B$ . The elementary lemma will be used in the following.

**Lemma.** *If  $(Y(t))$  and  $(Z(t)) \in \mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$  are adapted processes to the filtration  $(\mathcal{F}_t)$  then*

$$\begin{aligned} (I_Z(t)) &= \left( \int_0^t \int Y(s-) \mathbb{1}_{\{0 \leq z \leq Z(s-)\}} [\mathcal{N}(dz, ds) - dz ds] \right) \\ &= \left( \int_0^t Y(s-) [\mathcal{N}([0, Z(s-)], ds) - Z(s) ds] \right) \end{aligned}$$

is a local  $(\mathcal{F}_t)$ -martingale.

*Proof.* By localization with a stopping time,  $(Y(t))$  and  $(Z(t))$  can be assumed to be uniformly bounded by some constant  $C$ .

If, for  $T \geq 0$ ,  $(Z(t), 0 \leq t \leq T)$  has a finite set  $V$  of possible values then, for  $t \leq T$ ,

$$I_Z(t) = \sum_{v \in V} \int_0^t Y(s-) \mathbb{1}_{\{Z(s-)=v\}} (\mathcal{N}([0, v], ds) - v ds).$$

The classical properties of Poisson processes show that the martingale property holds on  $[0, T]$  for each term of the sum in the above expression and consequently for the process  $I_Z$ .

Returning to the general case, since

$$\sup_{0 \leq t \leq T} |I_Z(t)| \leq C(\mathcal{N}([0, C] \times [0, T]) + CT),$$

by approximating  $(Z(t), 0 \leq t \leq T)$  from below by processes with a finite number of possible values, and by using Lebesgue's Theorem, one obtains the desired martingale property.  $\square$

If  $W$  is a solution of Equation (4.1), for  $1 \leq k \leq K$ ,  $t \geq 0$  and  $f$  is a  $C^1$ -function from  $\mathbb{R}_+$  to  $\mathbb{R}$ , by defining

$$(4.2) \quad M_k^f(t) = f(W_k(t)) - f(W_k(0)) - \int_0^t a_k(W_k(s), u_W(s)) f'(W_k(s)) ds \\ - \int_0^t b_k(W_k(s), u_W(s)) (f(r_k W_k(s)) - f(W_k(s))) ds,$$

then  $(M_k^f(t))$  is a local martingale. For  $C^1$ -functions  $f$  and  $g$  and  $k \neq l$ , the local martingales  $M_k^f$  and  $M_l^g$  are independent and the Doob-Meyer brackets are given by

$$\langle M_k^f \rangle_t = \int_0^t b_k(W_k(s), u_W(s)) (f(r_k W_k(s)) - f(W_k(s)))^2 ds.$$

See Rogers and Williams [25].

A non-linear martingale problem formulation of the equations (4.1), is that there exists a probability  $P$  on the space  $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+^K)$  of sample paths  $(W(t)) = ((W_k(t), 1 \leq k \leq K), t \geq 0)$ , such that for any  $C^1$ -function  $f$ , the process  $M_k^f$  defined by Formula (4.2) is a local martingale for  $P$ .

**Existence and Uniqueness Results.** The following proposition is the central technical result used to establish the existence and uniqueness of solutions of Equation (4.1), as well as the mean-field convergence result.

**Proposition 4.1.** *If the functions  $(a_k)$  are bounded and  $(a_k)$  and  $(b_k)$  are Lipschitz, to any non-negative continuous  $\mathbb{R}_+^J$ -valued function  $u = (u(t))$  and a random variable  $X_0$  is associated the solution  $\phi(X_0, u) = (X_k(t))$  of the stochastic differential equation,*

$$(E_u) \quad dX_k(t) = a_k(X_k(t-), u(t)) dt \\ - (1 - r_k) X_k(t-) \int \mathbb{1}_{\{0 \leq z \leq b_k(X_k(t-), u(t))\}} \mathcal{N}_k(dz, dt), \quad 1 \leq k \leq K,$$

then, if  $(u(t))$  and  $(u'(t))$  are non-negative continuous functions and  $X_0$  and  $X'_0$  are random variables in  $\mathbb{R}_+^K$ ,

$$(4.3) \quad \mathbb{E} \left[ \|\phi(X_0, u) - \phi(X'_0, u')\|_t \mid X(0), X'(0) \right] \\ \leq \left[ \|X_0 - X'_0\| + \int_0^t C(s) \|u - u'\|_s ds \right] e^{tC(t)}, \quad t \geq 0,$$

where  $C(t) = A(1 + t + \|u\|_t + \|X_0\| + \|X'_0\|)$  and  $A$  is a constant depending only on the functions  $(a_k)$  and  $(b_k)$ .

*Proof.* The quantity  $L_b$  [resp.  $L_a$ ] denote the maximum of the Lipschitz constants for the function  $b_k$ , [resp.  $a_k$ ],  $1 \leq k \leq K$ . Denote by  $(X_k(t)) = (\phi(X_0, u)(t))$  and  $(X'_k(t)) = (\phi(X'_0, u')(t))$ .

Note that, since the functions  $(a_k)$  are bounded, by some constant  $\|a\|$  say, any solution  $(Z(t))$  of Equation (4.3) satisfies the relation

$$(4.4) \quad Z_k(t) \leq Z_k(0) + \|a\|t.$$

The quantity  $\mathbb{E}^0(\cdot)$  will refer to the conditional expectation given  $X(0)$  and  $X'(0)$ . For  $t \geq 0$ ,

$$(4.5) \quad \begin{aligned} \|X_k - X'_k\|_t &\leq |X_k(0) - X'_k(0)| \\ &\quad + \int_0^t |a_k(X_k(s), u(s)) - a_k(X'_k(s), u'(s))| ds \\ &\quad + \int_0^t \int \left| X_k(s-) \mathbb{1}_{\{z \leq b_k(X_k(s-), u(s))\}} - X'_k(s-) \mathbb{1}_{\{z \leq b_k(X'_k(s-), u'(s))\}} \right| \mathcal{N}(dz, ds). \end{aligned}$$

By using the above lemma and the upper bound on  $(X_k(t))$  which shows that the local martingale of this lemma is indeed a martingale, the conditional expected value of the last term of this expression is given by

$$(4.6) \quad \begin{aligned} \mathbb{E}^0 \left( \int_0^t \int \left| X_k(s) \mathbb{1}_{\{z \leq b_k(X_k(s), u(s))\}} - X'_k(s) \mathbb{1}_{\{z \leq b_k(X'_k(s), u'(s))\}} \right| dz ds \right) \\ \leq \mathbb{E}^0 \left( \int_0^t |X_k(s) - X'_k(s)| b_k(X_k(s), u(s)) ds \right) \\ + \mathbb{E}^0 \left( \int_0^t X'_k(s) |b_k(X_k(s), u(s)) - b_k(X'_k(s), u'(s))| ds \right). \end{aligned}$$

By using the growth conditions and the Lipschitz properties of  $(a_k)$  and  $(b_k)$ , one gets that

$$\begin{aligned} \mathbb{E}^0(\|X_k - X'_k\|_t) &\leq |X_k(0) - X'_k(0)| \\ &\quad + \int_0^t C_0(s) (\|u(s) - u'(s)\| + \mathbb{E}^0(\|X_k - X'_k\|_s)) ds, \end{aligned}$$

with

$$C_0(t) = L_a + L_b(2\|a\|t + \|u\|_t) + \sum_{k=1}^K b_k(0, 0) + L_b[X_k(0) + X'_k(0)] + \sum_{j=1}^J L_b u_j(0),$$

so that  $(C_0(t))$  can be replaced by  $(C(t))$  as in the statement of the proposition. Gronwall's Inequality gives the estimation

$$\begin{aligned} \mathbb{E}^0(\|X_k - X'_k\|_t) &\leq \left[ |X_k(0) - X'_k(0)| + \int_0^t C(s) \|u(s) - u'(s)\| ds \right] e^{\int_0^t C(s) ds} \\ &\leq \left[ |X_k(0) - X'_k(0)| + \int_0^t C(s) \|u - u'\|_s ds \right] e^{tC(t)}. \end{aligned}$$

The proposition is proved.  $\square$

One of the main motivations of this study is to obtain results which are valid for functions  $(a_k)$  and  $(b_k)$  of the form (3.7) and (3.8), which have basically a quadratic behavior. To control the evolution of the vector  $(W(t))$  and describe its stationary behavior, initial conditions cannot be assumed to be uniformly bounded, for this reason exponential and Gaussian moment assumptions are introduced.

**Condition (C)** *It is said to hold for a family of random variables  $\{X_0^\alpha, \alpha \in \mathcal{S}\}$  in  $\mathbb{R}_+^K$ , for  $(b_k)$ , and for  $\varepsilon > 0$  when at least one of the two conditions is satisfied:*

- (1) For any  $1 \leq k \leq K$ , the function  $b_k : \mathbb{R}_+ \times \mathbb{R}_+^J \rightarrow \mathbb{R}_+$  is Lipschitz and a uniform exponential moment of order  $\varepsilon$  holds:

$$\sup_{\alpha \in \mathcal{S}} \mathbb{E}(\exp(\varepsilon \|X_0^\alpha\|)) < \infty.$$

- (2) For any  $1 \leq k \leq K$ ,  $b_k(w, u) = w\beta_k(u)$  and  $\beta_k : \mathbb{R}_+^J \rightarrow \mathbb{R}_+$  is Lipschitz and a uniform Gaussian moment of order  $\varepsilon$ :

$$\sup_{\alpha \in \mathcal{S}} \mathbb{E}(\exp(\varepsilon \|X_0^\alpha\|^2)) < \infty.$$

The following theorem establishes the existence and uniqueness result of a solution of Equation (4.1).

**Theorem 4.2.** *If the functions  $a_k : \mathbb{R}_+ \times \mathbb{R}_+^J \rightarrow \mathbb{R}_+$ ,  $1 \leq k \leq K$  are bounded and Lipschitz and if Condition (C) holds for  $W_0$ ,  $b_K$  and  $\varepsilon > 0$ , then there is pathwise existence and uniqueness of a solution  $(W(t))$  of the non-linear stochastic differential equation (4.1) starting at  $W_0$ .*

*In this case, the solution depends continuously on the initial condition in the following way: if  $(W(t))$  and  $(W'(t))$  are solutions of Equation (4.1) with respective initial conditions  $W_0$  and  $W'_0$  having the same moment in Condition (C), for  $T \geq 0$ , there exists a constant  $A_T$  such that*

$$(4.7) \quad \mathbb{E}(\|W - W'\|_T) = \mathbb{E}\left(\sup_{s \leq T} \|W(s) - W'(s)\|\right) \\ \leq A_T \mathbb{E}\left(\|W_0 - W'_0\| e^{\varepsilon(\|W_0\|^\ell + \|W'_0\|^\ell)/2}\right) e^{\mathbb{E}(\exp[\varepsilon(\|W_0\|^\ell + \|W'_0\|^\ell)])}.$$

where  $\ell = 1$  or  $2$ , depending on the moment in Condition (C): exponential or Gaussian.

*Proof.* It is assumed for the moment that (1) of Condition (C) is satisfied. The functions  $(a_k)$  are bounded by the constant  $\|a\|$ . Recall that for an integrable process  $(Z(t)) = (Z_k(t))$ , for  $t \geq 0$ ,

$$u_Z(t) = (u_{Z,j}(t), 1 \leq j \leq J) = \left(\sum_{k=1}^K A_{jk} p_k \mathbb{E}(Z_k(t)), 1 \leq j \leq J\right).$$

Let  $\alpha$  denote the constant  $\max(A_{j1}p_1 + A_{j2}p_2 + \dots + A_{jK}p_K : 1 \leq j \leq J)$ .

The sequence  $(W^n)$  is constructed by induction as follows:

- $W^0 \equiv W(0)$ ;
- For  $n \geq 1$ ,  $W^n$  is the solution of Equation  $(\mathcal{E}_u)$ , see Proposition 4.1, with  $W^n(0) = W(0)$  and  $(u(t)) = (u_{W^{n-1}}(t))$ .

Note that by the growth condition  $W^n(t) \leq W^n(0) + \|a\|t$  for  $t \geq 0$  and  $n \geq 1$ , and consequently  $\|u_{W^n}\|_t \leq \alpha(\|W(0)\| + \|a\|t)$ . With the notation of Proposition 4.1,  $W^n = \phi(W(0), u_{W^{n-1}})$  for  $n \geq 1$ , so that Equation (4.3) gives the relation

$$\mathbb{E}\left[\|W^{n+1} - W^n\|_t \mid W(0)\right] \leq \left[\int_0^t C(s) \|u_{W^n} - u_{W^{n-1}}\|_s ds\right] e^{tC(t)}, \quad t \geq 0,$$

with  $C(t) = A_0(1 + t + \|W(0)\|)$  where  $A_0$  is a constant depending only on  $(a_k)$  and  $(b_k)$ . Because of the assumption on the existence of an exponential moment

for  $W(0)$ , by taking  $t_0 = \varepsilon/(2A_0)$ , then  $\delta = \alpha\mathbb{E}(C(t_0)\exp(t_0C(t_0))) < +\infty$ . One gets that for  $t \leq t_0$ ,

$$\|u_{W^{n+1}} - u_{W^n}\|_t \leq \delta \int_0^t \|u_{W^n} - u_{W^{n-1}}\|_s ds, \quad n \geq 1,$$

so that

$$\|u_{W^{n+1}} - u_{W^n}\|_t \leq \frac{(\delta t)^n}{n!} \|u_{W^1} - u_{W^0}\|_{t_0}, \quad n \geq 1,$$

with Relation (4.3). One gets therefore that

$$\mathbb{E}(\|W^{n+1} - W^n\|_{t_0}) \leq \frac{(\delta t_0)^n}{n!} \|u_{W^1} - u_{W^0}\|_{t_0}$$

holds almost surely. Consequently, the sequence of processes  $(W^n(t), t \leq t_0)$  converges uniformly to some process  $(W(t), t \leq t_0)$  and

$$\mathbb{E}(\|W^n - W\|_{t_0}) \leq \frac{(\delta t_0)^n}{n!} e^{\delta t_0} \|u_{W^1} - u_{W^0}\|_{t_0}$$

The above relation also shows that on the interval  $[0, t_0]$ , the sequence of functions  $(\mathbb{E}(W^n(t)))$  converges uniformly to  $(\mathbb{E}(W(t)))$ , in particular  $(u_{W^n}(t))$  converges uniformly to  $(u_W(t))$  on this interval.

By definition of  $W^{n+1}$ , for  $t \leq t_0$  and  $1 \leq k \leq K$ ,

$$\begin{aligned} W_k^{n+1}(t) &= W_k(0) + \int_0^t a_k(W_k^n(s), u_{W^n}(s)) ds \\ &\quad - (1 - r_k)W_k^n(t-) \int_0^t \mathcal{N}([0, b_k(W_k^n(s-), u_{W^n}(s))], ds), \end{aligned}$$

the uniform convergences, the Lipschitz assumption for  $a_k$  and  $b_k$  and continuity properties of the Poisson process  $\mathcal{N}_k$  show that, almost surely, the limit  $(W(t))$  satisfies indeed Relation (4.1) on  $[0, t_0]$ .

If  $(W'(t))$  is another solution of Equation (4.1) with  $W'(0) = W(0)$ , by using again Relation (4.3), one gets that, for  $t \leq t_0$ ,

$$\|u_W - u_{W'}\|_t \leq \delta \int_0^t \|u_W - u_{W'}\|_s ds,$$

so that  $u_W = u_{W'}$  on  $[0, t_0]$  and consequently,  $W = W'$  on this interval.

Since  $\|W(t_0)\| \leq \|W_0\| + \|a\|t_0$ , Condition (C) holds for  $W(t_0)$  and the same  $\varepsilon > 0$ , by taking it as an initial condition and by translating the Poisson processes by  $t_0$  (on the second coordinate), one can show existence and uniqueness results on the time interval  $[t_0, 2t_0]$ , and therefore by induction on  $\mathbb{R}_+$ .

If  $(W(t))$  and  $(W'(t))$  are the solutions of Equation (4.1) with respective initial conditions  $W_0$  and  $W'_0$ , Relation (4.3) gives that

$$\|u_W - u_{W'}\|_t \leq \alpha \mathbb{E} \left( \left[ \|W_0 - W'_0\| + \int_0^t \|u_W - u_{W'}\|_s C(s) ds \right] e^{tC(t)} \right),$$

then, for  $t < t_0$ , by using again Gronwall's Lemma, and the expression of  $C(t)$  one gets that there exists a constant  $A$  depending only on the functions  $(a_k)$  and  $(b_k)$  such that

$$\mathbb{E}(\|u_W - u_{W'}\|_t) \leq A \mathbb{E} \left( \|W_0 - W'_0\| e^{\varepsilon(\|W_0\| + \|W'_0\|/2)} \right) e^{\mathbb{E}(\exp(\varepsilon(\|W_0\| + \|W'_0\|)))}$$

for  $t \leq t_0$ . By plugging this inequality in Relation (4.3), one gets a similar relation for  $\mathbb{E}(\|W - W'\|_{t_0})$ . With the same procedure, an analogous inequality can be obtained on the interval  $[0, nt_0]$  for any  $n \geq 1$  (with a constant  $A$  depending on  $n$ ). The theorem has therefore been proved on  $\mathbb{R}_+$  in case (1) of Condition (C).

For the case (2) of Condition (C), when  $b_k(w, u) = w\beta_k(u)$  and  $\beta_k$  is Lipschitz, the corresponding inequality (4.6) in this setting is the relation

$$\begin{aligned} & |X_k(s)\beta_k(u(s)) - X'_k(s)\beta_k(u'(s))| \\ & \leq X_k(s)|\beta_k(u(s)) - \beta_k(u'(s))| + \beta_k(u'(s))|X_k(s) - X'_k(s)|, \end{aligned}$$

and the rest of the proof is analogous. The supplementary multiplication by  $W_k(s)$  of the Lipschitz bounds is handled by the Gaussian moment assumption.  $\square$

## 5. THE MEAN-FIELD LIMIT FOR CONVERGING INITIAL DATA

In this section, the convergence results to the limiting stochastic process analyzed in Section 4 are established. In the context of mean-field convergence, the notions of exchangeability and chaoticity play a fundamental role. If they are classical in the context of single-class systems, see Aldous [2] for example, these properties have to be extended to investigate the stochastic model of the multi-class network considered in this paper.

**Definition.** *The sequence of random variables  $(X_{n,k}, 1 \leq n \leq N_k, 1 \leq k \leq K)$  is said to be multi-exchangeable if its law is invariant under permutation of the indexes within the classes: for  $1 \leq k \leq K$  and any permutations  $\sigma_k$  of  $\{1, \dots, N_k\}$ , the equality in distribution*

$$(X_{\sigma_k(n),k}, 1 \leq n \leq N_k, 1 \leq k \leq K) \stackrel{\text{dist.}}{=} (X_{n,k}, 1 \leq n \leq N_k, 1 \leq k \leq K)$$

holds.

A sequence  $(X_{n,k}^N, 1 \leq n \leq N_k, 1 \leq k \leq K)$  of multi-class random variables indexed by  $N = (N_k) \in \mathbb{N}^K$  is  $P_1 \otimes \dots \otimes P_K$ -multi-chaotic if, for any  $m \geq 1$ , the convergence in distribution

$$\lim_{N \rightarrow \infty} (X_{n,k}^N, 1 \leq n \leq m, 1 \leq k \leq K) = P_1^{\otimes m} \otimes \dots \otimes P_K^{\otimes m}$$

holds for the topology of uniform convergence on compact sets, where  $P_k$  for  $1 \leq k \leq K$  is a probability distribution on  $\mathbb{R}_+$  and with the convention that  $N$  goes to infinity when  $\min_k N_k$  goes to infinity.

If a sequence of random variables is multi-chaotic, then any fixed finite sub-system is asymptotically independent, with particles of class  $k$  having the law  $P_k$ .

An interesting result, not used in this paper, is that a sequence of multi-exchangeable multi-class systems is multi-chaotic if and only if the restriction to each class is chaotic. See Theorem 3 in Graham [12].

**Mean Field Scaling: Initial conditions and definitions.** Equation (3.10) defines the growth rate of the number of class  $k$  connections when the total number of connections goes to infinity, one denotes

$$M^N = \min_{1 \leq k \leq K} N_k,$$

and the terminology “ $N$  goes to infinity” refers to the fact that  $M^N$  converges to infinity. Additionally, it will be assumed that the initial conditions of SDE (3.11)

$$(W_{n,k}^N(0), 1 \leq n \leq N_k, 1 \leq k \leq K)$$

are multi-exchangeable and  $P_{1,0} \otimes \cdots \otimes P_{K,0}$ -multi-chaotic where, for  $1 \leq k \leq K$ ,  $P_{k,0}$  is a probability distribution on  $\mathbb{R}_+$ . The process  $(W_{n,k}^N(t))$  is the solution of Equation (3.11) in this setting. For  $N \in \mathbb{N}^K$  and  $n \leq M^N$ , one defines

$$W_n^N(t) = (W_{n,k}^N(t), 1 \leq k \leq K).$$

Note that, because of exchangeability,  $(W_n^N(t), t \geq 0)$  and  $(W_1^N(t), t \geq 0)$  have the same distribution and, because of multi-chaoticity, the variable  $W_1^N(0)$  converges in distribution to  $P_{1,0} \otimes \cdots \otimes P_{K,0}$  as  $M^N$  goes to infinity.

The following theorem is the main result for the mean-field convergence. The underlying topology on the corresponding functional space is the uniform convergence on compact sets.

**Theorem 5.1.** *Under the conditions*

- (1) for  $1 \leq k \leq K$ ,

$$\lim_{N \rightarrow +\infty} \frac{N_k}{N_1 + \cdots + N_K} = p_k,$$

- (2) the variables  $(W_{n,k}^N(0), 1 \leq n \leq N_k, 1 \leq k \leq K)$  are multi-exchangeable and  $P_{1,0} \otimes \cdots \otimes P_{K,0}$ -multi-chaotic,  
(3) the function  $a_k : \mathbb{R}_+ \times \mathbb{R}_+^J \rightarrow \mathbb{R}_+$ ,  $1 \leq k \leq J$  are bounded and Lipschitz, and Condition (C) holds for the random variables  $\{W_1^N(0), N \in \mathbb{N}^K\}$ ,  $(b_k)$  and  $\varepsilon > 0$ ,

then, as  $N$  goes to infinity, the sequence of processes

$$((W_{n,k}^N(t), t \geq 0), 1 \leq n \leq N_k, 1 \leq k \leq K),$$

the solutions of SDE (3.11) with initial conditions  $(W_{n,k}^N(0))$ , is multi-exchangeable and  $P_W$ -multi-chaotic, where  $P_W = P_{W_1} \otimes \cdots \otimes P_{W_K}$  is the distribution of the process  $(W(t), t \geq 0) = ((W_k(t), t \geq 0), 1 \leq k \leq K)$ , the solution of the non-linear SDE (4.1) with initial distribution  $P_{1,0} \otimes \cdots \otimes P_{K,0}$ .

In particular, for any  $n \geq 1$ ,  $(W_n^N(t), t \geq 0) = ((W_{n,k}^N(t), 1 \leq k \leq K), t \geq 0)$  converges in distribution to  $(W(t), t \geq 0)$  as  $N$  goes to infinity.

*Proof.* We prove the result assuming that  $b_k$  is Lipschitz. The other case is similar, see the end of the proof of Theorem 4.2. A coupling method will be used.

Let  $(V_n^N(t)) = ((V_{n,k}^N(t), 1 \leq k \leq K))$  be the solution of Equation (4.1) with the initial conditions of the process  $(W_n^N(t))$ :

$$V_n^N(0) = (V_{n,k}^N(0), 1 \leq k \leq K) = (W_{n,k}^N(0), 1 \leq k \leq K) = W_n^N(0)$$

and Poisson processes  $(\mathcal{N}_{n,k}, 1 \leq k \leq K)$  driving the dynamic are the same as for  $(W_n^N(t))$  in Equation (3.11).

The key result to prove the theorem is to prove that the processes  $(W_n(t))$  and  $(V_n(t))$  are asymptotically close, that is, for any  $T > 0$  and  $n \geq 1$ ,

$$(5.1) \quad \lim_{N \rightarrow +\infty} \mathbb{E} \left( \sup_{0 \leq t \leq T} \|W_n^N(t) - V_n^N(t)\| \right) = 0.$$



Because of the multi-exchangeability property, one has to consider only the case  $n = 1$ . With the notation of Proposition 4.1, one has  $V_1^N = \phi(W_1^N(0), u_{V_1^N})$  by definition of  $V_1^N$  and  $W_1^N = \phi(W_1^N(0), \bar{U}^N(t))$  by Equation (3.11). By taking  $X_0 = X'_0 = W_1^N(0)$ ,  $u = u_{V_1^N}$ , and  $u' = \bar{U}^N$ , and the expectation of Inequality (4.3) with respect to the Poisson processes  $(\mathcal{N}_{n,k}, 1 \leq k \leq K, n \neq 1)$  gives

$$(5.2) \quad \mathbb{E}^N \left( \sup_{s \leq t} \|W_1^N(s) - V_1^N(s)\| \right) \leq C_1^N(t) e^{C_1^N(t)t} \int_0^t \mathbb{E}^N \left( \|\bar{U}^N(s) - u_{V_1^N}(s)\| \right) ds.$$

where  $\mathbb{E}^N(\cdot)$  denotes the expectation conditionally on the random vector  $W_1^N(0)$  and, since the function  $(a_k)$  are bounded by some  $\|a\|$ ,

$$(5.3) \quad V_{1,k}^N(t) \leq V_{1,k}^N(0) + t\|a\| = W_{1,k}^N(0) + t\|a\|, \quad 1 \leq k \leq K, t \geq 0,$$

and  $C_1^N(t)$  can be taken as  $A(1 + \|W_1^N(0)\| + t)$ , for some constant  $A$  depending only on the functions  $(a_k)$  and  $(b_k)$

We hereafter cope with the lack of symmetry of multi-exchangeable systems, whereas similar proofs exploit the strength of full exchangeability.

It is assumed that  $p_k > 0$  for all  $1 \leq k \leq K$  for ease of presentation (the proof extends easily), then, for  $t \geq 0$ ,

$$(5.4) \quad \mathbb{E}^N \left( \|\bar{U}^N(t) - u_{V_1^N}(t)\| \right) \leq \sum_{j=1}^J \sum_{k=1}^K A_{jk} \mathbb{E}^N \left| \frac{N_k}{|N|} \bar{W}_k^N(t) - p_k \mathbb{E}(V_{1,k}^N(t)) \right|,$$

with, for  $1 \leq k \leq K$ ,

$$\bar{W}_k^N(t) = \frac{1}{N_k} \sum_{n=1}^{N_k} W_{n,k}^N(t),$$

and

$$(5.5) \quad \mathbb{E}^N \left| \frac{N_k}{|N|} \bar{W}_k^N(t) - p_k \mathbb{E}(V_{1,k}^N(t)) \right| \leq p_k \mathbb{E}^N \left| \frac{1}{N_k} \sum_{n=1}^{N_k} [W_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right| + \Delta_k^N(t),$$

with

$$\Delta_k^N(t) \stackrel{\text{def.}}{=} \left| \frac{1}{|N|} - \frac{p_k}{N_k} \right| \mathbb{E}^N \left| \sum_{n=1}^{N_k} W_{n,k}^N(t) \right| \leq \left| \frac{N_k}{|N|} - p_k \right| \mathbb{E}^N |W_{1,k}^N(t)|,$$

and, consequently,

$$\lim_{N \rightarrow \infty} \mathbb{E} \left( \sup_{s \leq t} \Delta_k^N(s) \right) = 0$$

due to the uniform integrability bounds on the initial data and the growth bound of the solutions of Equation (3.11).

The finite set  $\{1, \dots, N_k\}$  is partitioned into  $\lfloor N_k/M^N \rfloor$  consecutive subsets with  $M^N$  elements and the remaining subset with cardinality  $R_k^N < M^N$ . Using again

the multi-exchangeability property, one gets

$$(5.6) \quad \mathbb{E}^N \left| \frac{1}{N_k} \sum_{n=1}^{N_k} [W_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right| \\ \leq \left[ \frac{N_k}{M^N} \right] \frac{M^N}{N_k} \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [W_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right| \\ + \frac{R_k^N}{N_k} \mathbb{E}^N \left| \frac{1}{R_k^N} \sum_{n=1}^{R_k^N} [W_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right|$$

where

$$\mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [W_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right| \\ \leq \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} (W_{n,k}^N(t) - V_{n,k}^N(t)) \right| + \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} (V_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))) \right| \\ (5.7) \quad \leq \mathbb{E}^N |W_{1,k}^N(t) - V_{1,k}^N(t)| + \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [V_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right|.$$

The Jensen Inequality yields

$$\mathbb{E} \left( \left( \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [V_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right| \right)^2 \right) \\ \leq \mathbb{E} \left( \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [V_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right|^2 \right)$$

and by developing the square, one gets

$$\mathbb{E} \left( \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [V_{n,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right|^2 \right) = \frac{1}{M^N} \mathbb{E} \left( [V_{1,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))]^2 \right) \\ + \frac{M^N - 1}{M^N} \mathbb{E} \left( [V_{1,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] [V_{2,k}^N(t) - \mathbb{E}(V_{1,k}^N(t))] \right),$$

and by exchangeability, the covariance term of the last expression is given by

$$\mathbb{E}(V_{1,k}^N(t)V_{2,k}^N(t)) - 2\mathbb{E}(V_{1,k}^N(t))\mathbb{E}(V_{1,k}^N(t)) + [\mathbb{E}(V_{1,k}^N(t))]^2.$$

Hence, the uniform integrability bounds yield

$$\lim_{N \rightarrow \infty} \mathbb{E} \left( \left( \sup_{s \leq t} \mathbb{E}^N \left| \frac{1}{M^N} \sum_{n=1}^{M^N} [V_{n,k}^N(s) - \mathbb{E}(V_{1,k}^N(s))] \right| \right)^2 \right) = 0.$$

For the remainder term containing  $R_k^N$  in (5.6) two cases have to be considered,

$$R_k^N \leq \sqrt{|N|} \Rightarrow \lim_{N \rightarrow \infty} \frac{R_k^N}{N_k} = 0, \quad \text{and } R_k^N > \sqrt{|N|} \Rightarrow \lim_{N \rightarrow \infty} R_k^N = \infty,$$

and the second case is treated as for  $M^N$ . Inequalities (5.4), (5.5), (5.6) and (5.7) yield

$$\mathbb{E}^N \|u_{V_1^N}(t) - \bar{U}^N(t)\| \leq \sum_{j=1}^J \sum_{k=1}^K A_{jk} p_k \mathbb{E}^N \|V_1^N(t) - W_1^N(t)\| + F^N(t),$$

with

$$(5.8) \quad \lim_{N \rightarrow \infty} \mathbb{E} \left( \sup_{s \leq t} |F^N(s)| \right) = 0.$$

By plugging this bound in Inequality (5.2), one gets

$$\mathbb{E}^N (\|W_1^N - V_1^N\|_t) \leq C_1^N(t) e^{C_1^N(t)t} \left( A \int_0^t \mathbb{E}^N (\|W_1^N - V_1^N\|_s) ds + F^N(t) \right).$$

for some constant  $A > 0$ . The Gronwall Lemma gives the relation

$$\mathbb{E}^N (\|W_1^N - V_1^N\|_t) \leq F^N(t) C_1^N(t) e^{C_1^N(t)t} \exp \left( At C_1^N(t) e^{C_1^N(t)t} \right),$$

the exponential moment assumption for the sequence  $(W_1^N(0), \in \mathbb{N}^K)$  shows that, for  $t < t_0$  with  $t_0$  depending only on the functions  $(a_k)$  and  $(b_k)$  and  $\varepsilon$ , the random variables  $C_1^N(t) \exp(C_1^N(t)t)$ ,  $N \geq 1$  are tight. Relation (5.8) implies therefore that the random variable  $\mathbb{E}^N (\|V_1^N - W_1^N\|_t)$  converges to 0 in probability when  $N$  goes to infinity. The growth bounds (5.3) (also valid for  $W_1^N$ ) and the uniform integrability property of  $(W_1^N(0), N \in \mathbb{N}^K)$  show that the convergence to 0 also holds for the expected value by Lebesgue's Theorem. Relation (5.1) has therefore been established for  $t \leq t_0$ . The extension to any arbitrary  $t$  is done as in the proof of Theorem 4.2.

The continuity property with respect to the initial conditions implied by Relation (4.7) and the uniqueness result of Theorem 4.2 show that the sequence of processes  $(V_{n,k}^N(t), 1 \leq n \leq N_k, 1 \leq k \leq K)$  is  $P_W = P_{W_1} \otimes \cdots \otimes P_{W_K}$ -multi-chaotic. Relation (5.1) gives that the same property also holds for the sequence of processes  $(W_{n,k}^N(t), 1 \leq n \leq N_k, 1 \leq k \leq K)$ . The theorem is proved.  $\square$

**Corollary 5.2.** *Under the assumptions of Theorem 5.1 and if, for  $1 \leq k \leq K$ ,  $P_{W_k}$  is the distribution of  $(W_k(t))$  such that the process  $(W(t)) = ((W_k(t)), 1 \leq k \leq K)$  is the solution of the non-linear SDE (4.1) with initial distribution  $P_{1,0} \otimes \cdots \otimes P_{K,0}$ , then the convergence in distribution of the empirical distributions*

$$\lim_{N \rightarrow \infty} \Lambda_k^N = P_{W_k}$$

*holds for the weak topology on  $\mathcal{P}(\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+))$  with  $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$  endowed with the Skorohod topology.*

*Proof.* For  $F$  a continuous bounded function on  $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$  then, with the same notations as in the proof of the above theorem and by using the exchangeability properties of the initial conditions,

$$|\Lambda_k^N(F) - P_{W_k}(F)| = |\mathbb{E}[F(W_{1,k}^N) - \mathbb{E}[F(V_{1,k}^N)]]|.$$

For the topology of the uniform convergence on compact sets, as  $N$  gets large,  $(V_{1,k}^N(t))$  and  $(W_{1,k}^N(t))$  have the same limit by Relation (5.1) and the processes  $(V_{1,k}^N(t))$  converge to  $(W_k(t))$  by Inequality (4.7) of Theorem 4.2, one gets that

$$\lim_{N \rightarrow +\infty} \Lambda_k^N(F) = P_{W_k}(F),$$

the weak convergence follows since there exists a convergence determining sequence  $(F_n)$  of  $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$  for the weak topology on  $\mathcal{P}(\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+))$ .  $\square$

## 6. FIXED POINTS AND INVARIANT DISTRIBUTIONS

This section investigates the stationary properties of the solutions of stochastic differential equations (4.1). Because of Poisson processes driving the corresponding random events, a solution  $(W(t))$  of the system of equations (4.1) with some deterministic initial state has clearly the Markov property. Since the function  $(u_W(t))$  has a role in these equations, it will imply that, in general, the Markov property is *non-homogeneous* with time. In the martingale problem formulation, the analogue of the infinitesimal generator *depends*, a priori, not only on time but also on  $\mathbb{E}(W(t))$  which complicates a lot the analysis of the asymptotic behavior of the process  $(W(t))$ . See the remark above in Section 4. As the following theorem shows, this has important implications, in particular for the existence and uniqueness of an invariant distribution for  $(W(t))$ .

**Theorem 6.1.** *In the case where, for  $1 \leq k \leq K$ ,*

$$a_k(w, u) = a_k(u), \text{ and } b_k(w, u) = w\beta_k(u), \quad w \geq 0, u \in \mathbb{R}_+^J,$$

*$\beta_k$  is a positive function and  $a_k$  and  $b_k$  are Lipschitz functions and  $a_k$  is bounded, then the invariant distributions for solutions  $(W(t))$  of Equation (4.1) are in one-to-one correspondence with the solutions  $u \in \mathbb{R}_+^J$  of the fixed point equation*

$$(6.1) \quad u_j = \sum_{k=1}^K A_{jk} p_k \psi(r_k) \sqrt{\frac{a_k(u)}{\beta_k(u)}}, \quad 1 \leq j \leq J,$$

where

$$\psi(r) = \sqrt{\frac{2}{\pi}} \prod_{n=1}^{+\infty} \frac{1 - r^{2n}}{1 - r^{2n-1}}.$$

*If  $u^*$  is such a solution, the corresponding invariant distribution has the density  $w \rightarrow \prod_{k=1}^K H_{r_k, \rho_k}(w_k)$  on  $\mathbb{R}_+^K$ , where  $\rho_k = a_k(u^*)/\beta_k(u^*)$  and  $H_{r, \rho}$  is defined in Proposition 2.1.*

*Proof.* Let  $u^* \in \mathbb{R}_+^J$  a solution of Equation (6.1) and  $W_0 = (W_{0,k}) \in \mathbb{R}_+^K$  a random variable such that, for  $1 \leq k \leq K$ ,  $W_{0,k}$  has the density (2.5) with  $a = a_k(u^*)$ ,  $b = \beta_k(u^*)$  and  $r = r_k$ . By invariance of this density, see Proposition 2.1, the solution  $(\widetilde{W}_k(t))$  of the stochastic differential equation

$$d\widetilde{W}_k(t) = a_k(u^*)dt - (1 - r_k)\widetilde{W}_k(t-)\mathcal{N}_k([0, \widetilde{W}_k(t-)\beta_k(u^*)], dt)$$

with initial condition  $\widetilde{W}_k(0) = W_{0,k}$  is a stationary process, in particular, for any  $t \geq 0$ ,  $\mathbb{E}(\widetilde{W}_k(t)) = \mathbb{E}(W_{0,k})$  so that, for  $1 \leq j \leq J$ ,

$$u_{\widetilde{W},j}(t) = \sum_{k=1}^K A_{jk} p_k \mathbb{E}(\widetilde{W}_k(t)) = \sum_{k=1}^K A_{jk} p_k \psi(r_k) \sqrt{\frac{a_k(u^*)}{\beta_k(u^*)}} = u_j^*$$

by Equation (2.6) and (6.1) so that  $u_{\widetilde{W}}(t) \equiv u^*$ . This implies that the stochastic process  $(\widetilde{W}_k(t), 1 \leq k \leq K)$  satisfies Equation (4.1). The variables  $W_{0,k}$  having a Gaussian moment, the above theorem shows that there exists a unique solution  $(W(t))$  of Equations (4.1) which is thus stationary and such that  $u_W(t) \equiv u^*$ .

Conversely, if  $\pi$  is a stationary distribution of  $(W(t))$ , then if the distribution of  $W(0)$  is  $\pi$ , the quantity  $u^* = u_W(t)$  does not depend on  $t$  and, consequently, for  $1 \leq k \leq K$ , the  $k$ th marginal of  $\pi$  is the invariant distribution of the Markov process whose infinitesimal generator is given by

$$\Omega(f)(x) = a_k(u^*)f'(x) + x\beta_k(u^*)(f(rx) - f(x)).$$

By Proposition 2.1, the density of the  $k$ th marginal of  $\pi$  is necessarily  $H_{r_k, \rho_k}$  with  $\rho_k = a_k(u^*)/\beta_k(u^*)$  and  $u^*$  satisfies Equation (6.1). The proposition is proved.  $\square$

The above theorem shows that if the fixed point equation (6.1) has several solutions then the limiting process  $(W(t))$  has several invariant distributions. Similarly, if equation (6.1) has no solution then, in particular,  $(W(t))$  cannot converge to an equilibrium. These possibilities have been suggested in the Internet literature through simulations, like the cyclic behavior of some nodes in the case of congestion. Obviously, there exist coefficients  $(a_k)$  and  $(\beta_k)$  with these possibilities for Equation (6.1). Under mild and natural assumptions, such that if the loss rate at any node is increasing with its utilization, for some networks there exists a unique fixed point. See the examples investigated below.

Several special cases are now considered under the assumptions of Theorem 6.1. It is assumed that  $A_{jk} = 1$  if node  $j \in \{1, \dots, J\}$  is used by connections of class  $k$  and 0 otherwise. In the same way, for  $k \in \{1, \dots, J\}$ , the function  $\beta_k$  for the loss rate for class  $k$  connections depends only on the sum of the utilizations of the nodes used by these connections, i.e.

$$\beta_k(u) = \beta_k \left( \sum_{j=1}^K A_{jk} u_j \right),$$

with a slight abuse of notations. Equation (6.1) becomes in this context

$$u_j = \sum_{k=1}^K A_{jk} p_k \psi(r_k) \frac{\sqrt{a_k}}{\sqrt{\beta_k \left( \sum_{j=1}^K A_{jk} u_j \right)}}, \quad 1 \leq j \leq J.$$

It is assumed that  $u \rightarrow \beta_k(u)$  is strictly non-decreasing and Lipschitz and that the function  $(a_k(u))$  is constant equal to  $a_k$ . Let  $\rho_k(u)$  denote  $a_k/\beta_k(u)$ .

**One single node.** Here  $J = 1$ , for  $k \in \{1, \dots, K\}$ , there is a unique solution  $u = U$  to Equation (6.1)

$$u = \sum_{k=1}^K \psi(r_k) p_k \sqrt{\frac{a_k}{\beta_k(u)}},$$

and consequently a unique invariant distribution for which the state of a class  $k$  connection has the density  $H_{r,\rho_k}$  and its expected value, the throughput, is

$$\psi(r_k) \sqrt{\frac{a_k}{\beta_k(U)}}.$$

In particular, if the effects of the congestion of the node on a connection are independent of its class, i.e.  $\beta_k \equiv \beta$  for  $1 \leq k \leq K$ , then, if the scaling factors ( $r_k$ ) are equal, the throughputs of the different classes of connections differ by the multiplicative factors ( $\sqrt{a_k}$ ). Recall that  $a_k$  is typically proportional to the inverse of the round trip time of a class  $k$  connection. See Section 3.

**A linear network.** A network with  $J$  nodes and  $J + 1 = K$  classes of connections is considered. For  $1 \leq j \leq J$  connections of class  $j$  uses only node  $j$  and the route of connections of class  $J + 1$  go along the  $J$  nodes. This network has been considered in Ben Fred *et al.* [9]. In this context, Equation (6.1) becomes




FIGURE 1. A network with  $J$  nodes and  $J + 1 = K$  classes of connections

$$(6.2) \quad u_j = \left( \frac{\alpha_j}{\sqrt{\beta(u_j)}} + \frac{\alpha_{J+1}}{\sqrt{\beta(\|u\|)}} \right), \quad 1 \leq j \leq J,$$

with  $\alpha_j = \psi(r_j)p_j\sqrt{a_j}$ ,  $1 \leq j \leq J$  and  $\|u\| = |u_1| + \dots + |u_J|$ . For  $\alpha > 0$ , the function  $\phi_{j,\alpha}(x) \stackrel{\text{def.}}{=} x - \alpha/\sqrt{\beta_j(x)}$  being strictly non-decreasing, the equation

$$u = \sum_{j=1}^J \phi_{j,\alpha_j}^{-1} \left( \frac{\alpha_{J+1}}{\sqrt{\beta(u)}} \right)$$

has a unique solution  $U$ . A solution ( $u_j$ ) of Equation (6.2) is necessarily given by

$$u_j = \phi_{j,\alpha_j}^{-1} \left( \frac{\alpha_{J+1}}{\sqrt{\beta(U)}} \right), \quad 1 \leq j \leq J.$$

Equation (6.2) has therefore a unique solution.

**Network on a Torus.** For this network, there are  $J$  nodes and  $K = J$  classes of connections and class  $j \in \{1, \dots, J\}$  uses two nodes: node  $j$  and  $j + 1$ . The assumptions on the rate functions are the same as before. For this network Equation (6.1) becomes

$$(6.3) \quad u_j = \left( \frac{\alpha_{j-1}}{\sqrt{\beta(u_{j-1} + u_j)}} + \frac{\alpha_j}{\sqrt{\beta(u_j + u_{j+1})}} \right), \quad 1 \leq j \leq J,$$

with the convention that  $J + 1 = 1$  and  $0 = J$  for the subscripts. For  $1 \leq j \leq J$ , define  $\alpha_j = \psi(r_j)p_j\sqrt{a_j}$  and  $\phi(x) = 1/\sqrt{\beta(x)}$ . If  $y_j = \alpha_j\phi(u_{j-1} + u_j)$ , then the above




FIGURE 2. Triangular Network

equation can be rewritten as follows

$$(6.4) \quad y_j = \alpha_j \phi(y_{j-1} + 2y_j + y_{j+1}), \quad 1 \leq j \leq J.$$

Take  $J = 3$ . Since  $\phi$  is strictly non-decreasing and continuous, for  $s > 0$  and  $\alpha > 0$ , there exists unique  $x_\alpha(s)$  such that,  $x_\alpha(s) = \alpha\phi(x_\alpha(s) + s)$  and the function  $s \rightarrow x_\alpha(s)$  is non-decreasing and continuous on  $\mathbb{R}_+$ . Similarly there exists a unique  $S > 0$  satisfying the relation

$$S = x_{\alpha_1}(S) + x_{\alpha_2}(S) + x_{\alpha_3}(S).$$

From this one gets that  $y = (x_{\alpha_i}(S), 1 \leq i \leq 3)$  is the unique solution of Equation (6.4). One concludes that Equation (6.3) has also a unique solution.

## REFERENCES

- [1] Cédric Adjih, Philippe Jacquet, and Nikita Vvedenskaya, *Performance evaluation of a single queue under multi-user TCP/IP connections*, Tech. Report RR-4141, INRIA, March 2001, <http://hal.archives-ouvertes.fr/docs/00/07/24/84/PDF/RR-4141.pdf>.
- [2] D.J. Aldous, *Exchangeability and related topics*, École d'été de Probabilités de Saint-Flour XIII, Lecture Notes in Mathematics, vol. 1117, Springer Verlag, 1985, pp. 1–198.
- [3] F. Baccelli, D.R. McDonald, and J. Reynier, *A mean-field model for multiple TCP connections through a buffer implementing RED*, *Performance Evaluation* **49** (2004), 77–97.
- [4] N. Bellomo and S. Stöcker, *Development of Boltzmann models in mathematical biology*, Modelling in Applied Sciences: A Kinetic Theory Approach (N. Bellomo and M. Pulvirenti, eds.), Birkhauser, 2000, pp. 225–262.
- [5] Patrick Billingsley, *Convergence of probability measures*, Wiley series in probability and mathematical statistics, John Wiley & Sons Ltd, New York, 1968.
- [6] C. Cercignani, R. Illner, and M. Pulvirenti, *The mathematical theory of dilute gases*, Applied Mathematical Sciences, no. 196, Springer Verlag, New-York, 1994.
- [7] M. H. A. Davis, *Markov models and optimization*, Chapman & Hall, London, 1993.
- [8] Vincent Dumas, Fabrice Guillemin, and Philippe Robert, *A Markovian analysis of Additive-Increase Multiplicative-Decrease (AIMD) algorithms*, *Advances in Applied Probability* **34** (2002), no. 1, 85–111.
- [9] S. Ben Fred, T. Bonald, A. Proutiere, G. Régnié, and J. W. Roberts, *Statistical bandwidth sharing: a study of congestion at flow level*, *SIGCOMM Comput. Commun. Rev.* **31** (2001), no. 4, 111–122.
- [10] Carl Graham, *McKean-Vlasov Itô-Skorohod equations, and nonlinear diffusions with discrete jump sets*, *Stochastic Processes and their Applications* **40** (1992), 69–82.
- [11] ———, *Kinetic limits for large communication networks*, Modelling in Applied Sciences: A Kinetic Theory Approach (N. Bellomo and M. Pulvirenti, eds.), Birkhauser, 2000, pp. 317–370.
- [12] ———, *Chaoticity for multi-class systems and exchangeability within classes*, Preprint, 2008.

- [13] Carl Graham and Sylvie Méléard, *Chaos hypothesis for a system interaction through shared resources*, Probability Theory and Related Fields **100** (1994), 157–173.
- [14] C.P. Grünfeld, *Nonlinear kinetic models with chemical reactions*, Modelling in Applied Sciences: A Kinetic Theory Approach (N. Bellomo and M. Pulvirenti, eds.), Birkhauser, 2000, pp. 173–224.
- [15] Fabrice Guillemin, Philippe Robert, and Bert Zwart, *AIMD algorithms and exponential functionals*, Annals of Applied Probability **14** (2004), no. 1, 90–117.
- [16] F. I. Karpelevich, E. A. Pechersky, and Yu. M. Suhov, *Dobrushin's approach to queueing network theory*, J. Appl. Math. Stochastic Anal. **9** (1996), no. 4, 373–397. MR **MR1429262 (98d:60182)**
- [17] F. P. Kelly, A.K. Maulloo, and D.K.H. Tan, *Rate control in communication networks: shadow prices, proportional fairness and stability*, Journal of the Operational Research Society **49** (1998), 237–252.
- [18] F.P. Kelly and R.J. Williams, *Fluid model for a network operating under a fair bandwidth-sharing policy*, The Annals of Applied Probability **14** (2004), no. 3, 1055–1083.
- [19] Laurent Massoulié, *Structural properties of proportional fairness: Stability and insensitivity*, Annals of Applied Probability **17** (2007), no. 3, 809–839.
- [20] Laurent Massoulié and James Roberts, *Bandwidth sharing: Objectives and algorithms*, INFOCOM '99. Eighteenth Annual Joint Conference of the IEEE Computer and Communications Societies, 1999, pp. 1395–1403.
- [21] Sylvie Méléard, *Asymptotic behaviour of some interacting particle systems, McKean-Vlasov and Boltzmann models*, CIME summer school Montecatini Terme 1995 (D. Talay and L. Tubaro, eds.), Lecture Notes in Mathematics, no. 1627, Springer Verlag, 1996, pp. 42–95.
- [22] Teunis J. Ott, J.H.B. Kemperman, and Matt Mathis, *The stationary behavior of ideal TCP congestion avoidance*, Unpublished manuscript, August 1996.
- [23] Vivek Raghunathan and P. R. Kumar, *A counterexample in congestion control of wireless networks*, Performance Evaluation **64** (2007), no. 5, 399–418.
- [24] Philippe Robert, *Stochastic networks and queues*, Stochastic Modelling and Applied Probability Series, vol. 52, Springer, New-York, June 2003.
- [25] L. C. G. Rogers and David Williams, *Diffusions, Markov processes, and martingales. Vol. 1: Foundations*, second ed., John Wiley & Sons Ltd., Chichester, 1994.
- [26] J. Struckmaier, *Numerical simulation of the Boltzmann equation by particle methods*, Modelling in Applied Sciences: A Kinetic Theory Approach (N. Bellomo and M. Pulvirenti, eds.), Birkhauser, 2000, pp. 371–419.
- [27] A.S. Sznitman, *Topics in propagation of chaos*, École d'été de Saint-Flour, Lecture Notes in Maths, vol. 1464, Springer Verlag, 1989, pp. 167–243.

(C. Graham) UMR 7641 CNRS — ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU, FRANCE

*E-mail address:* [carl@cmplx.polytechnique.fr](mailto:carl@cmplx.polytechnique.fr)

(Ph. Robert) INRIA PARIS — ROCQUENCOURT, DOMAINE DE VOLUCEAU, 78153 LE CHESNAY, FRANCE

*E-mail address:* [Philippe.Robert@inria.fr](mailto:Philippe.Robert@inria.fr)

*URL:* <http://www-rocq.inria.fr/~robert>