

## Symmetric tensor decomposition

Jérôme Brachat, Pierre Comon, Bernard Mourrain, Elias P. Tsigaridas

► **To cite this version:**

Jérôme Brachat, Pierre Comon, Bernard Mourrain, Elias P. Tsigaridas. Symmetric tensor decomposition. *Linear Algebra and Applications*, Elsevier - Academic Press, 2010, 433 (11-12), pp.1851-1872. <inria-00355713v2>

**HAL Id: inria-00355713**

**<https://hal.inria.fr/inria-00355713v2>**

Submitted on 25 Jan 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SYMMETRIC TENSOR DECOMPOSITION

JEROME BRACHAT, PIERRE COMON, BERNARD MOURRAIN, AND ELIAS P. TSIGARIDAS

ABSTRACT. We present an algorithm for decomposing a symmetric tensor, of dimension  $n$  and order  $d$  as a sum of rank-1 symmetric tensors, extending the algorithm of Sylvester devised in 1886 for binary forms.

We recall the correspondence between the decomposition of a homogeneous polynomial in  $n$  variables of total degree  $d$  as a sum of powers of linear forms (Waring's problem), incidence properties on secant varieties of the Veronese Variety and the representation of linear forms as a linear combination of evaluations at distinct points. Then we reformulate Sylvester's approach from the dual point of view.

Exploiting this duality, we propose necessary and sufficient conditions for the existence of such a decomposition of a given rank, using the properties of Hankel (and quasi-Hankel) matrices, derived from multivariate polynomials and normal form computations. This leads to the resolution of polynomial equations of small degree in non-generic cases.

We propose a new algorithm for symmetric tensor decomposition, based on this characterization and on linear algebra computations with these Hankel matrices.

The impact of this contribution is two-fold. First it permits an efficient computation of the decomposition of any tensor of sub-generic rank, as opposed to widely used iterative algorithms with unproved global convergence (e.g. Alternate Least Squares or gradient descents). Second, it gives tools for understanding uniqueness conditions, and for detecting the rank.

## CONTENTS

1. Introduction	2
1.1. The binary case	3
1.2. Notation and preliminaries	3
2. Problem formulations	4
2.1. Polynomial decomposition	4
2.2. Veronese and Secant Varieties	5
2.3. Decomposition using duality	5
3. Hankel operators and quotient algebra	6
4. Truncated Hankel operators	8
5. Symmetric tensor decomposition algorithm	11
5.1. First Example	11
5.2. Second Example	13
6. Conclusions and future work	16
Acknowledgments.	16
References	16
Appendix A. Ternary cubics	19
Appendix B. An example of extreme rank	19

## 1. INTRODUCTION

Symmetric tensors show up in applications mainly as high-order derivatives of multivariate functions. For instance in Statistics, cumulant tensors are derivatives of the second characteristic function [42].

Tensors have been widely utilized in Electrical Engineering since the nineties [51], and in particular in Antenna Array Processing [22] [9] or Telecommunications [54] [8] [48] [25] [19]. Even earlier in the seventies, tensors have been used in Chemometrics [4] or Psychometrics [33].

Arithmetic complexity is also an important field where the understanding of tensor decompositions has made a lot of progress, especially third order tensors, which represent bilinear maps [35] [3] [50] [37].

Another important application field is Data Analysis. For instance, Independent Component Analysis, originally introduced for symmetric tensors whose rank did not exceed dimension [12] [6]. Now, it has become possible to estimate more factors than the dimension [23] [32]. In some applications, tensors may be symmetric only in some modes [14], or may not be symmetric nor have equal dimensions [10] [49]. Further numerous applications of tensor decompositions may be found in [10] [49].

Note that in some cases, tensors are encountered in the form of a collection of symmetric matrices [21] [26] [54] [47] [53], in which case they may enjoy symmetries in some modes but not in others. Conversely, some algorithms treat symmetric tensors as a collection of symmetric matrix slices [55] [57] [20].

The problem of decomposition of a symmetric tensor, that we consider in this paper, is a rank determinant problem which extends the Singular Value Decomposition (SVD) problem for symmetric matrices. This former method is an important tool in (numerical) linear algebra, which received a lot of attention and which is routinely used in many applications [27]. As exhibited above, the extension to general symmetric tensors also appears in many application domains. However, many issues either theoretical or algorithmic remains unsolved. Among solved problems, let us mention the determination of the minimal number of terms in the decomposition of a generic tensor [2], which is stated there in terms of a dual interpolation problem. See [31, chap. 2] and section 2 for the link between these two points of view. Among open problems are the determination of the maximal rank of tensors of given degree and dimension, or the determination of the stratification of the set of symmetric tensors by the rank. See however [11] for an answer in the binary case. For a detailed presentation of the symmetric tensor decomposition problem, from a projective algebraic geometric point of view, we refer to [31]. The properties of so-called catalecticant matrices, related to the apolar duality induced by the symmetric tensor associated with homogeneous polynomials of a given degree, are extensively studied.

In a disconnected way, probably because of language barrier, investigations of this problem in numerical analysis have been developed, inspired by the successful work on order 2 tensors, i.e. matrices. However, despite their obvious practical interest, numerical algorithms presently used in most scientific communities are suboptimal, in the sense that they either do not fully exploit symmetries [1], minimize different successive criteria sequentially [55] [20], or are iterative and lack a guarantee of global convergence [30] [46]. In addition, they often request the rank to be much smaller than generic. Among these popular methods, we refer to PARAFAC techniques [4], heavily applied for solving an ill-posed problem. . . Indeed contrarily to the matrix case, the set of symmetric tensors of rank  $\leq r$  is not closed, and its closure has singularities corresponding to tensors of rank  $> r$ . This explains why iterative numerical methods encounter difficulties to compute a tensor decomposition. For more details on open problems on symmetric tensors, see [15].

The goal of this paper is to describe a new algorithm able to decompose a symmetric tensor of arbitrary order and dimension into a sum of rank-one terms. The algorithm proposed in this paper is inspired from Sylvester's theorem [36], and extends its principle to larger dimensions. Using apolar duality on polynomials, we show that the symmetric tensor decomposition problem reduces to the decomposition of a linear form as a linear combination of evaluations at distinct points. We give a necessary and sufficient condition for the existence of a decomposition of rank  $r$ , based on rank conditions of Hankel operators or commutation properties. Instead of working, degree by degree, as in [31], we consider affine situations in order to treat at the same time the various homogeneous components. In the binary case, the decomposition problem can be solved directly by computing ranks of catalecticant. In higher dimension, this is not so simple. An extension step is required to find the decomposition. This leads to the resolution of a polynomial system of small degree, from which we deduce the decomposition by solving a simple eigenvalue problem, thanks to linear algebra manipulations.

The algorithm is not restricted to strictly sub-generic ranks as for the method proposed in [31][chap. 5]. In sub-generic cases, the decomposition is essentially unique (i.e. up to scale and permutation) when some rank conditions are satisfied. Our algorithm fully exploits this symmetry and provides a complete answer to the questions of uniqueness and computation, for any order [13].

In the following section, we recall the method deduced from Sylvester's theorem to decompose a binary form. In section 2, we give three equivalent formulations of the same problem, used and studied in different communities. In section 3, we develop the duality point of view, extending the notion of generalized additive decomposition, introduced in [31], to any dimension. Section 4 is devoted to the algebraic characterization of the extension property of linear forms, in terms of rank condition on multivariate Hankel operators, or on commutation properties. Finally in section 5, we describe the algorithm and give examples.

**1.1. The binary case.** The present contribution is a generalization of Sylvester's algorithm devised to decompose homogeneous polynomials in two variables into a sum of powers of linear forms [52] [11]. It is hence convenient to first recall the latter algorithm.

**Theorem 1.1** (Sylvester, 1886). *A binary quantic  $p(x_1, x_2) = \sum_{i=0}^d \binom{d}{i} c_i x_1^i x_2^{d-i}$  can be written as a sum of  $d^{\text{th}}$  powers of  $r$  distinct linear forms in  $\mathbb{C}$  as:*

$$(1) \quad p(x_1, x_2) = \sum_{j=1}^r \lambda_j (\alpha_j x_1 + \beta_j x_2)^d,$$

if and only if (i) there exists a vector  $\mathbf{q}$  of dimension  $r + 1$ , with components  $q_\ell$ , such that

$$(2) \quad \begin{bmatrix} c_0 & c_1 & \cdots & c_r \\ \vdots & & & \vdots \\ c_{d-r} & \cdots & c_{d-1} & c_d \end{bmatrix} \mathbf{q} = \mathbf{0}.$$

and (ii) the polynomial  $q(x_1, x_2) = \sum_{\ell=0}^r q_\ell x_1^\ell x_2^{r-\ell}$  admits  $r$  distinct roots, i.e. it can be written as  $q(x_1, x_2) = \prod_{j=1}^r (\beta_j^* x_1 - \alpha_j^* x_2)$ .

The proof of this theorem is constructive [52] [13] [15] and yields Algorithm 1.1. Given a binary polynomial  $p(x_1, x_2)$  of degree  $d$  with coefficients  $a_i = \binom{d}{i} c_i$ ,  $0 \leq i \leq d$ , define the Hankel matrix  $H[r]$  of dimensions  $d - r + 1 \times r + 1$  with entries  $H[r]_{ij} = c_{i+j-2}$ . Note that step 5 is a specialization only if the dimension of the right kernel is strictly larger than 1.

**1.2. Notation and preliminaries.** Let  $\mathbb{K}$  be an algebraically closed field (e.g.  $\mathbb{K} = \mathbb{C}$  the field of complex numbers). For a vector space  $E$ , its associated projective space is denoted  $\mathbb{P}(E)$ . For  $\mathbf{v} \in E - \{0\}$  its class in  $\mathbb{P}(E)$  is denoted  $\bar{\mathbf{v}}$ . Let  $\mathbb{P}^n$  be the projective space of the field  $\mathbb{K}^n$ .

If  $\mathbf{a} = (a_1, \dots, a_n)$  is a vector in  $\mathbb{N}^n$ , then  $|\mathbf{a}|$  is the sum of its elements, i.e.  $|\mathbf{a}| = \sum_{i=1}^n a_i$ . We also use the greek letters  $\alpha$  and  $\beta$  for vectors in  $\mathbb{N}^n$ . We denote by  $\mathbf{x}^\alpha$  the monomial  $x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ . For a set  $B = \{b_1, \dots, b_m\}$ , we denote by  $\langle B \rangle$ , respectively  $(B)$ , the corresponding vector space, resp. ideal.

Let  $R$  be the ring of polynomials  $\mathbb{K}[x_1, \dots, x_n]$ , while  $R_d$  will denote the ring of polynomials of (total) degree at most  $d$ . The set  $\{\mathbf{x}^\alpha\}_{|\alpha| \leq d} = \{x_1^{\alpha_1} \cdots x_n^{\alpha_n}\}_{\alpha_1 + \dots + \alpha_n \leq d}$  represents the elements of the monomial basis of the vector space  $R_d$  and contains  $\binom{n+d}{d}$  elements. Hereafter, the superscript  $h$  denotes the homogenization of a polynomial. We denote by  $S_d$  the vector space of homogeneous polynomials in  $n + 1$  variables  $x_0, x_1, \dots, x_n$ . This is also the symmetric  $d$ -th power  $S^d(E)$  where  $E = \langle x_0, \dots, x_n \rangle$ . The dehomogenization of a polynomial  $f \in S_d$  with respect to the variable  $x_0$  is denoted  $f^a := f(1, x_1, \dots, x_n)$ .

Duality is an important ingredient of our approach. For a comprehensive treatment of duality on multivariate polynomials, we refer the reader to [44]. Hereafter, for a  $\mathbb{K}$ -vector space  $E$  its dual  $E^* = \text{Hom}_{\mathbb{K}}(E, \mathbb{K})$  is the set of  $\mathbb{K}$ -linear forms from  $E$  to  $\mathbb{K}$ .

A basis of the dual space  $R_d^*$ , is the set of linear forms that compute the coefficients of a polynomial in the primal basis. It is denoted by  $\{\mathbf{d}^\alpha\}_{|\alpha| \leq d}$ .

We may identify  $R^*$  with the (vector) space of formal power series, i.e.  $\mathbb{K}[[\mathbf{d}]] = \mathbb{K}[[d_1, \dots, d_n]]$ . Any element  $\Lambda \in R^*$  can be decomposed as  $\Lambda = \sum_{\mathbf{a}} \Lambda(\mathbf{x}^{\mathbf{a}}) \mathbf{d}^{\mathbf{a}}$ . Typical elements of  $R^*$  are the linear forms that

---

**Algorithm 1.1:** BINARY FORM DECOMPOSITION
 

---

**Input:** A binary form  $p(x_0, x_1)$  of degree  $d$ .

**Output:** A decomposition of  $p$  as  $p(x_1, x_2) = \sum_{j=1}^r \lambda_j \mathbf{k}_j(\mathbf{x})^d$  with  $r$  minimal.

- (1) Initialize  $r = 0$
- (2) Increment  $r \leftarrow r + 1$
- (3) If the column rank of  $H[r]$  is full, then go to step 2
- (4) Else compute a basis  $\{\mathbf{k}_1, \dots, \mathbf{k}_l\}$  of the right kernel of  $H[r]$ .
- (5) Specialization:
  - Take a generic vector  $\mathbf{q}$  in the kernel, e.g.  $\mathbf{q} = \sum_i \mu_i \mathbf{k}_i$
  - Compute the roots of the associated polynomial  $q(x_1, x_2) = \sum_{\ell=0}^r q_\ell x_1^\ell x_2^{d-\ell}$ . Denote them  $(\beta_j, -\alpha_j)$ , where  $|\alpha_j|^2 + |\beta_j|^2 = 1$ .
  - If the roots are not distinct in  $\mathbb{P}^2$ , try another specialization. If distinct roots cannot be obtained, go to step 2.
  - Else if  $q(x_1, x_2)$  admits  $r$  distinct roots then compute coefficients  $\lambda_j$ ,  $1 \leq j \leq r$ , by solving the linear system below, where  $a_i$  denotes  $\binom{d}{i} c_i$

$$\begin{bmatrix} \alpha_1^d & \dots & \alpha_r^d \\ \alpha_1^{d-1} \beta_1 & \dots & \alpha_r^{d-1} \beta_r \\ \alpha_1^{d-2} \beta_1^2 & \dots & \alpha_r^{d-2} \beta_r^2 \\ \vdots & \vdots & \vdots \\ \beta_1^d & \dots & \beta_r^d \end{bmatrix} \lambda = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_d \end{bmatrix}$$

- (6) The decomposition is  $p(x_1, x_2) = \sum_{j=1}^r \lambda_j \mathbf{k}_j(\mathbf{x})^d$ , where  $\mathbf{k}_j(\mathbf{x}) = (\alpha_j x_1 + \beta_j x_2)$ .
- 

correspond to the evaluation at a point  $\zeta \in \mathbb{K}^n$ :

$$\mathbb{1}_\zeta : R \rightarrow \mathbb{K} \\ p \mapsto p(\zeta)$$

The decomposition of  $\mathbb{1}_\zeta$  in the basis  $\{\mathbf{d}^{\mathbf{a}}\}_{|\mathbf{a}| \leq d}$  is  $\mathbb{1}_\zeta = \sum_{\mathbf{a}} \zeta^{\mathbf{a}} \mathbf{d}^{\mathbf{a}}$ . Such an evaluation form can be composed with differentiation. In fact, if  $\theta(\partial_1, \dots, \partial_n)$  is a differential polynomial, then

$$\mathbb{1}_\zeta \circ \theta(\partial_1, \dots, \partial_n) : R \rightarrow \mathbb{K} \\ p \mapsto \theta(\partial_1, \dots, \partial_n)(p)(\zeta).$$

The dual space  $R^*$  has a natural structure of  $R$ -module [24] which is defined as follows: for all  $p \in R$ , and for all  $\Lambda \in R^*$  consider the linear operator

$$p \star \Lambda : R \rightarrow \mathbb{K} \\ q \mapsto \Lambda(pq).$$

In particular, we have  $x_i \star \mathbf{d}^{\mathbf{a}} = \begin{cases} d_1^{a_1} \dots d_{i-1}^{a_{i-1}} d_i^{a_i-1} d_{i+1}^{a_{i+1}} \dots d_n^{a_n} & \text{if } a_i > 0, \\ 0 & \text{otherwise.} \end{cases}$

## 2. PROBLEM FORMULATIONS

In this section, we present three different formulations of the same problem, that we consider in this paper.

**2.1. Polynomial decomposition.** A symmetric tensor  $[a_{j_0, \dots, j_n}]$  of order  $d$  and dimension  $n$  can be associated with a homogeneous polynomial  $f(\mathbf{x}) \in S_d$ :

$$(3) \quad f(\mathbf{x}) = \sum_{j_0 + j_1 + \dots + j_n = d} a_{j_0, j_1, \dots, j_n} x_0^{j_0} x_1^{j_1} \dots x_n^{j_n}.$$

Our goal is to compute a decomposition of  $f$  as a sum of  $d^{\text{th}}$  powers of linear forms, i.e.

$$(4) \quad f(\mathbf{x}) = \sum_{i=1}^r \lambda_i (k_{i,0} x_0 + k_{i,1} x_1 + \dots + k_{i,n} x_n)^d = \lambda_1 \mathbf{k}_1(\mathbf{x})^d + \lambda_2 \mathbf{k}_2(\mathbf{x})^d + \dots + \lambda_r \mathbf{k}_r(\mathbf{x})^d,$$

where  $\lambda_i \neq 0$ ,  $\mathbf{k}_i \neq 0$ , and  $r$  is the smallest possible. This minimal  $r$  is called the *rank* of  $f$ .

The *direct* approach to solve this decomposition problem, is the following. Consider the relation

$$f(\mathbf{x}) = \sum_{i=1}^r (k_{i,0}x_0 + k_{i,1}x_1 + \cdots + k_{i,n}x_n)^d,$$

where  $\mathbf{k}_i \neq 0$ . We assume that  $r$ , the rank, is known and the smallest possible. We consider the  $r(n+1)$  coefficients  $k_{i,j}$  of the linear forms as unknowns. We expand (symbolically) the right hand side of the relation. The two polynomials of the left and right hand sides are equal, thus by equating the coefficients of the same monomials we get a polynomial system in the coefficients  $k_{i,j}$ . This is an over-constrained polynomial system of  $\binom{n+d}{d}$  equations and  $r(n+1)$  unknowns. The polynomials of the system are homogeneous of degree  $d$  and the magnitude of their coefficients is  $\leq r \binom{n+d}{d}$ . This approach describes the problem of decomposition in a non-optimal way. It introduces  $r!$  redundant solutions, since every permutation of the linear forms is also a solution. Another drawback of this approach is that the polynomials involved are of high degree, that is,  $d$ . The reader can compare this with the degree 2 polynomial system, described in Section 4, containing the polynomials that we have to solve in order to extend the matrix.

In the following sections, we are going to describe a new method, which is much more efficient to solve this decomposition problem.

**2.2. Veronese and Secant Varieties.** Let us recall the well-known correspondence between the symmetric outer product decomposition and secant varieties for symmetric tensors. The set of symmetric tensors or homogeneous polynomials of the form  $\mathbf{k}(\mathbf{x})^d = (k_0x_0 + k_1x_1 + \cdots + k_nx_n)^d$  for  $\mathbf{k} = (k_0, k_1, \dots, k_n) \in \mathbb{K}^n$  is a closed algebraic set. Scaling the vector  $\mathbf{k}$  by a non-zero scalar  $\lambda$  yields a homogeneous polynomial scaled by  $\lambda^d$ . Thus, we can also consider this construction as a map  $\mathbf{k} \mapsto \mathbf{k}(\mathbf{x})^d$  from the projective space  $\mathbb{P}^{n-1}$  to the projective space of symmetric tensors:

$$\begin{aligned} \nu &: \mathbb{P}(S_1) &\rightarrow & \mathbb{P}(S_d) \\ & \mathbf{k}(\mathbf{x}) &\mapsto & \mathbf{k}(\mathbf{x})^d. \end{aligned}$$

The image of  $\nu$  is called the Veronese variety  $\mathcal{V}_{n,d}$  [29, 56]. Following this point of view, a tensor is of rank 1 if it corresponds to a point on the Veronese variety. A tensor is of rank  $\leq r$  if it is a linear combination of  $r$  tensors of rank 1. In other words, it is in the linear space spanned by  $r$  points of the Veronese variety. The closure of the  $r$ -dimensional linear space spanned by  $r$  points of the Veronese variety  $\mathcal{V}_{n,d}$  is called the  $(r-1)$ -secant variety of  $\mathcal{V}_{n,d}$  and denoted  $\mathcal{S}^{r-1}(\mathcal{V}_{n,d})$ . We refer the reader to [29, 56] for examples and general properties of these algebraic sets. In the non-symmetric case, the so-called Segre variety of the projective space of tensors is considered instead of the Veronese variety. It corresponds to the set of (possibly non-symmetric) tensors of rank 1.

For any  $f \in S_d - \{0\}$ , the smallest  $r$  such that  $\bar{f} \in \mathcal{S}^{r-1}(\mathcal{V}_{n,d})$  is called the *typical rank* or *border rank* of  $f$  [5, 15, 53].

**2.3. Decomposition using duality.** Let  $f, g \in S_d$ , where  $f = \sum_{|\alpha|=d} f_\alpha x_0^{\alpha_0} \cdots x_n^{\alpha_n}$  and  $g = \sum_{|\alpha|=d} g_\alpha x_0^{\alpha_0} \cdots x_n^{\alpha_n}$ . We define the apolar inner product on  $S_d$  as

$$\langle f, g \rangle = \sum_{|\alpha|=d} f_\alpha g_\alpha \binom{d}{\alpha_0, \dots, \alpha_n}^{-1}.$$

Using this non-degenerate inner product, we can associate an element of  $S_d$  with an element  $S_d^*$ , through the following map:

$$\begin{aligned} \tau : S_d &\rightarrow S_d^* \\ f &\mapsto \Lambda_f, \end{aligned}$$

where the linear form  $\Lambda_f$  is defined as  $\Lambda_f : g \mapsto \langle f, g \rangle$ . A simple calculation shows that  $\langle f, \mathbf{k}(\mathbf{x})^d \rangle = f(\mathbf{k})$  so that under this duality it holds that  $\tau(\mathbf{k}(\mathbf{x})^d) = \mathbf{1}_{\mathbf{k}} \in S_d^*$ . Moreover, under  $\tau$ , the polynomial  $f = \sum_{|\alpha|=d} c_\alpha \binom{d}{\alpha} \mathbf{x}^\alpha \in S_d$  is mapped to  $f^* = \sum_{|\alpha|=d} c_\alpha \mathbf{d}^\alpha \in S_d^*$ .

The problem of decomposition of  $f$  can then be restated as follows:

Given  $f^* \in S_d^*$ , find the minimal number of non-zero vectors  $\mathbf{k}_1, \dots, \mathbf{k}_r \in \mathbb{K}^{n+1}$  and non-zero scalars  $\lambda_1, \dots, \lambda_r \in \mathbb{K} - \{0\}$  such that

$$f^* = \sum_{i=1}^r \lambda_i \mathbb{1}_{\mathbf{k}_i}.$$

By scaling  $\mathbf{k}_i$  and multiplying  $\lambda_i$  by the inverse of the  $d^{\text{th}}$  power of this scaling factor, we may assume that the first non-zero coordinate of  $\mathbf{k}_i$  is 1.

**Definition 2.1.** We say that  $f^*$  is an affine decomposition if for every  $\mathbf{k}_i$  in the decomposition,  $\mathbf{k}_{i,0} \neq 0$ .

By a generic change of coordinates, any decomposition of  $f^*$  can be transformed into an affine decomposition. To any  $f^* \in S_d^*$ , we can associate an element in  $R_d^*$ , defined by  $\tilde{\Lambda}_f : p \in R_d \mapsto f^*(p^h)$ , where  $p^h$  is the homogenization in degree  $d$  of  $p$ . If  $f^*$  admits an affine decomposition with  $\mathbf{k}_{i,0} = 1$  then we also have that  $\tilde{\Lambda}_f$  coincides with the linear form

$$\tilde{\Lambda} = \sum_{i=1}^r \lambda_i \mathbb{1}_{\tilde{\mathbf{k}}_i}$$

up to degree  $d$ , where  $\tilde{\mathbf{k}}_i$  is the vector made of the last  $n$  coordinates of  $\mathbf{k}_i$ .

### 3. HANKEL OPERATORS AND QUOTIENT ALGEBRA

In this section, we recall the algebraic tools we will need to describe and analyze our algorithm. For any  $\Lambda \in R^*$ , we define the bilinear form  $Q_\Lambda$ , such that

$$Q_\Lambda : R \times R \rightarrow \mathbb{K} \\ (a, b) \mapsto \Lambda(ab).$$

The matrix of  $Q_\Lambda$  in the monomial basis, of  $R$  is  $\mathbb{Q}_\Lambda = (\Lambda(\mathbf{x}^{\alpha+\beta}))_{\alpha,\beta}$ , where  $\alpha, \beta \in \mathbb{N}^n$ .

For any  $\Lambda \in R^*$ , we define the Hankel operator  $H_\Lambda$  from  $R$  to  $R^*$  as

$$H_\Lambda : R \rightarrow R^* \\ p \mapsto p \star \Lambda.$$

The matrix of the linear operator  $H_\Lambda$  in the monomial basis, and in the dual basis,  $\{\mathbf{d}^\alpha\}$ , is  $\mathbb{H}_\Lambda = (\Lambda(\mathbf{x}^{\alpha+\beta}))_{\alpha,\beta}$ , where  $\alpha, \beta \in \mathbb{N}^n$ . The following relates the Hankel operators with the bilinear forms. For all  $a, b \in R$ , thanks to the  $R$ -module structure, it holds

$$Q_\Lambda(a, b) = \Lambda(ab) = a \cdot \Lambda(b) = b \cdot \Lambda(a) = H_\Lambda(a)(b) = H_\Lambda(b)(a).$$

In what follows we will identify  $H_\Lambda$  and  $Q_\Lambda$ .

**Definition 3.1.** Given  $B = \{b_1, \dots, b_r\}, B' = \{b'_1, \dots, b'_{r'}\} \subset R$  we define

$$H_\Lambda^{B, B'} : \langle B \rangle \rightarrow \langle B' \rangle^*,$$

as the restriction of  $H_\Lambda$  to the vector space  $\langle B \rangle$  and inclusion of  $R^*$  in  $\langle B' \rangle^*$ . Let  $\mathbb{H}_\Lambda^{B, B'} = (\Lambda(b_i b'_j))_{1 \leq i \leq r, 1 \leq j \leq r'}$ . If  $B' = B$ , we also use the notation  $H_\Lambda^B$  and  $\mathbb{H}_\Lambda^B$ .

If  $B, B'$  are linearly independent, then  $\mathbb{H}_\Lambda^{B, B'}$  is the matrix of  $H_\Lambda^{B, B'}$  in this basis  $\{b_1, \dots, b_r\}$  of  $\langle B \rangle$  and the dual basis of  $B'$  in  $\langle B' \rangle^*$ . The *catalecticant* matrices of [31] correspond to the case where  $B$  and  $B'$  are respectively the set of monomials of degree  $\leq k$  and  $\leq d - k$  ( $k = 0, \dots, d$ ).

From the definition of the Hankel operators, we can deduce that a polynomial  $p \in R$  belongs to the kernel of  $H_\Lambda$  if and only if  $p \star \Lambda = 0$ , which in turn holds if and only if for all  $q \in R$ ,  $\Lambda(pq) = 0$ .

**Proposition 3.2.** Let  $I_\Lambda$  be the kernel of  $H_\Lambda$ . Then,  $I_\Lambda$  is an ideal of  $R$ .

*Proof.* Let  $p_1, p_2 \in I_\Lambda$ . Then for all  $q \in R$ ,  $\Lambda((p_1 + p_2)q) = \Lambda(p_1q) + \Lambda(p_2q) = 0$ . Thus,  $p_1 + p_2 \in I_\Lambda$ . If  $p \in I_\Lambda$  and  $p' \in R$ , then for all  $q \in R$ , it holds  $\Lambda(pp'q) = 0$ . Thus  $pp' \in I_\Lambda$  and  $I_\Lambda$  is an ideal.  $\square$

Let  $\mathcal{A}_\Lambda = R/I_\Lambda$  be the quotient algebra of polynomials modulo the ideal  $I_\Lambda$ , which, as Proposition 3.2 states is the kernel of  $H_\Lambda$ . The rank of  $H_\Lambda$  is the dimension of  $\mathcal{A}_\Lambda$  as a  $\mathbb{K}$ -vector space.

A quotient algebra  $\mathcal{A}$  is Gorenstein if there exists a non-degenerate bilinear form  $Q$  on  $\mathcal{A}$ , such that for all polynomials  $f, g, h \in \mathcal{A}$  it holds that  $Q(f, gh) = Q(fg, h)$  or equivalently if there exists  $\Lambda \in \mathcal{A}^*$  such that  $(f, g) \in \mathcal{A} \times \mathcal{A} \mapsto \Lambda(fg)$  is non-degenerate. Equivalently,  $\mathcal{A}$  is Gorenstein iff  $\mathcal{A}^*$  is a free  $\mathcal{A}$ -module generated by one element  $\Lambda \in \mathcal{A}^*$ :  $\mathcal{A}^* = \mathcal{A} \star \Lambda$ . See e.g. [24] for more details. The set  $\mathcal{A} \star \Lambda$  is also called the inverse system generated by  $\Lambda$  [41].

**Proposition 3.3.** *The dual space  $\mathcal{A}_\Lambda^*$  of  $\mathcal{A}_\Lambda$ , can be identified with the set  $D = \{q \star \Lambda \mid q \in R\}$  and  $\mathcal{A}_\Lambda$  is a Gorenstein algebra.*

*Proof.* Let  $D = \{q \star \Lambda; q \in R\}$  be the inverse system generated by  $\Lambda$ . By definition,

$$D^\perp = \{p \in R \mid \forall q \in R, q \star \Lambda(p) = \Lambda(pq) = 0\}.$$

Thus  $D^\perp = I_\Lambda$ , which is the ideal of the kernel of  $H_\Lambda$  (Proposition 3.2). Since  $\mathcal{A}_\Lambda^* = I_\Lambda^\perp$  is the set of linear forms in  $R^*$  which vanish on  $I_\Lambda$ , we deduce that  $\mathcal{A}_\Lambda^* = I_\Lambda^\perp = D^{\perp\perp} = D$ .

As  $p \star \Lambda = 0$  implies  $p \in I_\Lambda$  or  $p \equiv 0$  in  $\mathcal{A}_\Lambda$ , this shows that  $\mathcal{A}_\Lambda^*$  is free rank 1  $\mathcal{A}_\Lambda$ -module (generated by  $\Lambda$ ). Thus  $\mathcal{A}_\Lambda$  is Gorenstein.  $\square$

**Definition 3.4.** *For any  $B \subset R$ , let  $B^+ = B \cup x_1 B \cup \dots \cup x_n B$  and  $\partial B = B^+ - B$ .*

**Proposition 3.5.** *Assume that  $\text{rank}(H_\Lambda) = r < \infty$  and let  $B = \{b_1, \dots, b_r\} \subset R$  such that  $\mathbb{H}_\Lambda^B$  is invertible. Then  $b_1, \dots, b_r$  is a basis of  $\mathcal{A}_\Lambda$ . If  $1 \in \langle B \rangle$  the ideal  $I_\Lambda$  is generated by  $\ker H_\Lambda^{B^+}$ .*

*Proof.* Let us first prove that  $\{b_1, \dots, b_r\} \cap I_\Lambda = \{0\}$ . Let  $p \in \langle b_1, \dots, b_r \rangle \cap I_\Lambda$ . Then  $p = \sum_i p_i b_i$  with  $p_i \in \mathbb{K}$  and  $\Lambda(p b_j) = 0$ . The second equation implies that  $\mathbb{H}_\Lambda^B \cdot \mathbf{p} = \mathbf{0}$ , where  $\mathbf{p} = [p_1, \dots, p_r]^t \in \mathbb{K}^r$ . Since  $\mathbb{H}_\Lambda^B$  is invertible, this implies that  $\mathbf{p} = \mathbf{0}$  and  $p = 0$ .

As a consequence, we deduce that  $b_1 \star \Lambda, \dots, b_r \star \Lambda$  are linearly independent elements of  $R^*$ . This is so, because otherwise there exists  $\mathbf{m} = [\mu_1, \dots, \mu_r]^T \neq \mathbf{0}$ , such that  $\mu_1(b_1 \star \Lambda) + \dots + \mu_r(b_r \star \Lambda) = (\mu_1 b_1 + \dots + \mu_r b_r) \star \Lambda = 0$ . As  $\{b_1, \dots, b_r\} \cap \text{Kernel}(H_\Lambda) = \{0\}$ , this yields a contradiction.

Consequently,  $\{b_1 \star \Lambda, \dots, b_r \star \Lambda\}$  span the image of  $H_\Lambda$ . For any  $p \in R$ , it holds that  $p \star \Lambda = \sum_{i=1}^r \mu_i (b_i \star \Lambda)$  for some  $\mu_1, \dots, \mu_r \in \mathbb{K}$ . We deduce that  $p - \sum_{i=1}^r \mu_i b_i \in I_\Lambda$ . This yields the decomposition  $R = B \oplus I_\Lambda$ , and shows that  $b_1, \dots, b_r$  is a basis of  $\mathcal{A}_\Lambda$ .

If  $1 \in \langle B \rangle$ , the ideal  $I_\Lambda$  is generated by the relations  $x_j b_k - \sum_{i=1}^r \mu_i^{j,k} b_i \in I_\Lambda$ . These are precisely in the kernel of  $H_\Lambda^{B^+}$ .  $\square$

**Proposition 3.6.** *If  $\text{rank}(H_\Lambda) = r < \infty$ , then  $\mathcal{A}_\Lambda$  is of dimension  $r$  over  $\mathbb{K}$  and there exist  $\zeta_1, \dots, \zeta_d \in \mathbb{K}^n$  where  $d \leq r$ , and  $p_i \in \mathbb{K}[\partial_1, \dots, \partial_n]$ , such that*

$$(5) \quad \Lambda = \sum_{i=1}^d \mathbb{1}_{\zeta_i} \circ p_i(\partial)$$

*Moreover the multiplicity of  $\zeta_i$  is the dimension of the vector space spanned the inverse system generated by  $\mathbb{1}_{\zeta_i} \circ p_i(\partial)$ .*

*Proof.* Since  $\text{rank}(H_\Lambda) = r$ , the dimension of the vector space  $\mathcal{A}_\Lambda$  is also  $r$ . Thus the number of zeros of the ideal  $I_\Lambda$ , say  $\{\zeta_1, \dots, \zeta_d\}$  is at most  $r$ , viz.  $d \leq r$ . We can apply the structure Theorem [24, Th. 7.34, p. 185] in order to get the decomposition.  $\square$

In characteristic 0, the inverse system of  $\mathbb{1}_{\zeta_i} \circ p_i(\partial)$  by  $p_i$  is isomorphic to the vector space generated by  $p_i$  and its derivatives of any order with respect to the variables  $\partial_i$ . In general characteristic, we replace the derivatives by the product by the "inverse" of the variables [44], [24].

**Definition 3.7.** *For  $f \in S^d$ , we call generalized decomposition of  $f^*$  a decomposition such that  $f^* = \sum_{i=1}^d \mathbb{1}_{\zeta_i} \circ p_i(\partial)$  where the sum for  $i = 1, \dots, d$  of the dimensions of the vector spaces spanned by the inverse system generated by  $\mathbb{1}_{\zeta_i} \circ p_i(\partial)$  is minimal. This minimal sum of dimensions is called the length of  $f$ .*

This definition extends the definition introduced in [31] for binary forms. The length of  $f^*$  is the rank of the corresponding Hankel operator  $H_\Lambda$ .



**Theorem 3.8.** Let  $\Lambda \in R^*$ .  $\Lambda = \sum_{i=1}^r \lambda_i \mathbb{1}_{\zeta_i}$  with  $\lambda_i \neq 0$  and  $\zeta_i$  distinct points of  $\mathbb{K}^n$ , iff  $\text{rank } H_\Lambda = r$  and  $I_\Lambda$  is a radical ideal.

*Proof.* If  $\Lambda = \sum_{i=1}^r \lambda_i \mathbb{1}_{\zeta_i}$ , with  $\lambda_i \neq 0$  and  $\zeta_i$  distinct points of  $\mathbb{K}^n$ . Let  $\{e_1, \dots, e_r\}$  be a family of interpolation polynomials at these points:  $e_i(\zeta_j) = 1$  if  $i = j$  and 0 otherwise. Let  $I_\zeta$  be the ideal of polynomials which vanish at  $\zeta_1, \dots, \zeta_r$ . It is a radical ideal. We have clearly  $I_\zeta \subset I_\Lambda$ . For any  $p \in I_\Lambda$ , and  $i = 1, \dots, r$ , we have  $p \star \Lambda(e_i) = \Lambda(p e_i) = p(\zeta_i) = 0$ , which proves that  $I_\Lambda = I_\zeta$  is a radical ideal. As the quotient  $\mathcal{A}_\Lambda$  is generated by the interpolation polynomials  $e_1, \dots, e_r$ ,  $H_\Lambda$  is of rank  $r$ .

Conversely, if  $\text{rank } H_\Lambda = r$ , by Proposition 3.6  $\Lambda = \sum_{i=1}^r \mathbb{1}_{\zeta_i} \circ p_i(\partial)$  with a polynomial of degree 0, since the multiplicity of  $\zeta_i$  is 1. This concludes the proof of the equivalence.  $\square$

In order to compute the zeroes of an ideal  $I_\Lambda$  when we know a basis of  $\mathcal{A}_\Lambda$ , we exploit the properties of the operators of multiplication in  $\mathcal{A}_\Lambda$ :  $M_a : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_\Lambda$ , such that  $\forall b \in \mathcal{A}_\Lambda, M_a(b) = ab$  and its transposed operator  $M_a^t : \mathcal{A}_\Lambda^* \rightarrow \mathcal{A}_\Lambda^*$ , such that for  $\forall \gamma \in \mathcal{A}_\Lambda^*, M_a^t(\gamma) = a \star \gamma$ .

The following proposition expresses a similar result, based on the properties of the duality.

**Proposition 3.9.** For any linear form  $\Lambda \in R^*$  such that  $\text{rank } H_\Lambda < \infty$  and any  $a \in \mathcal{A}_\Lambda$ , we have

$$(6) \quad H_{a \star \Lambda} = M_a^t \circ H_\Lambda$$

*Proof.* By definition,  $\forall p \in R, H_{a \star \Lambda}(p) = a p \star \Lambda = a \star (p \star \Lambda) = M_a^t \circ H_\Lambda(p)$ .  $\square$

We have the following well-known theorem:

**Theorem 3.10.** Assume that  $\mathcal{A}_\Lambda$  is a finite dimensional vector space. Then  $\Lambda = \sum_{i=1}^d \mathbb{1}_{\zeta_i} \circ p_i(\partial)$  for  $\zeta_i \in \mathbb{K}^n$  and  $p_i(\partial) \in \mathbb{K}[\partial_1, \dots, \partial_n]$  and

- the eigenvalues of the operators  $M_a$  and  $M_a^t$ , are given by  $\{a(\zeta_1), \dots, a(\zeta_r)\}$ .
- the common eigenvectors of the operators  $(M_{x_i}^t)_{1 \leq i \leq n}$  are (up to scalar)  $\mathbb{1}_{\zeta_i}$ .

*Proof.* [16, 17, 24]  $\square$

Using the previous proposition, one can recover the points  $\zeta_i \in \mathbb{K}^n$  by eigenvector computation as follows. Assume that  $B \subset R$  with  $|B| = \text{rank}(H_\Lambda)$ , then equation (6) and its transposition yield

$$\mathbb{H}_{a \star \Lambda}^B = \mathbb{M}_a^t \mathbb{H}_\Lambda^B = \mathbb{H}_\Lambda^B \mathbb{M}_a,$$

where  $\mathbb{M}_a$  is the matrix of multiplication by  $a$  in the basis  $B$  of  $\mathcal{A}_\Lambda$ . By Theorem 3.10, the common solutions of the generalized eigenvalue problem

$$(7) \quad (\mathbb{H}_{a \star \Lambda} - \lambda \mathbb{H}_\Lambda) \mathbf{v} = \mathbb{0}$$

for all  $a \in R$ , yield the common eigenvectors  $\mathbb{H}_\Lambda^B \mathbf{v}$  of  $\mathbb{M}_a^t$ , that is the evaluation  $\mathbb{1}_{\zeta_i}$  at the roots. Therefore, these common eigenvectors  $\mathbb{H}_\Lambda^B \mathbf{v}$  are up to a scalar, the vectors  $[b_1(\zeta_i), \dots, b_r(\zeta_i)]$  ( $i = 1, \dots, r$ ). Notice that it is sufficient to compute the common eigenvectors of  $(\mathbb{H}_{x_i \star \Lambda}, \mathbb{H}_\Lambda)$  for  $i = 1, \dots, n$

If  $\Lambda = \sum_{i=1}^d \lambda_i \mathbb{1}_{\zeta_i}$  ( $\lambda_i \neq 0$ ), then the roots are simple, and one eigenvector computation is enough: for any  $a \in R$ ,  $\mathbb{M}_a$  is diagonalizable and the generalized eigenvectors  $\mathbb{H}_\Lambda^B \mathbf{v}$  are, up to a scalar, the evaluation  $\mathbb{1}_{\zeta_i}$  at the roots.

#### 4. TRUNCATED HANKEL OPERATORS

Coming back to our problem of symmetric tensor decomposition,  $f = \sum_{|\alpha| \leq d} c_\alpha \binom{d}{\alpha} \mathbf{x}^\alpha \in R_d$  admits an affine decomposition of rank  $r$ , iff  $\Lambda(\mathbf{x}^\alpha) = c_\alpha$  for all  $|\alpha| \leq d$  where

$$\Lambda = \sum_{i=1}^r \lambda_i \mathbb{1}_{\zeta_i},$$

for some distinct  $\zeta_1, \dots, \zeta_r \in \mathbb{K}^n$  and some  $\lambda_i \in \mathbb{K} - \{0\}$ .

Then, by theorem 3.8,  $H_\Lambda$  is of rank  $r$  and  $I_\Lambda$  is radical.

Conversely, given  $H_\Lambda$  of rank  $r$  with  $I_\Lambda$  radical which coincides up to degree  $d$  with  $\Lambda_d$ , by proposition 3.6,  $\Lambda = \sum_{i=1}^r \lambda_i \mathbb{1}_{\zeta_i}$  and  $f$  can be decomposed as a sum of  $r$   $d^{\text{th}}$ -powers of linear forms.

The problem of decomposition of  $f$  can thus be reformulated as follows:

Given  $f^* \in R_d^*$  find the smallest  $r$  such that there exists  $\Lambda \in R^*$  which extends  $f^*$  with  $H_\Lambda$  of rank  $r$  and  $I_\Lambda$  a radical ideal.

In this section, we are going to characterize under which conditions  $f^*$  can be extended to  $\Lambda \in R^*$  with  $H_\Lambda$  is of rank  $r$ .

We need the following technical property on the bases of  $\mathcal{A}_\Lambda$ , that we will consider:

**Definition 4.1.** Let  $B$  be a subset of monomials in  $R$ . We say that  $B$  is connected to 1 if  $\forall m \in B$  either  $m = 1$  or there exists  $i \in [1, n]$  and  $m' \in B$  such that  $m = x_i m'$ .

Let  $B \subset R_d$  be a set of monomials of degree  $\leq d$ , connected to 1. We consider the formal Hankel matrix

$$\mathcal{H}_\Lambda^B = (h_{\alpha+\beta})_{\alpha, \beta \in B},$$

with  $h_\alpha = f^*(\mathbf{x}^\alpha) = c_\alpha$  if  $|\alpha| \leq d$  and otherwise  $h_\alpha$  is a variable. The set of all these new variables is denoted  $\mathbf{h}$ .

Suppose that  $\mathcal{H}_\Lambda^B$  is invertible in  $\mathbb{K}(\mathbf{h})$ , then we define the formal multiplication operators

$$\mathcal{M}_i^B(\mathbf{h}) := (\mathcal{H}_\Lambda^B)^{-1} \mathcal{H}_{x_i \star \Lambda}^B.$$

The following result characterizes the cases where  $\mathbb{K}[\mathbf{x}] = B \oplus I_\Lambda$ :

**Theorem 4.2.** Let  $B = \{\mathbf{x}^{\beta_1}, \dots, \mathbf{x}^{\beta_r}\}$  be a set of monomials of degree  $\leq d$ , connected to 1 and  $\Lambda$  be a linear form in  $\langle B \cdot B^+ \rangle_d^*$ . Let  $\Lambda(\mathbf{h})$  be the linear form of  $\langle B \cdot B^+ \rangle^*$  defined by  $\Lambda(\mathbf{h})(\mathbf{x}^\alpha) = \Lambda(\mathbf{x}^\alpha)$  if  $|\alpha| \leq d$  and  $h_\alpha \in \mathbb{K}$  otherwise. Then,  $\Lambda(\mathbf{h})$  admits an extension  $\tilde{\Lambda} \in R^*$  such that  $H_{\tilde{\Lambda}}$  is of rank  $r$  with  $B$  a basis of  $\mathcal{A}_{\tilde{\Lambda}}$  iff

$$(8) \quad \mathcal{M}_i^B(\mathbf{h}) \circ \mathcal{M}_j^B(\mathbf{h}) - \mathcal{M}_j^B(\mathbf{h}) \circ \mathcal{M}_i^B(\mathbf{h}) = 0 \quad (1 \leq i < j \leq n)$$

and  $\det(\mathcal{H}_\Lambda^B)(\mathbf{h}) \neq 0$ . Moreover, such a  $\tilde{\Lambda}$  is unique.

*Proof.* If there exists  $\tilde{\Lambda} \in R^*$  which extends  $\Lambda(\mathbf{h})$ , with  $H_{\tilde{\Lambda}}$  of rank  $r$  then the tables of multiplications by the variables  $x_i$  are  $M_i = (\mathcal{H}_\Lambda^B)^{-1} \mathcal{H}_{x_i \star \Lambda}^B$  (proposition 3.9) and they commute.

Conversely suppose that these matrices commute. Then by [43], we have  $\mathbb{K}[\mathbf{x}] = \langle B \rangle \oplus (K)$ , where  $K$  is the vector space generated by the border relations  $x_i m - \mathcal{M}_i(m)$  for  $m \in B$  and  $i = 1, \dots, n$ . Let  $\pi_B$  be the projection of  $R$  on  $\langle B \rangle$  along  $(K)$ .

We define  $\tilde{\Lambda} \in R^*$  as follows:  $\forall p \in \mathbb{R}, \tilde{\Lambda}(p) = \Lambda(p(\mathcal{M})(1))$  where  $p(\mathcal{M})$  is the operator obtained by substitution of the variables  $x_i$  by the commuting operators  $\mathcal{M}_i$ . Notice that  $p(\mathcal{M})$  is also the operator of multiplication by  $p$  modulo  $(K)$ .

By construction,  $(K) \subset \ker H_{\tilde{\Lambda}}$  and  $B$  is a generating set of  $\mathcal{A}_{\tilde{\Lambda}}$ .

Let us prove by induction on the degree of  $b \in B$  that for all  $b' \in B$ , we have  $\Lambda(bb') = \Lambda(b(M)(b'))$ . The property is true for  $b = 1$ . As  $B$  is connected to 1, if  $b \neq 1$ , then  $b = x_i b'$  for some variable  $x_i$  and some element  $b' \in B$  of degree smaller than  $b$ . By construction of the operators  $M_i$ , we have  $\Lambda(x_i b'' b') = \Lambda(b'' M_i(b'))$ . By induction hypothesis, we deduce that  $\Lambda(bb') = \Lambda(b''(M) \circ M_i(b')) = \Lambda(b(M)(b'))$ . As  $b' = b'(M)(1)$  for all  $b' \in B$  (the multiplication of 1 by  $b$  is represented by  $b \in \langle B \rangle$  modulo  $(K)$ ), we deduce that

$$\Lambda(bb') = \Lambda(b''(M) \circ M_i(b')) = \Lambda(b(M) \circ b'(M)(1)) = \Lambda((bb')(M)(1)) = \tilde{\Lambda}(bb').$$

This shows that  $\Lambda = \tilde{\Lambda}$  on  $B \cdot B$ . As  $\det(\mathcal{H}_\Lambda^B) \neq 0$ , we deduce that  $B$  is a basis of  $\mathcal{A}_{\tilde{\Lambda}}$  and that  $H_{\tilde{\Lambda}}$  is of rank  $r$ .

If there exists another  $\Lambda' \in R^*$  which extends  $\Lambda(\mathbf{h}) \in \langle B \cdot B^+ \rangle^*$  with rank  $H_{\Lambda'} = r$ , by proposition 3.5,  $\ker H_{\Lambda'}$  is generated by  $\ker H_{\tilde{\Lambda}}^{B \cdot B^+}$  and thus coincides with  $\ker H_{\tilde{\Lambda}}$ . As  $\Lambda'$  coincides with  $\tilde{\Lambda}$  on  $B$ , the two elements of  $R^*$  must be equal. This ends the proof of the theorem.  $\square$

The degree of these commutation relations is at most 2 in the coefficients of the multiplications matrices  $\mathcal{M}_i$ . A direct computation yields the following, for  $m \in B$ :

- If  $x_i, m \in B, x_j m \in B$  then  $(\mathcal{M}_i^B \circ \mathcal{M}_j^B - \mathcal{M}_j^B \circ \mathcal{M}_i^B)(m) \equiv 0$  in  $\mathbb{K}(\mathbf{h})$ .
- If  $x_i m \in B, x_j m \notin B$  then  $(\mathcal{M}_i^B \circ \mathcal{M}_j^B - \mathcal{M}_j^B \circ \mathcal{M}_i^B)(m)$  is of degree 1 in the coefficients of  $\mathcal{M}_i, \mathcal{M}_j$ .
- If  $x_i m \notin B, x_j m \notin B$  then  $(\mathcal{M}_i^B \circ \mathcal{M}_j^B - \mathcal{M}_j^B \circ \mathcal{M}_i^B)(m)$  is of degree 2 in the coefficients of  $\mathcal{M}_i, \mathcal{M}_j$ .

We are going to give an equivalent characterization of the extension property, based on rank conditions.

**Theorem 4.3.** Let  $B = \{\mathbf{x}^{\beta_1}, \dots, \mathbf{x}^{\beta_r}\}$  be a set of monomials of degree  $\leq d$ , connected to 1. Then, the linear form  $f^* \in S_d^*$  admits an extension  $\Lambda \in R^*$  such that  $H_\Lambda$  is of rank  $r$  with  $B$  a basis of  $\mathcal{A}_\Lambda$  iff there exists an  $\mathbf{h}$  such that all  $(r+1) \times (r+1)$  minors of  $\mathcal{H}_\Lambda^{B^+}(\mathbf{h})$  vanish and  $\det(\mathcal{H}_\Lambda^B(\mathbf{h})) \neq 0$ .

*Proof.* Clearly, if there exists  $\Lambda \in R^*$  which extends  $f^* \in S_d^*$  with  $H_\Lambda$  of rank  $r$ , then all  $(r+1) \times (r+1)$  minors of  $\mathcal{H}_\Lambda^{B^+}(\mathbf{h})$  vanish.

Conversely, if  $\mathcal{H}_\Lambda^{B^+}(\mathbf{h})$  and  $\mathcal{H}_\Lambda^B(\mathbf{h})$  are of rank  $r$ , by [40, Theorem 1.4] there exists a unique  $\tilde{\Lambda} \in R^*$  such that  $H_{\tilde{\Lambda}}$  is of rank  $r$ , and which coincides with  $\Lambda$  on  $\langle B^+ \cdot B^+ \rangle$ .  $\square$

**Proposition 4.4.** Let  $B = \{\mathbf{x}^{\beta_1}, \dots, \mathbf{x}^{\beta_r}\}$  be a set of monomials of degree  $\leq d$ , connected to 1. Then, the linear form  $f^* \in S_d^*$  admits an extension  $\Lambda \in R^*$  such that  $H_\Lambda$  is of rank  $r$  with  $B$  a basis of  $\mathcal{A}_\Lambda$  iff

$$(9) \quad \mathbb{H}_\Lambda^{B^+} = \begin{pmatrix} \mathbb{H} & \mathbb{G} \\ \mathbb{G}^t & \mathbb{J} \end{pmatrix},$$

with  $\mathbb{H} = \mathbb{H}_\Lambda^B$  and

$$(10) \quad \mathbb{G} = \mathbb{H}\mathbb{W}, \mathbb{J} = \mathbb{W}^t \mathbb{H}\mathbb{W}.$$

for some matrix  $\mathbb{W} \in \mathbb{K}^{B \times \partial B}$ .

*Proof.* According to theorem 4.3,  $f^* \in S_d^*$  admits a (unique) extension  $\Lambda \in R^*$  such that  $H_\Lambda$  is of rank  $r$  with  $B$  a basis of  $\mathcal{A}_\Lambda$ , iff  $H^{B^+}$  is of rank  $r$ . Let us decompose  $H^{B^+}$  as (9) with  $\mathbb{H} = \mathbb{H}_\Lambda^B$ .

If we have  $\mathbb{G} = \mathbb{H}\mathbb{W}$ ,  $\mathbb{J} = \mathbb{W}^t \mathbb{H}\mathbb{W}$ , then

$$\begin{pmatrix} \mathbb{H} & \mathbb{H}\mathbb{W} \\ \mathbb{W}^t \mathbb{H} & \mathbb{W}^t \mathbb{H}\mathbb{W} \end{pmatrix}$$

is clearly of rank  $\leq \text{rank } \mathbb{H}$ .

Conversely, suppose that  $\mathbb{H}_\Lambda^{B^+} = \text{rank } \mathbb{H}$ . This implies that the image of  $\mathbb{G}$  is in the image of  $\mathbb{H}$ . Thus, there exists  $\mathbb{W} \in \mathbb{K}^{B \times \partial B}$  such that  $\mathbb{G} = \mathbb{H}\mathbb{W}$ . Without loss of generality, we can assume that the  $r$  first columns of  $\mathbb{H}$  ( $r = \text{rank } \mathbb{H}$ ) are linearly independent. Assume that we choose  $\mathbb{W}$  such that the  $i^{\text{th}}$  column of  $\mathbb{G}$  is the linear combination of the  $r$  first columns with coefficients corresponding to the  $i$  column  $\mathbb{W}_i$  of  $\mathbb{W}$ . As  $\text{rank } \mathbb{H}_\Lambda^{B^+} = r$  the same relation holds for the whole column of this matrix. Thus we have  $\mathbb{J} = \mathbb{G}^t \mathbb{W} = \mathbb{W}^t \mathbb{H}\mathbb{W}$ .  $\square$

Notice that if  $\mathbb{H}$  is invertible,  $\mathbb{W}$  is uniquely determined. In this case, we easily check that  $\ker \mathbb{H}_\Lambda^{B^+} = \begin{pmatrix} \mathbb{W} \\ -\mathbb{I} \end{pmatrix}$ .

This leads to the following system in the variables  $\mathbf{h}$  and the coefficients  $\mathbf{w}$  of matrix  $\mathbb{W}$ . It characterizes the linear forms  $f^* \in S_d^*$  that admit an extension  $\Lambda \in R^*$  such that  $H_\Lambda$  is of rank  $r$  with  $B$  a basis of  $\mathcal{A}_\Lambda$ .

$$(11) \quad \mathcal{H}_\Lambda^{B, \partial B}(\mathbf{h}) - \mathcal{H}_\Lambda^B(\mathbf{h}) \mathbb{W}(\mathbf{w}) = 0, \quad \mathcal{H}_\Lambda^{\partial B, \partial B}(\mathbf{h}) - \mathbb{W}^t(\mathbf{w}) \mathcal{H}_\Lambda^B(\mathbf{h}) \mathbb{W}(\mathbf{w}) = 0$$

with  $\det(\mathcal{H}_\Lambda^B(\mathbf{h})) \neq 0$ .

The matrix  $\mathcal{H}_\Lambda^{B^+}$  is a quasi-Hankel matrix [44], whose structure is imposed by equality (linear) constraints on its entries. If  $\mathbb{H}$  is known (ie.  $B \times B \subset R_d$ , the number of independent parameters in  $\mathcal{H}_\Lambda^{B, B^+}(\mathbf{h})$  or in  $\mathbb{W}$  is the number of monomials in  $B \times \partial B - R_d$ ). By Proposition 4.4, the rank condition is equivalent to the quadratic relations  $\mathbb{J} - \mathbb{W}^t \mathbb{H}^t \mathbb{W} = 0$  in these unknowns.

If  $\mathbb{H}$  is not completely known, the number of parameters in  $\mathbb{H}$  is the number of monomials in  $B \times B - R_d$ . The number of independent parameters in  $\mathcal{H}_\Lambda^{B, \partial B}(\mathbf{h})$  or in  $\mathbb{W}$  is then  $B \times \partial B - R_d$ .

The system (11) is composed of linear equations deduced from quasi-Hankel structure, quadratic relations for the entries in  $B \times \partial B$  and cubic relations for the entries in  $B \times \partial B$  in the unknown parameters  $\mathbf{h}$  and  $\mathbf{w}$ .

We are going to use explicitly these characterizations in the new algorithm we propose for minimal tensor decomposition.

## 5. SYMMETRIC TENSOR DECOMPOSITION ALGORITHM

The algorithm that we will present for decomposing a symmetric tensor as sum of rank 1 symmetric tensors generalizes the algorithm of Sylvester [52], devised for dimension 2 tensors, see also [11].

Consider the homogeneous polynomial  $f(\mathbf{x})$  in (3) that we want to decompose. We may assume without loss of generality, that for at least one variable, say  $x_0$ , all its coefficients in the decomposition are non zero, i.e.  $k_{i,0} \neq 0$ , for  $1 \leq i \leq r$ . We dehomogenize  $f$  with respect to this variable and we denote this polynomial by  $f^a := f(1, x_1, \dots, x_n)$ . We want to decompose the polynomial  $f^a(\mathbf{x}) \in R_d$  as a sum of powers of linear forms, i.e.

$$f(\mathbf{x}) = \sum_{i=1}^r \lambda_i (1 + k_{i,1}x_1 + \dots + k_{i,n}x_n)^d = \sum_{i=1}^r \lambda_i \mathbf{k}_i(\mathbf{x})^d$$

Equivalently, we want to decompose its corresponding dual element  $f^* \in R_d^*$  as a linear combination of evaluations over the distinct points  $\mathbf{k}_i := (k_{i,1}, \dots, k_{i,n})$ :

$$f^* = \sum_{i=1}^r \lambda_i \mathbb{1}_{\mathbf{k}_i}$$

(we refer the reader to the end of Section 2.3).

Assume that we know the value of  $r$ . As we have seen previously, knowing the value of  $\Lambda$  on polynomials of degree high enough, allows us to compute the table of multiplications modulo the kernel of  $\mathbb{H}_\Lambda$ . By Theorem 3.10, solving the generalized eigenvector problem  $(\mathbb{H}_{x_1 \star \Lambda} - \lambda \mathbb{H}_\Lambda)\mathbf{v} = \mathbb{O}$ , we will recover the points of evaluation  $\mathbf{k}_i$ . By solving a linear system, we will then deduce the value of  $\lambda_1, \dots, \lambda_r$ . Thus, the goal of the following algorithm is to extend  $f^*$  on a large enough set of polynomials, in order to be able to run this eigenvalue computation.

---

### Algorithm 5.1: SYMMETRIC TENSOR DECOMPOSITION

---

**Input:** A homogeneous polynomial  $f(x_0, x_1, \dots, x_n)$  of degree  $d$ .

**Output:** A decomposition of  $f$  as  $f = \sum_{i=1}^r \lambda_i \mathbf{k}_i(\mathbf{x})^d$  with  $r$  minimal.

- Compute the coefficients of  $f^*$ :  $c_\alpha = a_\alpha \binom{d}{\alpha}^{-1}$ , for  $|\alpha| \leq d$ ;
  - $r := 1$ ;
  - **Repeat**
    - (1) Compute a set  $B$  of monomials of degree  $\leq d$  connected to 1 with  $|B| = r$ ;
    - (2) Find parameters  $\mathbf{h}$  s.t.  $\det(\mathbb{H}_\Lambda^B) \neq 0$  and the operators  $\mathbb{M}_i = \mathbb{H}_{x_i \Lambda}^B (\mathbb{H}_\Lambda^B)^{-1}$  commute.
    - (3) If there is no solution, restart the loop with  $r := r + 1$ .
    - (4) Else compute the  $n \times r$  eigenvalues  $\zeta_{i,j}$  and the eigenvectors  $\mathbf{v}_j$  s.t.  $\mathbb{M}_i \mathbf{v}_j = \zeta_{i,j} \mathbf{v}_j$ ,  
 $i = 1, \dots, n, j = 1, \dots, r$ .
  - until** the eigenvalues are simple.
  - Solve the linear system in  $(\nu_j)_{j=1, \dots, k}$ :  $\Lambda = \sum_{j=1}^r \nu_j \mathbf{1}_{\zeta_j}$  where  $\zeta_j \in \mathbb{K}^n$  are the eigenvectors found in step 4.
- 

The critical part in this algorithm is the completion of step 2. Instead of the commutation relations, one can use the result of Proposition 4.4.

**5.1. First Example.** The example that follows will make the steps of the algorithm clearer.

- (1) Convert the symmetric tensor to the corresponding homogeneous polynomial.

Assume that a tensor of dimension 3 and order 5, or equivalently a 3-way array of dimension 5, corresponds to the following homogeneous polynomial

$$f = -1549440 x_0 x_1 x_2^3 + 2417040 x_0 x_1^2 x_2^2 + 166320 x_0^2 x_1 x_2^2 - 829440 x_0 x_1^3 x_2 - 5760 x_0^3 x_1 x_2 - 222480 x_0^2 x_1^2 x_2 + 38 x_0^5 - 497664 x_1^5 - 1107804 x_2^5 - 120 x_0^4 x_1 + 180 x_0^4 x_2 + 12720 x_0^3 x_1^2 + 8220 x_0^3 x_2^2 - 34560 x_0^2 x_1^3 - 59160 x_0^2 x_2^3 + 831840 x_0 x_1^4 + 442590 x_0 x_2^4 - 5591520 x_1^4 x_2 + 7983360 x_1^3 x_2^2 - 9653040 x_1^2 x_2^3 + 5116680 x_1 x_2^4.$$

The minimum decomposition of the polynomial as a sum of powers of linear forms is

$$(x_0 + 2x_1 + 3x_2)^5 + (x_0 - 2x_1 + 3x_2)^5 + \frac{1}{3}(x_0 - 12x_1 - 3x_2)^5 + \frac{1}{5}(x_0 + 12x_1 - 13x_2)^5,$$

that is, the corresponding tensor is of rank 4.

- (2) Compute the actual number of variables needed.

For algorithms computing the so-called number of *essential* variables, the reader may refer to the work of Oldenburger [45] or Carlini [7].

In our example the number of essential variable is 3, so we have nothing to do.

- (3) Compute the matrix of the quotient algebra.

We form a  $\binom{n+d-1}{d} \times \binom{n+d-1}{d}$  matrix, the rows and the columns of which correspond to the coefficients of the polynomial in the dual base. The map for this is

$$a_{j_0 j_1 \dots j_n} \mapsto c_{j_0 j_1 \dots j_n} := a_{j_0 j_1 \dots j_n} \binom{d}{j_0, \dots, j_n}^{-1},$$

where  $a_{d_0 d_1 \dots d_n}$  is the coefficient of the monomial  $x_0^{j_0} \dots x_n^{j_n}$  in  $f$ . Recall that, since the polynomial is homogeneous,  $\sum_{i=1}^n j_i = d$ .

This matrix is called quasi-Hankel [44] or Catalecticant [31].

Part of the corresponding matrix follows. The whole matrix is  $21 \times 21$ . We show only the  $10 \times 10$  principal minor.

	1	$x_1$	$x_2$	$x_1^2$	$x_1 x_2$	$x_2^2$	$x_1^3$	$x_1^2 x_2$	$x_1 x_2^2$	$x_2^3$
1	38	-24	36	1272	-288	822	-3456	-7416	5544	-5916
$x_1$	-24	1272	-288	-3456	-7416	5544	166368	-41472	80568	-77472
$x_2$	36	-288	822	-7416	5544	-5916	-41472	80568	-77472	88518
$x_1^2$	1272	-3456	-7416	166368	-41472	80568	-497664	-1118304	798336	-965304
$x_1 x_2$	-288	-7416	5544	-41472	80568	-77472	-1118304	798336	-965304	1023336
$x_2^2$	822	5544	-5916	80568	-77472	88518	798336	-965304	1023336	-1107804
$x_1^3$	-3456	166368	-41472	-497664	-1118304	798336	$h_{6,0,0}$	$h_{5,1,0}$	$h_{4,2,0}$	$h_{3,3,0}$
$x_1^2 x_2$	-7416	-41472	80568	-1118304	798336	-965304	$h_{5,1,0}$	$h_{4,2,0}$	$h_{3,3,0}$	$h_{2,4,0}$
$x_1 x_2^2$	5544	80568	-77472	798336	-965304	1023336	$h_{4,2,0}$	$h_{3,3,0}$	$h_{2,4,0}$	$h_{1,5,0}$
$x_2^3$	-5916	-77472	88518	-965304	1023336	-1107804	$h_{3,3,0}$	$h_{2,4,0}$	$h_{1,5,0}$	$h_{0,6,0}$

Notice that we do not know the elements in some positions of the matrix. In general we do not know the elements that correspond to monomials with (total) degree higher than 5.

- (4) Extract a principal minor of full rank.

We should re-arrange the rows and the columns of the matrix so that there is a principal minor of full rank,  $R$ . We call this minor  $\Delta_0$ . In order to do that we try to put the matrix in row echelon form, using elementary row and column operations.

In our example the  $4 \times 4$  principal minor is of full rank, so there is no need for re-arranging the matrix. The matrix  $\Delta_0$  is

$$\Delta_0 = \begin{bmatrix} 38 & -24 & 36 & 1272 \\ -24 & 1272 & -288 & -3456 \\ 36 & -288 & 822 & -7416 \\ 1272 & -3456 & -7416 & 166368 \end{bmatrix}$$

Notice that the columns of the matrix correspond to the monomials  $\{1, x_1, x_2, x_1^2\}$ .

- (5) We compute the “shifted” matrix  $\Delta_1 = x_1 \Delta_0$ .

The columns of  $\Delta_0$  correspond to set of some monomials, say  $\{\mathbf{x}^\alpha\}$  where  $\alpha \in \mathbb{N}^n$ . The columns of  $\Delta_1$  correspond to the set of monomials  $\{x_1 \mathbf{x}^\alpha\}$ .

The shifted matrix  $\Delta_1$  is

$$\Delta_1 = \begin{bmatrix} -24 & 1272 & -288 & -3456 \\ 1272 & -3456 & -7416 & 166368 \\ -288 & -7416 & 5544 & -41472 \\ -3456 & 166368 & -41472 & -497664 \end{bmatrix}$$

Notice that the columns correspond to the monomials  $\{x_1, x_1^2, x_1 x_2, x_1^3\}$ , which are just the corresponding monomials of the columns of  $\Delta_0$ , i.e.  $\{1, x_1, x_2, x_1^2\}$ , multiplied by  $x_1$ .

We assume for the moment that all the elements of the matrices  $\Delta_0$  and  $\Delta_1$  are known. If this is not the case, then we can compute the unknown entries of the matrix, using either necessary and sufficient conditions of the quotient algebra, e.g. it holds that  $M_{x_i} M_{x_j} - M_{x_j} M_{x_i} = 0$  [43] for any  $i, j \in \{1, \dots, n\}$ . There are other algorithms to extend a moment matrix, e.g. [18, 38, 39].

- (6) We solve the equation  $(\Delta_1 - \lambda \Delta_0)X = 0$ .

We solve the generalized eigenvalue/eigenvector problem using one of the well-known techniques [28]. We normalize the elements of the eigenvectors so that the first element is 1, and we read the solutions from the coordinates of the (normalized) eigenvectors.

The normalized eigenvectors of the generalized eigenvalue problem are

$$\begin{bmatrix} 1 \\ -12 \\ -3 \\ 144 \end{bmatrix}, \begin{bmatrix} 1 \\ 12 \\ -13 \\ 144 \end{bmatrix}, \begin{bmatrix} 1 \\ -2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$$

The coordinates of the eigenvectors correspond to the elements  $\{1, x_1, x_2, x_1^2\}$ . Thus, we can recover the coefficients of  $x_1$  and  $x_2$  in the decomposition from coordinates of the eigenvectors.

Recall that the coefficients of  $x_0$  are considered to be one. Thus, The polynomial admits a decomposition

$$f = \ell_1(x_0 - 12x_1 - 3x_2)^5 + \ell_2(x_0 + 12x_1 - 13x_2)^5 + \ell_3(x_0 - 2x_1 + 3x_2)^5 + \ell_4(x_0 + 2x_1 + 3x_2)^5$$

It remains to compute  $\ell_i$ 's. We can do this easily by solving an over-determined linear system, which we know that always has a solution, since the decomposition exists. Doing that, we deduce that  $\ell_1 = 3$ ,  $\ell_2 = 15$ ,  $\ell_3 = 15$  and  $\ell_4 = 5$ .

**5.2. Second Example.** One of the assumptions that the previous example fulfills is that all the entries of the matrices needed for the computations are known. However, this is not always the case as the following example shows.

- (1) Convert the symmetric tensor to the corresponding homogeneous polynomial.

Consider a tensor of dimension 3 and order 4, that corresponds to the following homogeneous polynomial

$$f = 79x_0x_1^3 + 56x_0^2x_2^2 + 49x_1^2x_2^2 + 4x_0x_1x_2^2 + 57x_0^3x_1,$$

the rank of which is 6.

- (2) Compute the actual number of variables needed.

In our example the number of essential variables is 3, so we have nothing to do.

- (3) Compute the matrix of the quotient algebra.

The matrix is  $15 \times 15$ .

	1	$x_1$	$x_2$	$x_1^2$	$x_1x_2$	$x_2^2$	$x_1^3$	$x_1^2x_2$	$x_1x_2^2$	$x_2^3$	$x_1^4$	$x_1^3x_2$	$x_1^2x_2^2$	$x_1x_2^3$	$x_2^4$
1	0	$\frac{57}{4}$	0	0	0	$\frac{28}{3}$	$\frac{79}{4}$	0	$\frac{1}{3}$	0	0	0	$\frac{49}{6}$	0	0
$x_1$	$\frac{57}{4}$	0	0	$\frac{79}{4}$	0	$\frac{1}{3}$	0	0	$\frac{49}{6}$	0	$h_{500}$	$h_{410}$	$h_{320}$	$h_{230}$	$h_{140}$
$x_2$	0	0	$\frac{28}{3}$	0	$\frac{1}{3}$	0	0	$\frac{49}{6}$	0	0	$h_{410}$	$h_{320}$	$h_{230}$	$h_{140}$	$h_{050}$
$x_1^2$	0	$\frac{79}{4}$	0	0	0	$\frac{49}{6}$	$h_{500}$	$h_{410}$	$h_{320}$	$h_{230}$	$h_{600}$	$h_{510}$	$h_{420}$	$h_{330}$	$h_{240}$
$x_1x_2$	0	0	$\frac{1}{3}$	0	$\frac{49}{6}$	0	$h_{410}$	$h_{320}$	$h_{230}$	$h_{140}$	$h_{510}$	$h_{420}$	$h_{330}$	$h_{240}$	$h_{150}$
$x_2^2$	$\frac{28}{3}$	$\frac{1}{3}$	0	0	$\frac{49}{6}$	0	$h_{320}$	$h_{230}$	$h_{140}$	$h_{050}$	$h_{420}$	$h_{330}$	$h_{240}$	$h_{150}$	$h_{060}$
$x_1^3$	$\frac{79}{4}$	0	0	$h_{500}$	$h_{410}$	$h_{320}$	$h_{600}$	$h_{510}$	$h_{420}$	$h_{330}$	$h_{700}$	$h_{610}$	$h_{520}$	$h_{430}$	$h_{340}$
$x_1^2x_2$	0	0	$\frac{49}{6}$	$h_{410}$	$h_{320}$	$h_{230}$	$h_{510}$	$h_{420}$	$h_{330}$	$h_{240}$	$h_{610}$	$h_{520}$	$h_{430}$	$h_{340}$	$h_{250}$
$x_1x_2^2$	$\frac{1}{3}$	$\frac{49}{6}$	0	$h_{320}$	$h_{230}$	$h_{140}$	$h_{420}$	$h_{330}$	$h_{240}$	$h_{150}$	$h_{520}$	$h_{430}$	$h_{340}$	$h_{250}$	$h_{160}$
$x_2^3$	0	0	0	$h_{230}$	$h_{140}$	$h_{050}$	$h_{330}$	$h_{240}$	$h_{150}$	$h_{060}$	$h_{430}$	$h_{340}$	$h_{250}$	$h_{160}$	$h_{070}$
$x_1^4$	0	$h_{500}$	$h_{410}$	$h_{600}$	$h_{510}$	$h_{420}$	$h_{700}$	$h_{610}$	$h_{520}$	$h_{430}$	$h_{800}$	$h_{710}$	$h_{620}$	$h_{530}$	$h_{440}$
$x_1^3x_2$	0	$h_{410}$	$h_{320}$	$h_{510}$	$h_{420}$	$h_{330}$	$h_{610}$	$h_{520}$	$h_{430}$	$h_{340}$	$h_{710}$	$h_{620}$	$h_{530}$	$h_{440}$	$h_{350}$
$x_1^2x_2^2$	$\frac{49}{6}$	$h_{320}$	$h_{230}$	$h_{420}$	$h_{330}$	$h_{240}$	$h_{520}$	$h_{430}$	$h_{340}$	$h_{250}$	$h_{620}$	$h_{530}$	$h_{440}$	$h_{350}$	$h_{260}$
$x_1x_2^3$	0	$h_{230}$	$h_{140}$	$h_{330}$	$h_{240}$	$h_{150}$	$h_{430}$	$h_{340}$	$h_{250}$	$h_{160}$	$h_{530}$	$h_{440}$	$h_{350}$	$h_{260}$	$h_{170}$
$x_2^4$	0	$h_{140}$	$h_{050}$	$h_{240}$	$h_{150}$	$h_{060}$	$h_{340}$	$h_{250}$	$h_{160}$	$h_{070}$	$h_{440}$	$h_{350}$	$h_{260}$	$h_{170}$	$h_{080}$

- (4) Extract a principal minor of full rank.

In our example the  $6 \times 6$  principal minor is of full rank. The matrix  $\Delta_0$  is

$$\Delta_0 = \begin{bmatrix} 0 & \frac{57}{4} & 0 & 0 & 0 & \frac{28}{3} \\ \frac{57}{4} & 0 & 0 & \frac{79}{4} & 0 & \frac{1}{3} \\ 0 & 0 & \frac{28}{3} & 0 & \frac{1}{3} & 0 \\ 0 & \frac{79}{4} & 0 & 0 & 0 & \frac{49}{6} \\ 0 & 0 & \frac{1}{3} & 0 & \frac{49}{6} & 0 \\ \frac{28}{3} & \frac{1}{3} & 0 & \frac{49}{6} & 0 & 0 \end{bmatrix}$$

The columns (and the rows) of the matrix correspond to the monomials  $\{1, x_1, x_2, x_1^2, x_1x_2, x_2^2\}$ .

- (5) We compute the “shifted” matrix  $\Delta_1 = x_1\Delta_0$ .

The shifted matrix  $\Delta_1$  is

$$\Delta_1 = \begin{bmatrix} \frac{57}{4} & 0 & 0 & \frac{79}{4} & 0 & \frac{1}{3} \\ 0 & \frac{79}{4} & 0 & 0 & 0 & \frac{49}{6} \\ 0 & 0 & \frac{1}{3} & 0 & \frac{49}{6} & 0 \\ \frac{79}{4} & 0 & 0 & h_{500} & h_{410} & h_{320} \\ 0 & 0 & \frac{49}{6} & h_{410} & h_{320} & h_{230} \\ \frac{1}{3} & \frac{49}{6} & 0 & h_{320} & h_{230} & h_{140} \end{bmatrix}$$

The columns of the matrix correspond to the monomials  $\{x_1, x_1^2, x_1x_2, x_1^3, x_1^2x_2, x_1x_2^2\}$  which are the monomials that correspond to the columns of  $\Delta_0$ , i.e.  $\{1, x_1, x_2, x_1^2, x_1x_2, x_2^2\}$ , multiplied by  $x_1$ . Since not all the entries of  $\Delta_1$  are known, we need to compute them in order to proceed.

Consider the following method to extend the matrix of a quotient algebra. In the quotient algebra it holds that  $\mathbb{M}_{x_i}\mathbb{M}_{x_j} - \mathbb{M}_{x_j}\mathbb{M}_{x_i} = \mathbb{O}$  [43] for any  $i, j \in \{1, \dots, n\}$ , i.e. the matrices of multiplications commute (cf. Section 4).

From Proposition 3.9 we know that  $\Delta_i = \mathbb{M}_{x_i}^t \Delta_0$ , and hence  $\mathbb{M}_{x_i}^t = \Delta_i \Delta_0^{-1}$ , for  $1 \leq i \leq n$ .

We form all the possible matrix equations,  $\mathbb{M}_{x_i} \mathbb{M}_{x_j} - \mathbb{M}_{x_j} \mathbb{M}_{x_i} = \mathbb{O}$ , there are  $\binom{n}{2}$ , and we equate their elements to zero. Since the dimension of the matrices is  $r \times r$ , this leads to at most  $\binom{n}{2} r^2$ , or  $\mathcal{O}(n^2 r^2)$  equations. Note that the equations are, at most of total degree 2.

In our example the matrix  $\Delta_2$  is

$$\Delta_2 = \begin{bmatrix} 0 & 0 & \frac{28}{3} & 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{3} & 0 & \frac{49}{6} & 0 \\ \frac{28}{3} & \frac{1}{3} & 0 & \frac{49}{6} & 0 & 0 \\ 0 & 0 & \frac{49}{6} & h_{410} & h_{320} & h_{230} \\ \frac{1}{3} & \frac{49}{6} & 0 & h_{320} & h_{230} & h_{140} \\ 0 & 0 & 0 & h_{230} & h_{140} & h_{050} \end{bmatrix}$$

Since we have only two variables, there is only one matrix equation,

$$\mathbb{M}_{x_i} \mathbb{M}_{x_j} - \mathbb{M}_{x_j} \mathbb{M}_{x_i} = \Delta_1 \Delta_0^{-1} \Delta_2 \Delta_0^{-1} - \Delta_2 \Delta_0^{-1} \Delta_1 \Delta_0^{-1} = \mathbb{O}.$$

Many of the resulting equations are trivial. After discarding them, we have 6 unknowns  $\{h_{500}, h_{410}, h_{320}, h_{230}, h_{140}, h_{050}\}$  and 15 equations.

A solution of the system is the following

$$\{h_{500} = 1, h_{410} = 2, h_{320} = 3, h_{230} = 1.5060, h_{140} = 4.960, h_{050} = 0.056\}.$$

We substitute these values to  $\Delta_1$  and we continue the algorithm as in the previous example.

Other algorithms to extend a moment matrix, e.g. [18, 38, 39], the so called *flat extensions*, are applicable when the  $\Delta_0$  is positive definite.

- (6) We solve the equation  $(\Delta_1 - \lambda \Delta_0)X = 0$ .

The normalized eigenvectors of the generalized eigenvalue problem are

$$\begin{bmatrix} 1 \\ -0.830 + 1.593i \\ -0.326 - 0.0501i \\ -1.849 - 2.645i \\ 0.350 - 0.478i \\ 0.103 + 0.0327i \end{bmatrix}, \begin{bmatrix} 1 \\ -0.830 - 1.593i \\ -0.326 + 0.050i \\ -1.849 + 2.645i \\ 0.350 + 0.478i \\ 0.103 - 0.032i \end{bmatrix}, \begin{bmatrix} 1.0 \\ 1.142 \\ 0.836 \\ 1.305 \\ 0.955 \\ 0.699 \end{bmatrix}, \\ \begin{bmatrix} 10.956 \\ -0.713 \\ 0.914 \\ -0.682 \\ 0.509 \end{bmatrix}, \begin{bmatrix} 1 \\ -0.838 + 0.130i \\ 0.060 + 0.736i \\ 0.686 - 0.219i \\ -0.147 - 0.610i \\ -0.539 + 0.089i \end{bmatrix}, \begin{bmatrix} 1 \\ -0.838 - 0.130i \\ 0.060 - 0.736i \\ 0.686 + 0.219i \\ -0.147 + 0.610i \\ -0.539 - 0.089i \end{bmatrix}.$$

The coordinates of the eigenvectors correspond to the elements  $\{1, x_1, x_2, x_1^2, x_1 x_2, x_2^2\}$  and we can recover the coefficients of  $x_1$  and  $x_2$  in the decomposition.

After, solving the over-constrained linear system for the coefficients of the linear forms we deduce the decomposition

$$\begin{aligned} & (0.517 + 0.044i)(x_0 - (0.830 - 1.593i)x_1 - (0.326 + 0.050i)x_2)^4 \\ & + (0.517 - 0.044i)(x_0 - (0.830 + 1.593i)x_1 - (0.326 - 0.050i)x_2)^4 \\ & \quad + 2.958(x_0 + (1.142)x_1 + 0.836x_2)^4 \\ & \quad + 4.583(x_0 + (0.956)x_1 - 0.713x_2)^4 \\ & - (4.288 + 1.119i)(x_0 - (0.838 - 0.130i)x_1 + (0.060 + 0.736i)x_2)^4 \\ & - (4.288 - 1.119i)(x_0 - (0.838 + 0.130i)x_1 + (0.060 - 0.736i)x_2)^4 \end{aligned}$$



## 6. CONCLUSIONS AND FUTURE WORK

We propose an algorithm that computes symmetric tensor decompositions, extending Sylvester's algorithm. The main ingredients are i) reformulate the problem in a dual space, ii) exploit the properties of multivariate Hankel operators and Gorenstein algebra, iii) devise an effective method to solve, when necessary, the truncated Hankel problem, iv) deduce the decomposition by solving a generalized eigenvalue problem.

There are several open questions that we are currently working on. What is the (arithmetic and Boolean) complexity of the algorithm? If we do not know all the elements of the tensor, can we still compute a decomposition?

**Acknowledgments.** This work is partially supported by contract ANR-06-BLAN-0074 "Decotes".

## REFERENCES

- [1] L. Albera, A. Ferreol, P. Comon, and P. Chevalier. Blind identification of overcomplete mixtures of sources (BIOME). *Lin. Algebra Appl.*, 391:1–30, November 2004.
- [2] J. Alexander and A. Hirschowitz. La méthode d'Horace éclatée : application à l'interpolation en degré quatre. *Inventh. math.*, 107:585–602, 1992.
- [3] D. Bini, M. Capovani, F. Romani, and G. Lotti.  $O(n^{2.77})$  Complexity for  $n \times n$  approximate matrix multiplication. *Inform. Process. Lett.*, 8(5):234–235, 1979.
- [4] R. Bro. Parafac, tutorial and applications. *Chemom. Intel. Lab. Syst.*, 38:149–171, 1997.
- [5] Peter Bürgisser, Michael Clausen, and M. Amin Shokrollahi. *Algebraic complexity theory*, volume 315 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1997. ISBN 3-540-60582-7. With the collaboration of Thomas Lickteig.
- [6] J. F. Cardoso. Blind signal separation: statistical principles. *Proc. of the IEEE*, 90:2009–2025, October 1998. special issue, R.W. Liu and L. Tong eds.
- [7] E. Carlini. Reducing the number of variables of a polynomial. In M. Elkadi, B. Mourrain, and R. Piene, editors, *Algebraic geometry and geometric modeling*, pages 237–247. Springer, 2005.
- [8] P. Chevalier. Optimal separation of independent narrow-band sources - concept and performance. *Signal Processing, Elsevier*, 73(1):27–48, February 1999. special issue on blind separation and deconvolution.
- [9] P. Chevalier, L. Albera, A. Ferreol, and P. Comon. On the virtual array concept for higher order array processing. *IEEE Trans. Sig. Proc.*, 53(4):1254–1271, April 2005.
- [10] A. Cichocki and S-I. Amari. *Adaptive Blind Signal and Image Processing*. Wiley, New York, 2002.
- [11] G. Comas and M. Seiguer. On the rank of a binary form, 2001. URL <http://arxiv.org/abs/math/0112311>.
- [12] P. Comon. Independent Component Analysis. In J-L. Lacoume, editor, *Higher Order Statistics*, pages 29–38. Elsevier, Amsterdam, London, 1992.
- [13] P. Comon and B. Mourrain. Decomposition of quantics in sums of powers of linear forms. *Signal Processing*, 53(2-3):93–107, 1996.
- [14] P. Comon and M. Rajih. Blind identification of under-determined mixtures based on the characteristic function. *Signal Processing*, 86(9):2271–2281, September 2006.
- [15] P. Comon, G. Golub, L-H. Lim, and B. Mourrain. Symmetric tensors and symmetric tensor rank. *SIAM Journal on Matrix Analysis Appl.*, 30(3):1254–1279, 2008.
- [16] D. Cox, J. Little, and D. O'Shea. *Ideals, Varieties, and Algorithms*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 2nd edition, 1997.
- [17] D. Cox, J. Little, and D. O'Shea. *Using Algebraic Geometry*. Number 185 in Graduate Texts in Mathematics. Springer, New York, 2nd edition, 2005.
- [18] R. E. Curto and L. A. Fialkow. Recursiveness, positivity, and truncated moment problems. *Houston J. Math*, 17(4):603–635, 1991.
- [19] L. de Lathauwer and J. Castaing. Tensor-based techniques for the blind separation of ds-cdma signals. *Signal Processing*, 87(2):322–336, February 2007.
- [20] L. de Lathauwer, J. Castaing, and J.F. Cardoso. Fourth-order cumulant-based identification of under-determined mixtures. *IEEE Trans. Sig. Proc.*, 55(6):2965–2973, June 2007.

- [21] J. de Leeuw. A new computational method to fit the weighted euclidean distance model. *Psychometrika*, 43(4):479–490, December 1978.
- [22] M. C. Dogan and J. Mendel. Applications of cumulants to array processing .I. aperture extension and array calibration. *IEEE Trans. Sig. Proc.*, 43(5):1200–1216, May 1995.
- [23] D. L. Donoho and X. Huo. Uncertainty principles and ideal atomic decompositions. *IEEE Trans. Inform. Theory*, 47(7):2845–2862, November 2001.
- [24] M. Elkadi and B. Mourrain. *Introduction à la résolution des systèmes polynomiaux*, volume 59 of *Mathématiques et Applications*. Springer, 2007. ISBN 978-3-540-71646-4.
- [25] A. Ferreol and P. Chevalier. On the behavior of current second and higher order blind source separation methods for cyclostationary sources. *IEEE Trans. Sig. Proc.*, 48:1712–1725, June 2000. erratum in vol.50, pp.990, Apr. 2002.
- [26] B. N. Flury. Common principal components in k groups. *Journal of the American Statistical Association*, 79(388):892–898, 1984.
- [27] G. H. Golub and C. F. Van Loan. *Matrix computations*. John Hopkins, Baltimore MD, 1983.
- [28] G.H. Golub and C.F. Van Loan. *Matrix Computations*. Johns Hopkins University Press, 1996.
- [29] J. Harris. *Algebraic Geometry: A First Course*. Number 133 in Graduate Texts in Mathematics. Springer-Verlag, 1992.
- [30] R. A. Harshman and M. Lundy. PARAFAC: Parallel factor analysis. *Computational Stat. Data Analysis*, 18:39–72, 1994.
- [31] A. Iarrobino and V. Kanev. *Power sums, Gorenstein algebras, and determinantal loci*, volume 1721 of *Lecture Notes in Computer Science*. Springer-Verlag, Berlin, 1999.
- [32] T. Jiang and N. Sidiropoulos. Kruskal’s permutation lemma and the identification of CANDECOMP/PARAFAC and bilinear models. *IEEE Trans. Sig. Proc.*, 52(9):2625–2636, September 2004.
- [33] H. A. L. Kiers and W. P. Krijnen. An efficient algorithm for Parafac of three-way data with large numbers of observation units. *Psychometrika*, 56:147, 1991.
- [34] I. A. Kogan and M. Moreno Maza. Computation of canonical forms for ternary cubics. In *Proc. Int. Symp. on Symbolic and Algebraic Computation*, pages 151–160, New York, NY, USA, 2002. ACM.
- [35] J. B. Kruskal. Three-way arrays: Rank and uniqueness of trilinear decompositions. *Linear Algebra and Applications*, 18:95–138, 1977.
- [36] J. Kung and G. Rota. The invariant theory of binary forms. *Bulletin of the American Mathematical Association*, pages 27–85, 1984.
- [37] J. Landsberg. Geometry and the complexity of matrix multiplication. *Bull. Amer. Math. Soc.*, 45(2): 247–284, April 2008.
- [38] M. Laurent. Revisiting two theorems of Curto and Fialkow on moment matrices. *Proc. Amer. Math. Soc.*, 133:2965–2976, 2005.
- [39] M. Laurent. Sums of squares, moment matrices and optimization over polynomials. In M. Putinar and S. Sullivant, editors, *IMA volume Emerging Applications of Algebraic Geometry*. IMA, 2008. (to appear).
- [40] M. Laurent and B. Mourrain. A sparse flat extension theorem for moment matrices. <http://fr.arxiv.org/abs/0812.2563>, 2008.
- [41] F.S. Macaulay. *The Algebraic Theory of Modular Systems*. Cambridge Univ. Press, 1916.
- [42] P. McCullagh. *Tensor Methods in Statistics*. Monographs on Statistics and Applied Probability. Chapman and Hall, 1987.
- [43] B. Mourrain. A new criterion for normal form algorithms. In M. Fossorier, H. Imai, S. Lin, and A. Poli, editors, *Proc. Applic. Algebra in Engineering, Communic. and Computing*, volume 1719 of *Lecture Notes in Computer Science*, pages 430–443. Springer, Berlin, 1999.
- [44] B. Mourrain and V.Y. Pan. Multivariate Polynomials, Duality, and Structured Matrices. *Journal of Complexity*, 16(1):110–180, 2000.
- [45] R. Oldenburger. Composition and rank of n-way matrices and multilinear forms. *The Annals of Mathematics*, 35(3):622–653, 1934.
- [46] P. Paatero. The multilinear engine: A table-driven, least squares program for solving multilinear problems, including the n-way parallel factor analysis model. *Journal of Computational and Graphical Statistics*, 8(4):854–888, December 1999.

- [47] D. T. Pham and J-F. Cardoso. Blind separation of instantaneous mixtures of nonstationary sources. *IEEE Trans. Sig. Proc.*, 49(9):1837–1848, September 2001.
- [48] N. D. Sidiropoulos, G. B. Giannakis, and R. Bro. Blind PARAFAC receivers for DS-CDMA systems. *IEEE Trans. on Sig. Proc.*, 48(3):810–823, March 2000.
- [49] A. Smilde, R. Bro, and P. Geladi. *Multi-Way Analysis*. Wiley, 2004.
- [50] V. Strassen. Rank and optimal computation of generic tensors. *Linear Algebra Appl.*, 52:645–685, July 1983.
- [51] A. Swami, G. Giannakis, and S. Shamsunder. Multichannel ARMA processes. *IEEE Trans. Sig. Proc.*, 42(4):898–913, April 1994.
- [52] J. J. Sylvester. Sur une extension d’un théorème de Clebsch relatif aux courbes du quatrième degré. *Comptes Rendus, Math. Acad. Sci. Paris*, 102:1532–1534, 1886.
- [53] J. M. F. ten Berge, N. D. Sidiropoulos, and R. Rocci. Typical rank and INDSCAL dimensionality for symmetric three-way arrays of order  $I \times 2 \times 2$  or  $I \times 3 \times 3$ . *Linear Algebra Appl.*, 388:363–377, 2004.
- [54] A. J. van der Veen and A. Paulraj. An analytical constant modulus algorithm. *IEEE Trans. Sig. Proc.*, 44(5):1136–1155, May 1996.
- [55] A. Yeredor. Non-orthogonal joint diagonalization in the LS sense with application in blind source separation. *IEEE Trans. Sig. Proc.*, 50(7):1545–1553, 2002.
- [56] F. L. Zak. *Tangents and secants of algebraic varieties*, volume 127 of *Translations of Mathematical Monographs*. AMS, Providence, RI, 1993.
- [57] A. Ziehe, G. Nolte, and K. R. Müller. A fast algorithm for joint diagonalization with non orthogonal transformations and its application to blind source separation. *Journal of Machine Learning Research*, 5:777–800, December 2004.

## APPENDIX A. TERNARY CUBICS

As an application, we present the decomposition of all the types of ternary cubics. The decomposition allows us to classify, up to projective transformations of the variables, homogeneous polynomials of degree three in three variables, for instance with the help of the algorithm described in [13]. For another algorithm for decomposing ternary cubics, based on the method of moving frames and on triangular decompositions of algebraic varieties, we refer the reader to [34]. Two polynomials are equivalent in this classification if there exists a variable invertible transformation which maps one polynomial to the other.

The classification algorithm goes as follows. Given a ternary cubic, we compute its decomposition as a sum of powers of linear forms. We have the following cases:

- If the rank is one then the polynomial is a 3<sup>rd</sup> power of a linear form, that is, it is equivalent to  $x_0^3$ .
- If the rank is 2, then the polynomial is equivalent to  $x_0^3 + x_1^3$  and is in the orbit of  $x_0x_1(x_0 + x_1)$ . In fact, the decomposition of the latter polynomial is

$$\begin{aligned} & \left(-260712 - \frac{1628000}{9}i\sqrt{3}\right) \left(\left(-\frac{125}{16492} - \frac{17}{16492}i\sqrt{3}\right)x_0 + \left(-\frac{22}{4123} + \frac{27}{8246}i\sqrt{3}\right)x_1\right)^3 \\ & + \left(-260712 + \frac{1628000}{9}i\sqrt{3}\right) \left(\left(-\frac{125}{16492} + \frac{17}{16492}i\sqrt{3}\right)x_0 + \left(-\frac{22}{4123} - \frac{27}{8246}i\sqrt{3}\right)x_1\right)^3. \end{aligned}$$

- If the rank is 3, then the polynomial is either in the orbit of  $x_0^2x_1$  or in the orbit of  $x_0^3 + x_1^3 + x_2^3$ . To identify the orbit, it suffices to check if the polynomial is square-free or not (that is, check whether the gcd between the polynomial and one of its derivatives is 1). If it is not square-free then it is in the orbit of  $x_0^2x_1$ . Otherwise it is in the orbit of  $x_0^3 + x_1^3 + x_2^3$ .

The decomposition of  $x_0^2x_1$  is

$$\begin{aligned} & -0.15563 - 0.54280i((-0.95437 - 0.08483i)x_0 + (-0.00212 + 0.28631i)x_1)^3 \\ & -0.15723 + 0.54841i((-0.95111 + 0.09194i)x_0 + (-0.00222 - 0.29484i)x_1)^3 \\ & -0.04320((-0.49451)x_0 + (-0.86917)x_1)^3. \end{aligned}$$

- If the rank is 4, then our polynomial is generic. As an example, consider the polynomial  $150x_0^2x_2 + x_1^2x_2 + x_2^3 - 12x_0^3$ ; a decomposition of which is

$$\begin{aligned} & 0.53629(+0.34496x_0 + 0.71403x_1 + 0.60923x_2)^3 \\ & -195.64389(-0.99227x_0 + 0.00286x_1 - 0.12403x_2)^3 \\ & +211.45588(-0.99282x_0 + 0.00311x_1 + 0.11962x_2)^3 \\ & +0.52875(-0.34600x_0 - 0.71671x_1 + 0.60549x_2)^3. \end{aligned}$$

- If the rank is 5, then the polynomial is of maximal rank and it is in the orbit of  $x_0^2x_1 + x_0x_2^2$ , a decomposition of which, is

$$\begin{aligned} & +0.28100(+0.06322x_0 - 0.99748x_1 + 0.03224x_2)^3 \\ & +0.97839(+0.14391x_0 + 0.50613x_1 + 0.85036x_2)^3 \\ & +0.44877(+0.73493x_0 + 0.56369x_1 - 0.37702x_2)^3 \\ & +(-0.97396 - 0.94535i)(0.45304x_0 + (-0.60752 + 0.14316i)x_1 + (-0.52915 + 0.35382i)x_2)^3 \\ & +(-0.97396 + 0.94535i)(0.45304x_0 + (-0.60752 + 0.14316i)x_1 + (-0.52915 - 0.35382i)x_2)^3. \end{aligned}$$

## APPENDIX B. AN EXAMPLE OF EXTREME RANK

In this section we present in detail the decomposition of a ternary cubic of maximal rank, that is 5. Consider the polynomial

$$x_0^2x_1 + x_0x_2^2.$$

The matrix of the quotient algebra is

$$\begin{bmatrix} 0 & \frac{1}{3} & 0 & 0 & 0 & \frac{1}{3} & 0 & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 0 & 0 & 0 & h_{4,0,0} & h_{3,1,0} & h_{2,2,0} & h_{1,3,0} \\ 0 & 0 & \frac{1}{3} & 0 & 0 & 0 & h_{3,1,0} & h_{2,2,0} & h_{1,3,0} & h_{0,4,0} \\ 0 & 0 & 0 & h_{4,0,0} & h_{3,1,0} & h_{2,2,0} & h_{5,0,0} & h_{4,1,0} & h_{3,2,0} & h_{2,3,0} \\ 0 & 0 & 0 & h_{3,1,0} & h_{2,2,0} & h_{1,3,0} & h_{4,1,0} & h_{3,2,0} & h_{2,3,0} & h_{1,4,0} \\ \frac{1}{3} & 0 & 0 & h_{2,2,0} & h_{1,3,0} & h_{0,4,0} & h_{3,2,0} & h_{2,3,0} & h_{1,4,0} & h_{0,5,0} \\ 0 & h_{4,0,0} & h_{3,1,0} & h_{5,0,0} & h_{4,1,0} & h_{3,2,0} & h_{6,0,0} & h_{5,1,0} & h_{4,2,0} & h_{3,3,0} \\ 0 & h_{3,1,0} & h_{2,2,0} & h_{4,1,0} & h_{3,2,0} & h_{2,3,0} & h_{5,1,0} & h_{4,2,0} & h_{3,3,0} & h_{2,4,0} \\ 0 & h_{2,2,0} & h_{1,3,0} & h_{3,2,0} & h_{2,3,0} & h_{1,4,0} & h_{4,2,0} & h_{3,3,0} & h_{2,4,0} & h_{1,5,0} \\ 0 & h_{1,3,0} & h_{0,4,0} & h_{2,3,0} & h_{1,4,0} & h_{0,5,0} & h_{3,3,0} & h_{2,4,0} & h_{1,5,0} & h_{0,6,0} \end{bmatrix},$$

and the matrices  $\Delta_0$ ,  $\Delta_1$  and  $\Delta_2$  are

$$\begin{bmatrix} 0 & 1/3 & 0 & 0 & 0 \\ 1/3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1/3 & 0 & 0 \\ 0 & 0 & 0 & h_{4,0,0} & h_{3,1,0} \\ 0 & 0 & 0 & h_{3,1,0} & h_{2,2,0} \end{bmatrix}, \begin{bmatrix} 1/3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & h_{4,0,0} & h_{3,1,0} \\ 0 & 0 & 0 & h_{3,1,0} & h_{2,2,0} \\ 0 & h_{4,0,0} & h_{3,1,0} & h_{5,0,0} & h_{4,1,0} \\ 0 & h_{3,1,0} & h_{2,2,0} & h_{4,1,0} & h_{3,2,0} \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1/3 & 0 & 0 \\ 0 & 0 & 0 & h_{3,1,0} & h_{2,2,0} \\ 1/3 & 0 & 0 & h_{2,2,0} & h_{1,3,0} \\ 0 & h_{3,1,0} & h_{2,2,0} & h_{4,1,0} & h_{3,2,0} \\ 0 & h_{2,2,0} & h_{1,3,0} & h_{3,2,0} & h_{2,3,0} \end{bmatrix}.$$

If we form the matrix equation

$$\mathbb{M}_{x_i} \mathbb{M}_{x_j} - \mathbb{M}_{x_j} \mathbb{M}_{x_i} = \Delta_1 \Delta_0^{-1} \Delta_2 \Delta_0^{-1} - \Delta_2 \Delta_0^{-1} \Delta_1 \Delta_0^{-1} = \mathbb{O}.$$

then we have a system of 8 equations in 8 unknowns. The unknowns are

$$\{h_{5,0,0}, h_{4,1,0}, h_{4,0,0}, h_{3,1,0}, h_{2,2,0}, h_{1,3,0}, h_{3,2,0}, h_{2,3,0}\}.$$

It turns out that the system is not zero dimensional, and that we can choose (randomly) the values of five of them, i.e.  $\{h_{1,3,0} = 3, h_{3,1,0} = 1, h_{2,2,0} = 2, h_{4,1,0} = 4, h_{4,0,0} = 5\}$ . Working as in the other examples we end up with the decomposition

$$\begin{aligned} &+0.000071(x_0 - 15.778x_1 + 0.510x_2)^3 \\ &+0.002916(x_0 + 3.517x_1 + 5.909x_2)^3 \\ &+0.178137(x_0 + 0.767x_1 - 0.513x_2)^3 \\ &(-0.09056 - 0.0879i)(x_0 + (-1.341 + 0.316i)x_1 + (-1.168 + 0.781i)x_2)^3 \\ &(-0.09056 + 0.0879i)(x_0 + (-1.341 + 0.316i)x_1 + (-1.168 - 0.781i)x_2)^3. \end{aligned}$$