



HAL
open science

A wireless sensor network to measure the health care workers exposure to tuberculosis

Guillaume Chelius, Eric Fleury, Antoine Fraboulet, Jean-Christophe Lucet

► **To cite this version:**

Guillaume Chelius, Eric Fleury, Antoine Fraboulet, Jean-Christophe Lucet. A wireless sensor network to measure the health care workers exposure to tuberculosis. International Workshop and Conference on Complex Networks and their Applications (NetSci 09), Jun 2009, Venice, Italy. inria-00385398

HAL Id: inria-00385398

<https://inria.hal.science/inria-00385398>

Submitted on 19 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A wireless sensor network to measure the health care workers exposure to tuberculosis

Guillaume Chelius

University of Lyon, INRIA, ENS Lyon
F-69364, France

Email: guillaume.chelius@inria.fr

Eric Fleury

University of Lyon, ENS Lyon, INRIA
F-69364, France

Email: eric.fleury@inria.fr

Antoine Fraboulet

University of Lyon, INSA Lyon, INRIA
F-69621, France

Email: antoine.fraboulet@insa-lyon.fr

Jean-Christophe Lucet

Bichat-Cl. Bernard Hospital, Paris VII University
France

Email: jean-christophe.lucet@bch.aphp.fr

May 19, 2009

In parallel to the advances in modern medicine, health sciences and public health policy, epidemic models aided by computer simulations and information technologies offer an important alternative for the understanding of transmission dynamics and epidemic patterns. In this paper, we focus on the first steps that may lead towards the design of epidemic models, i.e. the measure and analysis of interactions within a closed socio-professional context. More precisely, this project was motivated by the study of the Health Care Workers (HCWs) exposure to tuberculosis in their work environment.

Despite the progresses in treatment and prevention, tuberculosis remains a disease in expansion and represents the third cause of death by infectious pathologies in the world. In the health care context, if the transmission is globally controlled, the HCWs exposure remains obscure. Individual factors associated to the contamination of HCWs in their work environment are not precisely known. Our study focus on the evaluation of the intensity and the frequency of contacts between tuberculosis infected patients and HCWs. To gather this information, classical methods consist in performing audits, consulting medical and administrative files or using self-reports of conversations and trusting HCW souvenirs. All these methods are time-consuming, subject to memory failures and interpretations. As an alternate method, we have chosen to dedicate a Wireless Sensor Network (WSN) to gather these interactions inside a Service of Infectious and Tropical Diseases (Bichat-Claude Bernard Hospital, Paris) and a Service of Pneumology (La Pitié Salpêtrière Hospital, Paris). Within the two services, each room has been equipped with a sensor node and each HCW carries an autonomous sensor during his presence in the service. An important characteristic of this measurement campaign is that it was performed in a closed environment, over a closed population and during a large continuous period of time. That is, the presence of all HCWs of the units was monitored in all patient rooms, 24/7 during a three months period. In addition to the experimental measure system description, this paper main contributions are the analysis and characterization of this huge and unique data set describing a complex dynamic interaction network, and the impact study of the measurement process bias on the network dynamic. The analyze of large dynamic in situ interaction networks provides an opportunity to study dynamical processes occurring on dynamical networks, such as spreading or epidemical processes, taking into account the dynamics both on and of the network structure.