



**HAL**  
open science

## Directed acyclic graphs with the unique dipath property

Jean-Claude Bermond, Michel Cosnard, Stéphane Pérennes

► **To cite this version:**

Jean-Claude Bermond, Michel Cosnard, Stéphane Pérennes. Directed acyclic graphs with the unique dipath property. [Research Report] Inria. 2009, pp.19. inria-00387085v2

**HAL Id: inria-00387085**

**<https://hal.inria.fr/inria-00387085v2>**

Submitted on 10 Aug 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Directed acyclic graphs with the unique dipath property

Jean-Claude Bermond\*, Michel Cosnard, Stéphane Pérennes  
MASCOTTE, joint project CNRS-INRIA-UNSA,  
2004 Route des Lucioles, BP 93, F-06902 Sophia-Antipolis, France

## Abstract

Let  $\mathcal{P}$  be a family of dipaths of a DAG (Directed Acyclic Graph)  $G$ . The load of an arc is the number of dipaths containing this arc. Let  $\pi(G, \mathcal{P})$  be the maximum of the load of all the arcs and let  $w(G, \mathcal{P})$  be the minimum number of wavelengths (colors) needed to color the family of dipaths  $\mathcal{P}$  in such a way that two dipaths with the same wavelength are arc-disjoint.

There exist DAGs such that the ratio between  $w(G, \mathcal{P})$  and  $\pi(G, \mathcal{P})$  cannot be bounded. An internal cycle is an oriented cycle such that all the vertices have at least one predecessor and one successor in  $G$  (said otherwise every cycle contain neither a source nor a sink of  $G$ ). We prove that, for any family of dipaths  $\mathcal{P}$ ,  $w(G, \mathcal{P}) = \pi(G, \mathcal{P})$  if and only if  $G$  is without internal cycle.

We also consider a new class of DAGs, which is of interest in itself, those for which there is at most one dipath from a vertex to another. We call these digraphs UPP-DAGs. For these UPP-DAGs we show that the load is equal to the maximum size of a clique of the conflict graph. We prove that the ratio between  $w(G, \mathcal{P})$  and  $\pi(G, \mathcal{P})$  cannot be bounded (a result conjectured in an other article). For that we introduce “good labelings” of the conflict graph associated to  $G$  and  $\mathcal{P}$ , namely labelings of the edges such that for any ordered pair of vertices  $(x, y)$  there do not exist two paths from  $x$  to  $y$  with increasing labels.

## 1 Introduction

The problem we consider is motivated by routing, wavelength assignment and grooming in optical networks. But it can be of interest for other applications in parallel computing, where the graph will represent for example the precedence graph of a program or for scheduling complex operations on pipelined operators. A generic problem in the design of optical networks, [16, 19]), consists of satisfying a family of requests (or a traffic matrix) under various constraints like capacity constraints. The optimization problem associated consists in designing, for a given family of requests, a network optimizing some criteria, such as minimizing the number of wavelengths or the number of ADMs (Add Drop Multiplexers).

A request is satisfied by assigning to it a dipath in the network. A family of requests is satisfied, if we can route them in such a way that the capacity constraints of the network are satisfied. This is known as the routing problem. For a given routing let us define the load of an arc as the number of routes (dipaths) containing it and the load of the routing as the maximum load of the arcs. Typically one wants either to insure that the load of an arc does not exceed the capacity of this arc or to minimize the load of a routing satisfying a given family of requests.

Many backbone networks are now WDM optical ones. Indeed wavelength division multiplexing (WDM) enables to use the bandwidth of an optical fiber by dividing it in multiple non overlapping

---

\*Partially supported by the CRC CORSO with FRANCE TELECOM and by the European FET project AEOLUS

frequencies or wavelength channels. Satisfying a request in a WDM optical network consists in assigning to it a route (dipath), but also a wavelength, which shall stay unchanged if no conversion is allowed. Therefore the constraint is now that two requests, having the same wavelength, have to be routed by two arc disjoint dipaths or, equivalently, two requests whose associated dipaths share an arc, have to be assigned different wavelengths. Hence the scarce resource is the number of available wavelengths. For a given traffic matrix, either one wants to insure that the family of requests can be satisfied with the available number of wavelengths or one wants to minimize the number of wavelengths used. This problem is known in the literature as the RWA (Routing and Wavelength Assignment) problem.

Note that requests are satisfied on a virtual (logical) network which is itself embedded in the physical network (in fact there might be many layers). It is the case for example when considering SONET/WDM rings or in MPLS over WDM networks; in the latter case the RWA problem has to be considered for the lightpaths [9, 10]. Anyway, at the conceptual level of modeling of this article, the problems are the same and we will use the word request to indicate a connection at the upper level.

Minimizing the load or/and the number of wavelengths is a difficult problem and in general an NP-hard problem. These problems have been extensively studied in the literature for various topologies or special families of requests like multicast or all-to-all (see for example the survey [2] or [13, 18]). Many particular cases where the minimum number of wavelengths is equal to the minimum routing load have been given. For example, in [3] it is shown that for any digraph and for a multicast instance (all the requests have the same origin), there is equality and both problems can be solved in polynomial time. For some topologies the load might be easily computed, but the minimum number of wavelengths is NP-hard to compute as it is related to coloring problems. This is the case for symmetric trees (see the survey [8]). However, for symmetric trees it has been proved that there is equality for the all to all instance ([12]) and approximation algorithms have been given ([8, 11]).

As the RWA problem is very difficult to solve, it is often split into two separate problems. First one solves the routing problem by determining dipaths which minimize the load or are easy to compute like shortest paths. Then, the routing being given, the wavelength assignment problem is solved. In that case the input of the problem is not a family of requests but a family of dipaths  $\mathcal{P}$ . We will denote by  $\pi(G, \mathcal{P})$  the maximum of the load of all the arcs of the digraph  $G$  for the family  $\mathcal{P}$ . Determining the minimum number  $w(G, \mathcal{P})$  of wavelengths (colors) needed to color a family of dipaths  $\mathcal{P}$  in such a way that two dipaths with the same wavelength are arc-disjoint is still NP-hard in that case. Indeed it corresponds to finding the chromatic number of the **conflict graph** (also called the intersection graph) associated to the digraph  $G$  and the family of dipaths  $\mathcal{P}$  whose vertices represent the dipaths and where two vertices are joined if the corresponding dipaths are in conflict (that is share an arc).

There are examples of topologies and family of dipaths where there are at most 2 dipaths using an arc ( $\pi(G, \mathcal{P}) = 2$ ), but where we need as many wavelengths as we want. Figure 1 shows the example for  $k = 4$  wavelengths) In the example we consider  $k$  dipaths from  $s_i$  to  $t_i$ . The dipaths starts in  $s_i$ , then go alternatively right and down till they arrive at the bottom where they go right and up till they arrive at the destination  $t_i$ . Any two dipaths intersect so the conflict graph is complete and we need  $k$  colors. However the load of an arc is at most 2. Therefore the ratio between  $w(G, \mathcal{P})$  and  $\pi(G, \mathcal{P})$  is unbounded in general.

Here we consider the class of Directed Acyclic Graphs, DAGs, which plays a central role in Parallel

and Distributed Computing. Part of our motivation came when we tried to extend the results obtained in [6] for paths motivated by grooming problems for the paths ([4, 10]). In fact, we first proved that for rooted trees (directed trees where there is a unique dipath from the root to any vertex), for any family of requests, the minimum number of wavelengths is equal to the load.

The example given above in Figure 1 being a DAG there is no hope to bound ratio between  $w(G, \mathcal{P})$  and  $\pi(G, \mathcal{P})$ . In [5] we fully characterize when  $w(G, \mathcal{P}) = \pi(G, \mathcal{P})$  for a DAG. In fact the necessary and sufficient condition is that  $G$  does not contain what we call an internal cycle, i.e. an oriented cycle, such that all the vertices have at least one predecessor and one successor in  $G$  (said otherwise all cycles contain neither a source nor a sink). Here, we give a new shorter proof of this result deriving it from the case of trees where it is a known result as the conflict graph is a perfect graph (see for example [15]).




Figure 1: A pathological example

Then, we also consider a new (to our best knowledge) class of DAGs, which is of interest in itself, those for which there is at most one dipath from a vertex to another. Note that, if the original digraph has the property that for any request  $(x, y)$  there is a unique dipath from  $x$  to  $y$ , then it is equivalent to consider a family of requests or a family of dipaths. We call this property the UPP (Unique diPath Property) and call these digraphs UPP-DAGs. For these UPP-DAGs we show that the load is equal to the maximum size of a clique of the conflict graph. In [5] we proved that if an UPP-DAG has only one internal cycle, then for any family of dipaths  $w(G, \mathcal{P}) = \lceil \frac{4}{3} \pi(G, \mathcal{P}) \rceil$  and we exhibit an UPP-DAG and a family of dipaths reaching the bound.

In this article we prove that, for UPP-DAGs with load 2, the ratio between  $w(G, \mathcal{P})$  and  $\pi(G, \mathcal{P})$  cannot be bounded (a result conjectured in [5])

For that we introduce the notion of “good labelings” of the conflict graph associated to  $G$  and  $\mathcal{P}$ , namely labelings of the edges such that for any ordered pair of vertices  $(x, y)$  there do not exist two paths from  $x$  to  $y$  with increasing labels. We prove first that, if  $G$  is an UPP-DAG with load 2, then for any family of dipaths  $\mathcal{P}$ , the conflict graph  $C(G, \mathcal{P})$  has a good labeling. Then we also show that, if  $H$  is a graph with a good labeling, then there exists an UPP-DAG  $G$  with load 2 and a family of dipaths  $\mathcal{P}$  such that  $H = C(G, \mathcal{P})$ . Finally, we use that fact and a proof of the existence of graphs with good labelings but chromatic number as large as we want to conclude.




Figure 2: An oriented cycle (a) and an internal cycle (b)

## 2 Definitions

We model the network by a digraph  $G$ . The *outdegree* of a vertex  $x$  is the number of arcs with initial vertex  $x$  (that is the number of vertices  $y$  such that  $(x, y)$  is an arc of  $G$ ). The *indegree* of a vertex  $x$  is the number of arcs with terminal vertex  $x$  (that is the number of vertices  $y$  such that  $(y, x)$  is an arc of  $G$ ). A *source* is a vertex with indegree 0 and a *sink* a vertex with outdegree 0. A dipath is a sequence of vertices  $x_1, x_2, \dots, x_k$  such that  $(x_i, x_{i+1})$  is an arc of  $G$ . If  $x_k = x_1$  the dipath is called a directed cycle.

A **DAG (Directed Acyclic Graph)** is a digraph with no directed cycle. However the underlying (undirected) graph obtained by deleting the orientation can have cycles. An (oriented) cycle in a DAG consists therefore of an even sequence of dipaths  $P_1, P_2, \dots, P_{2k}$  alternating in direction (see Figure 2a). The vertices inside the dipaths have indegree and outdegree 1; those where there is a change of orientation have either indegree 2 and outdegree 0 or indegree 0 and outdegree 2.

An **internal cycle** of a DAG  $G$  is an oriented cycle, such that all its vertices have in  $G$  an indegree  $> 0$  and an outdegree  $> 0$ ; said otherwise no vertex is a source or a sink. Hence the vertices where there is a change of orientation in the cycle have a predecessor (resp. a successor) in  $G$ , if they are of indegree 0 (resp. outdegree 0) in the cycle (see Figure 2b).

We will say that a DAG has the **Unique Path Property** if between two vertices there is at most one dipath. A digraph satisfying this property will be called an **UPP-DAG**.

If  $G$  is an UPP-DAG, then any internal cycle contains at least  $2k \geq 4$  vertices where there is a change of orientation. Otherwise it would consist of a dipath from  $x$  to  $y$  and a reverse dipath from  $y$  to  $x$  and so there would be two dipaths from  $x$  to  $y$ .

Finally a DAG with no cycles is an **oriented tree** (its underlying graph has no cycles and so is a tree).

Given a digraph  $G$  and a family of dipaths  $\mathcal{P}$ , the **load of an arc**  $e$  is the number of dipaths of the family containing  $e$ :

$$\text{load}(G, \mathcal{P}, e) = |\{P : P \in \mathcal{P}; e \in P\}|$$

The **load of  $G$  for  $\mathcal{P}$**  will be the maximum over all the arcs of  $G$  and will be denoted by  $\pi(G, \mathcal{P})$ .

We will say that two dipaths are in conflict (or intersect) if they share at least one arc. We will denote by  $w(G, \mathcal{P})$  the **minimum number of colors** needed to color the dipaths of  $\mathcal{P}$  in such a way that two dipaths in conflict (sharing an arc) have different colors. Note that  $\pi(G, \mathcal{P}) \leq w(G, \mathcal{P})$ .

The **conflict graph** (also called the intersection graph) associated to the digraph  $G$  and the family of dipaths  $\mathcal{P}$  has as vertices the dipaths of  $\mathcal{P}$ , two vertices being joined if their associated dipaths are in conflict (that is intersect = share an arc). It will be denoted  $C(G, \mathcal{P})$ . Then  $w(G, \mathcal{P})$  is the chromatic number  $\gamma$  of the conflict graph: that is  $w(G, \mathcal{P}) = \gamma(C(G, \mathcal{P}))$ . Note that  $\pi$  is only upper bounded by the clique number of the conflict graph; indeed the  $\pi$  dipaths containing an arc  $e$  of maximum load are pairwise in conflict. The following property shows that if  $G$  is an UPP-DAG then  $\pi(G, \mathcal{P})$  is exactly the clique number  $\omega(C(G, \mathcal{P}))$  of the conflict graph.




Figure 3: Helly property

**Property 1** *If  $G$  is an UPP-DAG then the dipaths in conflict have the following Helly property : if a set of dipaths are pairwise in conflict, then their intersection is a dipath. Therefore  $\pi(G, \mathcal{P}) = \omega(C(G, \mathcal{P}))$*

**Proof:** If two dipaths intersect, then their intersection is a dipath. Indeed suppose their intersection contains two different dipaths  $(x_1, y_1)$  and  $(x_2, y_2)$  in this order. Then between  $y_1$  and  $x_2$  there are two dipaths, one via  $P_1$  and the other via  $P_2$  (see Figure 3 a).

So suppose  $P_1$  and  $P_2$  intersect in only one interval  $(x_1, y_1)$ , and  $P_3$  intersects  $P_1$  in an arc disjoint interval  $(u_1, v_1)$ . W.l.o.g. we may assume that  $v_1$  is before  $x_1$ . Let  $P_3$  intersects  $P_2$  in the interval  $(u_2, v_2)$ .

Case 1 :  $v_2$  is before  $u_1$  on  $P_3$ .  $v_2$  cannot be after  $y_1$  on  $P_2$  otherwise there will be a directed cycle. So  $v_2$  is before  $x_1$  on  $P_2$  and we have two dipaths from  $v_2$  to  $x_1$ , one via  $P_2$  and the other one via  $P_3$  till  $u_1$  and then via  $P_1$  (see Figure 3 b).

Case 2 :  $u_2$  is after  $v_1$  on  $P_3$ . If  $u_2$  is before  $x_1$  on  $P_2$  we have two dipaths from  $v_1$  to  $x_1$  one via  $P_1$

and the other going from  $v_1$  to  $u_2$  via  $P_3$  and to  $x_1$  via  $P_2$ . If  $u_2$  is after  $y_1$ , we have two dipaths from  $v_1$  to  $u_2$  one via  $P_3$  and the other via  $P_1$  till  $y_1$  and  $P_2$  (see Figure 3 c and d).  $\square$

### 3 Relations between $\pi(G, \mathcal{P})$ and $w(G, \mathcal{P})$

There exist DAGs  $G$  and a set of dipaths  $\mathcal{P}$  such that  $\pi(G, \mathcal{P}) = 2$  and  $w(G, \mathcal{P})$  is as big as we want (see Figure 1). These DAGs have many internal cycles.




Figure 4: Example for a DAG with an internal cycle

In Figure 4, we give an example of a DAG with one internal cycle and a set of 5 dipaths  $\mathcal{P}$  such that  $\pi(G, \mathcal{P}) = 2$  and  $w(G, \mathcal{P}) = 3$ . The dipaths are  $a_1, b_1, c_1$ ;  $b_1, c_1, d_1$ ;  $c_1, d_1, e_1$ ;  $b_1, d_1, e_1$  via the second dipath from  $b_1$  to  $d_1$ ;  $a_1, b_1, d_1$  also via this second dipath. The load is 2 and the conflict graph is a cycle of length 5 and so we need 3 colors to color its vertices.

In fact as shown by the following theorem, if a DAG  $G$  (which can be an UPP-DAG) contains an internal cycle there cannot be equality between  $\pi(G, \mathcal{P})$  and  $w(G, \mathcal{P})$  for all the set of dipaths  $\mathcal{P}$ .

**Theorem 2** *If a DAG  $G$  contains an internal cycle there exists a set  $\mathcal{P}$  of dipaths such that  $\pi(G, \mathcal{P}) = 2$  and  $w(G, \mathcal{P}) = 3$ .*

**Proof:** Let us consider an internal cycle consisting of  $2k$  dipaths  $k$  between  $b_i$  and  $c_i$  and  $k$  between  $b_i$  and  $c_{i-1}$  (the indices are taken modulo  $k$ ). So the  $b_i, i = 1, 2, \dots, k$ , have indegree 0 in the cycle and the  $c_i, i = 1, 2, \dots, k$ , have outdegree 0 in the cycle. As the cycle is internal, there exist  $k$  vertices  $a_i, i = 1, 2, \dots, k$  joined to the  $b_i$  and  $k$  vertices  $d_i, i = 1, 2, \dots, k$  to which are joined the  $c_i$ . Let us take as set  $\mathcal{P}$  of dipaths:  $a_1, b_1, c_1$ ;  $b_1, c_1, d_1$ ;  $a_i, b_i, c_{i-1}, d_{i-1}$  and  $a_i, b_i, c_i, d_i$  for  $i = 2, \dots, k$  and  $a_1, b_1, c_k, d_k$ . The vertices of the conflict graph associated to these dipaths form a cycle of odd length  $2k + 1$  in and so  $w = 3$  (see Figure 5).  $\square$

The example given in Theorem 2 above, for  $k = 2$ , gives an UPP-DAG  $G$  with  $\pi = 2$  and a set of 5 dipaths such that the conflict graph is a  $C_5$  and therefore  $w = 3$ . Replacing each of these dipaths with




Figure 5: Internal cycle and family of dipaths with  $\pi = 2$  and  $w = 3$ .

$h$  identical dipaths we obtain a family of  $5h$  dipaths with  $\pi = 2h$  and  $w = \lceil \frac{5h}{2} \rceil$  giving a ratio  $\frac{w}{\pi} = \frac{5}{4}$ . In fact the bound can be improved to  $\frac{4}{3}$  with the following example:

**Theorem 3** *There exists an UPP- DAG  $G$  with one internal cycle and a family  $\mathcal{P}$  of dipaths such that*

$$w(G, \mathcal{P}) = \left\lceil \frac{4}{3} \pi(G, \mathcal{P}) \right\rceil$$

**Proof:** The following example is due to Frédéric Havet (private communication). It consists of 8 dipaths generating the conflict graph consisting of a cycle of length 8 plus chords between the antipodal vertices (see Figure 6). Here again  $\pi = 2$  and  $w = 3$ ; but if we replace each of these dipaths with  $h$  identical dipaths we obtain a family  $\mathcal{P}$  of  $8h$  dipaths with  $\pi = 2h$  and  $w = \lceil \frac{8h}{3} \rceil$ ; indeed in the conflict graph an independent set has at most 3 vertices and so we need at least  $\frac{8h}{3}$  colors. Therefore this family satisfies the theorem (the reader can see the relation with fractional colouring). □

In [5] we showed that the DAGs for which for any family  $\mathcal{P}$  of dipaths,  $w(G, \mathcal{P}) = \pi(G, \mathcal{P})$  are exactly those with no internal cycles. We give here a simpler proof.

**Theorem 4** *Let  $G$  be a DAG. Then, for any family of dipaths  $\mathcal{P}$ ,  $w(G, \mathcal{P}) = \pi(G, \mathcal{P})$  if and only if  $G$  does not contain an internal cycle.*

**Proof:** Let  $G$  be a DAG without internal cycles. If  $G$  is an oriented tree, this is a known result as the conflict graph is a perfect graph (see for example [15] or for a polynomial algorithm in  $O(n, |\mathcal{P}|)$  [14]). If  $G$  is not an oriented tree, let  $G'$  be the digraph obtained as follows: replace each source  $s$  with  $d^+(s)$  neighbors  $v_i$  ( $i = 1, \dots, d^+(s)$ ) by  $d^+(s)$  sources  $s_i$  and join  $s_i$  to  $v_i$ . If a dipath  $P$  of  $\mathcal{P}$  contains the arc  $(s, v_i)$  associate in  $G'$  the dipath obtained by replacing  $(s, v_i)$  by  $(s_i, v_i)$ . Do also the same transformation for all the sinks replacing the sink  $t$  with  $d^-(t)$  neighbors  $w_j$  ( $j = 1, \dots, d^-(t)$ ) by  $d^-(t)$  sinks  $t_j$  and  $(w_j, t)$






Figure 6: An other UPP-DAG with  $\pi = 2$  and  $w = 3$ .

by  $(w_j, t_j)$ . Let  $\mathcal{P}'$  be the resulting family of dipaths obtained.  $G'$  is an oriented tree ; indeed, as there is no internal cycle, all the cycles in  $G$  contain either a source or a sink. So we have  $w(G', \mathcal{P}') = \pi(G', \mathcal{P}')$ .

By construction  $\pi(G, \mathcal{P}) = \pi(G', \mathcal{P}')$ . To conclude let us color the dipaths  $P$  of  $\mathcal{P}$  following the coloring of  $\mathcal{P}'$ . If  $P$  does not contain a source or a sink it belongs to  $\mathcal{P}'$  and we keep its color. If it contains  $(s, v_i)$  (resp  $(w_j, t)$ ) we give to it the color of the associated path in  $\mathcal{P}'$  obtained by replacing  $(s, v_i)$  by  $(s_i, v_i)$  (resp  $(w_j, t)$  by  $(w_j, t_j)$ ). We get a valid coloring as there are no conflicts between two arcs  $(s, v_i)$  and  $(s, v_j)$  (resp  $(w_i, t)$  and  $(w_j, t)$ ). So we also have  $w(G, \mathcal{P}) = w(G', \mathcal{P}')$  and therefore  $w(G, \mathcal{P}) = \pi(G, \mathcal{P})$ .

□

In [5] we asked the question whether for any UPP the ratio  $\frac{w}{\pi}$  was bounded by some constant. We were only able to prove (with an involved proof) that it was the case when there was only one internal cycle obtaining a ratio  $\frac{w}{\pi} = \frac{4}{3}$  which is the best possible in view of the example of theorem 3. We refer the reader to [5] for a proof.

**Theorem 5** [5] *Let  $G$  be an UPP-DAG with only one internal cycle. Then for any family of dipaths  $\mathcal{P}$ ,*

$$w(G, \mathcal{P}) \leq \left\lceil \frac{4}{3} \pi(G, \mathcal{P}) \right\rceil$$

*and the bound is the best possible.*

In the next section we show that the ratio is unbounded at least for UPP DAGs with load 2. For that we will characterize their conflict graphs.

## 4 UPP DAGS with load 2

We will now show that if  $G$  is an UPP-DAG with load 2, for some family of dipaths  $\mathcal{P}$  then its conflict graph  $H = C(G, \mathcal{P})$  admits a "good labeling" of its edges.

We give two equivalent definitions of a good labeling of the edges of a graph  $H$ .

**Definition 1:** Let us label the edges of a graph  $H$  with distinct labels (for example the integers from 1 to  $m = |E(G)|$ ). This labeling is said to be good if for any ordered pair of vertices  $(x, y)$  there do not exist 2 paths from  $x$  to  $y$  with increasing labels.

**Definition 2:** Let us label the edges of a graph  $H$  with non necessarily distinct labels. This labeling is said to be good if for any ordered pair of vertices  $(x, y)$  there do not exist 2 paths from  $x$  to  $y$  with non decreasing labels.

Clearly a labeling satisfying definition 1 satisfies definition 2.

Conversely let  $L$  be a labeling satisfying definition 2 and let the distinct labels used in  $L$  be  $a_i$   $1 \leq i \leq k$  with  $a_1 < a_2 < \dots < a_k$  and suppose  $a_i$  is repeated  $\lambda_i$  times. Let us define the labeling  $L'$  as follows: we label the  $\lambda_i$  edges having label  $a_i$  in  $L$  with the distinct labels  $a_i, a_i + \epsilon_i, \dots, a_i + (\lambda_i - 1)\epsilon_i$  where  $\lambda_i \epsilon_i \leq a_{i+1} - a_i$ . Now consider any ordered pair of vertices  $(x, y)$ . By definition of  $L$  there exists at most one path from  $x$  to  $y$  with non decreasing labels. So all the other paths contain two consecutive edges with labels  $a_j$  and  $a_i$  such that  $a_j > a_i$ . Then in the labeling  $L'$  these two edges have also decreasing labels as  $a'_j \geq a_j \geq a_{i+1} > a_i + (\lambda_i - 1)\epsilon_j \geq a'_i$ . So  $L'$  is a labeling with distinct integers satisfying definition 1.

Let us give now some examples of graphs having a good labeling or not:

**Property 6** *The cycles  $C_4$ ,  $C_5$  and the conflict graph of Figure 6 with 8 vertices have a good labeling*

**Proof:** Let the  $C_4$  be (a,b,c,d), then a good labeling with definition 2 is obtained by labeling the edges  $\{a, b\}$  and  $\{c, d\}$  with label 1 and the edges  $\{b, c\}$  and  $\{a, d\}$  with label 3. The labeling  $L'$  is obtained by giving label 1 to  $\{a, b\}$ , 2 to  $\{c, d\}$ , 3 to  $\{b, c\}$  and 4 to  $\{a, d\}$ .

Let the  $C_5$  be (a,b,c,d,e), then a good labeling with definition 2 is obtained by labeling the edges  $\{a, b\}$   $\{c, d\}$  and  $\{a, e\}$  with label 1 and the edges  $\{b, c\}$  and  $\{d, e\}$  with label 4. The labeling  $L'$  is obtained by giving label 1 to  $\{a, b\}$ , 2 to  $\{c, d\}$ , 3 to  $\{a, e\}$ , 4 to  $\{b, c\}$  and 5 to  $\{d, e\}$ .

For the conflict graph  $H$  of Figure 6 with 8 vertices, a good labeling with definition 2 is obtained by labeling the edges of the external cycle alternatively with labels 1 and 3 and the 4 diagonals with label 2.

□

**Property 7**  *$K_{2,3}$  does not admit a good labeling.*

**Proof:** This proof is due to J-S Sereni. Let the vertices of  $K_{2,3}$  be respectively  $a, b$  and  $c, d, e$ . Suppose it admits a good labeling  $L$  with definition 1. Wlog we can suppose that  $L(a, c) < L(a, d) < L(a, e)$ .

Then  $L(b, c) > L(a, c)$  otherwise there will be two increasing paths from b to d (b,d) and (b,c,a,d).  
Then  $L(a, d) > L(b, d)$  otherwise there will be two increasing paths from a to b (a,d,b) and (a,c,b).  
Then  $L(a, e) > L(b, e)$  otherwise there will be two increasing paths from a to b (a,e,b) and (a,c,b).  
But we get a contradiction as there are two increasing paths from b to a (b,d,a) and (b,e,a).  $\square$

**Theorem 8** *Let  $G$  be an UPP-DAG with load 2. Then for any family of dipaths  $\mathcal{P}$ , the conflict graph  $C(G, \mathcal{P})$  has a good labeling.*

**Proof:** Recall (see the proof of property 1) that, if two dipaths  $P$  and  $Q$  intersect, they intersect in an interval  $[x, y]$ . As the load of  $G$  is 2, the arcs of this interval belong only to these 2 dipaths. Therefore, if  $G'$  is the digraph obtained from  $G$  by replacing the interval  $[x, y]$  by a single arc  $(x, y)$ , then  $G'$  has the same conflict graph  $H$  as  $G$ . The edge of  $H = C(G, \mathcal{P})$  joining the two vertices  $P$  and  $Q$  will correspond to the intersection interval  $[x, y]$  of  $P$  and  $Q$  in  $G$  that is to the arc  $(x, y)$  of  $G'$ . (Note that if in  $G'$  we delete the arcs with load at most 1, that is covered by at most one path of  $\mathcal{P}$ , then there is a one to one mapping between the remaining arcs of  $G'$  and the edges of the conflict graph  $H$ ).

Now we label the arcs of  $G'$  according to the topological order; that is we label 1 the arcs leaving a source ; then we delete the arcs labeled 1 getting a digraph  $G'_2$  and label 2 the arcs leaving a source in  $G'_2$  and so on. As  $G$  and therefore  $G'$  is a DAG we can label all the arcs of  $G'$ . This induces a labeling of the edges of the conflict graph  $H$ , by giving to the edge joining the two vertices  $P$  and  $Q$  the label of the arc  $(x, y)$  of  $G'$  associated to the intersection interval  $[x, y]$  of  $P$  and  $Q$ .

Let us now show that it is a good labeling of  $H$ . Consider a non decreasing path in  $H$ , from  $P$  to  $Q$ ,  $P = P_1, P_2, \dots, P_k = Q$  and let  $(x_i, y_i)$  be the arc of  $G'$  associated to the intersection of  $P_i$  and  $P_{i+1}$ . As the labels are non decreasing, then  $(x_i, y_i)$  is in the topological order before  $(x_{i+1}, y_{i+1})$  and so  $y_i$  is before  $x_{i+1}$  in  $P_{i+1}$ . So this non decreasing path in  $H$  induces in  $G'$  a dipath  $x_1, y_1, x_2, y_2, \dots, x_{k-1}, y_{k-1}$  (in fact that implies that the labels are strictly increasing). Suppose we have two non decreasing paths in  $H$  from a vertex  $P$  to a vertex  $Q$ , then we have in  $G'$  two dipaths  $x_1, y_1, x_2, y_2, \dots, x_{k-1}, y_{k-1}$  and  $x'_1, y'_1, x'_2, y'_2, \dots, x'_{m-1}, y'_{m-1}$  with  $x_1, y_1$  and  $x'_1, y'_1$  belonging to  $P$  and  $x_{k-1}, y_{k-1}$  and  $x'_{m-1}, y'_{m-1}$  belonging to  $Q$ . Wlog we can suppose  $x'_1$  is after  $y_1$  on  $P$ .

If  $x'_{m-1}$  is after  $y_{k-1}$  on  $Q$ , we have two dipaths joining  $y_1$  and  $x'_{m-1}$  namely

$y_1, x_2, y_2, \dots, x_{k-1}, y_{k-1}, x'_{m-1}$  and  $y_1, x'_1, y'_1, x'_2, y'_2, \dots, x'_{m-2}, y'_{m-2}, x'_{m-1}$ .

If  $y'_{m-1}$  is before  $x_{k-1}$  on  $Q$ , we have two dipaths joining  $y_1$  and  $x_{k-1}$  namely

$y_1, x_2, y_2, \dots, x_{k-1}$  and  $y_1, x'_1, y'_1, x'_2, y'_2, \dots, x'_{m-1}, y'_{m-1}, x_{k-1}$ .

Therefore  $G'$  and so  $G$  cannot be an UPP digraph.  $\square$

So using the property 7 we obtain the following corollary proved in [5]:

**Corollary 9** *Let  $G$  be an UPP-DAG with load 2. Then its conflict graph cannot contain a  $K_{2,3}$ .*

**Theorem 10** *Let  $H$  be a graph with a good labeling. Then there exists an UPP-DAG  $G$  with load 2 and a family of dipaths  $\mathcal{P}$  such that  $H = C(G, \mathcal{P})$ .*

**Proof:** To each edge  $\{P, Q\}$  in  $H$  let us associate in  $G$  two vertices  $x_{PQ}$  and  $y_{PQ}$  joined by the arc  $(x_{PQ}, y_{PQ})$ . Now for each vertex  $P$  in  $H$ , order its neighbors  $Q_1, Q_2, \dots, Q_h$  according to the labels of the

edges  $\{P, Q_i\}$  that is  $L(P, Q_1) < L(P, Q_2) < \dots < L(P, Q_h)$ . Then, for  $i = 1, 2, \dots, h - 1$ , let us identify the vertex  $y_{PQ_i}$  to  $x_{PQ_{i+1}}$ . To the vertex  $P$  in  $H$  we associate in  $G$  the dipath  $P = (x_{PQ_1}, y_{PQ_1} = x_{PQ_2}, y_{PQ_2}, \dots, x_{PQ_h}, y_{PQ_h})$ . The family of dipaths  $\mathcal{P}$  consists of the dipaths associated to each vertex of  $H$ . The conflict graph associated to the graph  $G$  and the family of dipaths  $\mathcal{P}$  is exactly the graph  $H$ .  $G$  has load 2 as an arc  $x_{PQ}, y_{PQ}$  of  $G$  belongs exactly to the two dipaths  $P$  and  $Q$ .  $G$  is an UPP as a dipath in  $G$  corresponds to an increasing path in  $H$  and so if there were two dipaths in  $G$  joining some  $y_{PQ}$  to  $x_{P'Q'}$  there will be two increasing paths in  $H$  from  $P$  to  $P'$ .  $\square$

If we apply the construction of the proof to the graph  $H$  with 12 vertices of Figure 6 we get exactly the graph  $G$  and the dipaths of the example.

**Remark:** there is no equivalence between the two properties  $G$  being UPP with load 2 and its conflict graph having a good labeling. Indeed there exist digraphs which are not UPP but whose conflict graph has a good labeling : for example consider the graph of Figure 4 with the dipaths  $a_1, b_1, c_1$  ;  $b_1, c_1, d_1$  ;  $c_1, d_1, e_1$ ; and  $a_1, b_1, d_1, e_1$  via the second dipath from  $b_1$  to  $d_1$ ; it has  $C_4$  as conflict graph.

**Theorem 11** *There exists a family of graphs with a good labeling and a chromatic number as large as we want.*

**Proof:** Consider a regular graph  $H$  of degree at most  $k$ , girth  $> 2k + 2$  and with a large chromatic number. The existence of such graphs has been shown in [7, 17].

The edges of  $H$  can be partitioned in at most  $k + 1$  matchings (coloration of the edges of a graph with at most  $k + 1$  colors by Vizing's theorem). Let us give to the edges of each matching a different label  $1, 2, \dots, k + 1$ . Then any non decreasing (in fact increasing as there cannot be two consecutive edges with the same label) path in  $H$  has at most  $k + 1$  edges. Therefore there cannot exist two increasing paths otherwise there will be a cycle in  $H$  of length  $\leq 2k + 2$  contradicting the value of the girth.  $\square$

Now using theorems 10 and 11 we are able to answer the question asked in [5].

**Theorem 12** *There exist UPP- DAGS and a family of dipaths with load  $\pi(G, \mathcal{P}) = 2$  and  $w(G, \mathcal{P})$  as large as we want.*

## 5 Conclusions

Many questions are worth of being investigated. In this article we study the relations between the maximum of the load  $\pi(G, \mathcal{P})$  of all the arcs and  $w(G, \mathcal{P})$  the minimum number of wavelengths in particular for the class of UPP-DAGs. In particular we have the following conjecture (a specialization of a problem asked in [2]).

CONJECTURE 1: If  $G$  is a DAG and if we consider the all to all family of dipaths (i.e for each couple  $(x,y)$  connected by a dipath we have a request. Then  $w(G, ATA) = \pi(G, ATA)$ .

In case of binary trees we have obtained an explicit formula for coloring them in that case.

A natural question is to see when property 1 is valid.

QUESTION 2: When is  $\pi(G, \mathcal{P}) = \omega(C(G, \mathcal{P}))$ .

QUESTION 3: Given an undirected graph when is it possible to orient its edges such that the digraph obtained is UPP.

That is an NP-Hard problem but classes of such graphs could be exhibited.

For graphs with one or a small number of internal cycles we have the following questions.

QUESTION 4 Is the Theorem 5 true also for DAGS (not necessarily UPP) with exactly one internal cycle ?

QUESTION 5: is there a simple proof of the Theorem 5 with one internal cycle ?

QUESTION 6: What is the bound if we have exactly k internal cycles ?

In a preliminary version of this paper we asked for a characterization of graphs with a good labeling. In particular we asked if it was polynomial or not to decide if a graph has a good labeling. In a following paper [1], the authors exhibit infinite families of graphs for which no such edge-labelling can be found. They also show that deciding if a graph admits a good edge-labelling is NP-complete. Finally, they give large classes of graphs admitting a good edge-labelling like forests,  $C_3$ -free outerplanar graphs, planar graphs of girth at least 6.

A last question consists in extending the results to UPP-DAGS with load  $> 2$  perhaps by using hypergraphs.

QUESTION 7 Characterize UPP-DAGS with load 3 or load h.

Acknowledgement: We thank J. Araujo, D.Coudert and F.Peix for their helpful remarks.

## References

- [1] J. Araujo, N. Cohen, F. Giroire, and F. Havet. Good edge-labelling of graphs. In *proceedings of the Latin-American Algorithms, Graphs and Optimization Symposium (LAGOS'09)*, Electronic Notes in Discrete Mathematics, Gramado, Brazil, November 2009. Springer.
- [2] B. Beauquier, J-C. Bermond, L. Gargano, P. Hell, S. Pérennes, and U. Vaccaro. Graph problems arising from wavelength-routing in all-optical networks. In *IEEE Workshop on Optics and Computer Science*, Geneva, Switzerland, April 1997.
- [3] B. Beauquier, P. Hell, and S. Pérennes. Optimal wavelength-routed multicasting. *Discrete Applied Mathematics*, 84:15–20, 1998.

- [4] J.-C. Bermond, L. Braud, and D. Coudert. Traffic grooming on the path. *Theoretical Computer Science*, 384(2-3):139–151, October 2007.
- [5] J.-C. Bermond and M. Cosnard. Minimum number of wavelengths equals load in a dag without internal cycle. In *Parallel and Distributed Processing Symposium, 2007. IPDPS 2007. IEEE International*, pages 1–10, Long Beach, CA, U.S.A., March 2007.
- [6] J.-C. Bermond, M. Cosnard, D. Coudert, and S. Pérennes. Optimal solution of the maximum all request path grooming problem. In *Proceedings of the Advanced International Conference on Telecommunications, AICT06*, Guadeloupe, France, February 2006.
- [7] B. Bollobas. Chromatic number, girth and maximal degree. *Discrete Mathematics*, 24(3):311–314, 1978.
- [8] I. Caragiannis, C. Kaklamanis, and P. Persiano. Wavelength routing in all-optical tree networks: A survey. *Computers and Artificial Intelligence*, 20(2):95–120, 2001.
- [9] R. Dutta and N. Rouskas. A survey of virtual topology design algorithms for wavelength routed optical networks. *Optical Networks Magazine*, 1(1):73–89, 2000.
- [10] R. Dutta and N. Rouskas. Traffic grooming in WDM networks: Past and future. *IEEE Network*, 16(6):46–56, 2002.
- [11] T. Erlebach, K. Jansen, C. Kaklamanis, M. Mihail, and P. Persiano. Optimal wavelength routing on directed fiber trees. *Theoretical Computer Science*, 221(1–2):119–137, 1999.
- [12] L. Gargano, P. Hell, and S. Pérennes. Colouring all directed paths in a symmetric tree with an application to optical networks. *Journal of Graph Theory*, 38(4):183–196, 2001.
- [13] J. Kleinberg and A. Kumar. Wavelength conversion in optical networks. In *Proceedings of SODA '99*, pages 566–575, Baltimore,MD, USA, January 1999.
- [14] F. Meunier, C. Popa, and A. Romero. Directed paths on a tree: coloring and multicut. *submitted*, 2008.
- [15] C.L. Monma and V.K. Wei. Intersection graphs of paths in a tree. *Journal of Combinatorial Theory Ser.B*, 41:141–181, 1986.
- [16] B. Mukherjee. *Optical communication networks*. McGraw-Hill,New York, 1997.
- [17] J. Nešetřil and V. Rödl. A short proof of the existence of highly chromatic hypergraphs without short cycles. *Journal of Combinatorial Theory, Ser B*, 27(2):225–227, 2001.
- [18] R. Ramaswami and K.N. Sivajaran. Routing and wavelength assignment in all-optical networks. *IEEE Transactions on NetworkingNetwork*, 3(5):489–500, 1995.
- [19] R. Ramaswami and K.N. Sivajaran. *Optical networks : a practical perspective*. Morgan kaufmann, San Francisco, 1998.