

Local Existence and Uniqueness of a Mild Solution to the One Dimensional Vlasov-Poisson System with an Initial Condition of Bounded Variation

Simon Labrunie, Sandrine Marchal, Jean Rodolphe Roche

► **To cite this version:**

Simon Labrunie, Sandrine Marchal, Jean Rodolphe Roche. Local Existence and Uniqueness of a Mild Solution to the One Dimensional Vlasov-Poisson System with an Initial Condition of Bounded Variation. [Research Report] RR-6946, INRIA. 2009, pp.14. <inria-00389829>

HAL Id: inria-00389829

<https://hal.inria.fr/inria-00389829>

Submitted on 29 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Local Existence and Uniqueness of a Mild Solution
to the One Dimensional Vlasov-Poisson System with
an Initial Condition of Bounded Variation***

Simon Labrunie — Sandrine Marchal — Jean-Rodolphe Roche

N° 6946

May 28th 2009

Thème NUM



***Rapport
de recherche***

Local Existence and Uniqueness of a Mild Solution to the One Dimensional Vlasov-Poisson System with an Initial Condition of Bounded Variation

Simon Labrunie*, Sandrine Marchal†, Jean-Rodolphe Roche‡

Thème NUM — Systèmes numériques
Équipe-Projet Calvi

Rapport de recherche n° 6946 — May 28th 2009 — 14 pages

Abstract: We propose a result of local existence and uniqueness of a mild solution to the one-dimensional Vlasov–Poisson system. We establish the result for an initial condition lying in the Sobolev space of integrable functions with integrable derivatives, then we extend it to initial conditions lying in the space of functions of bounded variation, without any assumption of continuity, boundedness or compact support.

Key-words: Vlasov-Poisson system, bounded variation functions, Banach contraction mapping principle

* Institut Elie Cartan, Nancy

† Institut Elie Cartan, Nancy

‡ Institut Elie Cartan, Nancy

Existence et unicité locales d'une "bonne" solution du système de Vlasov-Poisson unidimensionnel avec une condition initiale à variation bornée

Résumé : Nous proposons un résultat d'existence et d'unicité locales d'une "bonne" solution du système de Vlasov-Poisson unidimensionnel. On établit le résultat pour une condition initiale dans l'espace de Sobolev des fonctions intégrables à dérivées intégrables, puis on étend le résultat à l'espace plus large des fonctions à variation bornée, mais non nécessairement continues, bornées, ou à support compact.

Mots-clés : système de Vlasov-Poisson, fonctions à variation bornée, théorème de la contraction

1 Introduction

1.1 Position of the problem

In this paper we study the one-dimensional Vlasov–Poisson system:

$$\forall(t, x, v) \in [0, T] \times \mathbb{R}^2, \quad \frac{\partial f}{\partial t}(t, x, v) + v \frac{\partial f}{\partial x}(t, x, v) + E(t, x) \frac{\partial f}{\partial v}(t, x, v) = 0, \quad (1)$$

$$\forall(t, x) \in [0, T] \times \mathbb{R}, \quad \frac{\partial E}{\partial x}(t, x) = \int_{\mathbb{R}} f(t, x, v) dv, \quad (2)$$

$$\forall(x, v) \in \mathbb{R}^2, \quad f(0, x, v) = f_0(x, v). \quad (3)$$

This system models the behaviour of a gas of protons in its self-consistent electrostatic field when the collisions between particles are neglected. In [6], Cooper and Klimas show the existence and uniqueness of a global mild solution to this system, i.e. a solution defined by characteristics, for a continuous initial condition which has its first two moments in v uniformly bounded in x . This was extended by Bostan [4] to the initial-boundary value problem, with similar assumptions on the boundary data. In [10], Guo shows that there exists a unique local weak solution to (1–3) in the space $L^\infty([0, T], BV(\mathbb{R}^2))$ for initial and boundary conditions with compact support and in the space $L^\infty(\mathbb{R}^2) \cap BV(\mathbb{R}^2)$.

In this article we extend the results of Guo to the initial value problem with initial data in the space $BV(\mathbb{R}^2)$, thus not necessarily compactly supported, bounded, or continuous. Our proof is based on the contraction mapping principle of Banach, and consists of two steps: first we establish the local existence and uniqueness of a mild solution for an initial data f_0 in $W^{1,1}(\mathbb{R}^2)$, then we extend the result to $f_0 \in BV(\mathbb{R}^2)$. As will appear in the course of the proof, the hypothesis $f_0 \in BV(\mathbb{R}^2)$ is close to the minimal assumption guaranteeing the well-posedness of the characteristic system as times goes by, and thus the possibility of the existence of a mild solution.

1.2 Notations and main results

We introduce the following notations (see [10]). Given $T > 0$, we denote

$$U_T = [0, T] \times \mathbb{R} \quad \text{and} \quad V_T = [0, T] \times \mathbb{R}^2.$$

For $s \in [0, T]$, we denote $\Pi_s = \{s\} \times \mathbb{R}^2$ the slice $t = s$ of V_T . Then we introduce the following functional spaces:

$$L(T) = L^\infty([0, T], W^{1,1}(\mathbb{R}^2)), \quad X(T) = L^\infty([0, T], W^{1,\infty}(\mathbb{R})).$$

The space $L(T)$ will be that of the solutions f to the Vlasov equation with initial data $f_0 \in W^{1,1}(\mathbb{R}^2)$; we equip it with its natural norm. As for $X(T)$, it is a space of electrostatic fields E for which the characteristic curves are globally well defined and Lipschitz-continuous in all their variables [4]. This can be shown by adapting the proof of the Cauchy–Lipschitz theorem: the only difference is that we integrate L^∞ functions instead of C^0 functions and so we get continuous solutions differentiable almost everywhere in the first variable and with bounded derivative. We equip it with the following norm:

$$\forall E \in X(T), \quad \|E\|_{X(T)} = \max(\|E\|_{L^\infty([0, T] \times \mathbb{R})}, \|\partial_x E\|_{L^\infty([0, T] \times \mathbb{R})}).$$

Moreover, for any $E \in X(T)$, we set

$$C(E) = \max(\|\partial_x E\|_{L^\infty([0, T] \times \mathbb{R})}, 1), \quad (4)$$

and we denote by Y_E the Vlasov differential operator:

$$Y_E = \frac{\partial}{\partial t} + v \frac{\partial}{\partial x} + E(t, x) \frac{\partial}{\partial v}. \quad (5)$$

We recall the definition of the total variation of a function $f \in L^1(\mathbb{R}^2)$ (see for example [7], p.39):

$$\forall f \in L^1(\mathbb{R}^2), \quad TV[f] = TV_x[f] + TV_v[f], \quad (6)$$

where:

$$TV_x[f] = \limsup_{\epsilon \rightarrow 0} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \left| \frac{f(x + \epsilon, v) - f(x, v)}{\epsilon} \right| dx dv \quad (7)$$

$$TV_v[f] = \limsup_{\epsilon \rightarrow 0} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \left| \frac{f(x, v + \epsilon) - f(x, v)}{\epsilon} \right| dx dv \quad (8)$$

The space of functions of bounded variation is defined as:

$$BV(\mathbb{R}^2) = \{f \in L^1(\mathbb{R}^2) : TV[f] < +\infty\}, \quad (9)$$

and equipped with the norm $\|f\|_{BV(\mathbb{R}^2)} = \|f\|_{L^1(\mathbb{R}^2)} + TV[f]$.

Finally, we denote by $L^{bv}(T)$ the space $L^\infty([0, T], BV(\mathbb{R}^2))$ equipped with its natural norm. We shall establish the following two theorems:

Theorem 1 (Local existence and uniqueness in $W^{1,1}$) *Let $f_0 \in W^{1,1}(\mathbb{R}^2)$, and let*

$$R \geq \max(\|f_0\|_1, |f_0|_{W^{1,1}(\mathbb{R}^2)}, 1) \quad \text{and} \quad T \in \left[0, \frac{1}{R} \ln \left(\frac{R}{|f_0|_{W^{1,1}(\mathbb{R}^2)}} \right) \right].$$

Then there exists a unique mild solution $(f, E) \in L(T) \times X(T)$ to (1-3).

Moreover, we have a lower bound on the existence time T_{ex} of the maximal solution to (1-3) with initial condition f_0 . Setting $R_0 := \max(1, \|f_0\|_1)$, there holds:

$$T_{\text{ex}} \geq \frac{1}{(e-1)|f_0|_{W^{1,1}}} \quad \text{if} \quad |f_0|_{W^{1,1}} \geq \frac{R_0}{e},$$

$$T_{\text{ex}} \geq \frac{1}{R_0} \left[\ln \frac{R_0}{|f_0|_{W^{1,1}}} + \frac{1}{(e-1)} \right] \quad \text{if} \quad |f_0|_{W^{1,1}} \leq \frac{R_0}{e}.$$

Theorem 2 (Local existence and uniqueness in BV) *Let $f_0 \in BV(\mathbb{R}^2)$, and let*

$$R \geq \max(\|f_0\|_1, TV[f_0], 1) \quad \text{and} \quad T \in \left[0, \frac{1}{R} \ln \left(\frac{R}{TV[f_0]} \right) \right].$$

Then there exists a unique mild solution $(f, E) \in L^{bv}(T) \times X(T)$ to (1-3). The existence time of the maximal solution is bounded as in the $W^{1,1}$ case, with $|f_0|_{W^{1,1}}$ replaced with $TV[f_0]$.

The proof is organised as follows. In §2, we recall the definitions of weak and mild solutions to the linear Vlasov equation (i.e. (1) and (3) with E a known function of (t, x)) and to the Vlasov–Poisson system (1-3). The next section (§3) we estimate the mild solutions to the linear Vlasov equation with initial data in $W^{1,1}(\mathbb{R}^2)$, and use these results to construct a contraction mapping on a suitable set, whose fixed point gives a mild solution to the Vlasov–Poisson system. Then we extend these results to initial conditions lying in $BV(\mathbb{R}^2)$ in §4.

2 Weak and mild solutions

2.1 Definition of a weak solution

We recall the definition of a weak solution to (1-3) by using the spaces of test functions and the functionals introduced by Guo in [9]. We define two spaces of test functions, one for the Vlasov equation and the other for the Poisson equation:

$$\mathcal{V} = C_c^\infty([0, T[\times\mathbb{R}^2), \quad \mathcal{M} = C_c^\infty([0, T[\times\mathbb{R}).$$

We define for $(E, f, f_0) \in L_{\text{loc}}^\infty(U_T) \times L_{\text{loc}}^1(V_T) \times L_{\text{loc}}^1(\mathbb{R}^2)$ and $\alpha \in \mathcal{V}$ (still like in [9]) the following functional:

$$A(f, E, f_0, \alpha) = \int_{\mathbb{R}^2} f_0(x, v) \alpha(0, x, v) dx dv + \int_0^T \int_{\mathbb{R}^2} [(Y_E \alpha) f](t, x, v) dx dv dt.$$

We define for $(E, f, E_0) \in L_{\text{loc}}^\infty(U_T) \times L^1(\mathbb{R}_v, L_{\text{loc}}^1(U_T)) \times L_{\text{loc}}^\infty(\mathbb{R})$ and $\psi \in \mathcal{M}$ the following functional:

$$\begin{aligned} C(f, E, E_0, \psi) &= \int_0^T \int_{\mathbb{R}} E(t, x) \partial_x \psi(t, x) dt dx + \int_{\mathbb{R}} E_0(x) \psi(0, x) dx \\ &\quad + \int_0^T \int_{\mathbb{R}} \psi(t, x) \int_{\mathbb{R}} f(t, x, v) dv dx dt. \end{aligned}$$

These functionals are well-defined.

A weak solution to the linear Vlasov equation associated to $E \in L_{\text{loc}}^\infty(U_T)$ with initial condition $f_0 \in L_{\text{loc}}^1(\mathbb{R}^2)$ is a function $f \in L_{\text{loc}}^1(V_T)$ which satisfies:

$$\forall \alpha \in \mathcal{V}, \quad A(f, E, f_0, \alpha) = 0.$$

A weak solution to the one-dimensional Vlasov-Poisson system with initial condition $f_0 \in L^1(\mathbb{R}_v, L^1_{\text{loc}}(\mathbb{R}_x))$ is a pair $(E, f) \in L_{\text{loc}}^\infty(V_T) \times L^1(\mathbb{R}_v, L^1_{\text{loc}}(U_T))$ which verifies:

$$\begin{aligned} \forall \alpha \in \mathcal{V}, \quad &A(f, E, f_0, \alpha) = 0, \\ \forall \phi \in \mathcal{M}, \quad &C(f, E, E_0, \phi) = 0, \end{aligned}$$

where E_0 is a fixed primitive of the density function $x \mapsto \int_{-\infty}^{+\infty} f_0(x, v) dv$; for instance, if $f_0 \in L^1(\mathbb{R}^2)$ one can take: $\forall x \in \mathbb{R}, E_0(x) = \int_{-\infty}^x \int_{-\infty}^{+\infty} f_0(y, v) dv dy$.

2.2 Characteristic curves associated to $E \in X(T)$

We recall the following results on the characteristic curves of a transport equation, see for example [6] or [11]. Given $E \in X(T)$ and $(t, x, v) \in V_T$, we consider the differential system :

$$\begin{aligned} \frac{dX}{ds}(s) &= V(s), \\ \frac{dV}{ds}(s) &= E(s, X(s)), \\ (X(t), V(t)) &= (x, v). \end{aligned} \tag{10}$$

As remarked above, this system admits a unique solution for all $(t, x, v) \in V_T$, which we denote $\Gamma(s; t, x, v) = (X(s; t, x, v), V(s; t, x, v))$ and is called the characteristic curve passing by (t, x, v) .

As E is bounded on $[0, T] \times \mathbb{R}$, every characteristic curve is defined from $s = 0$ to $s = T$; moreover, the characteristic curves form a partition of V_T . Thus for every characteristic $\Gamma(s; t, x, v)$, we can define an origin on Π_0 : $\Gamma(0; t, x, v) = (X(0; t, x, v), V(0; t, x, v))$.

Let $(t, s) \in [0, T]$. We denote by $\phi_{t,s}$ the function:

$$\begin{aligned} \phi_{t,s} : \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ (x, v) &\longmapsto \Gamma(s; t, x, v). \end{aligned} \quad (11)$$

$\phi_{t,s}$ transports a point (t, x, v) of the slice Π_t to a point (s, x', v') of the slice Π_s by following the characteristic curve passing by (t, x, v) . It is well-known that $\phi_{t,s}$ is a bijection (one-to-one and onto mapping) of \mathbb{R}^2 , which admits bounded partial derivatives and whose Jacobian is identically equal to 1.

2.3 Definition of a mild solution

Let $E \in X(T)$ and (X, V) be the associated characteristic curves. A mild solution to the linear Vlasov equation associated to E with initial condition $f_0 \in L^1_{loc}(\mathbb{R}^2)$ is a function $f \in L^1_{loc}(V_T)$ which satisfies:

$$f(t, x, v) = f_0(X(0; t, x, v), V(0; t, x, v)) \quad \text{for a.e. } (t, x, v) \in V_T.$$

We recall the following result (see for example [2]):

Proposition 1 *Let $E \in X(T)$ and $f_0 \in L^1(\mathbb{R}^2)$. Then $f \in L^1(V_T)$ is a weak solution to the linear Vlasov equation associated to E with initial condition f_0 if and only if it is a mild solution.*

This can be shown by using the characteristic change of variables: $(t, x, v) \mapsto (t, x_0, v_0) = (t, \phi_{t,0}(x, v))$, as e.g. in Guo [10]. We deduce the existence and uniqueness of a solution $f \in L^1(V_T)$ to the linear Vlasov equation associated to a field $E \in X(T)$:

Corollary 1 *Let $E \in X(T)$ and $f_0 \in L^1(\mathbb{R}^2)$. The linear Vlasov equation associated to E with initial condition f_0 admits a unique weak solution in $L^1(V_T)$ defined as: $\forall (t, x, v) \in V_T$, $f(t, x, v) = f_0(X(0; t, x, v), V(0; t, x, v))$.*

Finally, a mild solution to the Vlasov–Poisson system with initial condition $f_0 \in L^1(\mathbb{R}^2)$ is defined as a weak solution (E, f) , which belongs to $X(T) \times L^1(V_T)$, and such that f coincides a.e. with the mild solution to the linear Vlasov equation associated to E with initial condition f_0 .

3 Proof of Theorem 1

3.1 A priori estimates

The proof of Theorem 1 relies on the following two theorems whose version for a half space is given by Guo in [10].

Theorem 3 *Let $E \in X(T)$ and $p \in [1, +\infty[$. We suppose that $u \in L^p(V_T)$ and $Y_E u \in L^p(V_T)$. Then:*

1. *There exists $u_0 \in L^1_{loc}(\Pi_0) \simeq L^1_{loc}(\mathbb{R}^2)$, called the trace of u on Π_0 , such that $\forall \alpha \in C_c^\infty([0, T[\times \mathbb{R}^2)$,*

$$\int_{V_T} (Y_E u \alpha + u Y_E \alpha)(t, x, v) dx dv dt = - \int_{\mathbb{R}^2} u_0(x, v) \alpha(0, x, v) dx dv.$$

2. If $u_0 \in L^p(\mathbb{R}^2)$, then $\forall s \in [0, T]$, $u(s) \in L^p(\mathbb{R}^2)$ and

$$\int_{\mathbb{R}^2} |u(s)|^p dx dv = \int_{\mathbb{R}^2} |u_0|^p dx dv + p \int_0^s \int_{\mathbb{R}^2} (\operatorname{sgn} u |u|^{p-1} Y_E u)(\tau) dx dv d\tau.$$

Theorem 4 Let $E \in X(T)$. We suppose that $u \in L^1(V_T)$ and $Y_E u \in L^1(V_T)$. Let u_0 be the trace of u on Π_0 defined in Theorem 3. If K is a measurable set of \mathbb{R}^2 with non-vanishing Lebesgue measure, then:

$$\int_{\phi_{0,s}(K)} |u(s)| dx dv = \int_K |u_0| dx dv + \int_0^s \int_{\phi_{0,\tau}(K)} (\operatorname{sgn} u Y_E u)(\tau) dx dv d\tau.$$

The proofs rely on the characteristic change of variables and are entirely similar to those of [10].

With these results, we can prove the fundamental estimate on the solutions to the linear Vlasov equation. We introduce the semi-norm $|\cdot|_{W^{1,1}}$ defined by

$$\forall f \in W^{1,1}(\mathbb{R}^2), |f|_{W^{1,1}} = \|\partial_x f\|_1 + \|\partial_v f\|_1.$$

Theorem 5 Let $E \in X(T)$ and $f_0 \in W^{1,1}(\mathbb{R}^2)$. Let f be the unique mild solution in $L^1(V_T)$ of the linear Vlasov equation associated to E . Then $\forall s \in [0, T]$, $f(s) \in W^{1,1}(\mathbb{R}^2)$ and

$$|f(s)|_{W^{1,1}(\mathbb{R}^2)} \leq |f_0|_{W^{1,1}(\mathbb{R}^2)} \exp(C(E)s). \quad (12)$$

Thus, integrating from 0 to T :

$$\int_0^T |f(\tau)|_{W^{1,1}(\mathbb{R}^2)} d\tau \leq |f_0|_{W^{1,1}(\mathbb{R}^2)} \frac{\exp(C(E)T) - 1}{C(E)}. \quad (13)$$

Proof. The set of the indefinitely differentiable functions with compact support on \mathbb{R}^2 is dense in $W^{1,1}(\Pi_0)$ [1, p. 54]. Thus there exists a sequence $(f_0^n)_n$ of elements of $C_c^\infty(\mathbb{R}^2)$, such that $\|f_0^n - f_0\|_{W^{1,1}(\mathbb{R}^2)} \rightarrow 0$ when $n \rightarrow +\infty$.

Similarly, we regularise $E \in L^\infty([0, T], W^{1,\infty}(\mathbb{R}))$ in the following way. We define for all $t \in [0, T]$, $E_n(t, \cdot) = E(t, \cdot) * \rho_n$, where $(\rho_n) \in C_c^\infty(\mathbb{R}_x)$ is a mollifying sequence. The sequence $(E_n)_n$ satisfies: $E_n \in L^\infty([0, T], W^{1,\infty} \cap C^1(\mathbb{R}_x))$; $\|E_n\|_{L^\infty(U_T)} \leq \|E\|_{L^\infty(U_T)}$, $\|\partial_x E_n\|_{L^\infty(U_T)} \leq \|\partial_x E\|_{L^\infty(U_T)}$, and $\|E - E_n\|_{L^\infty(U_T)} \rightarrow 0$ when $n \rightarrow +\infty$.

Let f_n be the solution to the linear problem associated to E_n with initial condition f_0^n . We recall that this solution is given for a.e. $(t, x, v) \in V_T$ by $f_n(t, x, v) = f_0^n(X^n(0; t, x, v), V^n(0; t, x, v))$; as f_0^n is compactly supported, so is f_n , since f_n is different from 0 only on the characteristics that start on the support of f_0^n . Moreover, the characteristics associated to E_n are Lipschitz-continuous in all their variables (s, t, x, v) , therefore $f_n \in W^{1,\infty}(V_T)$.

All together, we have $\partial_x f_n$ and $\partial_v f_n \in L_c^\infty(V_T)$, thus $\partial_x f_n$ and $\partial_v f_n$ lie in $L^1(V_T)$. Moreover $Y_{E_n} \partial_x f_n = -\partial_x E_n \partial_v f_n$ in $\mathcal{D}'(\overset{\circ}{V}_T)$, thus $Y_{E_n} \partial_x f_n$ lie in $L^1(V_T)$. By an integration by parts, it can be shown that the trace of $\partial_x f_n$ on Π_0 is $\partial_x f_0^n$. If K is a measurable subset of \mathbb{R}^2 of non-vanishing Lebesgue measure, we get by Theorem 4:

$$\int_{\phi_{0,s}(K)} |\partial_x f_n(s)| = \int_K |\partial_x f_0^n| - \int_0^s \int_{\phi_{0,\tau}(K)} (\operatorname{sgn}(\partial_x f_n) \partial_x E_n \partial_v f_n)(\tau) d\tau;$$

for the sake of brevity we have omitted the kinetic integration element $dx dv$. Thus:

$$\begin{aligned} \int_{\phi_{0,s}(K)} |\partial_x f_n(s)| &\leq \int_K |\partial_x f_0^n| + \|\partial_x E_n\|_{L^\infty([0,s] \times \mathbb{R})} \int_0^s \int_{\phi_{0,\tau}(K)} |\partial_v f_n(\tau)| d\tau, \\ \int_{\phi_{0,s}(K)} |\partial_x f_n(s)| &\leq \int_K |\partial_x f_0^n| + \|\partial_x E\|_{L^\infty(U_T)} \int_0^s \int_{\phi_{0,\tau}(K)} |\partial_v f_n(\tau)| d\tau. \end{aligned} \quad (14)$$

In the same way, we have $\partial_v f_n \in L^1(V_T)$ and $Y_E \partial_v f_n = -\partial_x f_n \in \mathcal{D}'(\overset{\circ}{V}_T)$, thus $Y_E \partial_v f_n$ lie in $L^1(V_T)$; and one shows that the trace of $\partial_v f_n$ on Π_0 is $\partial_v f_0^n$. Reasoning as above, we obtain:

$$\int_{\phi_{0,s}(K)} |\partial_v f_n(s)| \leq \int_K |\partial_v f_0^n| + \int_0^s \int_{\phi_{0,\tau}(K)} |\partial_x f_n(\tau)| d\tau. \quad (15)$$

We add (14) and (15):

$$\begin{aligned} \int_{\phi_{0,s}(K)} \{|\partial_x f_n(s)| + |\partial_v f_n(s)|\} &\leq \int_K \{|\partial_x f_0^n| + |\partial_v f_0^n|\} \\ &+ \max(\|\partial_x E\|_{L^\infty(U_T)}, 1) \int_0^s \int_{\phi_{0,\tau}(K)} \{|\partial_x f_n(\tau)| + |\partial_v f_n(\tau)|\} d\tau. \end{aligned}$$

Then we utilize the Grönwall lemma, and we get:

$$\int_{\phi_{0,s}(K)} \{|\partial_x f_n(s)| + |\partial_v f_n(s)|\} \leq \exp(C(E)s) \int_K \{|\partial_x f_0^n| + |\partial_v f_0^n|\} \quad (16)$$

Therefore:

$$\int_0^T \int_{\phi_{0,s}(K)} |\nabla f_n(s)| ds \leq \frac{\exp(C(E)T) - 1}{C(E)} \int_K |\nabla f_0^n|. \quad (17)$$

Now we utilize the Dunford–Pettis weak compactness criterion in L^1 , that can be found for example in [5, p. 76] or [3, p. 167]:

Theorem 6 (Dunford–Pettis) *Let $(f_n)_n$ be a bounded sequence of $L^1(\Omega)$. The sequence is weakly compact if and only if $\{f_n\}_{n \in \mathbb{N}}$ is equiintegrable, that is to say:*

$$\forall \epsilon > 0, \exists K_\epsilon \text{ compact } \subset \Omega \text{ s.t. } \sup_n \int_{\Omega \setminus K_\epsilon} |f_n| d\Omega < \epsilon, \quad \text{and:}$$

$$\forall \epsilon > 0, \exists \eta > 0, \forall \mathcal{A} \subset \Omega \text{ measurable, } \text{meas}(\mathcal{A}) < \eta \implies \sup_n \int_{\mathcal{A}} |f_n| d\Omega < \epsilon.$$

Let $\epsilon > 0$. The sequences $(\partial_x f_0^n)_n$ and $(\partial_v f_0^n)_n$ converge in $L^1(\mathbb{R}^2)$, thus are weakly compact in $L^1(\mathbb{R}^2)$. By the Dunford–Pettis criterion, these sequences are equiintegrable. Thus, there exists a compact K_ϵ of \mathbb{R}^2 , and $\eta > 0$ such that:

$$\sup_n \int_{\mathbb{R}^2 \setminus K_\epsilon} \{|\partial_x f_0^n| + |\partial_v f_0^n|\} < e^{-C(E)T} \epsilon, \quad \text{and:}$$

$$\forall \mathcal{A} \subset \mathbb{R}^2 \text{ measurable, } \text{meas}(\mathcal{A}) < \eta \implies \sup_n \int_{\mathcal{A}} \{|\partial_x f_0^n| + |\partial_v f_0^n|\} < e^{-C(E)T} \epsilon.$$

Reporting these inequalities in (16), we see that the sequences $(\partial_x f_n(s))_n$ and $(\partial_v f_n(s))_n$ verify the Dunford–Pettis criterion in $L^1(\mathbb{R}^2)$, thus they are weakly compact in $L^1(\mathbb{R}^2)$. Remember that $\phi_{0,s}$ is a diffeomorphism and has a Jacobian identically equal to one; hence, $\text{meas}(\phi_{0,s}(\mathcal{A})) = \text{meas}(\mathcal{A})$. Therefore, $(\partial_x f_n(s))_n$ and $(\partial_v f_n(s))_n$ converge weakly (after extracting a subsequence) in $L^1(\mathbb{R}^2)$ toward some functions g and h of $L^1(\mathbb{R}^2)$.

On the other hand, we have $Y_E(f_n - f) = (E - E_n) \partial_v f_n$, thus $f_n - f$ and $Y_E(f_n - f)$ are in $L^1(V_T)$. Applying point 2 of Theorem 3 and then the bound (17), we find:

$$\begin{aligned} \int_{\mathbb{R}^2} |f(s) - f_n(s)| &\leq \int_{\mathbb{R}^2} |f_0 - f_0^n| + \int_0^s \int_{\mathbb{R}^2} |E(\tau) - E_n(\tau)| |\partial_v f_n(\tau)| d\tau \\ &\leq \int_{\mathbb{R}^2} |f_0 - f_0^n| + \|E - E_n\|_{L^\infty(U_T)} \frac{\exp(C(E)T) - 1}{C(E)} \int_{\mathbb{R}^2} |\nabla f_0^n|. \end{aligned}$$

Thus, $f_n(s)$ converges toward $f(s)$ in $L^1(\mathbb{R}^2)$. As a consequence, $\partial_x f_n(s)$ and $\partial_v f_n(s)$ converge toward $\partial_x f(s)$ and $\partial_v f(s)$ in $\mathcal{D}'(\mathbb{R}^2)$; therefore $g = \partial_x f(s)$ and $h = \partial_v f(s)$, i.e. $\partial_x f(s)$ and $\partial_v f(s)$ lie in $L^1(\mathbb{R}^2)$. In other words, $f(s)$ appears as the weak limit in $W^{1,1}(\mathbb{R}^2)$ of the sequence $(f_n(s))_n$. By passing to the limit in (16), we get:

$$\int_{\mathbb{R}^2} |\nabla f(s)| \leq \liminf_{n \rightarrow +\infty} \int_{\mathbb{R}^2} |\nabla f_n(s)| \leq \exp(C(E)s) \|\nabla f_0\|_{L^1(\mathbb{R}^2)},$$

which is (12), and yields (13) by integrating from 0 to T . \square

3.2 Construction of a contraction mapping

We now study the non-linear Vlasov–Poisson problem. We choose $f_0 \in W^{1,1}(\mathbb{R}^2)$, and we construct a contraction mapping from a closed set of a Banach space to itself. To this end, we define the following mappings:

- $\phi_1 : X(T) \rightarrow L^1(V_T)$ maps $E \in X(T)$ to the unique weak solution f to the linear Vlasov equation associated to E and with initial condition f_0 ;
- $\phi_2 : L^1(V_T) \rightarrow L^\infty(U_T)$ maps $f \in L^1(V_T)$ to the unique solution \mathcal{E} to the Poisson equation (2) satisfying $\forall t \in [0, T], \lim_{x \rightarrow -\infty} \mathcal{E}(t, x) = 0$, namely:

$$\mathcal{E}(t, x) = \int_{-\infty}^x \int_{-\infty}^{+\infty} f(t, y, v) dv dy.$$

We have in particular $\forall (t, x) \in [0, T] \times \mathbb{R}, \mathcal{E}(t, x) \geq 0$.

The following lemma will be crucial in our proof.

Lemma 1 *For $R \geq 0$, let B'_R be the closed ball of center 0 and radius R of the Banach space $(X(T), \|\cdot\|_{X(T)})$. Then, B'_R is a closed subset of the Banach space $(L^\infty(U_T), \|\cdot\|_\infty)$, hence it is complete for this norm.*

Proof. Let $(E_n)_n$ be a sequence of elements of B'_R which converges in $L^\infty(U_T)$ toward $E \in L^\infty(U_T)$. Of course, there holds: $\|E\|_\infty \leq R$.

Then, $(\partial_x E_n)_n$ is a sequence of elements of the closed ball of center 0 and radius R of $L^\infty(U_T)$. Thus, by the Banach–Alaoglu theorem, $(\partial_x E_n)_n$ converges for the weak-* topology of $L^\infty(U_T)$ toward some $g \in L^\infty(U_T)$ with $\|g\|_\infty \leq R$. In particular, $(\partial_x E_n)_n$ converges to g in $\mathcal{D}'(\overset{\circ}{U}_T)$; but as $(E_n)_n$ converges to $E \in \mathcal{D}'(\overset{\circ}{U}_T)$, $(\partial_x E_n)_n$ converges to $\partial_x E$ in $\mathcal{D}'(\overset{\circ}{U}_T)$. Thus $\partial_x E = g$, i.e. $\|\partial_x E\|_\infty \leq R$. This proves $E \in B'_R$. \square

3.2.1 Stability and Lipschitz continuity of $\phi_2 \circ \phi_1$

Let $E \in X(T)$. Theorem 5 gives $f = \phi_1(E) \in L(T)$; moreover we have:

$$\|\phi_1(E)\|_{L(T)} \leq \|f_0\|_1 + |f_0|_{W^{1,1}(\mathbb{R}^2)} \exp(C(E)T).$$

Let $f \in L(T)$. By the definition of ϕ_2 , we have: $\partial_x \phi_2(f)(t, x) = \int_{-\infty}^{+\infty} f(t, x, v) dv$. But, as $f(t) \in W^{1,1}(\mathbb{R}^2)$ for a.e. $t \in [0, T]$, we deduce by Fubini's theorem that, for a.e. $(t, v) \in [0, T] \times \mathbb{R}$, the mapping $x \mapsto f(t, x, v)$ is in $W^{1,1}(\mathbb{R})$, hence it satisfies $\lim_{x \rightarrow -\infty} f(t, x, v) = 0$. We have thus:

$$\begin{aligned} \partial_x \phi_2(f)(t, x) &= \int_{-\infty}^{+\infty} f(t, x, v) dv = \int_{-\infty}^{+\infty} \int_{-\infty}^x \partial_x f(t, y, v) dy dv; \\ |\partial_x \phi_2(f)(t, x)| &\leq \int_{-\infty}^{+\infty} \int_{-\infty}^x |\partial_x f(t, y, v)| dy dv \leq \|\partial_x f(t)\|_{L^1(\mathbb{R}^2)}; \\ \|\partial_x \phi_2(f)\|_{L^\infty([0, T] \times \mathbb{R})} &\leq \|\partial_x f\|_{L^\infty([0, T], L^1(\mathbb{R}^2))}. \end{aligned}$$

On the other hand: $\|\phi_2(f)(t)\|_{L^\infty(\mathbb{R})} = \|f(t)\|_{L^1(\mathbb{R}^2)}$; hence:

$$\|\phi_2(f)\|_{L^\infty([0, T] \times \mathbb{R})} = \|f\|_{L^\infty([0, T], L^1(\mathbb{R}^2))} \quad \text{and} \quad \|\phi_2(f)\|_{X(T)} \leq \|f\|_{L(T)}. \quad (18)$$

So, we finally have:

$$\forall E \in X(T), \quad \|\phi_2 \circ \phi_1(E)\|_{X(T)} \leq \max(\|f_0\|_1, |f_0|_{W^{1,1}(\mathbb{R}^2)} \exp(C(E)T)). \quad (19)$$

Now we show that $\phi_2 \circ \phi_1$ is a Lipschitz-continuous mapping in the norm of $L^\infty(U_T)$. Let $E_1, E_2 \in X(T)$; we denote $f_1 = \phi_1(E_1)$ and $f_2 = \phi_1(E_2)$. There holds: $Y_{E_1}(f_1 - f_2) = (Y_{E_2} - Y_{E_1})(f_2) = (E_2 - E_1) \partial_v f_2$. Thus, $(f_1 - f_2) \in L^1(V_T)$ and $Y_{E_1}(f_1 - f_2) \in L^1(V_T)$; we apply Theorem 3 and find:

$$\int_{\mathbb{R}^2} |f_1(s) - f_2(s)| \leq \int_0^s \int_{\mathbb{R}^2} |E_1(\tau) - E_2(\tau)| |\partial_v f_2(\tau)| d\tau.$$

Thus $\|f_1 - f_2\|_{L^\infty([0, T], L^1(\mathbb{R}^2))} \leq \|E_1 - E_2\|_{L^\infty(U_T)} \|\partial_v f_2\|_{L^1(V_T)}$; applying the bound (13), we obtain:

$$\begin{aligned} \|\phi_1(E_1) - \phi_1(E_2)\|_{L^\infty([0, T], L^1(\mathbb{R}^2))} &\leq \\ \|E_1 - E_2\|_{L^\infty(U_T)} |f_0|_{W^{1,1}(\mathbb{R}^2)} &\frac{\exp(C(E_2)T) - 1}{C(E_2)}. \end{aligned} \quad (20)$$

Now let $f_1, f_2 \in L(T)$. The linearity of the Poisson equation and the bound (18) allow one to write:

$$\|\phi_2(f_1) - \phi_2(f_2)\|_{L^\infty(U_T)} \leq \|f_1 - f_2\|_{L^\infty([0, T], L^1(\mathbb{R}^2))}.$$

Finally we arrive at:

$$\begin{aligned} \|\phi_2 \circ \phi_1(E_1) - \phi_2 \circ \phi_1(E_2)\|_{L^\infty(U_T)} &\leq \\ \|E_1 - E_2\|_{L^\infty(U_T)} |f_0|_{W^{1,1}(\mathbb{R}^2)} &\frac{\exp(C(E_2)T) - 1}{C(E_2)}. \end{aligned} \quad (21)$$

3.2.2 Local existence and uniqueness

We now give conditions on the parameters R and T in order to have: (i) the closed ball B'_R stable by $\phi_2 \circ \phi_1$, and (ii) $\phi_2 \circ \phi_1$ a contraction mapping on B'_R . The

stability estimate (19) implies (i) provided: $|f_0|_{W^{1,1}(\mathbb{R}^2)} \exp(\max(R, 1)T) \leq R$ and $\|f_0\|_1 \leq R$. Thus we choose:

$$R \geq \max(|f_0|_{W^{1,1}(\mathbb{R}^2)}, \|f_0\|_1) \quad \text{and} \quad T \leq \frac{1}{\max(R, 1)} \ln\left(\frac{R}{|f_0|_{W^{1,1}(\mathbb{R}^2)}}\right).$$

As for the point (ii), the Lipschitz estimate (21) yields the sufficient condition $R \geq 1$ and $|f_0|_{W^{1,1}(\mathbb{R}^2)} (\exp(RT) - 1)/R < 1$. We take for example:

$$R \geq \max(1, |f_0|_{W^{1,1}(\mathbb{R}^2)}) \quad \text{and} \quad T < \frac{1}{R} \ln\left(1 + \frac{R}{|f_0|_{W^{1,1}(\mathbb{R}^2)}}\right).$$

Considering the two conditions, we obtain that given

$$f_0 \in W^{1,1}(\mathbb{R}^2), \quad R \geq \max(1, |f_0|_{W^{1,1}(\mathbb{R}^2)}, \|f_0\|_1), \quad T \leq \frac{1}{R} \ln\left(\frac{R}{|f_0|_{W^{1,1}(\mathbb{R}^2)}}\right),$$

the mapping $\phi_2 \circ \phi_1$ goes from B'_R into B'_R and is a contraction for the norm $\|\cdot\|_{L^\infty(U_T)}$. By Lemma 1, B'_R is a complete space for this norm. Utilizing the contraction mapping principle, the mapping $\phi_2 \circ \phi_1$ admits a unique fixed point $E \in B'_R$. If we denote $f = \phi_1(E)$, the pair $(E, f) \in X(T) \times L(T)$ is a mild solution to (1-3).

3.2.3 Estimation of the existence time

Let $f_0 \in W^{1,1}(\mathbb{R}^2)$ be fixed; we define $R_0 = \max(1, \|f_0\|_1)$. The function $x \mapsto \ln(ax)/x$ admits a unique maximum at the point $x = e/a$, and its value is a/e . Thus, the greatest value of expression $\frac{1}{R} \ln\left(\frac{R}{|f_0|_{W^{1,1}}}\right)$ is attained at $R = e|f_0|_{W^{1,1}}$ and equal to $(e|f_0|_{W^{1,1}})^{-1}$.

There are two possibilities. If $|f_0|_{W^{1,1}} \geq R_0/e$, we can take $R = R_1 := e|f_0|_{W^{1,1}}$ and $T = T_1 := (e|f_0|_{W^{1,1}})^{-1}$ in §3.2.2. The estimate (12) then shows $|f(T_1)|_{W^{1,1}} = e|f_0|_{W^{1,1}}$. So, §3.2.2 proves the existence and uniqueness of the solution to the Vlasov-Poisson problem with initial data $f(T_1)$ during the time $T_2 := (e|f(T_1)|_{W^{1,1}})^{-1} = (e^2|f_0|_{W^{1,1}})^{-1}$. Thus, the solution generated by the initial data f_0 exists during $T_1 + T_2$. By induction, we obtain an existence time at least equal to:

$$\frac{1}{|f_0|_{W^{1,1}}} \left(\frac{1}{e} + \frac{1}{e^2} + \cdots + \frac{1}{e^n} + \cdots \right) = \frac{1}{(e-1)|f_0|_{W^{1,1}}}.$$

Now, if $|f_0|_{W^{1,1}} \leq R_0/e$, the existence time given by §3.2.2 is maximal for $R = R_0$ and equal to is equal to $T_0 := \frac{1}{R_0} \ln\left(\frac{R_0}{|f_0|_{W^{1,1}}}\right)$. Applying (12), we obtain $|f(T_0)|_{W^{1,1}} = e^{R_0 T_0} |f_0|_{W^{1,1}} = R_0 > R_0/e$. Thus we can use the previous argument to show that the solution to the Vlasov-Poisson problem with initial data $f(T_0)$ exists for a time at least equal to $((e-1)R_0)^{-1}$. Finally, the total existence time is no less than

$$\frac{1}{R_0} \left(\ln\left(\frac{R_0}{|f_0|_{W^{1,1}}}\right) + \frac{1}{(e-1)} \right).$$

4 Proof of Theorem 2

4.1 Preliminary results

Here we collect some well-known results on the functions of $W^{1,1}(\mathbb{R}^2)$ and $BV(\mathbb{R}^2)$. The following proposition can be found, for example, in [8, pp. 3-4]:

Proposition 2 $W^{1,1}(\mathbb{R}^2) \subset BV(\mathbb{R}^2)$ and $\forall f \in W^{1,1}(\mathbb{R}^2)$, $|f|_{W^{1,1}} = TV[f]$.

The following two theorems are taken from [8], p. 7 and p. 14:

Theorem 7 *Let $f \in L^1(\mathbb{R}^2)$ and $(f_n)_n$ be a sequence in $BV(\mathbb{R}^2)$ which converges to f in $L^1(\mathbb{R}^2)$. Then:*

$$TV[f] \leq \liminf_{n \rightarrow +\infty} TV[f_n].$$

Theorem 8 *Let $f \in BV(\mathbb{R}^2)$. There exists a sequence $(f_n)_n$ in $C^\infty(\mathbb{R}^2) \cap BV(\mathbb{R}^2)$ such that:*

$$\lim_{n \rightarrow +\infty} \|f_n - f\|_{L^1(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} TV[f_n] = TV[f].$$

4.2 A priori estimates

Theorem 9 *Let $f \in L^1(V_T)$ be the unique mild solution to the linear Vlasov equation associated to $E \in X(T)$ with initial condition $f_0 \in BV(\mathbb{R}^2)$. Then, $\forall s \in [0, T]$, $f(s) \in BV(\mathbb{R}^2)$ and*

$$TV[f(s)] \leq TV[f_0] \exp(C(E)s).$$

Thus, integrating from 0 to T :

$$\int_0^T TV[f(\tau)] d\tau \leq TV[f_0] \frac{\exp(C(E)T) - 1}{C(E)}.$$

Proof. Let E , f_0 , and f be as in the statement of the theorem. Theorem 8 yields the existence of a sequence $(f_0^n)_n$ in $C^\infty(\mathbb{R}^2) \cap BV(\mathbb{R}^2)$ such that:

$$\lim_{n \rightarrow +\infty} \|f_0^n - f_0\|_{L^1(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} TV[f_0^n] = TV[f_0].$$

In particular, we have: $\forall n \in \mathbb{N}$, $f_0^n \in W^{1,1}(\mathbb{R}^2)$.

Let f_n be the unique mild solution to the linear Vlasov equation associated to E with initial condition f_0^n . Using Theorem 5, we get that $\forall s \in [0, T]$, $f_n(s) \in W^{1,1}(\mathbb{R}^2)$, and

$$|f_n(s)|_{W^{1,1}(\mathbb{R}^2)} \leq |f_0^n|_{W^{1,1}(\mathbb{R}^2)} \exp(C(E)s),$$

thus, by Proposition 2:

$$TV[f_n(s)] \leq TV[f_0^n] \exp(C(E)s).$$

We have $Y_E(f_n - f) = Y_E(f_n) - Y_E(f) = 0$, so we can use Theorem 4 and obtain:

$$\int_{\mathbb{R}^2} |f_n(s) - f(s)| = \int_{\mathbb{R}^2} |f_0^n - f_0|.$$

Therefore, $\lim_{n \rightarrow +\infty} f_n(s) = f(s)$ in $L^1(\mathbb{R}^2)$, for almost every $s \in [0, T]$. Applying Theorem 7 then yields:

$$\begin{aligned} TV[f(s)] &\leq \liminf_{n \rightarrow +\infty} TV[f_n(s)] \leq \liminf_{n \rightarrow +\infty} TV[f_0^n] \exp(C(E)s) \\ &= TV[f_0] \exp(C(E)s), \end{aligned}$$

which implies $f(s) \in BV(\mathbb{R}^2)$. □

4.3 Construction of a contraction mapping

We now get down to the non-linear Vlasov–Poisson problem. We define the mappings ϕ_1 and ϕ_2 as in §3.2, and we find sufficient conditions for $\phi_2 \circ \phi_1$ to be a contraction mapping from B'_R to itself.

Let $f_0 \in BV(\mathbb{R}^2)$, $E \in X(T)$ and $f = \phi_1(E)$. By Theorem 9, $f \in L^{bv}(T)$ and

$$\|\phi_1(E)\|_{L^{bv}(T)} \leq \|f_0\|_{BV(\mathbb{R}^2)} \exp(C(E)T).$$

Let us examine the mapping ϕ_2 . As in §3.2.1, we find: $\|\phi_2(f)\|_{L^\infty([0,T] \times \mathbb{R})} = \|f\|_{L^\infty([0,T], L^1(\mathbb{R}^2))}$ and $\partial_x \phi_2(f)(t, x) = \int_{-\infty}^{+\infty} f(t, x, v) dv$. Then we state and prove the following lemma:

Lemma 2 *Let $f \in BV(\mathbb{R}^2)$. We denote by $\rho[f]$ the function of $L^1(\mathbb{R})$ defined by $\forall x \in \mathbb{R}$, $\rho[f](x) = \int_{-\infty}^{+\infty} f(x, v) dv$. Then, $\rho[f] \in L^\infty(\mathbb{R})$ and $\|\rho[f]\|_\infty \leq TV[f]$.*

Proof. According to Theorem 8, there exists a sequence $(f_n)_n$ of functions in $C^\infty(\mathbb{R}^2) \cap BV(\mathbb{R}^2) \subset W^{1,1}(\mathbb{R}^2)$ such that $\|f_n - f\|_1 \rightarrow 0$ and $TV[f_n] \rightarrow TV[f]$ when $n \rightarrow +\infty$.

From §3.2.1, we know that $\rho[f_n](x) \leq \|\partial_x f_n\|_1 \leq TV[f_n]$. But $\rho[f_n]$ converges to $\rho[f]$ in $L^1(\mathbb{R})$, thus there exists a subsequence $\rho[f_{\sigma(n)}]$ which converges almost everywhere to $\rho[f]$. We have $\rho[f_{\sigma(n)}](x) \leq TV[f_{\sigma(n)}]$ and passing to the limit we get for a.e. $x \in \mathbb{R}$, $\rho[f](x) \leq TV[f]$. Therefore, $\rho[f] \in L^\infty(\mathbb{R})$ and $\|\rho[f]\|_\infty \leq TV[f]$. \square

Lemma 2 gives: $\|\phi_2(f)\|_{X(T)} \leq \|f\|_{L^{bv}(T)}$. Thus we have:

$$\forall E \in X(T), \quad \|\phi_2 \circ \phi_1(E)\|_{X(T)} \leq \|f_0\|_{BV(\mathbb{R}^2)} \exp(C(E)T).$$

Now we establish that the mapping $\phi_2 \circ \phi_1$ is Lipschitz continuous in the norm of $L^\infty([0, T] \times \mathbb{R})$. Let $E_1, E_2 \in X(T)$; we denote $f_1 = \phi_1(E_1)$ and $f_2 = \phi_1(E_2)$. Moreover, as we did in the proof of Theorem 9, we approximate f_0 by a sequence $(f_0^n)_n$ whose terms lie in $W^{1,1}(\mathbb{R}^2)$, and such that

$$\lim_{n \rightarrow +\infty} \|f_0^n - f_0\|_{L^1(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} TV[f_0^n] = TV[f_0].$$

The solutions to the linear Vlasov equation with field E_1 (resp. E_2) and initial condition f_0^n will be denoted f_1^n (resp. f_2^n). Applying the $W^{1,1}$ estimate (20) to these functions yields:

$$\|f_1^n - f_2^n\|_{L^\infty([0,T], L^1(\mathbb{R}^2))} \leq \|E_1 - E_2\|_{L^\infty(U_T)} TV[f_0^n] \frac{\exp(C(E_2)T) - 1}{C(E_2)}. \quad (22)$$

As seen in the proof of Theorem 9, we have

$$\|f_i^n(s) - f_i(s)\|_{L^1(\mathbb{R}^2)} = \|f_0^n - f_0\|_{L^1(\mathbb{R}^2)}, \quad \text{for a.e. } s \in [0, T], \text{ and } i = 1, 2.$$

Thus, f_i^n converges toward f_i in $L^\infty([0, T], L^1(\mathbb{R}^2))$. Passing to the limit in (22), we obtain:

$$\|f_1 - f_2\|_{L^\infty([0,T], L^1(\mathbb{R}^2))} \leq \|E_1 - E_2\|_{L^\infty(U_T)} TV[f_0] \frac{\exp(C(E_2)T) - 1}{C(E_2)}.$$

Then, the linearity of ϕ_2 and the bound (18) imply:

$$\begin{aligned} \|\phi_2 \circ \phi_1(E_1) - \phi_2 \circ \phi_1(E_2)\|_{L^\infty(U_T)} &\leq \\ &\|E_1 - E_2\|_{L^\infty(U_T)} TV[f_0] \frac{\exp(C(E_2)T) - 1}{C(E_2)}. \end{aligned} \quad (23)$$

Reasoning like in §3.2.2, we infer that $\phi_2 \circ \phi_1$ admits a unique fixed point in B'_R for suitable values of R and T (using the contraction mapping principle of Banach), then we deduce the local existence and uniqueness of a mild solution to (1–3). The existence time is estimated as in §3.2.3.

References

- [1] R.A. ADAMS. *Sobolev spaces*. Pure and applied mathematics, **65**, Academic Press, 1975.
- [2] R. BEALS, V. PROTOPODESCU. *Abstract time-dependent transport equations*. J. Math. Anal. Appl. **121** (1987), pp. 370–405.
- [3] B. BEAUZAMY. *Introduction to Banach spaces and their geometry*. Mathematics studies **68**, North-Holland, 1985.
- [4] M. BOSTAN. *Existence and uniqueness of the mild solution for the 1D Vlasov–Poisson initial-boundary value problem*. SIAM J. Math. Anal. **37** (2005), 156–188.
- [5] H. BREZIS. *Analyse fonctionnelle*. Mathématiques appliquées pour la maîtrise, Dunod, 1999.
- [6] J. COOPER, A. KLIMAS. *Boundary Value Problems for the Vlasov–Maxwell Equation in One Dimension*. J. Math. Anal. Appl. **75**, 306–329 (1980).
- [7] F. FILBET. *Contribution à l’analyse et à la simulation numérique de l’équation de Vlasov*. Thèse de doctorat, Université Nancy 1, 2001.
- [8] E. GIUSTI. *Minimal surfaces and functions of bounded variation*. Monographs in mathematics, Birkhäuser, 1984.
- [9] Y. GUO. *Global weak solutions of the Vlasov–Maxwell system with boundary conditions*. Commun. Math. Phys. **174**, 245–263, 1993.
- [10] Y. GUO. *Singular Solutions of the Vlasov–Maxwell System on a Half Line*. Arch. Rational Mech. Anal. **131** (1995) 241–304. Springer-Verlag 1995.
- [11] S. UKAI, T. OKABE. *On classical solutions in the large in time of two-dimensional Vlasov’s equation*. Osaka J. Math. **15** (1978), 245–261.



Centre de recherche INRIA Nancy – Grand Est
LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Centre de recherche INRIA Bordeaux – Sud Ouest : Domaine Universitaire - 351, cours de la Libération - 33405 Talence Cedex
Centre de recherche INRIA Grenoble – Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier
Centre de recherche INRIA Lille – Nord Europe : Parc Scientifique de la Haute Borne - 40, avenue Halley - 59650 Villeneuve d'Ascq
Centre de recherche INRIA Paris – Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex
Centre de recherche INRIA Rennes – Bretagne Atlantique : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex
Centre de recherche INRIA Saclay – Île-de-France : Parc Orsay Université - ZAC des Vignes : 4, rue Jacques Monod - 91893 Orsay Cedex
Centre de recherche INRIA Sophia Antipolis – Méditerranée : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399