

## Extension of LCAO to excited states: non interacting response and molecular spectra

Petr Koval, Olivier Coulaud, Dietrich Foerster

► **To cite this version:**

Petr Koval, Olivier Coulaud, Dietrich Foerster. Extension of LCAO to excited states: non interacting response and molecular spectra. ETSF 2009: Ab-initio tools for the characterization of nanostructures, Sep 2009, Evora, Portugal. pp.20. <inria-00437608>

**HAL Id: inria-00437608**

**<https://hal.inria.fr/inria-00437608>**

Submitted on 1 Dec 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Extension of LCAO to excited states:  
non interacting response and molecular spectra

Peter Koval, Olivier Coulaud, Dietrich Foerster

ANR



Evora 16/09/2009

## Petersilka–Gossmann–Gross equation: reminder

- ▶ Interacting response  $\chi$  satisfies<sup>1</sup>

$$\chi^{-1} = \chi_0^{-1} - \Sigma$$

$$\Sigma = \frac{\delta(t - t')}{|\mathbf{r} - \mathbf{r}'|} + \frac{\delta V_{xc}(\mathbf{r}, t)}{\delta n(\mathbf{r}', t')} \text{ still to be determined.}$$

- ▶ Interacting polarizability  $P^{ik}(\omega) = d^i \chi d^k$  is the goal.

---

<sup>1</sup>Petersilka, Gossmann and Gross, Phys. Rev. Lett., **76** (1996) 1212

Construction of  $\chi_0$  from its spectral function

- ▶ Ansatz  $f_a(\mathbf{r})f_b(\mathbf{r}) = \sum_\mu V_{ab}^\mu F_\mu(\mathbf{r})$  leads to

$$\chi_{\mu\nu}^0(\omega) = \sum_{E < 0, F > 0} \sum_{pqrs} \chi_E^p \chi_F^q \chi_E^r \chi_F^s \frac{V_{pq}^\mu V_{rs}^\nu}{\omega - (E - F) + i\epsilon}$$

- ▶ Spectral function  $\rho_{\mu\nu} = \text{Im}\chi_{\mu\nu}^0$

$$\rho_{\mu\nu}(\lambda) = \sum_{E < 0, F > 0} \delta(\lambda - (E - F)) \sum_{pqrs} \chi_F^q \chi_F^s \chi_E^p \chi_E^r V_{pq}^\mu V_{rs}^\nu$$

- ▶  $\rho_{\mu\nu}(\lambda)$  via convolution

$$\rho_{\mu\nu}(\lambda) = \sum_{pqrs} \rho_{qs}^-(F) * \rho_{pr}^+(\lambda - F) V_{pq}^\mu V_{rs}^\nu$$

electronic spectral function  $\rho_{qs}^-(F < 0) = \sum_\nu \delta(\nu - F) \chi_F^q \chi_F^s$

- ▶ Discretise  $\lambda = E - F \Rightarrow \chi_{\mu\nu}^0(\omega)$  is again convolution

$$\chi_{\mu\nu}^0(\omega) = \sum_\lambda \rho_{\mu\nu}(\lambda) \cdot (\omega - \lambda + i\epsilon)^{-1}$$

## Spectral function: discretisation

- ▶ FFT needs equidistant grid, but energy differences do not fit any equidistant grid



- ▶ Solution: 'occupy' adjacent nodes



- ▶ Result:  $\chi_0$  exact up to discretisation

Coulomb self energy  $\Sigma_{\text{H}}$ : sketch of basis

- ▶ In the basis of dominant products  $\Sigma_{\text{H}}$  reads

$$\Sigma_{\text{H}}^{\mu\nu} = \int dr dr' F^{\mu}(r) |r - r'|^{-1} F^{\nu}(r').$$

Dominant functions  $F^{\mu}(r)$  are either local or bilocal.



- ▶ Generally  $F^{\mu}(r) = \sum_j F_j^{\mu}(r') S_{jm}(\mathbf{R}_{\mu} r')$ ,  $r' = r - \mathbf{C}_{\mu}$ .
- ▶ Radial functions  $F_j^{\mu}(r')$  are given on a logarithmic grid.

Coulomb self energy  $\Sigma_{\text{H}}$ : method

- ▶  $\Sigma_{\text{H}}^{\mu\nu}$  can be reduced to a sum over elementary Coulomb interactions

$$E_{jm,j'm'}(\mathbf{c}, \mathbf{c}') = \int d\mathbf{r} d\mathbf{r}' \frac{g_{jm}(\mathbf{r} - \mathbf{c}) g_{j'm'}(\mathbf{r}' - \mathbf{c}')}{|\mathbf{r} - \mathbf{r}'|}$$

$$g_{jm}(\mathbf{r}) = g_j(r) S_{jm}(\mathbf{r})$$

- ▶  $E_{jm,j'm'}(\mathbf{c}, \mathbf{c}') = \langle g_{jm} | p^{-2} | g_{j'm'} \rangle$  in momentum space.
- ▶ Conversion to momentum space by Talman's transform<sup>2</sup>.
- ▶ No 4-center integrals involved :-)

---

<sup>2</sup>Talman J. D., Comput. Phys. Commun. **180** 332 (2009)



## Dynamical polarizability: CPU costs

- ▶ Non interacting response  $\chi_0$   $O(N^2 N_\omega)$
  - ▶ Self energies  $\Sigma_H, \Sigma_{xc}$   $O(N)$
  - ▶ Solving  $\chi^{-1} = \chi_0^{-1} - \Sigma$   $O(N^3 N_\omega)$
  - ☺ Fortunately  $P^{ik} = d^i \chi d^k$  can be calculated iteratively,  
thanks to a special Lanczos method  $O(N^2 N_\omega)$
- 
- ✓ Interacting polarizability  $P^{ik}$ , total CPU  $O(N^2 N_\omega)$



# Performance of iterative method: naphthalene $C_{10}H_8$

How well does the iterative method work?




- ▶ 898 dominant products  $\rightarrow$  6 Krylov vectors.
- ▶ CPU cost is dramatically reduced.

# Performance of dominant products basis: benzene $C_6H_6$

How many dominant products  $F^\mu(\mathbf{r})$  needed?

- ▶ Compare with exact result in the conventional basis.
- ▶  $\chi_{EF,PQ}^0$  is diagonal ☺, but interaction kernel costs  $O(N^3)$  ☹



interaction kernel  $K$  via  
 self energy  $\Sigma$ , vertex  $V$   
 and density matrix  $D$   
 $K = (DV)\Sigma(V^T D)$



- ▶ Original basis 5832 products  $f_a(\mathbf{r})f_b(\mathbf{r})$   
 Dominant basis 1236 products  $F^\mu(\mathbf{r})$ .

# Comparison with experiment: indigo dye $C_{16}H_{10}N_2O_2$

How good can be comparison with experiment<sup>3</sup> ?





- ▶ Original basis 48050 products  $f_a(\mathbf{r})f_b(\mathbf{r})$   
Dominant basis 2831 products  $F^\mu(\mathbf{r})$ .
- ▶ One hour on 8 core machine; 11 GByte RAM.

<sup>3</sup>Ross Brown, IPREM unpublished (2008)

Scaling of CPU time: carbon chains  $C_n$ 

Does the approach really scale as  $N^2 N_\omega$  ?




- ▶ Size of matrix is proportional to the number of atoms  $N$ ,  $O(N^2)$  non zero elements.
- ▶ Each element costs  $O(N_\omega)$  because of FFT.

# Parallelization approach versus runtime

Timing for benzene example, sequential version

Task	Time, s	Parallelization
Basis of dominant products	10	OMP
Self energies	160	OMP+MPI
Response function	1100	OMP+MPI
Excitation spectrum	120	OMP+MPI

- ▶ Easy to parallelize
- ▶ OMP mostly done
- ▶ MPI for response function  $\chi_0$  and polarizability  $P_{ik}(\omega)$  together



## Conclusions and Outlook

- ✓  $\chi^0$  is constructed in  $O(N^2 N_\omega)$  operations.
- ✓  $\Sigma$  is cheap in dominant products: no 4-center integrals
- ✓  $P^{ik}$  in  $O(N^2 N_\omega)$  operations due to Lanczos.
- ▶ The algorithm is easily parallelizable
  - ⊕ Memory bottleneck made less severe with MPI.
- ▶  $\chi^0$  in the basis of dominant products will be useful in applications to GW and Bethe–Salpeter.

# Acknowledgements

- ▶ James Talman, London, Canada
- ▶ Mark Casida, Grenoble, France
- ▶ Andrei Postnikov, Metz, France
- ▶ Ross Brown, Pau, France

ANR



\$\$ Work supported by ANR-project **NOSSI**

**Nouveaux Outils pour la Simulation des Solides et des Interfaces**