

Unsatisfiable Linear CNF Formulas Are Large, and Difficult to Construct Explicitly

Dominik Scheder

► **To cite this version:**

Dominik Scheder. Unsatisfiable Linear CNF Formulas Are Large, and Difficult to Construct Explicitly. Jean-Yves Marion and Thomas Schwentick. 27th International Symposium on Theoretical Aspects of Computer Science - STACS 2010, Mar 2010, Nancy, France. pp.621-632, 2010, Proceedings of the 27th Annual Symposium on the Theoretical Aspects of Computer Science. <inria-00455809>

HAL Id: inria-00455809

<https://hal.inria.fr/inria-00455809>

Submitted on 11 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNSATISFIABLE LINEAR CNF FORMULAS ARE LARGE AND COMPLEX

DOMINIK SCHEDER

ETH Zürich, Institute of Theoretical Computer Science
Universitätstrasse 6, CH-8092 Zürich, Switzerland
E-mail address: dscheder@inf.ethz.ch

ABSTRACT. We call a CNF formula *linear* if any two clauses have at most one variable in common. We show that there exist unsatisfiable linear k -CNF formulas with at most $4k^2 4^k$ clauses, and on the other hand, any linear k -CNF formula with at most $\frac{4^k}{8e^{2k^2}}$ clauses is satisfiable. The upper bound uses probabilistic means, and we have no explicit construction coming even close to it. One reason for this is that unsatisfiable linear formulas exhibit a more complex structure than general (non-linear) formulas: First, any treelike resolution refutation of any unsatisfiable linear k -CNF formula has size at least $2^{2^{\frac{k}{2}-1}}$. This implies that small unsatisfiable linear k -CNF formulas are hard instances for Davis-Putnam style splitting algorithms. Second, if we require that the formula F have a *strict* resolution tree, i.e. every clause of F is used only once in the resolution tree, then we need at least $a^{a^{\dots^a}}$ clauses, where $a \approx 2$ and the height of this tower is roughly k .

1. Introduction

How can CNF formulas become unsatisfiable? Roughly speaking, there are two ways: Either some constraint (clause) is itself impossible to satisfy – the empty clause; or, every clause can be satisfied individually, but one cannot satisfy all of them simultaneously. In the latter case, the clauses have to somehow overlap. How much? For example, take k boolean variables x_1, \dots, x_k . The conjunction of all 2^k possible clauses of size k is the *complete k -CNF formula* and denote by \mathcal{K}_k . It is unsatisfiable, and as small as possible: Any k -CNF formula with less than 2^k clauses is satisfiable. Clearly, the clauses of \mathcal{K}_k overlap a lot. What if we require that any two distinct clauses share at most one variable? We call such a formula *linear*. There are unsatisfiable linear k -CNF formulas, but they are significantly larger and have a much more complex structure than \mathcal{K}_k .

A CNF formula is a conjunction (AND) of *clauses*, and a clause is a disjunction (OR) of *literals*. A literal is either a boolean variable x or its negation \bar{x} . We require that a clause does not contain the same literal twice, and does not contain complementary literals, i.e.,

1998 ACM Subject Classification: Computational and structural complexity.

Key words and phrases: Extremal Combinatorics, Proof Complexity, Probabilistic Method.

Research is supported by the SNF Grant 200021-118001/1.

both x and \bar{x} . To simplify notation, we also regard formulas as sets of clauses and clauses as sets of literals. A clause with k literals is a k -clause, and a k -CNF formula is a CNF formula consisting of k -clauses. For a clause C , we denote by $\text{vbl}(C)$ set of variables x with $x \in C$ or $\bar{x} \in C$. Consequently, a CNF formula F is *linear* if $|\text{vbl}(C) \cap \text{vbl}(D)| \leq 1$ for any two distinct clauses $C, D \in F$. As a relaxation of this notion, we call F *weakly linear* if $|C \cap D| \leq 1$ for any distinct $C, D \in F$.

Example. The formula $(\bar{x}_1 \vee x_2) \wedge (\bar{x}_2 \vee x_3) \wedge (x_3 \vee x_4) \wedge (\bar{x}_4 \vee \bar{x}_1)$ is linear, whereas $(\bar{x}_1 \vee x_2) \wedge (x_1 \vee x_2) \wedge (x_2 \vee x_3)$ is weakly linear, but not linear, and finally $(x_1 \vee x_2 \vee x_3) \wedge (x_1 \vee x_2 \vee \bar{x}_3)$ is not weakly linear (and not linear, either).

It is not very difficult to construct an unsatisfiable linear 2-CNF formula, but significantly more effort is needed for a 3-CNF formula. It is not obvious whether unsatisfiable linear k -CNF formulas exist for every k . These questions have been asked first by Porschen, Speckenmeyer and Randerath [15], who also proved that for any $k \geq 3$, if an unsatisfiable linear k -CNF formula exists, then deciding satisfiability of linear k -CNF formulas is NP-complete. Later, Porschen, Speckenmeyer and Zhao [16] and, independently, myself [18] gave a construction of unsatisfiable linear k -CNF formulas, for every $k \in \mathbb{N}_0$:

Theorem 1.1 ([16], [18]). *For every $k \geq 0$, there exists an unsatisfiable linear k -CNF formula F_k , with F_0 containing one clause and F_{k+1} containing $|F_k|2^{|F_k|}$ clauses.*

The $|F_k|$ are extremely large. Here, we will give an almost optimal construction.

Theorem 1.2. *All weakly linear k -CNF formulas with at most $\frac{4^k}{8e^2(k-1)^2}$ clauses are satisfiable. There exists an unsatisfiable linear k -CNF formula with $4k^24^k$ clauses.*

It is a common phenomenon in extremal combinatorics that by probabilistic means one can show that a certain object exists (in our case, a “small” linear unsatisfiable k -CNF formula), but one cannot explicitly construct it. We have no explicit construction avoiding the tower-like growth in Theorem 1.1. We give some arguments why this is so, and show that small linear unsatisfiable k -CNF formulas have a more complex structure than their non-linear relatives. To do so, we speak about *resolution*.

1.1. Resolution Trees

If C and D are clauses and there is unique literal u such that $u \in C$ and $\bar{u} \in D$, then $(C \setminus \{u\}) \cup (D \setminus \{\bar{u}\})$ is called the *resolvent* of C and D . It is easy to check that every assignment satisfying C and D also satisfies the resolvent.

Definition 1.3. A *resolution tree* for a CNF formula F is a tree T whose vertices are labeled with clauses, such that

- each leaf of T is labeled with a clause of F ,
- the root of T is labeled with the empty clause,
- if vertex a has children b and c , and these are labeled with clauses C_a, C_b, C_c , respectively, then C_a is the resolvent of C_b and C_c .

It is well-known that a CNF formula F is unsatisfiable if and only if it has a resolution tree (which can be exponentially large in $|F|$). Proving lower bounds on the size of resolution

trees (and general resolution proofs, which we will not introduce here) has been and still is an area of intensive research. See for example Ben-Sasson and Wigderson [2].

Theorem 1.4. *Let $k \geq 2$. Every resolution tree of an unsatisfiable weakly linear k -CNF formula has at least $2^{2^{\frac{k}{2}-1}}$ leaves.*

A large ratio between the size of F and the size of a smallest resolution tree is an indication that F has a complex structure. For example, it is well-known that the running time of so-called Davis-Putnam procedures on a formula F is lower bounded by the size of the smallest resolution tree of F (actually those procedures were introduced by Davis, Logeman and Loveland [3]). Such a procedure tries to find a satisfying assignment for a formula F (or to prove that none exists) by choosing a variable x , and then recursing on the formulas $F^{[x \rightarrow 0]}$ and $F^{[x \rightarrow 1]}$, obtained from F by fixing the value of x to 0 or 1, respectively. If F is unsatisfiable, the procedure implicitly constructs a resolution tree.

A CNF formula F is *minimal unsatisfiable* if it is unsatisfiable, and for every clause $C \in F$, $F \setminus \{C\}$ is satisfiable. The complete k -CNF formula introduced above is minimal unsatisfiable, and has a resolution tree with 2^k leaves, one for every clause. This is as small as possible, since for a minimal unsatisfiable formula, every clause must appear as label of at least one leaf of any resolution tree. We call a resolution tree *strict* if no two leaves are labeled by the same clause, and a formula F *strictly treelike* if it has a strict resolution tree. In some sense, strictly treelike formulas are the least complex formulas possible. For example, the complete formula \mathcal{K}_k and the formulas constructed in the proof of Theorem 1.1 are strictly treelike.

Theorem 1.5. *For any $\epsilon > 0$, there exists a constant c such that for any $k \in \mathbb{N}$, any strictly treelike weakly linear k -CNF formula has at least $\text{tower}_{2-\epsilon}(k - c)$ clauses, where $\text{tower}_a(n)$ is defined by $\text{tower}_a(0) = 1$ and $\text{tower}_a(n + 1) = a^{\text{tower}_a(n)}$.*

Strictly treelike formulas appear in other contexts, too. Consider $\text{MU}(1)$, the class of minimal unsatisfiable formulas whose number of variables is one less than the number of clauses. A result of Davydov, Davydova and Kleine Büning ([4], Theorem 12) implies that every $\text{MU}(1)$ -formula is strictly treelike. Also, $\text{MU}(1)$ -formulas serve as “universal patterns” for unsatisfiable formulas: Szeider [19] shows that a formula F is unsatisfiable if and only if it can be obtained from a $\text{MU}(1)$ -formula G by renaming the variables of G (in a possibly non-injective manner). It is not difficult to show that a strictly treelike linear k -CNF formula can be transformed into a linear $\text{MU}(1)$ -formula with the same number of clauses.

1.2. Related Work

For a CNF formula F and a variable x , let $d_F(x)$ denote the *degree* of x , i.e. the number of clauses of F containing x or \bar{x} , and let $d(F) := \max_x d_F(x)$ denote the *maximum degree* of F . For the complete k -CNF formula \mathcal{K}_k , we have $d(\mathcal{K}_k) = 2^k$. Intuitively, in an unsatisfiable k -CNF formula, some variables should occur in many clauses. In other words, the following function should be large:

$$f(k) := \max\{d \mid \text{every } k\text{-CNF formula } F \text{ with } d(F) \leq d \text{ is satisfiable}\} . \tag{1.1}$$

The function $f(k)$ has first been investigated by Tovey [20], who showed $f(k) \geq k$, using Hall's Theorem. Using the famous Lovász Local Lemma (see [5] for the original proof, or [1] for several generalized versions), Kratochvíl, Savický and Tuza [12] proved that $f(k) \geq \frac{2^k}{ek}$, and that while all k -CNF formulas F with $d(F) \leq f(k)$ are trivially satisfiable, deciding satisfiability of k -CNF formulas F with $d(F) \leq f(k) + 1$ is already NP-complete, for $k \geq 3$. For $k = 3$, this is already observed in [20]. For an upper bound, the complete k -CNF formula witnesses that $f(k) \leq 2^k - 1$. Savický and Sgall [17] showed $f(k) \in O(k^{-0.26}2^k)$. This was improved by Hoory and Szeider [9] to $f(k) \in O\left(\frac{\ln(k)2^k}{k}\right)$, and recently Gebauer [7] proved that $f(k) \leq \frac{2^{k+2}}{k}$. Thus, $f(k)$ is known up to a constant factor. The best upper bounds on $f(k)$ come from MU(1)-formulas. This is true for large values of k , since the formulas constructed in [7] are MU(1), as for small values: Hoory and Szeider [8] show that the function $f(k)$, when restricted to MU(1)-formulas, is computable (in general this is not known), and derive the currently best-known bounds on $f(k)$ for small k ($k \leq 9$). To summarize: When we try to find unsatisfiable k -CNF formulas minimizing a certain parameter, like number of clauses or maximum degree, strictly treelike formulas do an excellent job. However, if we try to construct a small unsatisfiable linear k -CNF formula, they perform horribly. Just compare our upper bound in Theorem 1.2 with the lower bound for strictly treelike formulas in Theorem 1.5

While interest in linear CNF formulas is rather young, *linear hypergraphs* have been studied for quite some time. A hypergraph $H = (V, E)$ is linear if $|e \cap f| \leq 1$ for any two distinct hyperedges $e, f \in E$. A k -uniform hypergraph is a hypergraph where every hyperedge has cardinality k . We ask when a hypergraph is 2-colorable, i.e., admits a 2-coloring of its vertices such that no hyperedge becomes monochromatic. Bounds on the number of edges in such a hypergraph were given by Erdős and Lovász [5] (interestingly, this is the paper where the Local Lemma has been proven). They show that there are non-2-colorable linear k -uniform hypergraphs with ck^44^k hyperedges, but not with less than $\frac{c'4^k}{k^3}$. The proof of the lower bound directly translates into our lower bound for linear k -CNF formulas. For the number of edges in linear k -uniform hypergraphs that are not 2-colorable, the currently best upper bound is ck^24^k by Kostochka and Rödl [11], and the best lower bound is $k^{-\epsilon}4^k$, for any $\epsilon > 0$ and sufficiently large k , due to Kostochka and Kumbhat [10].

2. Existence and Upper and Lower Bounds

Proof of Theorem 1.1. Choose F_0 to be the formula consisting of only the empty clause. Suppose we have constructed F_k , and want to construct F_{k+1} . Let $m = |F_k|$. We create m new variables x_1, \dots, x_m , and let $\mathcal{K}_m = \{D_1, D_2, \dots, D_{2^m}\}$ be the complete m -CNF formula over x_1, \dots, x_m . It is unsatisfiable, but not linear. We take 2^m variable disjoint copies of F_k , denoted by $F_k^{(1)}, F_k^{(2)}, \dots, F_k^{(2^m)}$. For each $1 \leq i \leq 2^m$, we build a linear $(k+1)$ -CNF formula $\tilde{F}_k^{(i)}$ from $F_k^{(i)}$ by adding, for each $1 \leq j \leq m$, the j^{th} literal of D_i to the j^{th} clause of $F_k^{(i)}$. Note that every assignment satisfying $\tilde{F}_k^{(i)}$ also satisfies D_i . Finally, we set $F_{k+1} := \bigcup_{i=1}^{2^m} \tilde{F}_k^{(i)}$. This is an unsatisfiable linear $(k+1)$ -CNF formula with $m2^m$ clauses. ■

Using induction, it is not difficult to see that the formulas F_k are strictly treelike. We will prove the upper bound in Theorem 1.2 by giving a probabilistic construction of

a comparably small unsatisfiable linear k -CNF formula. Our construction consists of two steps. First, we construct a linear k -uniform hypergraph H that is “dense” in the sense that $\frac{m}{n}$ is large, where m and n are the number of hyperedges and vertices, respectively, and then transform it randomly into a linear k -CNF formula F that is unsatisfiable with high probability.

Lemma 2.1. *If there is a linear k -uniform hypergraph H with n vertices and m edges such that $\frac{m}{n} \geq 2^k$, then there is an unsatisfiable linear k -CNF formula with m clauses.*

Proof. Let $H = (V, E)$. By viewing V as a set of variables and E as a set of clauses (each containing only positive literals), this is a (satisfiable) linear k -CNF formula. We replace each literal in each clause by its complement with probability $\frac{1}{2}$, independently in each clause. Let F denote the resulting (random) formula. For any fixed truth assignment α , it holds that $\Pr[\alpha \text{ satisfies } F] = (1 - 2^{-k})^m$. Hence the expected number of satisfying assignments of F is

$$2^n(1 - 2^{-k})^m < 2^n e^{-2^{-k}m} = e^{\ln(2)n - 2^{-k}m} \leq 1,$$

where the last inequality follows from $\frac{m}{n} \geq 2^k$. Hence some formula F has fewer than one satisfying assignment, i.e., none. ■

How can we construct a dense linear hypergraph? We use a construction by Kuzjurin [13]. Our application of this construction is motivated by Kostochka and Rödl [11], who use it to construct linear hypergraphs of large chromatic number.

Lemma 2.2. *For any prime power q and any $k \in \mathbb{N}$, there exists a k -uniform linear hypergraph with kq vertices and q^2 edges.*

With $n = kq$, this hypergraph has n^2/k^2 hyperedges. This is almost optimal, since any linear k -uniform hypergraph on n vertices has at most $\binom{n}{2}/\binom{k}{2}$ hyperedges: The n vertices provide us with $\binom{n}{2}$ vertex pairs. Each hyperedge occupies $\binom{k}{2}$ pairs, and because of linearity, no pair can be occupied by more than one hyperedge.

Proof. Choose the vertex set $V = V_1 \uplus \dots \uplus V_k$, where each V_i is a disjoint copy of the finite field $GF(q)$. The hyperedges consist of all k -tuples (x_1, \dots, x_k) with $x_i \in V_i, 1 \leq i \leq k$, such that

$$\begin{pmatrix} 1 & 1 & & 1 & & 1 \\ 1 & 2 & \dots & i & \dots & k \\ 1 & 4 & & i^2 & & k^2 \\ \vdots & \vdots & & \vdots & & \vdots \\ 1 & 2^{k-3} & \dots & i^{k-3} & \dots & k^{k-3} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_i \\ \vdots \\ x_k \end{pmatrix} = \mathbf{0}. \tag{2.1}$$

Consider two distinct vertices $x \in V_i, y \in V_j$. How many hyperedges contain both of them? If $i = j$, none. If $i \neq j$, we can find out by plugging the fixed values x, y into (2.1). We obtain a (possibly non-uniform) $(k - 2) \times (k - 2)$ linear system with a Vandermonde matrix, which has a unique solution. In other words, x and y are in exactly one hyperedge, and the hypergraph is linear. By the same argument, there are exactly q^2 hyperedges. ■

Proof of the upper bound in Theorem 1.2. Choose a prime power $q \in \{k2^k, \dots, 2k2^k - 1\}$. By Lemma 2.2, there is a linear k -uniform hypergraph H with $n = qk$ vertices and $m = q^2$ hyperedges. Since $\frac{m}{n} = \frac{q}{k} \geq 2^k$, Lemma 2.1 shows that there is an unsatisfiable linear k -CNF formula with $q^2 \leq 4k^24^k$ clauses. ■

Let us prove the lower bound of Theorem 1.2. For a literal u and a CNF formula F , we write $\text{occ}_F(u) := |\{C \in F \mid u \in C\}|$, the *degree* of the literal u . Thus $d_F(x) = \text{occ}_F(x) + \text{occ}_F(\bar{x})$. We write $\text{occ}(F) = \max_u \text{occ}_F(u)$. In analogy to $f(k)$, we define $f_{\text{occ}}(k)$ to be the largest integer d such that any k -CNF formula F with $\text{occ}(F) \leq d$ is satisfiable. Clearly $f_{\text{occ}}(k) \geq \frac{f(k)}{2}$, and thus from [12] it follows that $f_{\text{occ}}(k) \geq \frac{2^k}{2ek}$. Actually, an application of the *Lopsided Lovász Local Lemma* [6, 1, 14] yields $f_{\text{occ}}(k) \geq \frac{2^k}{ek} - 1$.

Lemma 2.3. *Let F be a linear k -CNF formula with at most $1 + f_{\text{occ}}(k - 1)$ variables of degree at least $1 + f_{\text{occ}}(k - 1)$. Then F is satisfiable.*

Proof. Transform F into a $(k - 1)$ -CNF formula F' by removing in every clause in F a literal of maximum degree. We claim that $\deg_{F'}(u) \leq f_{\text{occ}}(k - 1)$ for every literal u . Therefore F' is satisfiable, and F is, as well.

For the sake of contradiction, suppose there is a literal u such that $t := \text{occ}_{F'}(u) \geq 1 + f_{\text{occ}}(k - 1)$. Let C'_i , $i = 1, 2, \dots, t$, be the clauses in F' containing u . C'_i is obtained by removing some literal v_i from some clause $C_i \in F$. By construction of F' , $\text{occ}_F(v_i) \geq \text{occ}_F(u) \geq f_{\text{occ}}(k - 1) + 1$ for all $1 \leq i \leq t$. The v_i are pairwise distinct: If $v_i = v_j$, then $\{u, v_i\} \subseteq C_i \cap C_j$. Since F is weakly linear, this can only mean $i = j$. Now u, v_1, v_2, \dots, v_t are $t + 1 \geq 2 + f_{\text{occ}}(k - 1)$ variables of degree at least $1 + f_{\text{occ}}(k - 1)$ in F , a contradiction. ■

We see that an unsatisfiable weakly linear k -CNF formula has at least $f_{\text{occ}}(k - 1) + 2 \geq \frac{2^k}{2e(k-1)} + 1$ literals of degree at least $f_{\text{occ}}(k - 1) + 1 \geq \frac{2^k}{2e(k-1)}$. Double counting yields $k|F| = \sum_u \text{occ}_F(u) > \frac{4^k}{4e^2(k-1)^2}$, thus $|F| > \frac{4^k}{16e^2k^3}$. By a more careful argument, we can improve this by a factor of k . We call a hypergraph (j, d) -rich if at least j vertices have degree at least d . The following lemma is due to Welzl [22].

Lemma 2.4. *For $d \in \mathbb{N}_0$, every linear (d, d) -rich hypergraph has at least $\binom{d+1}{2}$ edges. This bound is tight for all $d \in \mathbb{N}_0$.*

Proof. We proceed by induction over d . Clearly, the assertion of the lemma is true for $d = 0$. Now let $H = (V, E)$ be a linear (d, d) -rich hypergraph for $d \geq 1$. Choose some vertex v of degree at least d in H and let $H' = (V, E')$ be the hypergraph with $E' := E \setminus \{e \in E \mid e \ni v\}$. We have (i) $|E| \geq |E'| + d$, (ii) H' is linear, since this property is inherited when edges are removed, and (iii) H' is $(d - 1, d - 1)$ -rich, since for no vertex other than v the degree decreases by more than 1 due to the linearity of H . It follows that $|E| \geq \binom{d}{2} + d = \binom{d+1}{2}$. The complete 2-uniform hypergraph (graph, so to say) on $d + 1$ vertices shows that the bound given is tight for all $d \in \mathbb{N}_0$. ■

Proof of the lower bound in Theorem 1.2. A weakly linear k -CNF formula F is a linear k -uniform hypergraph, with literals as vertices. If F is unsatisfiable, then by Lemma 2.3, it is $(f_{\text{occ}}(k - 1) + 1, f_{\text{occ}}(k - 1) + 1)$ -rich. By Lemma 2.4, F has at least $\binom{f_{\text{occ}}(k-1)+2}{2} > \frac{4^k}{8e^2(k-1)^2}$ clauses. ■




Figure 1: A resolution tree, with its edges labeled in the obvious way. Every clause is unsatisfied when applying the assignments on the path to the root.

There is an obvious generalization of the notion of being linear. We say a CNF formula is *b-linear*, if any two distinct clauses $C, D \in F$ fulfill $|\text{vbl}(C) \cap \text{vbl}(D)| \leq b$, and *weakly b-linear* if $|C \cap D| \leq b$ holds for all distinct $C, D \in F$. Thus, a (weakly) 1-linear formula is (weakly) linear. We can generalize Theorem 1.2 for $b \geq 2$. However, the proofs do not introduce new ideas and goes along the lines of the proofs presented above.

Theorem 2.5. *Let $b \geq 2$. Every weakly b -linear k -CNF formula with at most $\frac{2^{k(1+\frac{1}{b})}}{2^{b+2}e^2k^{2+\frac{1}{b}}}$ clauses is satisfiable. There exists an unsatisfiable b -linear k -CNF formula with at most $2^{b+1}(k2^k)^{1+\frac{1}{b}}$ clauses.*

3. Proof of Theorem 1.4

Let F be an unsatisfiable weakly linear k -CNF formula, and let T be a resolution tree of minimal size of F . We want to show that T has a large number of nodes. It is not difficult to see that a resolution tree of minimal size is *regular*, meaning that no variable is resolved more than once on a path from a leaf to the root. See Urquhart [21], Lemma 5.1, for a proof of this fact. We take a random walk of length ℓ in T starting at the root, in every step choosing randomly to go to one of the two children of the current node. If we arrive at a leaf, we stay there. We claim that if $\ell \leq \sqrt{2^{k-2}}$, then with probability at least $\frac{1}{2}$, our walk does not end at a leaf. Thus, T has at least $2^{\ell-1}$ inner vertices at distance ℓ from the root, thus at least $2^{2^{\frac{k}{2}-1}}$ leaves.

As illustrated in Figure 1, we label each edge in T with an assignment. If C is the resolvent of D_1 and D_2 , $x \in D_1$ and $\bar{x} \in D_2$, we label the edge from C to D_1 by $x \mapsto 0$ and from C to D_2 by $x \mapsto 1$. Each path from the root to a node gives a partial assignment α . If that node is labeled with clause C , then C evaluates to **false** under α . In our random walk, let α_i denote the partial assignment associated with the first i steps. α_0 is the empty assignment, and α_i assigns exactly i variables (if we are not yet at a leaf). We set $F_i := F^{[\alpha_i]}$, i.e., the formula obtained from F by fixing the variables according to the partial assignment α_i . For a formula G , we define the *weight* $w(G)$ to be

$$w(G) := \sum_{C \in G, |C| \leq k-2} 2^{k-|C|}. \tag{3.1}$$

Since F is a k -CNF formula, $w(F) = 0$. If some formula G contains the empty clause, then $w(G) \geq 2^k$. In our random walk, $w(F_i)$ is a random variable.

Lemma 3.1. $\mathbb{E}[w(F_{i+1})] \leq \mathbb{E}[w(F_i)] + 4i$.

Since $w(F_0) = 0$, this implies $\mathbb{E}[w(F_\ell)] \leq 4\binom{\ell}{2} \leq 2\ell^2$. If our random walk ends at a leaf, then F_ℓ contains the empty clause, thus $w(F_\ell) \geq 2^k$. Therefore $2\ell^2 \geq \mathbb{E}[w(F_\ell)] \geq 2^k \Pr[\text{the random walk ends at a leaf}]$. We conclude that at least half of all paths of length $\ell^* = \sqrt{2^{k-2}}$ starting at the root do not end at a leaf. Thus T has at least 2^{ℓ^*-1} internal nodes at distance ℓ^* from the root, and thus at least 2^{ℓ^*} leaves, which proves the theorem. It remains to prove the lemma.

Proof of the lemma. For a formula G and a variable x , let $d_{k-1}(x, G)$ denote the number of $(k-1)$ -clauses containing x or \bar{x} . Since F_0 is a k -CNF formula, $d_{k-1}(x, F_0) = 0$, for all variables x . We claim that $d_{k-1}(x, F_{i+1}) \leq d_{k-1}(x, F_i) + 2$ for every variable x . To see this, note that in step i , some variable y is set to $b \in \{0, 1\}$, say to 0. At most one k -clause of F_i contains y and x , and at most one contains y and \bar{x} , since F_i is weakly linear, thus $d_{k-1}(x, F_{i+1}) \leq d_{k-1}(x, F_i) + 2$. It follows immediately that $d_{k-1}(x, F_i) \leq 2i$.

Consider $w(F_i)$, which was in (3.1). F_{i+1} is obtained from F_i by setting some variable y randomly to 0 or 1. Consider a clause C . How does its contribution to (3.1) change when setting y ? If (i) $y \notin \text{vbl}(C)$ or $|C| = k$, it does not change. If (ii) $y \in \text{vbl}(C)$ and $|C| \leq k-2$, then with probability $\frac{1}{2}$ each, its contribution to (3.1) doubles or vanishes. Hence on expectation, it does not change. If (iii) $y \in \text{vbl}(C)$ and $|C| = k-1$, then C contributes nothing to $w(F_i)$, and with probability $\frac{1}{2}$, it contributes 4 to $w(F_{i+1})$. On expectation, its contribution to (3.1) increases by 2. Case (iii) applies to at most $d_{k-1}(y, F_i) \leq 2i$ clauses. Hence $\mathbb{E}[w(F_{i+1})] \leq \mathbb{E}[w(F_i)] + 4i$. ■

4. Proof of Theorem 1.5

Let F be a strictly treelike weakly linear k -CNF formula F , and let T be a strict resolution tree of F . Letters a, b, c denote nodes of T , and u, v, w denote literals. Every node a of T is labeled with a clause C_a . We define a graph G_a with vertex set C_a , connecting $u, v \in C_a$ if $u, v \in D$ for some clause $D \in F$ that occurs as a label of a leaf in the subtree of a . Since T is a strict resolution tree and F is weakly linear, every edge in G_a comes from a unique leaf of T . Resolution now has a simple interpretation as a "calculus on graphs", see Figure 2. If a is a leaf, then $G_a = K_k$. Since the root of a resolution tree is labeled with the empty clause, we have $G_{\text{root}} = (\emptyset, \emptyset)$. For a graph G , let $\kappa_i(G)$ denote the minimum size of a set $U \subseteq V(G)$ such that $G - U$ contains no i -clique. Here, $G - U$ is the subgraph of G induced by $V(G) \setminus U$. Thus, $\kappa_1(G) = |V(G)|$, and $\kappa_2(G)$ is the size of a minimum vertex cover of G . For the complete graph K_k , $\kappa_i(K_k) = k - i + 1$. We write $\kappa_i(a) := \kappa_i(G_a)$. The tuple $(\kappa_1(a), \dots, \kappa_k(a))$ can be viewed as the complexity measure for a . We observe that if a is a leaf, then $\kappa_i(a) = k - i + 1$, and $\kappa_i(\text{root}) = 0$, for all $1 \leq i \leq k$. If a is an ancestor of b in T , let $\text{dist}(a, b)$ denote the number of edges in the T -path from a to b . Since one resolution step deletes one literal (and may add several), the next proposition is immediate:

Proposition 4.1. *If b is a descendant of a in T , then $\kappa_i(b) \leq \kappa_i(a) + \text{dist}(a, b)$.*




Figure 2: Resolution as a calculus on graphs. A resolution step amounts to deleting the resolved vertex and taking the union of the two graphs.

At this point we want to give an intuition of the proofs that follow. Our goal is to show that if the values $\kappa_i(a)$ are small for some node a in the tree, then the subtree of a is big. The proof goes roughly as follows: If the subtree of a is small, then there are many descendants b of a that are not too far from a and have even smaller subtrees. By induction, we will be able to show that $\kappa_{i+1}(b)$ is fairly large. Thus, on the path from b to a , not all $(i + 1)$ -cliques are destroyed, and every such descendant b of a provides G_a with an $(i + 1)$ -clique. These cliques need not be vertex-disjoint, but they are edge-disjoint. This implies that G_a has many vertex-disjoint i -cliques, a contradiction to $\kappa_i(a)$ being small. To make this intuition precise, we have to define what small and big actually means in this context: We fix a value $1 \leq \ell \leq k$ and define ν_i and θ_i for $1 \leq i \leq \ell$ as follows: $\theta_\ell := \lfloor \frac{k-\ell+1}{2} \rfloor - 1$ and $\nu_\ell := 1$, and for $1 \leq i < \ell$, we inductively define $\theta_i := \lfloor \frac{2^{\nu_{i+1}\theta_{i+1}-2}}{\theta_{i+1}} \rfloor - 1$ and $\nu_i := \frac{\nu_{i+1}\theta_{i+1}-1}{\theta_i} \lfloor \frac{\theta_i}{\theta_{i+1}} \rfloor$. One should not worry about these ugly expressions too much, they are only chosen that way to make the induction go through. For the right value of ℓ , one checks that θ_1 is a tower function in k . More precisely, for any $\epsilon > 0$, there exists a $c \in \mathbb{N}$ such that when choosing $\ell = k - c$, then $\theta_1 \geq \text{tower}_{2-\epsilon}(k - c)$. The following theorem is a more precise version of Theorem 1.5.

Theorem 4.2. *Let F be a strictly treelike linear k -CNF formula. Then F has at least $2^{\nu_1\theta_1}$ clauses.*

Proof. A node a in T is i -extendable if $\kappa_j(a) \leq \theta_j$ for each $i \leq j \leq \ell$. We observe that if a is i -extendable, it is also $(i + 1)$ -extendable. For $i = \ell + 1$, the condition is void, so every node is $(\ell + 1)$ -extendable. Also, the root is 1-extendable, since $\kappa_1(\text{root}) = 0$.

Definition 4.3. A set A of descendants of a in T such that (i) no vertex in A is an ancestor of any other vertex in A and (ii) $\text{dist}(a, b) \leq d$ for all $b \in A$ is called an *antichain of a at distance at most d* . If furthermore every $b \in A$ is i -extendable, we call A an *i -extendable antichain*.

Lemma 4.4. *Let $1 \leq i \leq \ell$, and let a be a node in T . If a is i -extendable, then there is an $(i + 1)$ -extendable antichain A of a at distance at most θ_i such that $|A| = 2^{\nu_i\theta_i}$.*

Proof. We use induction on $\ell - i$. For the base case $i = \ell$, we have $\kappa_\ell(a) \leq \theta_\ell$, as a is ℓ -extendable. Since each leaf b of T has $\kappa_\ell(b) = k - \ell + 1 \geq 2\theta_\ell + 2$, Proposition 4.1 tells us that every leaf in the subtree of a has distance at least $\theta_\ell + 2$ from a . Since T is a complete binary tree, there are 2^{θ_ℓ} descendants of a at distance exactly θ_ℓ from a . This is the desired antichain A of a . Since every node is $(\ell + 1)$ -extendable, the base case holds. For the step,




Figure 3: Illustration of the claim in the proof of Lemma 4.4. If node a is i -extendable, and b is a close $(i + 1)$ -extendable descendant of a , then b itself has many close descendants A , at least half of which are $(i + 1)$ -extendable themselves.

let a be i -extendable, for $1 \leq i < \ell$.

Claim: Let b be a descendant of a with $\text{dist}(a, b) \leq \theta_i$. If b is $(i + 1)$ -extendable, then there is an $(i + 1)$ -extendable antichain A of b at distance at most θ_{i+1} of size $2^{\nu_{i+1}\theta_{i+1}-1}$.

Proof of the claim. By applying the induction hypothesis of the lemma to b , there is an $(i + 2)$ -extendable antichain A of b at distance at most θ_{i+1} of size $2^{\nu_{i+1}\theta_{i+1}}$. Let $A_{\text{good}} := \{c \in A \mid \kappa_{i+1}(c) \leq \theta_{i+1}\}$. This is an $(i + 1)$ -extendable antichain. If A_{good} contains at least half of A , we are done. See Figure 3 for an illustration. Write $A_{\text{bad}} := A \setminus A_{\text{good}}$ and suppose for the sake of contradiction that $|A_{\text{bad}}| > 2^{\nu_{i+1}\theta_{i+1}-1}$. Consider any $c \in A_{\text{bad}}$. On the path from c to b , in each step some literal gets removed (and others may be added). Let P denote the set of the removed literals. Then $C_c \setminus \{P\} \subseteq C_b$, and $G_c - P$ is a subgraph of G_b . Node c is not $(i + 1)$ -extendable, thus $\kappa_{i+1}(c) \geq \theta_{i+1} + 1$. Since $|P| = \text{dist}(b, c) \leq \theta_{i+1}$, the graph $G_c - P$ contains at least one $(i + 1)$ -clique, which is also contained in G_b . This holds for every $c \in A_{\text{bad}}$, and by weak linearity, G_b contains at least $|A_{\text{bad}}|$ edge disjoint $(i + 1)$ -cliques. Since b is $(i + 1)$ -extendable, there exists a set $U \subseteq V(G_b)$, $|U| = \kappa_{i+1}(b)$ such that $G_b - U$ contains no $(i + 1)$ -clique. Each of the $|A_{\text{bad}}|$ edge-disjoint $(i + 1)$ -cliques G_b contains some vertex of U , thus some vertex $v \in U$ is contained in at least $\frac{|A_{\text{bad}}|}{|U|} \geq \frac{2^{\nu_{i+1}\theta_{i+1}-1}}{\theta_{i+1}} \geq 2\theta_i + 1$ edge-disjoint $(i + 1)$ -cliques. Two such cliques overlap in no vertex besides v , hence G_b contains at least $2\theta_i + 1$ vertex-disjoint i -cliques, thus $\kappa_i(b) \geq 2\theta_i + 1$. By Proposition 4.1, $\kappa_i(a) \geq \kappa_i(b) - \text{dist}(a, b) \geq \theta_i + 1$. This contradicts the assumption of Lemma 4.4 that a is i -extendable. We conclude that $|A_{\text{bad}}| \leq \frac{1}{2}|A|$, which proves the claim. ■

Let us continue with the proof of the lemma. If A is an $(i + 1)$ -extendable antichain of a at distance $d \leq \theta_i$, then by the claim for each vertex $b \in A$ there exists an $(i + 1)$ -extendable antichain of b at distance at most θ_{i+1} , of size $2^{\nu_{i+1}\theta_{i+1}-1}$. Their union is an $(i + 1)$ -extendable antichain A' of a at distance at most $d + \theta_{i+1}$, of size $|A|2^{\nu_{i+1}\theta_{i+1}-1}$. Hence we can “inflate” A to A' , as long as $d \leq \theta_i$. Starting with the $(i + 1)$ -extendable antichain $\{a\}$ and inflate it $\lfloor \frac{\theta_i}{\theta_{i+1}} \rfloor$ times, and obtain a final $(i + 1)$ -extendable antichain of a at distance at most θ_i of size at least $(2^{\nu_{i+1}\theta_{i+1}-1})^{\lfloor \frac{\theta_i}{\theta_{i+1}} \rfloor} = 2^{\nu_i\theta_i}$. ■

Applying Lemma 4.4 to the root of T , which is 1-extendable, we obtain an antichain A of size $2^{\nu_1 \theta_1}$ nodes. Since T has at least $|A|$ leaves, this proves the theorem. ■

5. Open Problems

Let $f_{\text{LIN}}(k)$ be the largest integer d such that any linear k -CNF formula F with $d(F) \leq d$ is satisfiable. Clearly $f_{\text{LIN}}(k) \geq f(k)$, and from the proof of the upper bound in Theorem 1.2 it follows that $f_{\text{LIN}}(k) \leq 2k2^k$. Is there a significant gap between $f(k)$ and $f_{\text{LIN}}(k)$? It is not difficult to show that $f(2) = f_{\text{LIN}}(2) = 2$, but we do not know the value of $f_{\text{LIN}}(k)$ for any $k \geq 3$. How do unsatisfiable linear k -CNF formulas look like? Can one find an explicit construction of an unsatisfiable linear k -CNF formula whose size is singly exponential in k ? We suspect one has to come up with some algebraic construction. What is the resolution complexity of linear k -CNF formulas? Tree resolution complexity is doubly exponential in k . We suspect the same to be true for general resolution.

Acknowledgments

My thanks go to Emo Welzl, Robin Moser, Heidi Gebauer, Andreas Razen and Philipp Zumstein for very helpful and fruitful discussions.

References

- [1] N. Alon and J. H. Spencer. *The probabilistic method*. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience [John Wiley & Sons], New York, second edition, 2000. With an appendix on the life and work of Paul Erdős.
- [2] E. Ben-Sasson and A. Wigderson. Short proofs are narrow—resolution made simple. *J. ACM*, 48(2):149–169, 2001.
- [3] M. Davis, G. Logemann, and D. Loveland. A machine program for theorem-proving. *Comm. ACM*, 5:394–397, 1962.
- [4] G. Davydov, I. Davydova, and K. Büning. An efficient algorithm for the minimal unsatisfiability problem for a subclass of CNF. *Ann. Math. Artificial Intelligence*, 23(3-4):229–245, 1998.
- [5] P. Erdős and L. Lovász. Problems and results on 3-chromatic hypergraphs and some related questions. In A. Hajnal, R. Rado, and V. T. Sós, editors, *Infinite and Finite Sets (to Paul Erdős on his 60th birthday)*, Vol. II, pages 609–627. North-Holland, 1975.
- [6] P. Erdős and J. Spencer. Lopsided Lovász Local Lemma and Latin transversals. *Discrete Appl. Math.*, 30(2-3):151–154, 1991. ARIDAM III (New Brunswick, NJ, 1988).
- [7] H. Gebauer. Disproof of the neighborhood conjecture with implications to SAT. In A. Fiat and P. Sanders, editors, *17th Annual European Symposium on Algorithms (ESA 2009)*, volume 5757 of *Lecture Notes in Computer Science*, pages 764–775. Springer, 2009.
- [8] S. Hoory and S. Szeider. Computing unsatisfiable k -SAT instances with few occurrences per variable. *Theoretical Computer Science*, 337(1-3):347–359, 2005.
- [9] S. Hoory and S. Szeider. A note on unsatisfiable k -CNF formulas with few occurrences per variable. *SIAM Journal on Discrete Mathematics*, 20(2):523–528, 2006.
- [10] A. V. Kostochka and M. Kumbhat. Coloring uniform hypergraphs with few edges, *manuscript*.
- [11] A. V. Kostochka and V. Rödl. Constructions of sparse uniform hypergraphs with high chromatic number, *manuscript*.
- [12] J. Kratochvíl, P. Savický, and Z. Tuza. One more occurrence of variables makes satisfiability jump from trivial to NP-complete. *SIAM Journal of Computing*, 22(1):203–210, 1993.
- [13] N. N. Kuzjurin. On the difference between asymptotically good packings and coverings. *European J. Combin.*, 16(1):35–40, 1995.

- [14] L. Lu and L. Székely. Using Lovász Local Lemma in the space of random injections. *Electron. J. Combin.*, 14(1):Research Paper 63, 13 pp. (electronic), 2007.
- [15] S. Porschen, E. Speckenmeyer, and B. Randerath. On linear CNF formulas. In *SAT*, pages 212–225, 2006.
- [16] S. Porschen, E. Speckenmeyer, and X. Zhao. Linear CNF formulas and satisfiability. *Discrete Appl. Math.*, 157(5):1046–1068, 2009.
- [17] P. Savický and J. Sgall. DNF tautologies with a limited number of occurrences of every variable. *Theoret. Comput. Sci.*, 238(1–2):495–498, 2000.
- [18] D. Scheder. Unsatisfiable linear k -CNFs exist, for every k . *CoRR*, abs/0708.2336, 2007.
- [19] S. Szeider. Homomorphisms of conjunctive normal forms. *Discrete Appl. Math.*, 130(2):351–365, 2003.
- [20] C. A. Tovey. A simplified NP-complete satisfiability problem. *Discrete Appl. Math.*, 8(1):85–89, 1984.
- [21] A. Urquhart. The complexity of propositional proofs. *Bulletin of Symbolic Logic*, 1:425–467, 1995.
- [22] E. Welzl. personal communication.