

An optimal error estimate in stochastic homogenization of discrete elliptic equations

Antoine Gloria, Felix Otto

► **To cite this version:**

Antoine Gloria, Felix Otto. An optimal error estimate in stochastic homogenization of discrete elliptic equations. *Annals of Applied Probability*, Institute of Mathematical Statistics (IMS), 2012, 22 (1), pp.1-28. 10.1214/10-AAP745 . inria-00457020v3

HAL Id: inria-00457020

<https://hal.inria.fr/inria-00457020v3>

Submitted on 18 Dec 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AN OPTIMAL ERROR ESTIMATE IN STOCHASTIC HOMOGENIZATION OF DISCRETE ELLIPTIC EQUATIONS¹

BY ANTOINE GLORIA AND FELIX OTTO

*INRIA Lille-Nord Europe and Max Planck Institute for
 Mathematics in the Sciences*

This paper is the companion article to [Ann. Probab. **39** (2011) 779–856]. We consider a discrete elliptic equation on the d -dimensional lattice \mathbb{Z}^d with random coefficients A of the simplest type: They are identically distributed and independent from edge to edge. On scales large w.r.t. the lattice spacing (i.e., unity), the solution operator is known to behave like the solution operator of a (continuous) elliptic equation with constant deterministic coefficients. This symmetric “homogenized” matrix $A_{\text{hom}} = a_{\text{hom}}\text{Id}$ is characterized by $\xi \cdot A_{\text{hom}}\xi = \langle (\xi + \nabla\phi) \cdot A(\xi + \nabla\phi) \rangle$ for any direction $\xi \in \mathbb{R}^d$, where the random field ϕ (the “corrector”) is the unique solution of $-\nabla^* \cdot A(\xi + \nabla\phi) = 0$ in \mathbb{Z}^d such that $\phi(0) = 0$, $\nabla\phi$ is stationary and $\langle \nabla\phi \rangle = 0$, $\langle \cdot \rangle$ denoting the ensemble average (or expectation).

In order to approximate the homogenized coefficients A_{hom} , the corrector problem is usually solved in a box $Q_L = [-L, L]^d$ of size $2L$ with periodic boundary conditions, and the space averaged energy on Q_L defines an approximation A_L of A_{hom} . Although the statistics is modified (independence is replaced by periodic correlations) and the ensemble average is replaced by a space average, the approximation A_L converges almost surely to A_{hom} as $L \uparrow \infty$. In this paper, we give estimates on both errors. To be more precise, we do not consider periodic boundary conditions on a box of size $2L$, but replace the elliptic operator by $T^{-1} - \nabla^* \cdot A\nabla$ with (typically) $T \sim L^2$, as standard in the homogenization literature. We then replace the ensemble average by a space average on Q_L , and estimate the overall error on the homogenized coefficients in terms of L and T .

1. Introduction.

1.1. *Motivation.* In this article, we continue the analysis we began in [6] on stochastic homogenization of discrete elliptic equations. More precisely, we consider real functions u of the sites x in a d -dimensional Cartesian lattice \mathbb{Z}^d . Every edge e of the lattice is endowed with a “conductivity” $a(e) > 0$. This defines a discrete elliptic differential operator $-\nabla^* \cdot A\nabla$ via

$$-\nabla^* \cdot (A\nabla u)(x) := \sum_{y \in \mathbb{Z}^d, |x-y|=1} a(e)(u(x) - u(y)),$$

Received February 2010; revised August 2010.

¹Supported in part of the Hausdorff Center for Mathematics, Bonn, Germany.
MSC2010 subject classifications. 35B27, 39A70, 60H25, 60F99.

Key words and phrases. Stochastic homogenization, effective coefficients, difference operator.

where the sum is over the $2d$ sites y which are connected by an edge $e = [x, y]$ to the site x . It is sometimes more convenient to think in terms of the associated Dirichlet form, that is,

$$\begin{aligned} \sum \nabla v \cdot A \nabla u &:= \sum_{x \in \mathbb{Z}^d} v(x) (-\nabla^* \cdot (A \nabla u))(x) \\ &= \sum_e (v(x) - v(y)) a(e) (u(x) - u(y)), \end{aligned}$$

where the last sum is over all edges e and (x, y) denotes the two sites connected by e , that is, $e = [x, y] = [y, x]$ (with the convention that an edge is not oriented). We assume the conductivities a to be uniformly elliptic in the sense of

$$\alpha \leq a(e) \leq \beta \quad \text{for all edges } e$$

for some fixed constants $0 < \alpha \leq \beta < \infty$.

We are interested in random coefficients. To fix ideas, we consider the simplest situation possible:

$$\{a(e)\}_e \text{ are independently and identically distributed (i.i.d.).}$$

Hence, the statistics are described by a distribution on the finite interval $[\alpha, \beta]$. We'd like to see this discrete elliptic operator with random coefficients as a good model problem for continuum elliptic operators with random coefficients of correlation length unity.

Classical results in stochastic homogenization of linear elliptic equations (see [8] and [13] for the continuous case, and [10] and [9] for the discrete case) state that there exist *homogeneous and deterministic* coefficients A_{hom} such that the solution operator of the continuum differential operator $-\nabla \cdot A_{\text{hom}} \nabla$ describes the large scale behavior of the solution operator of the discrete differential operator $-\nabla^* \cdot A \nabla$. As a by product of this homogenization result, one obtains a characterization of the homogenized coefficients A_{hom} : It is shown that for every direction $\xi \in \mathbb{R}^d$, there exists a unique scalar field ϕ such that $\nabla \phi$ is stationary [stationarity means that the fields $\nabla \phi(\cdot)$ and $\nabla \phi(\cdot + z)$ have the same statistics for all shifts $z \in \mathbb{Z}^d$] and $\langle \nabla \phi \rangle = 0$, solving the equation

$$(1.1) \quad -\nabla^* \cdot (A(\xi + \nabla \phi)) = 0 \quad \text{in } \mathbb{Z}^d$$

and normalized by $\phi(0) = 0$. As in periodic homogenization, the function $\mathbb{Z}^d \ni x \mapsto \xi \cdot x + \phi(x)$ can be seen as the A -harmonic function which macroscopically behaves as the affine function $\mathbb{Z}^d \ni x \mapsto \xi \cdot x$. With this ‘‘corrector’’ ϕ , the homogenized coefficients A_{hom} (which in general form a symmetric matrix and for our simple statistics in fact a multiple of the identity: $A_{\text{hom}} = a_{\text{hom}} \text{Id}$) can be characterized as follows:

$$(1.2) \quad \xi \cdot A_{\text{hom}} \xi = \langle (\xi + \nabla \phi) \cdot A(\xi + \nabla \phi) \rangle.$$

Since the scalar field $(\xi + \nabla\phi) \cdot A(\xi + \nabla\phi)$ is stationary, it does not matter (in terms of the distribution) at which site x it is evaluated in the formula (1.2), so that we suppress the argument x in our notation.

When one is interested in explicit values for A_{hom} , one has to solve (1.1). Since this is not possible in practice, one has to make approximations. For a discussion of the literature on error estimates, in particular the pertinent work by Yurinskii [15] and Naddaf and Spencer [12], we refer to [6], Section 1.2. A standard approach used in practice consists in solving (1.1) in a box $Q_L = [-L, L]^d \cap \mathbb{Z}^d$ with periodic boundary conditions

$$(1.3) \quad -\nabla^* \cdot (A(\xi + \nabla\phi_{L,\#})) = 0 \quad \text{in } Q_L,$$

and replacing (1.2) by a space average

$$(1.4) \quad \xi \cdot A_{L,\#}\xi = \int_{Q_L} (\xi + \nabla\phi_{L,\#}) \cdot A(\xi + \nabla\phi_{L,\#}) dx.$$

Such an approach is consistent in the sense that

$$\lim_{L \rightarrow \infty} A_{L,\#} = A_{\text{hom}}$$

almost surely, as proved, for instance, in [1] for the continuous case, and in [2] for the discrete case. Numerical experiments tend to show that the use of periodic boundary conditions gives better results than other choices such as homogeneous Dirichlet boundary conditions, see [14].

An important question for practical purposes is to quantify the dependence of the error $\langle |A_{\text{hom}} - A_{L,\#}|^2 \rangle^{1/2}$ in terms of L . Let us give another interpretation of (1.3): This equation on Q_L is equivalent to (1.1) on \mathbb{Z}^d with a modified conductivity matrix \tilde{A}_L , that is the periodization of $A|_{Q_L}$ on \mathbb{Z}^d . Doing this, we have replaced independent coefficients A by Q_L -periodically correlated coefficients \tilde{A} . Since A and \tilde{A} are not jointly stationary (see Definition 4), it may be difficult to compare $\nabla\phi$ to $\nabla\phi_{L,\#}$. To circumvent this difficulty, and following the route of [10, 13, 15] and [12], and as in [6], we slightly depart from (1.3) by introducing a zero-order term in (1.1):

$$(1.5) \quad T^{-1}\phi_T - \nabla^* \cdot (A(\xi + \nabla\phi_T)) = 0 \quad \text{in } \mathbb{Z}^d.$$

As for the periodization, this localizes the dependence of $\phi_T(z)$ upon $A(z')$ to those points $z' \in \mathbb{Z}^d$ such that $|z - z'| \lesssim \sqrt{T}$ (at first order). Yet, unlike the periodization, $\nabla\phi_T$ and $\nabla\phi$ are jointly stationary. In terms of random walk interpretation, the lifetime of the random walker is of order T , and the distance to the origin of order \sqrt{T} . Hence, up to taking $T \sim L^2$, in first approximation, the function $\phi_T|_{Q_L}$ only depends on the coefficients $A(z)$ for $z \in Q_L$, as it is the case for $\phi_{L,\#}$.

We'd like to view $\phi_T|_{Q_L}$ as a variant of $\phi_{L,\#}$ which is convenient for our analysis. We then define

$$(1.6) \quad \xi \cdot A_{T,L}\xi = \int_{\mathbb{Z}^d} (\xi + \nabla\phi_T) \cdot A(\xi + \nabla\phi_T)\eta_L dx,$$

where η_L is a smooth mask with unit mass and support Q_L . The aim of this paper is to determine the scaling of the error $\langle |A_{\text{hom}} - A_{T,L}|^2 \rangle^{1/2}$ in terms of L and T . Eventually this will allow us to make a reasonable choice for T and L at fixed computational complexity.

1.2. *Informal statement of the results.* When approximating A_{hom} by $A_{T,L}$, we make two types of errors: A “systematic error” and a “random error.” In particular, as shown in [6],

$$\langle (\xi \cdot A_{\text{hom}} \xi - \xi \cdot A_{T,L} \xi)^2 \rangle = \langle \xi \cdot (A_{\text{hom}} - A_T) \xi \rangle^2 + \text{var}[A_{T,L}].$$

The first term is the square of the systematic error (see [6], (1.10))

$$(1.7) \quad \begin{aligned} \text{Error}_{\text{sys}}(T) &:= |\langle (\xi + \nabla \phi_T) \cdot A(\xi + \nabla \phi_T) \rangle - \langle (\xi + \nabla \phi) \cdot A(\xi + \nabla \phi) \rangle| \\ &= \langle (\nabla \phi_T - \nabla \phi) \cdot A(\nabla \phi_T - \nabla \phi) \rangle. \end{aligned}$$

It measures the fact that the coefficient $a(e)$ at bond e does (up to exponentially small terms) not influence $\phi_T(x)$ if $|x - e| \gg \sqrt{T}$. This error vanishes for $T = L^2 \uparrow \infty$. The second term is the square of the random error,

$$(1.8) \quad \text{Error}_{\text{rand}}(T, L) = \text{var} \left[\int_{\mathbb{Z}^d} (\xi + \nabla \phi_T) \cdot A(\xi + \nabla \phi_T) \eta_L dx \right]^{1/2}.$$

It measures the fluctuations of the energy density. This error vanishes as $L \uparrow \infty$.

In [6], Theorem 1, we have proved that

$$(1.9) \quad \text{var} \left[\int_{\mathbb{Z}^d} (\xi + \nabla \phi_T) \cdot A(\xi + \nabla \phi_T) \eta_L dx \right]^{1/2} \lesssim \begin{cases} d = 2, & L^{-1} \ln^q T, \\ d > 2, & L^{-d/2}, \end{cases}$$

for some q depending only on α, β , where “ \lesssim ” stands for “ \leq ” up to a multiplicative constant depending only on α, β and d . We have also identified the systematic error in the limit of vanishing conductivity contrast, that is, $1 - \beta/\alpha \ll 1$, and found

$$\text{Error}_{\text{sys}}(T) \sim \begin{cases} d = 2, & T^{-1}, \\ d = 3, & T^{-3/2}, \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}, \end{cases}$$

where “ \sim ” means that both terms have the same scaling (in T). In this paper, we shall actually prove that for general α and β (see Theorem 1)

$$(1.10) \quad \text{Error}_{\text{sys}}(T) \lesssim \begin{cases} d = 2, & T^{-1} \ln^q T, \\ d = 3, & T^{-3/2}, \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}, \end{cases}$$

where there is a logarithmic correction for $d = 2$ when compared to the vanishing conductivity asymptotics.

Assuming that ϕ_T can be well approximated on domains of size L if we choose $T \sim L^2$, the combination of (1.10) and (1.9) yields

$$\langle |A_{\text{hom}} - A_{T,L}|^2 \rangle^{1/2} \lesssim \begin{cases} d = 2, & L^{-1} \ln^q L, \\ 2 < d \leq 7, & L^{-d/2}, \\ d = 8, & L^{-4} \ln L, \\ d > 8, & L^{-4}. \end{cases}$$

Hence, the numerical strategy converges at the rate of the central limit theorem for $2 \leq d \leq 8$ (up to logarithmic corrections for $d = 2$ and $d = 8$).

Up to dimension 4, the systematic error for $T \sim L^2$ scales as the square of the random error. In particular, this leaves room for the choice T . If we take $T \sim L$, then the systematic error is of the same order as the random error. What we have gained is that ϕ_T can now be well-approximated on domains of size $R \sim \sqrt{T} \sim \sqrt{L}$, and not only L . Note also that the random error is unchanged if instead of taking the average of one realization of ϕ_T on Q_L (with the mask μ_L) we take the empirical average of the averages of N independent realizations of ϕ_T on a domain $Q_{L/N^{1/d}}$ (with the according mask $\mu_{L/N^{1/d}}$). Hence, since ϕ_T can be well-approximated on domains of size $R \gtrsim \sqrt{L}$, considering $N = 1$ realization of ϕ_T approximated on Q_L or $N = \sqrt{L^d}$ independent realizations of ϕ_T approximated on $Q_{\sqrt{L}}$ yields the same scaling for the error between the homogenized coefficients and their approximations. Since the computational cost of solving a linear problem is superlinear in the number of unknowns, it seems best to choose N as large as possible, and therefore taking $N = \sqrt{L^d}$ seems a reasonable strategy at first order. Yet, we do not make precise in this paper the relation between R and \sqrt{L} in terms of absolute values (we only consider the scaling), which may make the optimal choice for N more subtle in practice than this general principle. A complete numerical analysis of the numerical method (including the influence of R and the optimization of N) will be presented in [4].

We conclude this introduction by mentioning the very recent contribution [11] by Mourrat. The equation under investigation is the same as above, namely a discrete elliptic equation on \mathbb{Z}^d with i.i.d. coefficients. The object under study is the spectral measure associated with the generator of the environment viewed by the particle. Without entering into details, there exists some nonnegative measure e_∂ associated with the elliptic operator and direction $\xi \in \mathbb{R}^d$, such that the homogenized coefficient is given by

$$\xi \cdot A_{\text{hom}} \xi = \langle \xi \cdot A \xi \rangle - \int_{\mathbb{R}^+} \frac{1}{\lambda} d e_\partial(\lambda).$$

As recalled in [11], we also have

$$(1.11) \quad \xi \cdot A_T \xi = \langle \xi \cdot A \xi \rangle - \int_{\mathbb{R}^+} \frac{\lambda - 2/T}{(1/T + \lambda)^2} d e_\partial(\lambda).$$

In particular, the systematic error can be written as

$$\xi \cdot A_T \xi - \xi \cdot A_{\text{hom}} \xi = \frac{1}{T^2} \int_{\mathbb{R}^+} \frac{1}{\lambda(1/T + \lambda)^2} de_\delta(\lambda),$$

so that information on the scaling of the systematic error in terms of T yields information on the spectral behavior and conversely. The interplay between the strategy used in the present paper and the spectral measure is further investigated by Mourrat and the first author in [5]. In what follows, we do not make use of the spectral measure, which makes our approach self-contained.

The article is organized as follows: In Section 2, we introduce the general framework and state the main results of this paper, that is, the systematic error actually scales as in (1.10). The last two sections are dedicated to its proof.

Throughout the paper, we make use of the following notation:

- $d \geq 2$ is the dimension;
- $\int_{\mathbb{Z}^d} dx$ denotes the sum over $x \in \mathbb{Z}^d$, and $\int_D dx$ denotes the sum over $x \in \mathbb{Z}^d$ such that $x \in D$, D subset of \mathbb{R}^d ;
- $\langle \cdot \rangle$ is the ensemble average, or equivalently the expectation in the underlying probability space;
- $\text{var}[\cdot]$ is the variance associated with the ensemble average;
- $\text{cov}[\cdot; \cdot]$ is the covariance associated with the ensemble average;
- \lesssim and \gtrsim stand for \leq and \geq up to a multiplicative constant which only depends on the dimension d and the constants α, β (see Definition 1 below) if not otherwise stated;
- when both \lesssim and \gtrsim hold, we simply write \sim ;
- we use \gg instead of \gtrsim when the multiplicative constant is (much) larger than 1;
- $(\mathbf{e}_1, \dots, \mathbf{e}_d)$ denotes the canonical basis of \mathbb{Z}^d .

2. Main result.

2.1. General framework.

DEFINITION 1. We say that a is a conductivity function if there exist $0 < \alpha \leq \beta < \infty$ such that for every edge e of \mathbb{Z}^d , one has $a(e) \in [\alpha, \beta]$. We denote by $\mathcal{A}_{\alpha\beta}$ the set of such conductivity functions.

DEFINITION 2. The elliptic operator $L : L_{\text{loc}}^2(\mathbb{Z}^d) \rightarrow L_{\text{loc}}^2(\mathbb{Z}^d)$, $u \mapsto Lu$ associated with a conductivity function $a \in \mathcal{A}_{\alpha\beta}$ is defined for all $x \in \mathbb{Z}^d$ by

$$(2.1) \quad (Lu)(x) = -\nabla^* \cdot A(x) \nabla u(x),$$

where

$$\nabla u(x) := \begin{bmatrix} u(x + \mathbf{e}_1) - u(x) \\ \vdots \\ u(x + \mathbf{e}_d) - u(x) \end{bmatrix}, \quad \nabla^* u(x) := \begin{bmatrix} u(x) - u(x - \mathbf{e}_1) \\ \vdots \\ u(x) - u(x - \mathbf{e}_d) \end{bmatrix}$$

and

$$A(x) := \text{diag}[a(e_1), \dots, a(e_d)],$$

$$e_1 = [x, x + \mathbf{e}_1], \dots, e_d = [x, x + \mathbf{e}_d].$$

We now turn to the definition of the statistics of the conductivity function.

DEFINITION 3. A conductivity function is said to be independent and identically distributed (i.i.d.) if the coefficients $a(e)$ are i.i.d. random variables.

DEFINITION 4. The conductivity matrix A is obviously stationary in the sense that for all $z \in \mathbb{Z}^d$, $A(\cdot + z)$ and $A(\cdot)$ have the same statistics, so that for all $x, z \in \mathbb{Z}^d$,

$$\langle A(x + z) \rangle = \langle A(x) \rangle.$$

Therefore, any translation invariant function of A , such as the modified corrector ϕ_T (see Lemma 2), is jointly stationary with A . In particular, not only are ϕ_T and its gradient $\nabla\phi_T$ stationary, but also any function of A , ϕ_T and $\nabla\phi_T$. A useful such example is the energy density $(\xi + \nabla\phi_T) \cdot A(\xi + \nabla\phi_T)$, which is stationary by joint stationarity of A and $\nabla\phi_T$.

Another translation invariant function of A is the Green functions G_T of Definition 6. In this case, stationarity means that $G_T(\cdot + z, \cdot + z)$ has the same statistics as $G_T(\cdot, \cdot)$ for all $z \in \mathbb{Z}^d$, so that in particular, for all $x, y, z \in \mathbb{Z}^d$,

$$\langle G_T(x + z, y + z) \rangle = \langle G_T(x, y) \rangle.$$

LEMMA 1 (Corrector; [10], Theorem 3). *Let $a \in \mathcal{A}_{\alpha\beta}$ be an i.i.d. conductivity function, then for all $\xi \in \mathbb{R}^d$, there exists a unique random function $\phi: \mathbb{Z}^d \rightarrow \mathbb{R}$ which satisfies the corrector equation*

$$(2.2) \quad -\nabla^* \cdot A(x)(\xi + \nabla\phi(x)) = 0 \quad \text{in } \mathbb{Z}^d,$$

and such that $\phi(0) = 0$, $\nabla\phi$ is stationary and $\langle \nabla\phi \rangle = 0$. In addition, $\langle |\nabla\phi|^2 \rangle \lesssim |\xi|^2$.

We also define an ‘‘approximation’’ of the corrector as follows.

LEMMA 2 (Approximate corrector; [10], Proof of Theorem 3). *Let $a \in \mathcal{A}_{\alpha\beta}$ be an i.i.d. conductivity function, then for all $T > 0$ and $\xi \in \mathbb{R}^d$, there exists a unique stationary random function $\phi_T: \mathbb{Z}^d \rightarrow \mathbb{R}$ which satisfies the ‘‘approximate’’ corrector equation*

$$(2.3) \quad T^{-1}\phi_T(x) - \nabla^* \cdot A(x)(\xi + \nabla\phi_T(x)) = 0 \quad \text{in } \mathbb{Z}^d,$$

and such that $\langle \phi_T \rangle = 0$. In addition, $T^{-1}\langle \phi_T^2 \rangle + \langle |\nabla\phi_T|^2 \rangle \lesssim |\xi|^2$.

DEFINITION 5 (Homogenized coefficients). Let $a \in \mathcal{A}_{\alpha\beta}$ be an i.i.d. conductivity function and let $\xi \in \mathbb{R}^d$ and ϕ be as in Lemma 1. We define the homogenized $d \times d$ -matrix A_{hom} as

$$(2.4) \quad \xi \cdot A_{\text{hom}} \xi = \langle (\xi + \nabla \phi) \cdot A(\xi + \nabla \phi)(0) \rangle.$$

Note that (2.4) fully characterizes A_{hom} since A_{hom} is a symmetric matrix (it is actually of the form $a_{\text{hom}} \text{Id}$ for an i.i.d. conductivity function).

2.2. *Statement of the main results.* The main result of the article is the following estimate of the systematic error introduced in Section 1.

THEOREM 1. Let $a \in \mathcal{A}_{\alpha\beta}$ be an i.i.d. conductivity function, and let ϕ_T denote the approximate corrector associated with the conductivity function a and direction $\xi \in \mathbb{R}^d$, $|\xi| = 1$. We then define for all $T \gg 1$ the symmetric matrix A_T characterized by

$$(2.5) \quad \xi \cdot A_T \xi := \langle (\xi + \nabla \phi_T) \cdot A(\xi + \nabla \phi_T) \rangle.$$

Then, there exists an exponent $q > 0$ depending only on α, β such that

$$(2.6) \quad \begin{aligned} d = 2: & \quad |A_{\text{hom}} - A_T| \lesssim T^{-1} (\ln T)^q, \\ d = 3: & \quad |A_{\text{hom}} - A_T| \lesssim T^{-3/2}, \\ d = 4: & \quad |A_{\text{hom}} - A_T| \lesssim T^{-2} \ln T, \\ d > 4: & \quad |A_{\text{hom}} - A_T| \lesssim T^{-2}. \end{aligned}$$

As a by-product of the proof of Theorem 1, we obtain the following corollary.

COROLLARY 1. Let $a \in \mathcal{A}_{\alpha\beta}$ be an i.i.d. conductivity function, $d > 2$, $T > 0$, and let ϕ_T and $\tilde{\phi}$ denote the approximate corrector and stationary corrector (see [6], Corollary 1) associated with the conductivity function a and direction $\xi \in \mathbb{R}^d$, $|\xi| = 1$, respectively. Then

$$(2.7) \quad T^{-1} \langle (\phi_T - \tilde{\phi})^2 \rangle + \langle |\nabla \phi_T - \nabla \tilde{\phi}|^2 \rangle \lesssim \begin{cases} d = 3, & T^{-3/2}, \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}. \end{cases}$$

In particular,

$$\lim_{T \rightarrow \infty} \langle (\phi_T - \tilde{\phi})^2 \rangle + \langle |\nabla \phi_T - \nabla \tilde{\phi}|^2 \rangle = 0.$$

This corollary gives a full characterization of the convergence of the regularized corrector to the exact corrector for $d > 2$.

REMARK 1. Note that the definition (2.5) of A_T does not include the zero-order term $T^{-1}\langle\phi_T^2\rangle$, so that $\xi \cdot A_T \xi$ does not coincide with the energy associated with the equation. Surprisingly, the addition of the zero-order term in the definition of A_T would make the estimate (2.6) saturate at T^{-1} for $d > 2$.

REMARK 2. For $d = 2$, although we lose control of ϕ_T we may still quantify the rate of convergence of $\nabla\phi_T$ to $\nabla\phi$, the gradient of the corrector of Definition 1. In particular, (2.7) is replaced by

$$\langle|\nabla\phi_T - \nabla\phi|^2\rangle \lesssim T^{-1} \ln^q T$$

for some $q > 0$ depending only on α, β .

2.3. *Auxiliary lemmas.* In order to prove Theorem 1 and Corollary 1, we need three auxiliary lemmas in addition to the results of [6]: The first one is a covariance estimate very similar to the variance estimate in [6], Lemma 2.3, the next one is a refined version of the decay estimates of [6], Lemma 2.8, whereas the last one is a generalization of the convolution estimate of [6], Lemma 2.10.

LEMMA 3 (Covariance estimate). *Let $a = \{a_i\}_{i \in \mathbb{N}}$ be a sequence of i.i.d. random variables with range $[\alpha, \beta]$. Let X and Y be two Borel measurable functions of $a \in \mathbb{R}^{\mathbb{N}}$ (i.e., measurable w.r.t. the smallest σ -algebra on $\mathbb{R}^{\mathbb{N}}$ for which all coordinate functions $\mathbb{R}^{\mathbb{N}} \ni a \mapsto a_i \in \mathbb{R}$ are Borel measurable, cf. [7], Definition 14.4).*

Then we have

$$(2.8) \quad \text{cov}[X; Y] \leq \sum_{i=1}^{\infty} \left\langle \sup_{a_i} \left| \frac{\partial X}{\partial a_i} \right|^2 \right\rangle^{1/2} \left\langle \sup_{a_i} \left| \frac{\partial Y}{\partial a_i} \right|^2 \right\rangle^{1/2} \text{var}[a_1],$$

where $\sup_{a_i} \left| \frac{\partial Z}{\partial a_i} \right|$ denotes the supremum of the modulus of the i th partial derivative

$$\frac{\partial Z}{\partial a_i}(a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots)$$

of Z with respect to the variable $a_i \in [\alpha, \beta]$, for $Z = X, Y$.

The proof of this lemma is standard. As for [6], Lemma 2.3, it relies on a martingale difference decomposition.

We define discrete Green's functions in the following definition.

DEFINITION 6 (Discrete Green's function). Let $d \geq 2$. For all $T > 0$, the Green function $G_T : \mathcal{A}_{\alpha\beta} \times \mathbb{Z}^d \times \mathbb{Z}^d \rightarrow \mathbb{Z}^d$, $(a, x, y) \mapsto G_T(x, y; a)$ associated with the conductivity function a is defined for all $y \in \mathbb{Z}^d$ and $a \in \mathcal{A}_{\alpha\beta}$ as the unique solution $G(\cdot, y; a) \in L^2(\mathbb{Z}^d)$ to

$$(2.9) \quad \int_{\mathbb{Z}^d} T^{-1} G_T(x, y; a) v(x) dx + \int_{\mathbb{Z}^d} \nabla v(x) \cdot A(x) \nabla_x G_T(x, y; a) dx = v(y)$$

$$\forall v \in L^2(\mathbb{Z}^d),$$

where A is as in (2.1).

Throughout this paper, when no confusion occurs, we use the shorthand notation $G_T(x, y)$ for $G_T(x, y; a)$. We need a decay of the Green function $G_T(x, y)$ and its (discrete) gradient $\nabla_x G_T(x, y)$ in $|x - y| \gg 1$ that is *uniform* in a but nevertheless coincides (in terms of scaling) with the decay of the *constant-coefficient* Green function. The constant-coefficient Green function in the continuous case is known to decay as

$$\begin{aligned} |x - y|^{2-d} \exp\left(-\text{const.} \frac{|x - y|}{\sqrt{T}}\right) & \quad \text{for } d > 2 \quad \text{and} \\ \left(\ln \frac{\sqrt{T}}{|x - y|}\right) \exp\left(-\text{const.} \frac{|x - y|}{\sqrt{T}}\right) & \quad \text{for } d = 2; \end{aligned}$$

its gradient decays as the first derivative of these expressions. Note the cross-over of the decay at distances $|x - y|$ of the order of the intrinsic length scale $\sqrt{T} \gg 1$ from algebraic (or logarithmic in case of $d = 2$) to exponential.

In the class of a -uniform estimates, these decay properties survive as *pointwise* in (x, y) estimates on the level of the discrete Green function $G_T(x, y)$ itself, but only as *averaged* estimates on the level of its discrete gradient $\nabla_x G_T(x, y)$. More precisely, $\nabla_x G_T(x, y)$ has to be averaged in x on dyadic annuli centered at $x = y$. It will be important that the average can be (at least slightly) stronger than a *square* average (see [6], Lemma 2.9). On the other hand, we do not need the exponential decay: Super algebraic decay is sufficient for our purposes.

LEMMA 4 (Pointwise decay estimate on G_T). *Let $a \in \mathcal{A}_{\alpha\beta}$, and G_T be the associated Green function. For $d > 2$, we have for all $k > 0$, and all $x, y \in \mathbb{Z}^d$*

$$(2.10) \quad G_T(x, y) \lesssim (1 + |x - y|)^{2-d} \min\left\{1, \left(\frac{|x - y|}{\sqrt{T}}\right)^{-k}\right\},$$

where the constant in “ \lesssim ” depends on k . For $d = 2$, we have for all $k > 0$

$$(2.11) \quad G_T(x, y) \lesssim \begin{cases} \ln\left(\frac{\sqrt{T}}{1 + |x - y|}\right) & \text{for } |x - y| \ll \sqrt{T} \\ \left(\frac{|x - y|}{\sqrt{T}}\right)^{-k} & \text{for } |x - y| \gtrsim \sqrt{T} \end{cases},$$

where the constant in “ \lesssim ” depends on k .

Finally, for the proof of Theorem 1, we need to know that also the *convolution* of the gradient of the Green’s function with itself decays at the optimal rate, that is, with the following lemma.

LEMMA 5 (Convolution estimate). *Let $h_T, g_T : \mathbb{Z}^d \rightarrow \mathbb{R}^+$ satisfy the following properties.*

Assumptions on h_T [estimate of $|\nabla_x G_T(y+z, y)|$]: For all $R \gg 1$ and $T > 0$,

$$(2.12) \quad d > 2: \quad \int_{R < |z| \leq 2R} h_T(z)^2 dz \lesssim R^{2-d},$$

$$(2.13) \quad d = 2: \quad \int_{R < |z| \leq 2R} h_T(z)^2 dz \lesssim \min\{1, \sqrt{T}R^{-1}\}^2,$$

and for $R \sim 1$

$$(2.14) \quad d \geq 2: \quad \int_{|z| \leq R} h_T(z)^2 dz \lesssim 1.$$

Assumptions on g_T [estimate of $G_T(y+z, y)$]: For $d > 2$, and for all $z \in \mathbb{Z}^d$,

$$(2.15) \quad g_T(z) = (1 + |z|)^{2-d} \min\left\{1, \left(\frac{|z|}{\sqrt{T}}\right)^{-3}\right\},$$

and for $d = 2$,

$$(2.16) \quad g_T(z) = \begin{cases} \ln\left(\frac{1+|z|}{\sqrt{T}}\right) & \text{for } |z| \leq \sqrt{T} \\ \left(\frac{|z|}{\sqrt{T}}\right)^{-3} & \text{for } |z| > \sqrt{T} \end{cases}.$$

Then we have

$$(2.17) \quad \int_{\mathbb{Z}^d} g_T(z) \int_{\mathbb{Z}^d} h_T(w) h_T(z-w) dw dz \lesssim \begin{cases} d = 2, & T, \\ d = 3, & \sqrt{T}, \\ d = 4, & \ln T, \\ d > 4, & 1. \end{cases}$$

3. Proof of the main results. Throughout this section, we let $\xi \in \mathbb{R}^d$ be such that $|\xi| = 1$.

3.1. *Proof of Theorem 1.* In view of (1.7), in order to estimate $|A_T - A_{\text{hom}}|$, we need to estimate how close the modified corrector ϕ_T is to the original corrector ϕ [in terms of $\langle |\nabla \phi_T - \nabla \phi|^2 \rangle$]. Therefore, it is natural to introduce $\psi_T = T^2 \frac{\partial \phi_T}{\partial T}$ (the prefactor T^2 is such that ψ_T is properly renormalized in the limit $T \uparrow \infty$ at least for large d). Considering ψ_T is also convenient since for $d = 2$, the corrector ϕ is not known to be stationary (only its gradient is known to be stationary) so that working with the modified correctors ϕ_T , which are known to be stationary, avoids technical subtleties. In fact, we opt for a dyadically discrete version of ψ_T defined via

$$(3.1) \quad \psi_T := T(\phi_{2T} - \phi_T).$$

This discrete version has the technical advantage that we do not have to think about the differentiability of ϕ_T in T . Moreover, its dyadic nature is in line with the dyadic decomposition of the T -axis according to

$$(3.2) \quad |A_T - A_{\text{hom}}| \leq \sum_{i=0}^{\infty} |A_{2^i T} - A_{2^{i+1} T}|$$

forced upon us in the case of $d = 2$. In order to get (3.2), we used the fact that

$$(3.3) \quad \lim_{T \rightarrow \infty} A_T = A_{\text{hom}},$$

which is proved in [6], Proof of Theorem 1, Step 8. We shall also use that ψ_T solves

$$(3.4) \quad T^{-1} \psi_T - \nabla^* \cdot A \nabla \psi_T = \frac{1}{2} \phi_{2T}.$$

We split the proof in eight steps.

Step 1. Derivation of

$$(3.5) \quad |\xi \cdot (A_{2T} - A_T) \xi| \leq T^{-2} |\langle \phi_T \psi_T \rangle| + \frac{T^{-2}}{2} |\langle \phi_{2T} \psi_T \rangle|.$$

Although this could be directly inferred from the spectral formula (1.11) for A_T , we give an elementary argument relying only on the corrector equation. We recall the following consequence of (2.3) which is proved in [6], Proof of Theorem 1, Step 8:

$$(3.6) \quad T^{-1} \langle \phi_T \chi \rangle + \langle (\xi + \nabla \phi_T) \cdot A \nabla \chi \rangle = 0$$

for every field $\chi : \mathbb{Z}^d \rightarrow \mathbb{R}$ that is jointly stationary with A and such that $\langle \chi^2 \rangle < \infty$. From formally differentiating the definition (2.5) of A_T w.r.t. T and using (3.6) for $\chi = \frac{\partial \phi_T}{\partial T}$, we obtain

$$\xi \cdot \frac{\partial A_T}{\partial T} \xi = -2T^{-1} \left\langle \frac{\partial \phi_T}{\partial T} \phi_T \right\rangle.$$

We claim that the corresponding discrete-in- T version reads

$$(3.7) \quad \xi \cdot (A_{2T} - A_T) \xi = -T^{-2} (\langle \psi_T \phi_T \rangle + \frac{1}{2} \langle \psi_T \phi_{2T} \rangle).$$

Indeed, by definition of A_T , by expanding the square, by symmetry of A , by definition of ψ_T , and (3.6), we have

$$\begin{aligned} & \xi \cdot (A_{2T} - A_T) \xi \\ &= \langle (\xi + \nabla \phi_{2T}) \cdot A (\xi + \nabla \phi_{2T}) \rangle - \langle (\xi + \nabla \phi_T) \cdot A (\xi + \nabla \phi_T) \rangle \\ &= \langle (\nabla \phi_{2T} - \nabla \phi_T) \cdot A (\xi + \nabla \phi_{2T}) \rangle + \langle (\nabla \phi_{2T} - \nabla \phi_T) \cdot A (\xi + \nabla \phi_T) \rangle \\ &\stackrel{(3.1)}{=} T^{-1} (\langle \nabla \psi_T \cdot A (\xi + \nabla \phi_{2T}) \rangle + \langle \nabla \psi_T \cdot A (\xi + \nabla \phi_T) \rangle) \\ &\stackrel{(3.6)}{=} -T^{-1} ((2T)^{-1} \langle \psi_T \phi_{2T} \rangle + T^{-1} \langle \psi_T \phi_T \rangle). \end{aligned}$$

In the next four steps, we focus on the first term of the r.h.s. of (3.5). The second term will be dealt with the same way in Step 7.

Step 2. Proof of

$$(3.8) \quad |\langle \phi_T \psi_T \rangle| \lesssim \sum_e \left\langle \sup_{a(e)} \left| \frac{\partial \phi_T(0)}{\partial a(e)} \right|^2 \right\rangle^{1/2} \left\langle \sup_{a(e)} \left| \frac{\partial \psi_T(0)}{\partial a(e)} \right|^2 \right\rangle^{1/2},$$

where the sum runs over the edges e , and proof of the representation formulas

$$(3.9) \quad \frac{\partial \phi_T(0)}{\partial a(e)} = -(\xi_i + \nabla_i \phi_T(z)) \nabla_{z_i} G_T(z, 0),$$

$$(3.10) \quad \begin{aligned} \frac{\partial \psi_T(0)}{\partial a(e)} &= -\nabla_i \psi_T(z) \nabla_{z_i} G_T(z, 0) \\ &\quad - \frac{1}{2} \int_{\mathbb{Z}^d} G_T(0, w) (\xi_i + \nabla_i \phi_{2T}(z)) \nabla_{z_i} G_{2T}(z, w) dw, \end{aligned}$$

where the edge is $e = [z, z + \mathbf{e}_i]$.

Due to [6], Lemma 2.6, the functions ϕ_T and ψ_T are measurable with respect to the coefficients a . Hence, (3.8) is a consequence of the covariance estimate of Lemma 3: Since $\langle \phi_T \rangle = \langle \psi_T \rangle = 0$,

$$\begin{aligned} \langle \phi_T \psi_T \rangle &= \langle (\phi_T - \langle \phi_T \rangle) (\psi_T - \langle \psi_T \rangle) \rangle \\ &= \text{cov}[\phi_T; \psi_T]. \end{aligned}$$

Formula (3.9) is identical to [6], Lemma 2.4, (2.12). To prove (3.10), we first make use of the Green representation formula for the solution to (3.4):

$$(3.11) \quad \psi_T(x) = \frac{1}{2} \int_{\mathbb{Z}^d} G_T(x, w) \phi_{2T}(w) dw$$

for all $x \in \mathbb{Z}^d$. Since $a(e) \mapsto \phi_T(\cdot; a(e))$ and $a(e) \mapsto \phi_{2T}(\cdot; a(e))$ are continuously differentiable by [6], Lemma 2.4, we deduce by formula (3.1) that $a(e) \mapsto \psi_T(\cdot; a(e))$ is also continuously differentiable. Using then the formulas [6], Lemma 2.5, (2.15), and [6], Lemma 2.4, (2.12), for the derivatives of G_T and ϕ_T with respect to $a(e)$, and the fact that $G_T \in L^1(\mathbb{Z}^d)$ (see [6], Corollary 2.2), we may switch the order of the differentiation and the integration to obtain for all $x \in \mathbb{Z}^d$

$$(3.12) \quad \begin{aligned} \frac{\partial \psi_T(x)}{\partial a(e)} &= \frac{1}{2} \int_{\mathbb{Z}^d} \frac{\partial G_T(x, w)}{\partial a(e)} \phi_{2T}(w) dw \\ &\quad + \frac{1}{2} \int_{\mathbb{Z}^d} G_T(x, w) \frac{\partial \phi_{2T}(w)}{\partial a(e)} dw \\ &\stackrel{[6], (2.12) \text{ and } (2.15)}{=} -\frac{1}{2} \int_{\mathbb{Z}^d} \nabla_{z_i} G_T(x, z) \nabla_{z_i} G_T(z, w) \phi_{2T}(w) dw \end{aligned}$$

$$\begin{aligned}
& - \frac{1}{2} \int_{\mathbb{Z}^d} G_T(x, w) (\xi_i + \nabla_i \phi_{2T}(z)) \nabla_{z_i} G_{2T}(z, w) dw \\
& \stackrel{(3.11)}{=} - \nabla_{z_i} G_T(x, z) \nabla_i \psi_T(z) \\
& - \frac{1}{2} \int_{\mathbb{Z}^d} G_T(x, w) (\xi_i + \nabla_i \phi_{2T}(z)) \nabla_{z_i} G_{2T}(z, w) dw,
\end{aligned}$$

which is (3.10) taking $x = 0$.

From now on in the proof, we let g_T be defined as in Lemma 5 (i.e., g_T decays as the Green function G_T).

Step 3. In this step, we shall prove that

$$(3.13) \quad |\langle \phi_T \psi_T \rangle| \lesssim \mathcal{L} + \mathcal{N},$$

where

$$\begin{aligned}
(3.14) \quad \mathcal{L} & := \int_{\mathbb{Z}^d} \langle (1 + |\nabla \phi_T(z)|^2) |\nabla_z G_T(z, 0)|^2 \rangle^{1/2} \\
& \quad \times \left\langle \left(\int_{\mathbb{Z}^d} g_T(w) (1 + |\nabla \phi_{2T}(z)|) |\nabla_z G_{2T}(z, w)| dw \right)^2 \right\rangle^{1/2} dz,
\end{aligned}$$

and $\mathcal{N} = \mathcal{N}_1 + \mathcal{N}_2$,

$$\begin{aligned}
(3.15) \quad \mathcal{N}_1 & := \int_{\mathbb{Z}^d} \langle (1 + |\nabla \phi_T(z)|^2) |\nabla_z G_T(z, 0)|^2 \rangle^{1/2} \\
& \quad \times \langle |\nabla \psi_T(z)|^2 |\nabla_z G_T(z, 0)|^2 \rangle^{1/2} dz,
\end{aligned}$$

$$\begin{aligned}
(3.16) \quad \mathcal{N}_2 & := \mu_d(T) \int_{\mathbb{Z}^d} \langle (1 + |\nabla \phi_T(z)|^2) |\nabla_z G_T(z, 0)|^2 \rangle^{1/2} \\
& \quad \times \langle (1 + |\nabla \phi_{2T}(z)|^2) |\nabla_z G_T(z, 0)|^2 \rangle^{1/2} dz
\end{aligned}$$

with

$$\mu_d(T) := \begin{cases} d = 2, & \ln T, \\ d > 2, & 1. \end{cases}$$

The term \mathcal{L} is a linear error: It is of the same type as for the analysis in the limit of vanishing ellipticity contrast (see [6], the Appendix). On the contrary, the term \mathcal{N} is nonlinear and does not appear in the limit of vanishing ellipticity contrast. As we shall prove, it is of lower order. The terms \mathcal{L} and \mathcal{N}_1 in estimate (3.13) would be direct consequences of (3.8), and (3.9) and (3.10), disregarding the suprema in $a(e)$ in (3.8). Taking the suprema in $a(e)$ into account actually brings the second nonlinear term \mathcal{N}_2 , which turns out to be of lower order than \mathcal{N}_1 .

According to [6], Lemma 2.4, (2.13), we have for (3.9)

$$(3.17) \quad \sup_{a(e)} \left| \frac{\partial \phi_T(0)}{\partial a(e)} \right| \lesssim (1 + |\nabla_i \phi_T(z)|) |\nabla_z G_T(z, 0)|.$$

It remains to deal with (3.10). Using the pointwise decay of G_T in Lemma 4 combined with the susceptibility estimates [6], Lemma 2.4, (2.14), and [6], Lemma 2.5, (2.16), of $\nabla\phi_T$ and ∇G_T w.r.t. $a(e)$, we obtain

$$(3.18) \quad \sup_{a(e)} \left| \frac{1}{2} \int_{\mathbb{Z}^d} G_T(0, w) (\xi_i + \nabla_i \phi_{2T}(z)) \nabla_{z_i} G_{2T}(z, w) dw \right| \\ \lesssim \int_{\mathbb{Z}^d} g_T(w) (1 + |\nabla_i \phi_{2T}(z)|) |\nabla_z G_{2T}(z, w)| dw,$$

which together with (3.17) gives the linear term \mathcal{L} .

To treat the first term of the r.h.s. of (3.10), we need to deal with the supremum of $|\nabla_i \psi_T(z)|$ over $a(e)$. We appeal to (3.12) that we rewrite in the form

$$\frac{\partial \psi_T(x)}{\partial a(e)} = -\nabla_i \psi_T(z) G_T(x, e) \\ - \frac{1}{2} (\xi_i + \nabla_i \phi_{2T}(z)) \int_{\mathbb{Z}^d} G_T(x, w) G_T(e, w) dw,$$

where $G_T(x, e) := G_T(x, z + \mathbf{e}_i) - G_T(x, z)$ and $G_T(e, w) := G_T(z + \mathbf{e}_i, w) - G_T(z, w)$. Hence,

$$(3.19) \quad \frac{\partial \nabla_i \psi_T(z)}{\partial a(e)} = -\nabla_i \psi_T(z) G_T(e, e) \\ - \frac{1}{2} (\xi_i + \nabla_i \phi_{2T}(z)) \int_{\mathbb{Z}^d} G_T(e, w) G_{2T}(e, w) dw,$$

where $G_T(e, e) := G_T(z + \mathbf{e}_i, z + \mathbf{e}_i) + G_T(z, z) - G_T(z + \mathbf{e}_i, z) - G_T(z, z + \mathbf{e}_i)$. On the one hand, the uniform bound [6], Corollary 2.3, on ∇G_T yields $|G_T(e, e)| \lesssim 1$. On the other hand, as we shall argue, the integrability of ∇G_T and ∇G_{2T} from [6], Lemma 2.9 (combined with the uniform bound [6], Corollary 2.3, on gradients) implies

$$(3.20) \quad \int_{\mathbb{Z}^d} G_T(e, w) G_{2T}(e, w) dw \lesssim \mu_d(T) = \begin{cases} d = 2, & \ln T, \\ d > 2, & 1. \end{cases}$$

Hence, if we regard (3.19) as an ordinary differential equation for $\nabla_i \psi_T(z)$ in the variable $a(e)$, we obtain

$$(3.21) \quad \sup_{a(e)} |\nabla_i \psi_T(z)| \lesssim |\nabla_i \psi_T(z)| + \mu_d(T) (1 + |\nabla_i \phi_{2T}(z)|)$$

since $a(e)$ lies in a bounded domain $[\alpha, \beta]$, and $\sup_{a(e)} |\nabla_i \phi_{2T}(z)| \lesssim 1 + |\nabla_i \phi_{2T}(z)|$ according to [6], Lemma 2.4, (2.14), with $2T$ instead of T . Note that (3.17), (3.21) and $\sup_{a(e)} |\nabla_{z_i} G_T(z, 0)| \lesssim |\nabla_{z_i} G_T(z, 0)|$ give the nonlinear terms \mathcal{N}_1 and \mathcal{N}_2 .

We now give the argument for (3.20). We first use the Cauchy–Schwarz inequality

$$\begin{aligned} & \int_{\mathbb{Z}^d} G_T(e, w) G_{2T}(e, w) dw \\ & \leq \left(\int_{\mathbb{Z}^d} G_T(e, w)^2 dw \right)^{1/2} \left(\int_{\mathbb{Z}^d} G_{2T}(e, w)^2 dw \right)^{1/2} \\ & \leq \left(\int_{\mathbb{Z}^d} |\nabla_z G_T(z, w)|^2 dw \right)^{1/2} \left(\int_{\mathbb{Z}^d} |\nabla_z G_{2T}(z, w)|^2 dw \right)^{1/2} \end{aligned}$$

and then make a decomposition of \mathbb{Z}^d into the ball of radius $R \sim 1$, and dyadic annuli $\{w : 2^i R < |z - w| \leq 2^{i+1} R\}$ for $i \in \mathbb{N}$. On the ball of radius R , we use the uniform estimate of [6], Corollary 2.3, on ∇G_T , whereas on the dyadic annuli we appeal to the decay estimate in [6], Lemma 2.9, for the gradient of the Green function, which requires R to be sufficiently large although still of order 1. Both terms in the r.h.s. scale the same way and we only treat the first one:

$$\begin{aligned} & \int_{\mathbb{Z}^d} |\nabla_z G_T(z, w)|^2 dw \\ & = \int_{|z-w| \leq R} |\nabla_z G_T(z, w)|^2 dw + \sum_{i=0}^{\infty} \int_{2^i R < |z-w| \leq 2^{i+1} R} |\nabla_z G_T(z, w)|^2 dw \\ & \lesssim 1 + \sum_{i=1}^{\infty} (2^i)^{d+2(1-d)} \min\{1, \sqrt{T}(2^i R)^{-1}\}^2 \\ & \lesssim \mu_d(T), \end{aligned}$$

using [6], Corollary 2.3, and [6], Lemma 2.9, for $k = 2$, respectively. This concludes Step 3.

Step 4. Suboptimal estimate of the nonlinear term \mathcal{N} :

$$(3.22) \quad \mathcal{N}_1 \lesssim \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \begin{cases} d = 2, & \sqrt{T} \ln^q T, \\ d = 3, & \ln T, \\ d > 3, & 1, \end{cases}$$

$$(3.23) \quad \mathcal{N}_2 \lesssim \mu_d(T)^q,$$

where q is a generic exponent which only depends on α, β . We first deal with \mathcal{N}_1 , and begin with the second factor of the r.h.s. of (3.15). The pointwise estimate (2.10) of Lemma 4 for $d > 2$ on the Green function gives the *suboptimal pointwise* estimate on the gradient of the Green function

$$(3.24) \quad |\nabla G_T(z, 0)| \leq G_T(z, 0) + \sum_{i=1}^d G_T(z + \mathbf{e}_i, 0) \lesssim (1 + |z|)^{2-d}.$$

This estimate coincides for $d = 2$ with the uniform bound of [6], Corollary 2.3. The coercivity of A thus yields

$$\begin{aligned} & \langle |\nabla G_T(z, 0)|^2 |\nabla \psi_T(z)|^2 \rangle^{1/2} \\ & \lesssim (1 + |z|)^{2-d} \langle \nabla \psi_T(z) \cdot A(z) \nabla \psi_T(z) \rangle^{1/2} \\ & = (1 + |z|)^{2-d} \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \end{aligned}$$

by joint stationarity of $\nabla \psi_T$ and A . Hence, (3.15) turns into

$$\mathcal{N}_1 \lesssim \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle (1 + |\nabla \phi_T(z)|^2) |\nabla G_T(z, 0)|^2 \rangle^{1/2} dz.$$

We then let $p > 2$ be a Meyers' exponent as in [6], Lemma 2.9 and use Hölder's inequality in probability with exponents $(p/(p-2), p/2)$, the stationarity of $\nabla \phi_T$, the fact that the gradient of ϕ_T is estimated by ϕ_T as in (3.24), and the bounds on the stochastic moments of ϕ_T in [6], Proposition 1,

$$\begin{aligned} & \mathcal{N}_1 \lesssim \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \\ & \quad \times \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle 1 + |\nabla \phi_T(z)|^{2p/(p-2)} \rangle^{(p-2)/(2p)} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \\ & = \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \langle 1 + |\nabla \phi_T|^{2p/(p-2)} \rangle^{(p-2)/(2p)} \\ (3.25) \quad & \times \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \\ & \lesssim \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \langle 1 + |\phi_T|^{2p/(p-2)} \rangle^{(p-2)/(2p)} \\ & \quad \times \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \\ & \lesssim \mu_d(T)^q \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle^{1/2} \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz, \end{aligned}$$

for some generic q depending only on α, β . Hölder's inequality with exponents $(p, p/(p-1))$ in \mathbb{Z}^d , combined with the same dyadic decomposition of \mathbb{Z}^d as for the proof of (3.20) (and the uniform bound on ∇G_T from [6], Corollary 2.3) yields

$$\begin{aligned} & \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \\ & \lesssim 1 + \sum_{i=0}^{\infty} \left(\left\langle \int_{2^i R < |z| \leq 2^{i+1} R} |\nabla G_T(z, 0)|^p dz \right\rangle \right)^{1/p} \\ & \quad \times \left(\int_{2^i R < |z| \leq 2^{i+1} R} (1 + |z|)^{(2-d)p/(p-1)} dz \right)^{(p-1)/p}. \end{aligned}$$

Using the optimal decay of ∇G_T on dyadic annuli in L^p norm from [6], Lemma 2.9, with $k = 2p$, this turns into

$$\begin{aligned} & \int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \\ & \lesssim 1 + \sum_{i=0}^{\infty} \left((2^i R)^d (2^i R)^{(1-d)p} \min \left\{ 1, \frac{\sqrt{T}}{2^i R} \right\}^{2p} \right)^{1/p} \\ & \quad \times ((2^i R)^d (2^i R)^{(2-d)p/(p-1)})^{(p-1)/p} \\ & = 1 + \sum_{i=0}^{\infty} (2^i R)^{3-d} \min \left\{ 1, \frac{\sqrt{T}}{2^i R} \right\}^2. \end{aligned}$$

Recalling that $R \sim 1$, this implies

$$\int_{\mathbb{Z}^d} (1 + |z|)^{2-d} \langle |\nabla G_T(z, 0)|^p \rangle^{1/p} dz \lesssim \begin{cases} d = 2, & \sqrt{T}, \\ d = 3, & \ln T, \\ d > 3, & 1. \end{cases}$$

Combined with (3.25) it proves (3.22).

We now turn to \mathcal{N}_2 . Proceeding as above to deal with the terms $\nabla \phi_T$ and $\nabla \phi_{2T}$ in \mathcal{N}_2 , we obtain as desired

$$\begin{aligned} \mathcal{N}_2 & \lesssim \mu_d(T) \mu_d(T)^{2q} \int_{\mathbb{Z}^d} \langle |\nabla_z G_T(z, 0)|^p \rangle^{2/p} dz \\ & \lesssim \mu_d(T)^{2q+2}, \end{aligned}$$

using the same dyadic decomposition of \mathbb{Z}^d as for the proof of (3.20) together with the higher integrability of gradients of [6], Lemma 2.9 and [6], Corollary 2.3.

Step 5. Estimate of the linear term \mathcal{L} :

$$(3.26) \quad \mathcal{L} \lesssim \begin{cases} d = 2, & T \ln^q T, \\ d = 3, & \sqrt{T}, \\ d = 4, & \ln T, \\ d > 4, & 1. \end{cases}$$

We first treat the second factor of (3.14). We proceed as in Step 4 to deal with the expectation of the corrector term, and let $p > 2$ be a Meyers' exponent as in [6], Lemma 2.9. We obtain by Hölder's inequality in probability with exponents $(p/(p-2), p, p)$ and the bounds on the stochastic moments of ϕ_T from [6], Proposition 1:

$$\begin{aligned} & \left\langle \left(\int_{\mathbb{Z}^d} g_T(w) (1 + |\nabla \phi_{2T}(z)|) |\nabla_{z_i} G_{2T}(z, w)| dw \right)^2 \right\rangle \\ & = \int_{\mathbb{Z}^d} \int_{\mathbb{Z}^d} g_T(w) g_T(w') \end{aligned}$$

$$\begin{aligned}
& \times \langle (1 + |\nabla\phi_{2T}(z)|)^2 |\nabla_{z_i} G_{2T}(z, w)| |\nabla_{z_i} G_{2T}(z, w')| \rangle dw dw' \\
& \lesssim (1 + \langle |\phi_{2T}|^{2p/(p-2)} \rangle^{(p-2)/p}) \\
& \quad \times \int_{\mathbb{Z}^d} \int_{\mathbb{Z}^d} g_T(w) g_T(w') \langle |\nabla_{z_i} G_{2T}(z, w)|^p \rangle^{1/p} \\
& \quad \quad \times \langle |\nabla_{z_i} G_{2T}(z, w')|^p \rangle^{1/p} dw dw' \\
& \lesssim \mu_d(T)^q \left(\int_{\mathbb{Z}^d} g_T(w) \langle |\nabla_z G_{2T}(z, w)|^p \rangle^{1/p} dw \right)^2.
\end{aligned}$$

We thus have

$$\begin{aligned}
\mathcal{L} & \lesssim \mu_d(T)^q \int_{\mathbb{Z}^d} \langle (1 + |\nabla\phi_T(z)|^2) |\nabla G_T(z, 0)|^2 \rangle^{1/2} \\
& \quad \times \int_{\mathbb{Z}^d} g_T(w) \langle |\nabla_z G_{2T}(z, w)|^p \rangle^{1/p} dw dz.
\end{aligned}$$

Appealing once more to Hölder's inequality in probability with exponents $(p/(p-2), p/2)$ and to [6], Proposition 1, this turns into

$$\begin{aligned}
\mathcal{L} & \lesssim \mu_d(T)^{2q} \int_{\mathbb{Z}^d} g_T(w) \int_{\mathbb{Z}^d} \langle |\nabla_z G_{2T}(z, w)|^p \rangle^{1/p} \langle |\nabla_z G_T(z, 0)|^p \rangle^{1/p} dz dw \\
& = \mu_d(T)^{2q} \int_{\mathbb{Z}^d} g_T(w) \int_{\mathbb{Z}^d} h_{2T}(z-w) h_T(z) dz dw,
\end{aligned}$$

where, by stationarity, we have set

$$\begin{aligned}
h_T(w) & = \langle |\nabla_w G_T(w, 0)|^p \rangle^{1/p}, \\
h_{2T}(w) & = \langle |\nabla_w G_{2T}(w, 0)|^p \rangle^{1/p}.
\end{aligned}$$

By the optimal decay estimate of ∇G_T on dyadic annuli from [6], Lemma 2.9 (and by the uniform bounds on ∇G_T from [6], Corollary 2.3), and by definition of g_T , we are in position to apply Lemma 5. Estimate (3.26) is thus proved.

Step 6. Proof of

$$(3.27) \quad \langle \nabla\psi_T \cdot A\nabla\psi_T \rangle \leq |\langle \phi_T \psi_T \rangle|.$$

Using (3.1), we rewrite (3.4) as

$$\begin{aligned}
(3.28) \quad (2T)^{-1}\psi_T - \nabla^* \cdot A\nabla\psi_T & = \frac{1}{2}\phi_{2T} - (2T)^{-1}\psi_T \\
& = \frac{1}{2}\phi_T.
\end{aligned}$$

We now multiply (3.28) by ψ_T :

$$(2T)^{-1}\psi_T^2 - (\nabla^* \cdot A\nabla\psi_T)\psi_T = \frac{1}{2}\phi_T\psi_T.$$

By integration by parts and joint stationarity of ψ_T , $\nabla\psi_T$ and A (see [6], Proof of Theorem 1, Step 8, for details), this turns into

$$(2T)^{-1}\langle \psi_T^2 \rangle + \langle \nabla\psi_T \cdot A\nabla\psi_T \rangle = \frac{1}{2}\langle \phi_T\psi_T \rangle.$$

We then conclude by the nonnegativity of the first term.

Step 7. Proof of

$$(3.29) \quad |\langle \phi_T \psi_T \rangle| \lesssim \begin{cases} d=2, & T \ln^q T, \\ d=3, & \sqrt{T}, \\ d=4, & \ln T, \\ d>4, & 1, \end{cases}$$

and

$$(3.30) \quad |\langle \phi_{2T} \psi_T \rangle| \lesssim \begin{cases} d=2, & T \ln^q T, \\ d=3, & \sqrt{T}, \\ d=4, & \ln T, \\ d>4, & 1. \end{cases}$$

From Steps 3, 4 and 5, and Young's inequality, we deduce that

$$|\langle \phi_T \psi_T \rangle| - \frac{1}{2} \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle \lesssim \begin{cases} d=2, & T \ln^q T, \\ d=3, & \sqrt{T}, \\ d=4, & \ln T, \\ d>4, & 1. \end{cases}$$

Combined with Step 6, this shows (3.29).

For (3.30), we proceed exactly as for (3.29) in Steps 2–6. In particular, with obvious notation, we have

$$|\langle \phi_{2T} \psi_T \rangle| \lesssim \mathcal{N}' + \mathcal{L}',$$

where

$$\mathcal{N}' - \frac{1}{2} \langle \nabla \psi_T \cdot A \nabla \psi_T \rangle \lesssim \begin{cases} d=2, & T \ln^q T, \\ d=3, & \ln^2 T, \\ d>3, & 1, \end{cases}$$

and

$$\mathcal{L}' \lesssim \begin{cases} d=2, & T \ln^q T, \\ d=3, & \sqrt{T}, \\ d=4, & \ln T, \\ d>4, & 1. \end{cases}$$

We then conclude as above.

Step 8. Proof of (2.6).

Steps 1 and 7 yield

$$(3.31) \quad \begin{aligned} |\xi \cdot (A_T - A_{2T}) \xi| &\leq T^{-2} |\langle \phi_T \psi_T \rangle| + (2T^2)^{-1} |\langle \phi_{2T} \psi_T \rangle| \\ &\lesssim T^{-2} \begin{cases} d=2, & T \ln^q T, \\ d=3, & \sqrt{T}, \\ d=4, & \ln T, \\ d>4, & 1. \end{cases} \end{aligned}$$

We finally appeal to the dyadic decomposition of the T -axis (3.2), which, combined with (3.31), turns into

$$\begin{aligned} |\xi \cdot (A_T - A_{\text{hom}})\xi| &\lesssim \sum_{i=1}^{\infty} \begin{cases} d = 2, & (2^i T)^{-1} \ln^q(2^i T), \\ d = 3, & (2^i T)^{-3/2}, \\ d = 4, & (2^i T)^{-2} \ln(2^i T), \\ d > 4, & (2^i T)^{-2}, \end{cases} \\ &\lesssim \begin{cases} d = 2, & T^{-1} \ln^q T, \\ d = 3, & T^{-3/2} \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}. \end{cases} \end{aligned}$$

This concludes the proof of the theorem.

3.2. Proof of Corollary 1. By Steps 6 and 7 in the proof of Theorem 1 and by the definition (3.1) of ψ_T , we learn that

$$\begin{aligned} \langle |\nabla\phi_{2T} - \nabla\phi_T|^2 \rangle &\stackrel{(3.1)}{=} T^{-2} \langle |\nabla\psi_T|^2 \rangle \\ &\stackrel{(3.27) \text{ and } (3.29)}{\lesssim} \begin{cases} d = 3, & T^{-3/2}, \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}. \end{cases} \end{aligned}$$

In particular, $\nabla\phi_T$ is a Cauchy sequence in L^2 in probability. Hence, $\nabla\phi_T$ converges in L^2 to its weak limit $\nabla\phi$, and by a dyadic decomposition of the T -axis the above estimate yields

$$\langle |\nabla\phi_T - \nabla\phi|^2 \rangle \lesssim \begin{cases} d = 3, & T^{-3/2}, \\ d = 4, & T^{-2} \ln T, \\ d > 4, & T^{-2}, \end{cases}$$

which gives the second term of the l.h.s. of (2.7).

Likewise, from Step 7 in the proof of Theorem 1, we learn that

$$\begin{aligned} \langle (\phi_{2T} - \phi_T)^2 \rangle &\stackrel{(3.1)}{=} T^{-1} \langle (\phi_{2T} - \phi_T)\psi_T \rangle \\ &\leq T^{-1} (\langle |\phi_{2T}\psi_T| \rangle + \langle |\phi_T\psi_T| \rangle) \\ &\stackrel{(3.29) \text{ and } (3.30)}{\lesssim} \begin{cases} d = 3, & T^{-1/2}, \\ d = 4, & T^{-1} \ln T, \\ d > 4, & T^{-1}, \end{cases} \end{aligned}$$

so that ϕ_T is a Cauchy sequence in L^2 in probability and ϕ_T converges in L^2 to its weak limit $\tilde{\phi}$ provided by [6], Corollary 1. In particular, by a dyadic decomposition of the T -axis the above estimate yields

$$\langle (\phi_T - \tilde{\phi})^2 \rangle = \begin{cases} d = 3, & T^{-1/2}, \\ d = 4, & T^{-1} \ln T, \\ d > 4, & T^{-1}, \end{cases}$$

which is the first term of the l.h.s. of (2.7). This concludes the proof of the corollary.

4. Proof of the auxiliary lemmas.

4.1. *Proof of Lemma 3.* Without loss of generality we may assume

$$(4.1) \quad \sum_{i=1}^{\infty} \left\langle \sup_{a_i} \left| \frac{\partial X}{\partial a_i} \right|^2 \right\rangle, \sum_{i=1}^{\infty} \left\langle \sup_{a_i} \left| \frac{\partial Y}{\partial a_i} \right|^2 \right\rangle < \infty.$$

Let Z_n denote the expected value of Z conditioned on a_1, \dots, a_n , that is

$$Z_n(a_1, \dots, a_n) := \langle Z | a_1, \dots, a_n \rangle.$$

From [6], (5.2) and (5.3), in the proof of [6], Lemma 2.3, we learn that

$$\lim_{n \uparrow \infty} \langle (Z - Z_n)^2 \rangle = 0$$

for $Z = X$, $Z_n = X_n$ and $Z = Y$, $Z_n = Y_n$, respectively, so that, by the Cauchy–Schwarz inequality in probability,

$$\begin{aligned} \lim_{n \uparrow \infty} \langle X_n \rangle &= \langle X \rangle, \\ \lim_{n \uparrow \infty} \langle Y_n \rangle &= \langle Y \rangle, \\ \lim_{n \uparrow \infty} \langle X_n Y_n \rangle &= \langle XY \rangle. \end{aligned}$$

Hence,

$$(4.2) \quad \begin{aligned} \lim_{n \uparrow \infty} \text{cov}[X_n; Y_n] &= \lim_{n \uparrow \infty} (\langle X_n Y_n \rangle - \langle X_n \rangle \langle Y_n \rangle) \\ &= \langle XY \rangle - \langle X \rangle \langle Y \rangle \\ &= \text{cov}[X; Y]. \end{aligned}$$

Note also that

$$(4.3) \quad \text{cov}[X_n; Y_n] = \sum_{i=1}^n (\langle X_i Y_i \rangle - \langle X_{i-1} Y_{i-1} \rangle)$$

with the notation $X_0 = \langle X \rangle$ and $Y_0 = \langle Y \rangle$, so that $\langle X_n \rangle = X_0$ and $\langle Y_n \rangle = Y_0$. Inequality (2.8) then follows from (4.1), (4.2), (4.3), and

$$(4.4) \quad \langle X_i Y_i \rangle - \langle X_{i-1} Y_{i-1} \rangle \lesssim \left\langle \sup_{a_i} \left| \frac{\partial X}{\partial a_i} \right|^2 \right\rangle^{1/2} \left\langle \sup_{a_i} \left| \frac{\partial Y}{\partial a_i} \right|^2 \right\rangle^{1/2},$$

that we prove now. By our assumption that $\{a_i\}_{i \in \mathbb{N}}$ are i.i.d., we have

$$\begin{aligned} Z_{i-1}(a_1, \dots, a_{i-1}) &= \int Z_i(a_1, \dots, a_{i-1}, a_i'') \beta(da_i''), \\ \langle X_i(a_1, \dots, a_i) Y_i(a_1, \dots, a_i) \rangle \\ &= \left\langle \int X_i(a_1, \dots, a_{i-1}, a_i') Y_i(a_1, \dots, a_{i-1}, a_i') \beta(da_i') \right\rangle, \end{aligned}$$

where β denotes the distribution of a_1 . Hence, we obtain

$$\begin{aligned} \langle X_i Y_i \rangle - \langle X_{i-1} Y_{i-1} \rangle &= \left\langle \int X_i(a_1, \dots, a_{i-1}, a_i') Y_i(a_1, \dots, a_{i-1}, a_i') \beta(da_i') \right\rangle \\ &\quad - \left\langle \int X_i(a_1, \dots, a_{i-1}, a_i') \beta(da_i') \int Y_i(a_1, \dots, a_{i-1}, a_i'') \beta(da_i'') \right\rangle \\ &= \left\langle \int \int \frac{1}{2} (X_i(a_1, \dots, a_{i-1}, a_i') - X_i(a_1, \dots, a_{i-1}, a_i'')) \right. \\ &\quad \left. \times (Y_i(a_1, \dots, a_{i-1}, a_i') - Y_i(a_1, \dots, a_{i-1}, a_i'')) \beta(da_i') \beta(da_i'') \right\rangle \\ &\leq \left\langle \int \int \frac{1}{2} (X_i(a_1, \dots, a_{i-1}, a_i') - X_i(a_1, \dots, a_{i-1}, a_i''))^2 \beta(da_i') \beta(da_i'') \right\rangle^{1/2} \\ &\quad \times \left\langle \int \int \frac{1}{2} (Y_i(a_1, \dots, a_{i-1}, a_i') - Y_i(a_1, \dots, a_{i-1}, a_i''))^2 \beta(da_i') \beta(da_i'') \right\rangle^{1/2}. \end{aligned}$$

We then conclude the proof of (4.4) as in the proof of [6], Lemma 2.3.

4.2. Proof of Lemma 4. We divide the proof in two main parts and deal with $|z| \leq \sqrt{T}$ and $|z| > \sqrt{T}$ separately. The proof relies on the Harnack inequality on graphs. We refer to Zhou [16] for \mathbb{Z}^d , and to Delmotte [3] for other graphs. We recall here the easy part of Harnack's inequality (see [3], Proposition 5.3, or [16], Proof of Theorem 3.3, (3.11)).

LEMMA 6 (Harnack's inequality). *Let $a \in \mathcal{A}_{\alpha\beta}$ and $R \gg 1$. If $g: \mathbb{Z}^d \rightarrow \mathbb{R}^+$ satisfies*

$$(4.5) \quad -\nabla^* \cdot A \nabla g(x) \leq 0$$

in the annulus $\{R/2 < |x| \leq 4R\}$ (i.e., g is a nonnegative subsolution), then

$$(4.6) \quad \sup_{R < |x| \leq 2R} g(x) \lesssim \left(R^{-d} \int_{R/2 < |x| \leq 4R} g(x)^2 dx \right)^{1/2}.$$

Step 1. Proof of (2.10) for $|x - y| \leq \sqrt{T}$.

Since G_T satisfies

$$(4.7) \quad -\nabla_x^* \cdot A \nabla_x G_T(x, y) = -T^{-1} G_T(x, y) \leq 0$$

for $|x - y| \gg 1$, one may apply Lemma 6. For $R \gg 1$, we then have

$$\sup_{x: R < |x-y| \leq 2R} G_T(x, y) \lesssim \left(R^{-d} \int_{R/2 < |x-y| \leq 4R} G_T(x, y)^2 dx \right)^{1/2}.$$

Combined with [6], Lemma 2.8, (2.21), for $q = 2$ (which is uniform in $T > 0$ and $y \in \mathbb{Z}^d$), this yields

$$\sup_{R < |x-y| \leq 2R} G_T(x, y) \lesssim R^{2-d},$$

from which we deduce (2.10) for $\sqrt{T} \geq |x - y| \gg 1$. For $|x - y| \sim 1$, we appeal to [6], Proof of Lemma 2.8, (4.4), with $R \sim 1$ and $q = 1$, which yields $\sup_{|x-y| \leq R} G_T(x, y) \lesssim 1$ by the discrete $L^1 - L^\infty$ estimate.

Step 2. Proof of (2.11) for $|x - y| \leq \sqrt{T}$.

Let N be a positive integer such that $2^N \sim \sqrt{T}$ and $2^{-N} \sqrt{T} \gg 1$. For all $i \in \{1, \dots, N\}$, we first show that

$$(4.8) \quad \begin{aligned} & \left((2^{-i} \sqrt{T})^{-2} \int_{2^{-i-1} \sqrt{T} < |x-y| \leq 2^{-i+2} \sqrt{T}} G_T(x, y)^2 dx \right)^{1/2} \\ & \lesssim i \sim \ln \left(\frac{\sqrt{T}}{1 + 2^{-i} \sqrt{T}} \right). \end{aligned}$$

Estimate (4.8) follows from the triangle inequality and the BMO estimate of [6], Lemma 2.8 (2.20), provided we show that

$$(4.9) \quad \overline{G_T}_{\{|x-y| \leq 2^{-i+2} \sqrt{T}\}} \lesssim i,$$

where $\overline{G_T}_{\{|x-y| \leq 2^{-i+2} \sqrt{T}\}}$ denotes the average of $G_T(x, y)$ on the set $\{|x - y| \leq 2^{-i+2} \sqrt{T}\}$. By the triangle inequality and the BMO estimate of [6], Lemma 2.8, (2.20), we have

$$\begin{aligned} & \overline{G_T}_{\{|x-y| \leq 2^{-i+2} \sqrt{T}\}} \\ & \leq \overline{G_T}_{\{|x-y| \leq 2^{-i+3} \sqrt{T}\}} \\ & \quad + 2 \left(\frac{1}{|\{|x - y| \leq 2^{-i+3} \sqrt{T}\}|} \right. \\ & \quad \left. \times \int_{|x-y| \leq 2^{-i+3} \sqrt{T}} (G_T(x, y) - \overline{G_T}_{\{|x-y| \leq 2^{-i+3} \sqrt{T}\}})^2 dx \right)^{1/2} \\ & \leq \overline{G_T}_{\{|x-y| \leq 2^{-i+3} \sqrt{T}\}} + C, \end{aligned}$$

where C is a universal constant independent of i . Combined with the estimate for $i = 1$

$$\overline{G_T}_{\{|x-y| \leq 4\sqrt{T}\}} \lesssim 1,$$

which is a consequence of [6], Lemma 2.8, (2.22), this implies (4.9) by induction.

We are now in position to prove (2.11) for $|x - y| \leq \sqrt{T}$. Since $x \mapsto G_T(x, y)$ satisfies

$$-\nabla_x^* \cdot A \nabla_x G_T(x, y) = -T^{-1} G_T(x, y) \leq 0$$

in the annulus $\{x, 2^{-i-1}\sqrt{T} < |x - y| \leq 2^{-i+2}\sqrt{T}\}$, Lemma 6 implies

$$\begin{aligned} & \sup_{x: 2^{-i}\sqrt{T} < |x-y| \leq 2^{-i+1}\sqrt{T}} G_T(x, y) \\ & \lesssim \left((2^{-i}\sqrt{T})^{-2} \int_{2^{-i-1}\sqrt{T} < |x-y| \leq 2^{-i+2}\sqrt{T}} G_T(x, y)^2 dx \right)^{1/2} \\ & \lesssim \ln \left(\frac{\sqrt{T}}{1 + 2^{-i}\sqrt{T}} \right) \end{aligned}$$

using (4.8) for $2^{-i}\sqrt{T} \gg 1$. For $|x - y| \leq R \sim 1$, we appeal to (4.9) and to the discrete $L^1 - L^\infty$ estimate

$$G_T(x, y) \leq R^2 \overline{G_T}_{\{|x-y| \leq R\}} \lesssim \ln T.$$

This completes the proof of (2.11) for $|x - y| \leq \sqrt{T}$.

Step 3. Proof of (2.10) and (2.11) for $|x - y| > \sqrt{T}$.

Let $R \geq \sqrt{T}$. Since G_T satisfies

$$-\nabla_x^* \cdot A \nabla_x G_T(x, y) = -T^{-1} G_T(x, y) \leq 0 \quad \text{for } |x - y| \geq 1,$$

Lemma 6 implies

$$\sup_{x: R < |x-y| \leq 2R} G_T(x, y) \lesssim \left(R^{-d} \int_{R/2 < |x-y| \leq 4R} G_T(x, y)^2 dx \right)^{1/2}.$$

Combined with [6], Lemma 2.8, (2.23), for $q = 2$ and $r = k$, that is,

$$\int_{R/2 < |x-y| \leq 4R} G_T(x, y)^2 dx \lesssim R^{d+(2-d)2} (\sqrt{T} R^{-1})^k,$$

this yields the desired pointwise bound.

4.3. *Proof of Lemma 5.* First note that by symmetry

$$\begin{aligned} \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz &= \int_{|z| \geq |z-x|} h_T(z) h_T(z-x) dz \\ &\geq \frac{1}{2} \int_{\mathbb{Z}^d} h_T(z) h_T(z-x) dz. \end{aligned}$$

Hence, it is enough to consider

$$\int_{\mathbb{Z}^d} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx.$$

In this proof, we essentially combine the pointwise decay of g_T with the results of [6], Lemma 2.10, that we recall here for the reader's convenience (see [6], Proof of Lemma 2.10, Steps 1, 2 and 4): There exists $\tilde{R} \sim 1$ such that for all $R \geq \tilde{R}/2$,

$$(4.10) \quad \int_{R < |x| \leq 2R} \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \lesssim \begin{cases} d = 2, & R^2 \max\{1, \ln(\sqrt{T} R^{-1})\}, \\ d > 2, & R^2, \end{cases}$$

$$(4.11) \quad \int_{|x| \leq 4\tilde{R}} \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \lesssim \begin{cases} d = 2, & \ln T, \\ d > 2, & 1. \end{cases}$$

In view of (4.11) and (4.10), it will be convenient to make a dyadic decomposition of space. In order to also benefit from the decay of $g_T(x)$ for $|x| \gg \sqrt{T}$, we make the following decomposition of \mathbb{Z}^d :

$$(4.12) \quad \mathbb{Z}^d = \{|x| \leq 2^{-I} \sqrt{T}\}$$

$$(4.13) \quad \cup \bigsqcup_{i=-I, \dots, -1} \{2^i \sqrt{T} < |x| \leq 2^{i+1} \sqrt{T}\}$$

$$(4.14) \quad \cup \bigsqcup_{i \in \mathbb{N}} \{2^i \sqrt{T} < |x| \leq 2^{i+1} \sqrt{T}\},$$

where I is characterized by $2\tilde{R} < 2^{-I} \sqrt{T} \leq 4\tilde{R}$.

For the integral over the r.h.s. of (4.12), we appeal to (4.11) and to the definitions (2.15) and (2.16) of $g_T(x)$ for $|x| \lesssim 1$:

$$(4.15) \quad \int_{|x| \leq 2^{-I} \sqrt{T}} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \lesssim \begin{cases} d = 2, & \ln^2 T, \\ d > 2, & 1. \end{cases}$$

For the integral over (4.14), we use this time (4.10) for $R \geq \sqrt{T}$ and the definitions (2.15) and (2.16) of $g_T(x)$ for $|x| \geq \sqrt{T}$, so that for all $i \in \mathbb{N}$ we have

$$\begin{aligned} &\int_{2^i \sqrt{T} < |x| \leq 2^{i+1} \sqrt{T}} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \\ &\lesssim (2^i \sqrt{T})^{2-d} (2^i)^{-3} (2^i \sqrt{T})^2 \\ &= \sqrt{T}^{4-d} (2^i)^{1-d}. \end{aligned}$$

Summing this inequality on $i \in \mathbb{N}$ then yields the estimate

$$(4.16) \quad \int_{\sqrt{T} < |x|} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \lesssim \sqrt{T}^{4-d}.$$

We now deal with the integral over the last part (4.13) of \mathbb{Z}^d . To this aim, we combine (4.10) for $R \leq \sqrt{T}$ with the definitions (2.15) and (2.16) of $g_T(x)$ for $|x| \leq \sqrt{T}$. In particular, for all $i \in \{-I, \dots, -1\}$, we have

$$\begin{aligned} & \int_{2^i \sqrt{T} < |x| \leq 2^{i+1} \sqrt{T}} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \\ & \lesssim \begin{cases} d = 2, & \ln(2^{-i}) (2^i \sqrt{T})^2 \ln(2^{-i}) \sim i^2 (2^i \sqrt{T})^2, \\ d > 2, & (2^i \sqrt{T})^2 (2^i \sqrt{T})^{2-d} = (2^i \sqrt{T})^{4-d}. \end{cases} \end{aligned}$$

Summing this inequality over $i \in \{-I, \dots, -1\}$ and using that $2^I \sim \sqrt{T}$ then yield

$$(4.17) \quad \begin{aligned} & \int_{2^{-I} \sqrt{T} < |x| \leq \sqrt{T}} g_T(x) \int_{|z| \leq |z-x|} h_T(z) h_T(z-x) dz dx \\ & \lesssim \begin{cases} d = 2, & 1 + T \sum_{i=-I}^{-1} i^2 4^i, \\ d > 2, & 1 + \sqrt{T}^{4-d} \sum_{i=-I}^{-1} (2^{4-d})^i, \end{cases} \\ & \lesssim \begin{cases} d = 2, & T, \\ d = 3, & \sqrt{T}, \\ d = 4, & \ln T, \\ d > 4, & 1. \end{cases} \end{aligned}$$

The combination of (4.15), (4.16) and (4.17) finally proves (2.17).

REFERENCES

- [1] BOURGEAT, A. and PIATNITSKI, A. (2004). Approximations of effective coefficients in stochastic homogenization. *Ann. Inst. H. Poincaré Probab. Statist.* **40** 153–165. [MR2044813](#)
- [2] CAPUTO, P. and IOFFE, D. (2003). Finite volume approximation of the effective diffusion matrix: The case of independent bond disorder. *Ann. Inst. H. Poincaré Probab. Statist.* **39** 505–525. [MR1978989](#)
- [3] DELMOTTE, T. (1997). Inégalité de Harnack elliptique sur les graphes. *Colloq. Math.* **72** 19–37. [MR1425544](#)
- [4] GLORIA, A. Numerical approximation of effective coefficients in stochastic homogenization of discrete elliptic equations. Preprint.
- [5] GLORIA, A. and MOURRAT, J. C. (2011). Spectral measure and approximation of homogenized coefficients. *Probab. Theory Related Fields*. To appear.
- [6] GLORIA, A. and OTTO, F. (2011). An optimal variance estimate in stochastic homogenization of discrete elliptic equations. *Ann. Probab.* **39** 779–856.
- [7] KLENKE, A. (2006). *Wahrscheinlichkeitstheorie*. Springer, Berlin. English version appeared as: *Probability Theory. A Comprehensive Course* (2008). Springer, London.

- [8] KOZLOV, S. M. (1979). The averaging of random operators. *Mat. Sb. (N.S.)* **109** 188–202, 327. [MR0542557](#)
- [9] KOZLOV, S. M. (1987). Averaging of difference schemes. *Math. USSR Sbornik* **57** 351–369.
- [10] KÜNNEMANN, R. (1983). The diffusion limit for reversible jump processes on \mathbf{Z}^d with ergodic random bond conductivities. *Comm. Math. Phys.* **90** 27–68. [MR0714611](#)
- [11] MOURRAT, J. C. (2011). Variance decay for functionals of the environment viewed by the particle. *Ann. Inst. H. Poincaré Probab. Statist.* **47** 294–327. [MR2779406](#)
- [12] NADDAF, A. and SPENCER, T. (1998). Estimates on the variance of some homogenization problems. Preprint.
- [13] PAPANICOLAOU, G. C. and VARADHAN, S. R. S. (1981). Boundary value problems with rapidly oscillating random coefficients. In *Random Fields, Vol. I, II (Esztergom, 1979). Colloquia Mathematica Societatis János Bolyai* **27** 835–873. North-Holland, Amsterdam. [MR0712714](#)
- [14] YUE, X. and E, W. (2007). The local microscale problem in the multiscale modeling of strongly heterogeneous media: Effects of boundary conditions and cell size. *J. Comput. Phys.* **222** 556–572. [MR2313415](#)
- [15] YURINSKIĬ, V. V. (1986). Averaging of symmetric diffusion in a random medium. *Sibirsk. Mat. Zh.* **27** 167–180, 215. [MR0867870](#)
- [16] ZHOU, X. Y. (1993). Green function estimates and their applications to the intersections of symmetric random walks. *Stochastic Process. Appl.* **48** 31–60. [MR1237167](#)

PROJECT-TEAM SIMPAF
INRIA LILLE-NORD EUROPE
PARC SCIENTIFIQUE DE LA HAUTE BORNE
40, AVENUE HALLEY BAT. A
PARK PLAZA
59650 VILLENEUVE D'ASCQ
FRANCE
E-MAIL: antoine.gloria@inria.fr

MAX PLANCK INSTITUTE
FOR MATHEMATICS IN THE SCIENCES
INSELSTR. 22
04103 LEIPZIG
GERMANY
E-MAIL: otto@mis.mpg.de