



Cop and robber games when the robber can hide and ride

Jérémie Chalopin, Victor Chepoi, Nicolas Nisse, Yann Vaxès

► **To cite this version:**

Jérémie Chalopin, Victor Chepoi, Nicolas Nisse, Yann Vaxès. Cop and robber games when the robber can hide and ride. 8th French Combinatorial Conference, Jun 2010, Orsay, France. 2010. <inria-00482117>

HAL Id: inria-00482117

<https://hal.inria.fr/inria-00482117>

Submitted on 9 May 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cop and robber games when the robber can hide and ride

JÉRÉMIE CHALOPIN¹, VICTOR CHEPOI¹, NICOLAS NISSE², and YANN VAXÈS¹

¹ LIF, Aix-Marseille Université and CNRS,

² MASCOTTE, INRIA, I3S, CNRS,
UNS, Sophia Antipolis, France

In the classical cop and robber game, two players, the cop \mathcal{C} and the robber \mathcal{R} , move alternatively along edges of a finite graph $G = (V, E)$. The cop captures the robber if both players are on the same vertex at the same moment of time. A graph G is called *cop win* if the cop always captures the robber after a finite number of steps. Nowakowski, Winkler (1983) and Quilliot (1983) characterized the cop-win graphs as dismantlable graphs. In this talk, we will characterize in a similar way the class $\mathcal{CWFR}(s, s')$ of cop-win graphs in the game in which the cop and the robber move at different speeds s' and s , $s' \leq s$. We also establish some connections between cop-win graphs for this game with $s' < s$ and Gromov's hyperbolicity. In the particular case $s' = 1$ and $s = 2$, we prove that the class of cop-win graphs is exactly the well-known class of dually chordal graphs. We show that all classes $\mathcal{CWFR}(s, 1)$, $s \geq 3$, coincide and we provide a structural characterization of these graphs. We also investigate several dismantling schemes necessary or sufficient for the cop-win graphs (which we call *k-winnable* and denote by $\mathcal{CWW}(k)$) in the game in which the robber is visible only every k moves for a fixed integer $k > 1$. We characterize the graphs which are *k-winnable* for any value of k .

We present now our main results (for the full version, see [1]). A graph $G = (V, E)$ is (s, s') -*dismantlable* if its vertices can be ordered v_1, \dots, v_n so that for each $v_i, 1 \leq i < n$, there exists v_j with $j > i$, such that $N_s(v_i, G \setminus \{v_j\}) \cap \{v_i, v_{i+1}, \dots, v_n\} \subseteq N_{s'}(v_j)$. A graph G is *dually chordal* if its clique hypergraph is a hypertree. Dually chordal graphs are exactly the graphs $G = (V, E)$ admitting a *maximum neighborhood ordering*.

Theorem 1. *For any $s, s' \in \mathbb{N} \cup \{\infty\}$, $s' \leq s$, a graph $G = (V, E)$ belongs to $\mathcal{CWFR}(s, s')$ iff G is (s, s') -dismantlable.*

Theorem 2. *For a graph $G = (V, E)$, the following conditions are equivalent: (i) $G \in \mathcal{CWFR}(2)$; (ii) G is $(2, 1)$ -dismantlable; (iii) G is dually chordal.*

A graph G is a *big brother graph* if its blocks can be ordered B_1, \dots, B_r such that B_1 is dominated and, for any $i > 1$, the block B_i is a leaf in the block-decomposition of $\cup_{j < i} B_j$ and is dominated by the articulation point connecting B_i to $\cup_{j < i} B_j$. A graph G is a *big two-brother graph* if G can be represented as a union of its subgraphs G_1, \dots, G_r labeled in such a way that G_1 has a dominating vertex, and for any $i > 1$, either the subgraph G_i intersects $\cup_{j < i} G_j$ in two adjacent vertices x_i, y_i belonging to a common subgraph $G_j, j < i$, so that y_i dominates G_i , or G_i has a dominating vertex y_i and intersects $\cup_{j < i} G_j$ in a single vertex x_i (that may coincide with y_i).

Theorem 3. *For a graph $G = (V, E)$ the following conditions are equivalent: (i) $G \in \mathcal{CWFR}(3)$; (i') G is $(3, 1)$ -dismantlable; (ii) $G \in \mathcal{CWFR}(\infty)$; (ii') G is $(\infty, 1)$ -dismantlable; (iii) G is a big brother graph. In particular, the classes of graphs $\mathcal{CWFR}(s)$, $s \geq 3$, coincide.*

Theorem 4. *A graph $G = (V, E)$ is k -winnable for all $k \geq 1$ if and only if G is a big two-brother graph.*

A graph $G = (V, E)$ is *k-bidismantlable* if its vertices can be ordered v_1, \dots, v_n in such a way that for each vertex $v_i, 1 \leq i < m$, there exist two adjacent or coinciding vertices x, y with $y = v_j, x = v_\ell$ and $j, \ell > i$ such that $N_k(v_i, G \setminus \{x, y\}) \cap X_i \subseteq N_1(y)$, where $X_i := \{v_i, v_{i+1}, \dots, v_m\}$.

Theorem 5. *Any graph $G = (V, E)$ of $\mathcal{CWW}(2)$ is 2-bidismantlable, however there exist 2-bidismantlable graphs G with $G \notin \mathcal{CWW}(2)$. For any odd integer $k \geq 3$, if a graph G is k -bidismantlable, then $G \in \mathcal{CWW}(k)$.*

[1] J. Chalopin, V. Chepoi, N. Nisse, Y. Vaxès, Cop and robber games when the robber can hide and ride, arXiv:1001.4457.