

# LES DIMENSIONS FRACTALS : MYTHES ET RÉALITÉS ÉCOLOGIQUES

Nicolas Bez

Institut de Recherche pour le Développement (IRD), Centre de Recherche Halieutique Méditerranéenne et Tropicale, Avenue Jean Monnet, BP 171, 34203 Sète Cedex, France

Mots clefs : Biostatistique, Statistique Spatiale

## Français

La notion de fractal proposée par Mandelbrot a séduit la communauté écologique pour les notions d'indépendance d'échelle et de transfert d'échelle qu'elle recèle. L'objectif de ce travail est de préciser qu'en réalité, la notion de fractal cache deux concepts différents de rugosité d'un part et d'auto-homothétie d'autre part. Les cas où ces notions se rejoignent sont très particuliers (processus Gaussiens) et rarement rencontrés dans la nature. En s'appuyant sur la définition rigoureuse de la dimension fractal qui repose sur la dimension de Hausdorff-Besicovitch, on montre que la dimension fractal quantifie une propriété locale des processus à savoir la rugosité. De plus, lorsque son estimation est basée sur le variogramme, on montre que la formule retenue par la bibliographie écologique qui utilise la demi-pente du log-variogramme est erronée. L'utilisation de la pente du log-madogramme est recommandée. La différence est illustrée à l'aide de données réelles.

La dimension d'auto-homothétie doit être distinguée de la dimension fractal. On montre des cas où la dimension fractal est non entière alors que le processus n'est pas auto-homothétique. Il est donc important, dans la pratique, de ne pas confondre ces deux notions souvent cachées derrière le même vocable de dimension fractal.

Enfin, parce que précisément la dimension fractal peut être associée à des indépendances d'échelles, il importe de préciser l'effet du support d'information, qui ici, comme dans l'ensemble des statistiques spatiales existe.


## Anglais

Fractals have been widely used in ecology. Among other, their scale independence seems interesting for describing scale transfer in ecology. We show here that the original definition based on the Hausdorff-Besicovitch dimension (DHB), as well as the estimators based on Minkowski-Boulingand dimension (DMB) actually quantify a local property called roughness. As a consequence, the scale independence property of fractals, which is very attractive and powerful in ecological modelling, is left out by the mathematical definition which refers to a local property, the roughness. The primary objective of the present paper is to clarify the dual aspect of fractals and to show that those two properties may be quantified by a single dimension (DHB) only in some specific cases. We show that for natural objects in ecology, a non-integer DHB quantifies per se only the roughness and self-similarity needs to be evidenced or postulated by other means. Second, we revisit some aspects of the practical estimation of DHB. We show that when geostatistics is used, the formula based on the slope at the origin of the variogram, widely found in the literature, indeed only holds for Gaussian situations. We recommend the use of a more general formula based on the first order variogram (madogram) which holds for any type of variable distribution. We discuss then other aspects related to the estimation of DHB on non self-similar objects, its simplification for 2D fields, and its possible relationship with self-similarity. In resonance with earlier works, we point out the problem of scale and resolution when estimating a fractal dimension. Some of our considerations and developments are illustrated with field data recorded during a scientific acoustic survey on the North Sea herring. I conclude on a synthesis of practical

recommendations and warnings to ecologists when using fractal dimension..

## **Bibliographie**

- [1] Bruno, R., and G., Raspa, 1989. Geostatistical characterization of fractal models of surfaces. Kluwer academics publisher, M.Armstrong (Ed.), Geostatistics, Vol. 1, 77-89.
- [2] Chilès J.P. and P. Delfiner, 1999. Geostatistics: Modeling spatial uncertainty. J. Wiley & Sons, New York, 695 p.
- [3] Dubuc B., S.W., Zucker, C. Tricot, J.F., Quiniou and D., Wehbi, 1989. Evaluating the fractal dimension of surfaces. Proc. R. Soc. Lond. A 425, 113-127
- [4] Dungan, J. L., J. N. Perry, M. R. T. Dale, P. Legendre, S. Citron-Pousty, M.-J. Fortin, A. Jakomulska, M. Miriti, and M. S. Rosenberg. 2002. A balanced view of scale in spatial statistical analysis. *Ecography* 25:626-640.
- [5] Falconer, K.L. 1990. Fractal geometry. Mathematical foundations and applications. J. Wiley & Sons, New York.
- [6] Fortin, M.-J., T. H. Keitt, B. A. Maurer, M. L. Taper, D. M. Kaufman, and T. M. Blackburn. 2005. Species' geographic ranges and distributional limits: pattern analysis and statistical issues. *OIKOS* 108:7-17.
- [7] Halley, J.M., S., Hartley, A.S., Kallimanis, W.E., Kunin, J.J., Lennon, and S.P., Sgardelis, 2004. Uses and abuses of fractal methodology in ecology. *Ecology Letters*, 7, 254-271.
- [8] Leduc, A., Y.T., Prairie, and Y., Bergeron, 1994. Fractal dimension estimates of a fragmented landscape: source of variability. *Landscape Ecology*, vol. 9, no 4, 279-286.
- [9] Mandelbrot, B.B., 1975. Les objets fractals: forme, hasard et dimension. Paris & Montréal, Flammarion.
- [10] Mandelbrot, B.B. 1977. Fractals: forms, chance and dimension. Freeman, California, USA, 365 pp.
- [11] Matheron, G. 1963. Principles of geostatistics. *Economic Geology*, 58, n°8, 1246-1266
- [11] Matheron, G. 1973. The intrinsic random functions and their applications. *Adv. Appl. Prob.*, Vol. 11, n°1, 184-189.
- [12] Seuront, L. 1998. Fractals and multifractals: new tools to characterize space-time heterogeneity in marine ecology. *Océanis* 24: 123-158.
- [13] Stoyan, D., and H. Stoyan, 1994. Fractals, random shapes and point fields; methods of geometrical statistics. John Wiley sons, Chichester, England, 389 pp.
- [14] Sugihara, G., and R. M. May. 1990. Applications of fractals in ecology. *Trends in Ecology and Evolution* 5:79-86.
- [15] Taylor C.C., and S.J. Taylor, 1991. Estimating the dimension of a fractal. *Journal of the Royal Statistical Society, Series B (Methodology)*, Vol. 53, No 2, 353-364.
- [16] Tricot C., 1982. Two definitions of fractional dimension. *Math. Proc. Camb. Phil. Soc.*, 91, 57-74.
- [17] Tricot C., J.F., Quiniou, D., Wehbi, C., Roques-Carmes et B., Dubuc, 1988. Evaluation de la dimension fractale d'un graphe. *Revue Phys. Appl.* 23, 111-124.
- [18] Turchin, P., 1996. Fractal analyses of animal movement: a critique. *Ecology*, 77(7), 2086-2090.
- [19] Viswanathan, G. M., V. Afanasyev, S. V. Buldyrev, E. J. Murphy, P. A. Prince, and H. E. Stanley. 1996. Lévy flight search patterns of wandering albatrosses. *Nature* 381:413-415.



Field acoustic data collected along one scientific survey transect in the North Sea. These data were available at the smallest possible resolution (i.e. one ping  $\sim 3$ m) up to any concatenation of pings: 10 pings  $\sim 30$  m, 100 pings  $\sim 300$  m, and 1000 pings  $\sim 3000$  m. They should be considered as exhaustive along the survey transect. For each of the four resolutions, we computed the empirical madogram (right panel) and variogram (left panel) and considered their log-log representations together with their linear regression for small distances. The slopes at the origin of the log-variogram are not twice the slopes of the log-madogram. In addition, they vary with the support indicating a modification of the fractal dimension with the support.