

## On removable edges in 3-connected cubic graphs

Jean-Luc Fouquet, Henri Thuillier

► **To cite this version:**

Jean-Luc Fouquet, Henri Thuillier. On removable edges in 3-connected cubic graphs. *Discrete Mathematics*, Elsevier, 2012, Article in press, pp.9. <10.1016/j.disc.2011.11.025>. <inria-00516060>

**HAL Id: inria-00516060**

**<https://hal.inria.fr/inria-00516060>**

Submitted on 8 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON REMOVABLE EDGES IN 3-CONNECTED CUBIC GRAPHS

J.L. FOUQUET AND H. THULLIER

ABSTRACT. A removable edge in a 3-connected cubic graph  $G$  is an edge  $e = uv$  such that the cubic graph obtained from  $G \setminus \{u, v\}$  by adding an edge between the two neighbours of  $u$  distinct from  $v$  and an edge between the two neighbours of  $v$  distinct from  $u$  is still 3-connected. Li and Wu [3] showed that a spanning tree in a 3-connected cubic graph avoids at least two removable edges, and Kang, Li and Wu [4] showed that a spanning tree contains at least two removable edges. We show here how to obtain these results easily from the structure of the sets of non removable edges and we give a characterization of the extremal graphs for these two results.

## 1. INTRODUCTION

In 1961 Tutte [5] gave a structural characterization for 3-connected graphs by using the existence of contractible or removable edges. A *cubic graph* is a simple 3-regular graph. From now on, all graphs considered here are cubic graphs. An edge  $e$  of a 3-connected cubic graph  $G$  is said to be *removable* when the cubic graph obtained from  $G$  by the following operations remains to be 3-connected.

- Delete  $u$  and  $v$  from  $V(G)$  and their incident edges from  $E(G)$
- Add one edge between the two neighbours of  $u$  distinct from  $v$  as well as between the two neighbours of  $v$  distinct from  $u$

An edge which is not removable is said to be *non removable*. The set of removable edges of  $G$  is denoted by  $R(G)$  and the set of non removable edges is denoted by  $N(G)$ .

Conversely, we can get a new 3-connected cubic graph from a 3-connected cubic graph  $G$  by *inserting* one edge between two existing edges. More formally, let  $uv$  and  $u'v'$  be two edges of a 3-connected cubic graph  $G$ , we get a new 3-connected cubic graph  $G'$  when the three following operations are performed.

- Delete  $uv$  and  $u'v'$  from  $E(G)$
- Add two new adjacent vertices  $x$  and  $y$  to  $V(G)$
- Join  $x$  to  $u$  and  $v$  and  $y$  to  $u'$  and  $v'$ .

We shall say that we have proceeded to the *insertion* (of the edge  $xy$ ). Obviously the new edge  $xy$  is removable in the obtained graph.

Li and Wu [3] showed that a spanning tree in a 3-connected cubic graph avoids at least two removable edges:

**Theorem 1.1.** [3] *Let  $G$  be a 3-connected cubic graph with at least six vertices. Then every spanning tree of  $G$  avoids at least two removable edges.*

---

1991 *Mathematics Subject Classification.* 035 C.

*Key words and phrases.* Cubic graphs, Removable edges, spanning trees.

Kang, Li and Wu [4] showed that a spanning tree contains at least two removable edges:

**Theorem 1.2.** [4] *Let  $G$  be a 3-connected cubic graph with at least six vertices. Then every spanning tree of  $G$  contains at least two removable edges.*

We shall show in Section 3 how to obtain these results easily from the structure of the set of non removable edges (Corollaries 3.4 and 3.7) and we give a characterization of the extremal graphs for these two theorems. More precisely, we shall exhibit two infinite families of 3-connected cubic graphs, the *PR*-graphs and the *3T*-graphs (defined below in Subsection 1.2) and we shall prove that a 3-connected cubic graph having a spanning tree avoiding exactly two removable edges is a *PR*-graph (Corollary 3.6), and that a 3-connected cubic graph having a spanning tree containing exactly two removable edges is a *3T*-graph (Corollary 3.8).

**1.1. Edge cut.** Let  $\{V_1, V_2\}$  be a partition of the vertex set  $V(G)$  of  $G$ . The set  $F$  of edges joining  $V_1$  to  $V_2$  denoted by  $(V_1, V_2)$  is an *edge cut* and the partition  $\{V_1, V_2\}$  of  $V(G)$  is the *associated partition*. An edge cut  $F$  of  $k$  edges is a *k-edge cut*. An edge cut  $F$  is *minimal* if no proper subset of  $F$  is an edge cut, it is *trivial* if it is minimal and one component of  $G \setminus F$  is a single vertex.

Obviously, a 3-connected cubic graph has no 2-edge cut. Moreover, any non trivial 3-edge cut  $F$  is a matching of three edges and the edges of this edge cut are contained in  $N(G)$  (non removable edges). By deleting the edges of  $F$ , we get two connected graphs (the subgraphs  $G[V_1]$  and  $G[V_2]$  of  $G$  induced respectively by  $V_1$  and  $V_2$ ) and we remark that these two subgraphs are 2-connected. By contracting  $G[V_2]$  in a single new vertex  $u$  and  $G[V_1]$  in a single new vertex  $v$ , we get two smaller 3-connected cubic graphs  $G_1$  and  $G_2$ . Conversely, let  $G_1$  and  $G_2$  be two 3-connected cubic graphs and  $u \in V(G_1)$ ,  $v \in V(G_2)$  with  $N_u = \{u_1, u_2, u_3\}$  and  $N_v = \{v_1, v_2, v_3\}$ . We construct a new 3-connected cubic graph  $G$  where  $V(G) = (V(G_1) \setminus \{u\}) \cup (V(G_2) \setminus \{v\})$  and  $E(G) = (E(G_1) \setminus \{uu_1, uu_2, uu_3\}) \cup (E(G_2) \setminus \{vv_1, vv_2, vv_3\}) \cup \{u_1v_1, u_2v_2, u_3v_3\}$  having  $\{u_1v_1, u_2v_2, u_3v_3\}$  as a non trivial 3-edge cut (note that  $G$  may contain other non trivial 3-edge cuts).

## 1.2. Two special families of 3-connected cubic graphs.

**1.2.1. The family of *PR*-graphs.** Let  $PR_{0,0}$  be the 3-connected cubic graph on six vertices formed by two triangles joined by a matching of three edges. Let us remark that these three edges are not removable. Starting from  $PR_{0,0}$  we proceed to successive insertions between edges of non trivial 3-edge cuts or insertions of claws (by adding three vertices of degree 2 on the edges of a non trivial 3-edge cut and joining these 3 vertices to a fourth vertex). To proceed to an insertion of an edge, we choose two edges of a 3-edge cut  $F$  and we insert an edge between these two chosen edges. To proceed to an insertion of a claw, we proceed first to the insertion of an edge as previously (let  $xy$  be the new edge obtained) and we insert a new edge between  $xy$  and the last edge of the considered 3-edge cut  $F$ . Let  $k_1$  and  $k_2$  be two integers such that  $k_1 \geq 0, k_2 \geq 0$  and  $k_1 + k_2 \geq 1$ . A cubic graph obtained from  $PR_{0,0}$  by  $k_1$  insertions of edges and  $k_2$  insertions of claws is said to be a *graph of type  $PR_{k_1, k_2}$*  (or simply, a  $PR_{k_1, k_2}$ ). More precisely, a graph of type  $PR_{k_1+1, k_2}$  is obtained from a  $PR_{k_1, k_2}$  by insertion of an edge and a graph of type  $PR_{k_1, k_2+1}$  is obtained from a  $PR_{k_1, k_2}$  by insertion of a claw. It must be clear that given  $k_1$  and  $k_2$ , we may obtain several non isomorphic cubic graphs of type

$PR_{k_1, k_2}$ . Since the operation of insertion of an edge preserves the 3-connectivity, it is easy to see that a  $PR_{k_1, k_2}$  is a 3-connected cubic graph. A  $PR$ -graph is a graph  $G$  such that there exist integers  $k_1$  and  $k_2$  and  $G$  is of type  $PR_{k_1, k_2}$ . In Figure 1, we give example of a graph of type  $PR_{2,1}$  and a graph of type  $PR_{1,2}$ .




FIGURE 1. Type  $PR_{2,1}$  and Type  $PR_{1,2}$

It can be easily verified that the only non removable edges of a graph of type  $PR_{k_1, k_2}$  are the edges of the three disjoint paths  $P_1, P_2$  and  $P_3$  joining the two triangles (drawn in bold in Figure 1). Then a graph  $G$  of type  $PR_{k_1, k_2}$  has  $n = 2k_1 + 4k_2 + 6$  vertices and it verifies  $|R(G)| = k_1 + 3k_2 + 6$  and  $|N(G)| = 2k_1 + 3k_2 + 3$ .

1.2.2. *The family of 3T-graphs.* A *fundamental*  $3T_{k+2}$  (with  $k \geq 0$ ) is a cubic graph obtained from three isomorphic trees  $T_1, T_2$  and  $T_3$  of maximum degree 3 and no vertex of degree 2 with  $k + 2$  vertices of degree one and  $k$  vertices of degree three each. Each triple of pendent vertices (one in each tree) mapped by the isomorphism are joined by a triangle. It must be clear that a fundamental  $3T_{k+2}$  is 3-connected.

A  $p$ -extended  $3T_{k+2}$  is a cubic graph obtained from a fundamental  $3T_{k+2}$  by insertion of  $p$  edges. The family of  $p$ -extended  $3T_{k+2}$  shall be denoted by  $3T_{k+2, p}$ . As above, to proceed to the insertion of an edge, we choose a non trivial 3-edge cut  $F$  and two distinct edges of  $F$  in the graph in construction. Note that given  $k \geq 0$  and  $p \geq 2$ , the family  $3T_{k+2, p}$  may contain several non isomorphic cubic graphs. Since the operation of insertion of an edge preserves the 3-connectivity, a  $p$ -extended  $3T_{k+2}$  is a 3-connected cubic graph. In Figure 2 we give a fundamental  $3T_4$  and in Figure 3 a 3-extended  $3T_4$ .

A fundamental  $3T_{k+2}$  can be seen as a 0-extended  $3T_{k+2}$ . A  $3T$ -graph is a graph that belongs to the union  $\bigcup_{k \geq 0, p \geq 0} 3T_{k+2, p}$ .




FIGURE 2. A fundamental  $3T_4$



also an edge of  $F$  ( $|F \cap F'| = 1$ ). It is easy to see that, one of the sets  $V_1'$  or  $V_2'$  (say  $V_1'$ ) contains the whole set  $V_1$ , so  $V_1 \cap V_2' = \emptyset$ . We let  $A = V_1 \cap V_1'$  (hence  $A = V_1$ ),  $B = V_2 \cap V_1'$  and  $C = V_2 \cap V_2'$  (hence  $C = V_2'$ ) and we can check that the first item is verified when  $F \cap F' = \emptyset$  while the second item is verified when  $|F \cap F'| = 1$ .  $\square$

An *end 3-edge cut* is a 3-edge cut such that every edge of the subgraph induced on one of the two sets of the associated partition is removable. This subgraph without any non removable edge will be called an *extremity* (it may happens that the two sets of the associated partition are extremities). Let us remark that an extremity of a 3-connected cubic graph  $G$  is a 2-connected induced subgraph of  $G$ .

**Lemma 2.4.** *Each set of the associated partition of any 3-edge cut  $F$  contains an extremity.*

**Proof** If every edge of  $G[V_2]$  is a removable edge then  $V_2$  is an extremity. If  $G[V_2]$  contains a non removable edge  $e$  then let  $F'$  be a 3-edge cut containing  $e$ . Clearly,  $F'$  is distinct from  $F$ . By Lemma 2.3, we have the partition  $A = V_1 \cap V_1' = V_1$ ,  $B = V_2 \cap V_1'$  and  $C = V_2 \cap V_2' = V_2'$ . We have thus obtained a refining of the partition  $\{V_1, V_2\}$ . If every edge of  $G[V_2']$  is removable then  $V_2'$  is an extremity, otherwise we can proceed to a new refinement of  $V_2'$ . Since the number of 3-edge cuts is finite, we shall be left with an extremity in  $V_2$ . The same holds for  $V_1$  and the Lemma follows.  $\square$

**Lemma 2.5.** *Let  $P = u_1u_2 \dots u_k$  ( $k \geq 3$ ) be a path contained in  $N(G)$  and let  $F$  be a 3-edge cut of  $G$ . Then  $F$  has at most one edge in  $P$ .*

**Proof** Assume to the contrary that there exists a 3-edge cut  $F$  containing two edges of  $P$ ,  $u_iu_{i+1}$  and  $u_ju_{j+1}$  ( $i \neq j$ ,  $1 \leq i \leq k-2$ ,  $i+1 \leq j \leq k-1$ ). Since  $F$  is a matching, the edge  $u_{i+1}u_{i+2}$  is distinct from  $u_ju_{j+1}$ . Assume moreover that the subpath  $P' = u_{i+1}u_{i+2} \dots u_j$  of  $P$  does not contained the third edge of  $F$ . We can suppose that  $F$  has been chosen in such a way that the distance on  $P$  between  $u_iu_{i+1}$  and  $u_ju_{j+1}$  is as short as possible.

Let  $F'$  be a 3-edge cut containing  $u_{i+1}u_{i+2}$ . The choice of  $F$  forces  $F'$  to have no other edge between  $u_iu_{i+1}$  and  $u_ju_{j+1}$ . We consider that  $u_{i+1}$  and  $u_j$  are in  $V_1$  (hence,  $P'$  is a path in  $G[V_1]$  and  $u_i$  and  $u_{j+1}$  are in  $V_2$ ). Let  $Q$  be a path in  $G[V_2]$  joining  $u_i$  to  $u_{j+1}$  and consider the cycle obtained by concatenation of  $u_iu_{i+1}$ ,  $P'$ ,  $u_ju_{j+1}$  and  $Q$ . By Lemma 2.1, this cycle contains an edge  $e$  of  $F'$  distinct from  $u_{i+1}u_{i+2}$ . By the choice of  $F$ , this edge  $e$  must be on  $Q$ . We do not know the exact position of the third edge of  $F'$ , but we are certain that at least one of the two 2-connected subgraphs  $G[V_1]$  or  $G[V_2]$  contains exactly one edge of  $F'$ . Hence  $G[V_1]$  or  $G[V_2]$  has an isthmus, a contradiction.  $\square$

**Lemma 2.6.** *Let  $P = u_1u_2 \dots u_k$  ( $k \geq 3$ ) be a path contained in  $N(G)$ . Then  $P$  is an induced path of  $G$ .*

**Proof** Assume to the contrary that  $u_iu_j$  is an edge of  $G$  ( $i \neq j$ ,  $1 \leq i \leq k-1$ ,  $i+2 \leq j \leq k$ ). Then the concatenation of the subpath  $P'$  of  $P$  with ends  $u_i$  and  $u_j$  together with the edge  $u_iu_j$  gives a cycle of  $G$ . This cycle intersects a 3-cut edge

containing the edge  $u_i u_{i+1}$ . By Lemma 2.1, a second edge of this 3-edge cut must be contained in  $P$ , a contradiction with Lemma 2.5.  $\square$

### 3. ON THE SET OF NON REMOVABLE EDGES

**Theorem 3.1.** [1] *The subgraph of a 3-connected cubic graph  $G$  induced by the set  $N(G)$  of non removable edges is an induced forest with at least three trees. Each 3-edge cut intersects three distinct trees of that forest.*

**Proof** Assume that the edge-induced subgraph on  $N(G)$  (denoted also  $N(G)$ ) contains a cycle  $C$  and let  $e \in C$ . By Lemma 2.1, any 3-edge cut containing  $e$  must intersect  $C$  at least twice. Then two edges of this 3-edge cut are contained in a path  $P$  of  $N(G)$ , a contradiction with Lemma 2.5. Hence,  $N(G)$  is a forest as claimed and, by Lemma 2.6, it is clear that this forest is an induced forest.

Let  $F$  be a 3-edge cut. If two edges of  $F$  are contained in the same tree of  $N(G)$  then we can find a path contained in  $N(G)$  joining these two edges, again a contradiction with Lemma 2.5. The theorem follows.  $\square$

*Remark 3.2.* Since  $N(G)$  has at most  $n - 3$  edges (with  $n = |V(G)|$ ), the graph  $G$  contains at least  $\frac{n+6}{2}$  removable edges. By Remark 1.3, we see that the 3T-graphs are extremal for these numbers. More precisely, we have proved in [2] that the family of 3T-graphs is exactly the family of 3-connected cubic graphs having the minimum number of removable edges.

**Corollary 3.3.** *Let  $G$  be a 3-connected cubic graph and let  $C$  be  $C = u_0 u_1 \dots u_k u_0$  be a cycle of  $G$ . Then  $C$  contains at least two removable edges.*

**Proof** Since by Theorem 3.1  $N(G)$  is a forest,  $C$  contains at least one removable edge. Assume that  $C$  contains only one removable edge. Let  $P$  be the path obtained from  $C$  by deleting this edge and let  $e$  be an edge of  $P$ . Since  $P$  is contained in  $N(G)$  there is a 3-edge cut  $F$  containing  $e$ . By Lemma 2.1,  $F$  contains exactly one other edge of  $F$ , a contradiction with Lemma 2.5.  $\square$

**Corollary 3.4.** [3] *Let  $G$  be a 3-connected cubic graph with at least six vertices. Then every spanning tree of  $G$  avoids at least two removable edges.*

**Proof** Let  $n$  be the number of vertices of  $G$ . Since  $G$  has  $3\frac{n}{2}$  edges, a spanning tree avoids  $\frac{n+2}{2} \geq 4$  edges. If every edge of  $G$  is removable, the result is immediate.

Now, assume that  $N(G) \neq \emptyset$ . By Lemma 2.4,  $G$  contains at least two extremities. Let  $F$  be an end 3-edge cut with associated partition  $\{V_1, V_2\}$  such that  $G[V_1]$  is an extremity. The subgraph  $G[V_1]$  contains  $2p + 1$  vertices (with  $p \geq 1$ ) and  $3p$  edges. The trace  $T_1$  on  $G[V_1]$  of a spanning tree  $T$  of  $G$  is a spanning forest of this extremity having  $k$  trees (with  $1 \leq k \leq 3$ ). Hence,  $T_1$  has  $2p - k + 1$  edges and avoids  $p + k - 1 \geq p$  edges of  $G[V_1]$ . Thus,  $T$  must avoid at least one edge in each extremity and the theorem follows.  $\square$

**Lemma 3.5.** *Let  $G$  be a 3-connected cubic graph with at least six vertices having a spanning tree avoiding exactly two removable edges. Then  $G$  has exactly two extremities and these extremities are isomorphic to a triangle.*

**Proof** Let  $n$  be the number of vertices of  $G$ . Since we know that there are  $\frac{n+2}{2} \geq 4$  edges outside any spanning tree, if a spanning tree  $T$  avoids exactly two removable edges then  $N(G)$  is not empty. By Lemma 2.4,  $G$  has  $k \geq 2$  extremities  $H_1, H_2, \dots, H_k$ . We have seen in the proof of Corollary 3.4 that if  $H_i$  ( $i = 1, \dots, k$ ) is an extremity having  $2p_i + 1$  vertices then a spanning tree  $T$  of  $G$  avoids at least  $p_i \geq 1$  edges of  $H_i$ . Hence,  $T$  must avoid at least  $p_1 + p_2 + \dots + p_k$  removable edges. Since  $T$  avoids exactly two removable edges,  $p_1 + p_2 + \dots + p_k = 2$ . Hence  $k = 2$  and  $p_1 = p_2 = 1$ , that is the graph  $G$  has exactly two extremities and each extremity has three vertices.  $\square$

**Corollary 3.6.** *Let  $G$  be a 3-connected cubic graph. Then  $G$  has a spanning tree  $T$  avoiding exactly two removable edges if and only if  $G$  is a PR-graph.*

**Proof** Assume that  $G$  is isomorphic to some 3-connected cubic graph of type  $PR_{k_1, k_2}$  ( $k_1 + k_2 \geq 0$ ). Let  $M$  be the set of edges involved in the insertions operated from  $PR_{0,0}$  in order to obtain  $G$ . Assume that the two triangles are  $a, b, c$  and  $a', b', c'$ . Let  $M' = M \cup \{ab, bc, a'b', b'c'\}$ . We can easily find a spanning tree  $T$  containing the edges of  $M'$  (perform the greedy Kruskal's algorithm to find a minimum spanning tree of  $G$  when the edges of  $M'$  are placed at the beginning of the ordering of  $E(G)$ ). Since the removable edges of  $G$  are the edges of  $M$  and the six edges contained in the two triangles, exactly two removable edges are outside this spanning tree.

We prove now by induction on the number of vertices  $n \geq 6$ , that whenever  $G$  is a 3-connected cubic graph having a spanning tree avoiding exactly two removable edges then  $G$  is isomorphic to some graph of type  $PR_{k_1, k_2}$  ( $k_1 + k_2 \geq 0$ ).

When  $n = 6$ ,  $PR_{0,0}$  is the only graph with that property. Assume that the result holds for any 3-connected cubic graph with  $6 \leq n' < n$  vertices having a spanning tree avoiding exactly two removable edges.

Let  $G$  be a 3-connected cubic graph with  $n$  vertices having a spanning tree avoiding exactly two removable edges. By lemma 3.5,  $G$  has exactly two extremities isomorphic to a triangle. Assume that these triangles are  $\Delta_1$  and  $\Delta_2$ . If  $T$  is a spanning tree of  $G$  avoiding exactly two removable edges, one of this edge (say  $e_1$ ) must be in  $\Delta_1$  and the other (say  $e_2$ ) is in  $\Delta_2$ .

When there is no 3-edge cut distinct from the 3-edge cut incident to  $\Delta_1$  or to  $\Delta_2$ , it is not difficult to see that  $G$  is isomorphic to  $PR_{0,0}$  or to  $PR_{1,0}$  or to  $PR_{0,1}$  (see  $PR_{0,1}$  in Figure 4).




FIGURE 4.  $PR_{0,1}$

Let  $F = \{x_1x_2, y_1y_2, z_1z_2\}$  be a 3-edge cut of  $G$  distinct from the end 3-edge cuts  $F_1$  and  $F_2$  respectively incident to  $\Delta_1$  and to  $\Delta_2$ . Let  $\{V_1, V_2\}$  be the associated partition of  $F$ . We can construct a new 3-connected cubic graph  $G_1$  by replacing



in  $G$  the subgraph  $G[V_2]$  by the triangle  $\Delta'_1 = \{x_2, y_2, z_2\}$ . In the same way, we construct  $G_2$  by replacing  $G[V_1]$  by the triangle  $\Delta'_2 = \{x_1, y_1, z_1\}$ . Clearly, for  $i = 1, 2$   $R(G_i) = (R(G) \cap E(G[V_i])) \cup E(\Delta'_i)$ .

Let  $U_i$  be the trace of the spanning tree  $T$  on  $G[V_i]$  ( $i = 1, 2$ ). Note that  $U_i$  is a spanning forest of  $G[V_i]$  having at most three trees and that  $U_i$  avoids exactly one removable edge in  $E(G[V_i])$  (the edge  $e_i$  in  $\Delta_i$ ). By using the trace  $U_i$  we will construct a spanning tree  $T_i$  of  $G_i$  avoiding exactly two removable edges in  $G_i$ .

Following the number of edges of  $F$  in  $E(T)$  there are three cases :

**Case 1 :**  $|E(T) \cap F| = 1$ . Assume that  $E(T) \cap F = \{x_1x_2\}$ . We see that  $U_1$  and  $U_2$  are trees. Hence,  $T_1 = U_1 + \{x_1x_2, x_2y_2, x_2z_2\}$  is a spanning tree of  $G_1$  and  $T_2 = U_2 + \{x_1x_2, x_1y_1, x_1z_1\}$  is a spanning tree of  $G_2$ .

**Case 2 :**  $|E(T) \cap F| = 2$ . Assume that  $E(T) \cap F = \{x_1x_2, y_1y_2\}$ . Consider the unique path  $P$  in  $T$  connecting  $x_1x_2$  to  $y_1y_2$ . Then either  $P$  is a subpath of  $G[V_1]$  having  $x_1$  and  $y_1$  as end vertices or  $P$  is a subpath of  $G[V_2]$  having  $x_2$  and  $y_2$  as end vertices. If  $P$  is a subpath of  $G[V_1]$  then  $U_1$  is a tree and there is no path in  $U_2$  connecting  $x_2$  to  $y_2$ . Then  $U_2$  is a forest of two trees, one of them containing  $x_2$  and the other containing  $y_2$ . We see that  $T_1 = U_1 + \{x_1x_2, x_2y_2, x_2z_2\}$  and  $T_2 = U_2 + \{x_1x_2, y_1y_2, x_1y_1, x_1z_1\}$  are respectively spanning trees of  $G_1$  and  $G_2$ . Analogously, if  $P$  is a subpath of  $G[V_2]$  then  $U_2$  is a tree and  $U_1$  is a forest of two trees, one of them containing  $x_1$  and the other containing  $y_1$ . Hence,  $T_1 = U_1 + \{x_1x_2, y_1y_2, x_2y_2, x_2z_2\}$  and  $T_2 = U_2 + \{x_1x_2, x_1y_1, x_1z_1\}$  are spanning trees of  $G_1$  and  $G_2$ .

**Case 3 :**  $F \subset E(T)$ . Up to symmetries, there are two subcases:

*Subcase 3.1 :*  $U_1$  is a tree and  $U_2$  is a forest of three trees (the first containing  $x_2$ , the second containing  $y_2$  and the third containing  $z_2$ ). We consider  $T_1 = U_1 + \{x_1x_2, x_2y_2, x_2z_2\}$  and  $T_2 = U_2 + \{x_1x_2, y_1y_2, z_1z_2, x_1y_1, x_1z_1\}$ .

*Subcase 3.2 :*  $U_1$  is a forest of two trees (one of them containing  $x_1$  and the other containing  $y_1$  and  $z_1$ ) and  $U_2$  is a forest of two trees (one of them containing  $x_2$  and  $z_2$  and the other containing  $y_2$ ). We consider  $T_1 = U_1 + \{x_1x_2, y_1y_2, x_2y_2, x_2z_2\}$  and  $T_2 = U_2 + \{x_1x_2, y_1y_2, x_1y_1, x_1z_1\}$ .

In every case, we have constructed a spanning tree  $T_1$  of  $G_1$  (respectively  $T_2$  of  $G_2$ ) avoiding exactly two removable edges in  $G_1$  (respectively  $G_2$ ), the edges  $e_1$  and  $y_2z_2$  (resp.  $e_2$  and  $y_1z_1$ ).

By the induction hypothesis,  $G_1$  is isomorphic to a graph of type  $PR_{p_1, q_1}$  and  $G_2$  is isomorphic to a graph of type  $PR_{p_2, q_2}$ . At last,  $G$  itself is isomorphic to a graph of type  $PR_{p_1+p_2, q_1+q_2}$ .  $\square$

**Corollary 3.7.** [4] *Let  $G$  be a 3-connected cubic graph with at least six vertices. Then every spanning tree contains at least two removable edges.*

**Proof** A spanning tree  $T$  of  $G$  containing at most one removable edge  $e$  contains only edges in  $N(G) \cup \{e\}$ . Since  $N(G)$  has at most  $n - 3$  edges, this is impossible.  $\square$

**Corollary 3.8.** *Let  $G$  be a 3-connected cubic graph with at least six vertices. Then there is a spanning tree containing exactly two removable edges if and only if  $G$  is a 3T-graph.*

**Proof** Assume that  $G$  is isomorphic to some  $p$ -extended  $3T_{k+2}$  ( $k \geq 0, p \geq 0$ ). Following the notation of Remark 1.3, let  $T_1, T_2$  and  $T_3$  be the three trees of  $N(G)$ . By adding to  $N(G)$  two edges of any given triangle of  $G$  we get a spanning tree containing exactly two removable edges.

We prove now by induction on  $n \geq 6$  that, if  $G$  a 3-connected cubic graph on  $n$  vertices spanned by a tree  $T$  containing exactly two removable edges, then it is isomorphic to some  $p$ -extended  $3T_{k+2}$  ( $k \geq 0, p \geq 0$ ).

When  $n = 6$ ,  $G$  is isomorphic to  $3T_{2,0}$  (that is,  $PR_{0,0}$ ) and the result is obvious. Assume that the result holds for any 3-connected cubic graph with  $6 \leq n' < n$  vertices having a spanning tree containing exactly two removable edges.

Since  $|T| = n - 1$  and  $|N(G)| \leq n - 3$ , we need to have  $|N(G)| = n - 3$  (that is  $N(G)$  is a spanning forest and is formed of exactly three trees,  $T_1, T_2$  and  $T_3$ ) and every edge of  $N(G)$  must be contained in  $T$ . If no 3-edge cut distinct from an end 3-edge cut exists then  $G$  is isomorphic either to  $PR_{0,0}$  (that is,  $3T_{2,0}$ ) or to  $PR_{1,0}$  (that is,  $3T_{2,1}$ ) or to the graph  $3T_{3,0}$  depicted in Figure 5.




FIGURE 5.  $3T_{3,0}$

Let  $F$  be a 3-edge cut distinct from an end 3-edge cut. Let  $\{V_1, V_2\}$  be the associated partition of  $F$ . We can construct a new 3-connected cubic graph  $G_1$  by replacing in  $G$  the subgraph  $G[V_2]$  by a triangle  $\Delta'_1$ . In the same way, we construct  $G_2$  by replacing  $G[V_1]$  by a triangle  $\Delta'_2$ . The trace of the forest  $N(G)$  in  $G_1$  gives a spanning forest of three trees of non removable edges. If we add two edges of  $\Delta'_1$  to these trees, we get a spanning tree of  $G_1$  containing exactly two removable edges. By the induction hypothesis,  $G_1$  is isomorphic to a  $p_1$ -extended  $3T_{k_1+2}$  and, in the same way  $G_2$  is isomorphic to a  $p_2$ -extended  $3T_{k_2+2}$ . The reconstruction of  $G$  gives a  $(p_1 + p_2)$ -extended  $3T_{k_1+k_2+2}$ , and the result follows.  $\square$

#### REFERENCES

1. J.-L. Fouquet and H. Thuillier, *Cycles through given vertices in planar 3-connected cubic graphs*, *Ars Combinatoria* **20B** (1985), 75–105.
2. ———, *k-minimal 3-connected cubic graphs*, *Ars Combinatoria* **25A** (1988), 149–190.
3. Wu Jichang and Li Xueliang, *Removable Edges Outside a Spanning Tree of a 3-connected 3-regular Graph*, *Journal of Mathematical Study* **36** (2003), no. 3, 223–229.
4. Haiyan Kang, Jichang Wu, and Guojun Li, *Removable Edges of a Spanning Tree in 3-Connected 3-Regular Graphs*, LNCS, no. 4613, pp. 337–345, Springer Verlag Berlin Heidelberg, 2007.
5. W. T. Tutte, *A theory of 3-connected graphs*, *Indag Math* **23** (1961), 441–455.

L.I.F.O., FACULTÉ DES SCIENCES, B.P. 6759, UNIVERSITÉ D'ORLÉANS, 45067 ORLÉANS CEDEX 2, FR