

## Sampling per mode for rare event simulation in switching diffusions

Jaroslav Krystul, François Le Gland, Pascal Lezaud

► **To cite this version:**

Jaroslav Krystul, François Le Gland, Pascal Lezaud. Sampling per mode for rare event simulation in switching diffusions. [Research Report] RR-7499, INRIA. 2010, pp.19. <inria-00550716>

**HAL Id: inria-00550716**

**<https://hal.inria.fr/inria-00550716>**

Submitted on 29 Dec 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

*Sampling per Mode for Rare Event Simulation  
in Switching Diffusions*

Jaroslav Krystul — François Le Gland — Pascal Lezaud

N° 7499

Décembre 2010

---

A large, light gray stylized 'R' logo is positioned to the left of the text. A horizontal gray brushstroke is located below the text.

*R*apport  
de recherche



## Sampling per Mode for Rare Event Simulation in Switching Diffusions

Jaroslav Krystul\* , François Le Gland\*\* , Pascal Lezaud<sup>||</sup>

Thème : Modèles et méthodes stochastiques  
Équipe-Projet ASPI

Rapport de recherche n° 7499 — Décembre 2010 — 19 pages

**Abstract:** An interacting particle system (IPS) approach is virtually applicable to estimate rare event for switching diffusions, since these processes own the strong Markov property. Nevertheless, in practice the straightforward application of this approach to switching diffusions fails to produce reasonable estimates within a reasonable amount of simulation time. This happens because each resampling step tends to sample more "heavy" particles from modes with higher probabilities, thus "light" particles in the modes with small probability tend to be discarded. To avoid this, a conditional "sampling per mode" algorithm has been proposed by Krystul (2006): instead of starting the algorithm with  $N$  particles randomly distributed, we draw in each mode  $j$ , a fixed number  $N^j$  of particles and at each resampling step, the same number of particles is sampled for each visited mode. In this paper, we establish a law of large numbers theorem as well as a central limit theorem (CLT) for the estimate of the rare event probability.

**Key-words:** rare event simulation, switching diffusion, multilevel splitting, stratification, central limit theorem (CLT).

This work was partially supported by the European Commission under the project *Safety, Complexity and Responsibility-Based Design and Validation of Highly Automated Air Traffic Management (iFLY)* (contract number TREN/07/FP6AE/S07.71574/037180) within the 6th Framework Programme FP6-2005-Aero-4 (Aeronautics and Space). The authors gratefully acknowledge Henk Blom, for his continuing support and interest in this work.

\* University of Twente, Department of Applied Mathematics, 7500 AE ENSCHEDE, The Netherlands.

— [krystulj@ewi.utwente.nl](mailto:krystulj@ewi.utwente.nl)

\*\* INRIA Rennes — Bretagne Atlantique, Campus de Beaulieu, 35042 RENNES Cédex, France. — [legland@irisa.fr](mailto:legland@irisa.fr)

<sup>||</sup> Direction Générale de l'Aviation Civile, DSNA/DTI/R&D, 7 avenue Edouard-Belin, 31055 TOULOUSE Cédex 4, France.

— [lezaud@recherche.enac.fr](mailto:lezaud@recherche.enac.fr)

# Échantillonnage par mode pour la simulation d'évènements rares dans les diffusions à paramètre markovien

**Résumé :** Un algorithme de branchement multi-niveaux peut en principe être utilisé pour estimer des évènements rares dans des diffusions à paramètre markovien, puisque ces processus vérifient la propriété de Markov forte. En pratique, l'application directe de cet algorithme peut échouer à produire une estimation raisonnable en un temps de simulation raisonnable. En effet, chaque étape de ré-échantillonnage tend à favoriser les particules dans les modes de forte probabilité, de sorte que les particules dans les modes de faible probabilité tendent à être éliminées. Pour éviter cet écueil, un algorithme de ré-échantillonnage par mode a été proposé par Krystul (2006) : au lieu de démarrer l'algorithme avec  $N$  particules distribuées aléatoirement, on génère dans chaque mode  $j$  un nombre fixé  $N^j$  de particules, et à chaque étape de ré-échantillonnage, le même nombre de particules est généré dans chaque mode visité. Dans cet article, on établit une loi des grands nombres et un théorème central limite (TCL) pour l'estimation de la probabilité de l'évènement rare.

**Mots-clés :** simulation d'évènement rare, diffusion à paramètre markovien, branchement multi-niveaux, stratification, théorème central limite (TCL).

## 1 Introduction

Rare event simulation requires acceleration techniques to speed up the occurrence of the rare events under consideration, otherwise it may take unacceptably large sample sizes to get enough positive realizations, or even a single one, on average. A well known technique is *importance sampling*, whose idea is to change the probability laws driving the model in order to make the events of interest more likely, and to correct the bias by multiplying the estimator by the suitable likelihood ratio [13]. An alternative technique called *multilevel splitting* does not need to modify the probability laws that drive the system; this means that the computer program that implements the simulation model can just be a black box [12]. The idea of the splitting is to express the small probability of rare event to be estimated as the product of a certain number of larger probabilities, which can be efficiently estimated by the Monte Carlo methods. This can be achieved by introducing sets of intermediate states that are visited by stochastic process one after the other, in an ordered sequence, before reaching the final set of rare event states. The probability of rare event is then given by the product of the conditional probabilities of reaching a set of intermediate states given that the previous set of intermediate states have been reached. Each conditional probability is estimated by simulating in parallel several copies of the system, i.e. each copy is considered as a particle following the trajectory generated through the system dynamics. Each particle branches (i.e. the trajectory splits into a number of independent subpaths, which subsequently evolve independently of each other) as soon as it enters the intermediate states, which is usually characterized by the crossing of a threshold defined by an *importance function*. Reaching intermediate states is more likely than reaching the rare event states, and by splitting at each threshold the chances to reach the rare event states are increasing. Several strategies have been designed to determine the importance function, to decide the number of splits at each level, and to handle the trajectories that tend to go in the wrong direction (away from the rare event of interest). The most difficulty is to find an appropriate importance function, a poor choice can easily lead to bad results [6, 7, 10]. One of the best-know versions of splitting is the RESTART method [14, 15, 16].

The multilevel splitting technique can also be considered as an interacting particle interpretation (IPS) of the Feynman-Kac models, a general framework presented in [4]. This abstract Feynman-Kac formulation gives a powerful tool which in particular allows to establish a Strong law of large numbers and a central limit theorem for the estimate of the rare event probability [4, 5, 3].

Owing to the increasing demands for modelling large-scale and complex systems, switching diffusions (a subclass of hybrid processes) are lately receiving growing attention [1, 9, 17, 2]. A distinctive feature of these systems is the coexistence of continuous dynamics and discrete events; they also satisfy the strong Markov property. While in theory the IPS approach is virtually applicable to any strong Markov process, in practice the straightforward application of this approach to switching diffusions fails to produce reasonable estimates within a reasonable amount of simulation time. The reason is that there may be few if no particles in modes with small probabilities (i.e. "light" modes). This happens because each resampling step tends to sample more "heavy" particles from modes with higher probabilities, thus, "light" particles in the "light" modes tend to be discarded. By increasing the number of particles the IPS estimates should improve but only at the cost of substantially increased simulation time which makes the performance of IPS approach in switching diffusions similar to one of the standard Monte Carlo. To avoid this, a conditional "sampling per mode" algorithm has been proposed in [9]; instead of starting the algorithm with  $N$  particles randomly distributed, we draw in each mode  $j$ , a fixed number  $N^j$  particles and at each resampling step, the same number of particles is sampled for each visited mode. Using the techniques introduced in [4, 11], we recently established a law of large numbers theorem as well as a central limit theorem for the estimate of the rare event probability as the number of particles increase to infinity.

The rest of the paper is arranged as follows. In Section 2, we introduce the abstract of Feynman-Kac and particle theory in the context of multilevel splitting and we adapt this framework to take into account the discrete modes of the switching diffusion. In particular, we detail the conditional "sampling per mode" algorithm. The Section 3 is devoted to the asymptotic behaviour of algorithm as the number of particles tends to infinity. Using an approach based on a martingale decomposition, as presented in [4], we establish a law of large numbers and a central limit theorem for the particle approximation of the rare event probability. Finally, Section 4 summarizes the paper and outlines a number of directions for future research.

## 2 Multilevel Feynman-Kac distributions

### 2.1 Formulation

We consider a switching diffusion  $Z = \{(X_t, \theta_t); t \geq 0\}$  taking values in  $\mathbb{R}^d \times \mathbb{M}$  for some  $d > 0$  and  $\mathbb{M} = \{1, \dots, M\}$ . More precisely,  $Z_t$  is a two-component Markov process such that  $(X_t)$  is a continuous component taking values in  $\mathbb{R}^d$  and  $(\theta_t)$  is a jump process taking values in the finite set  $\mathbb{M}$ ; it can be described by

$$dX_t = b(X_t, \theta_t) dt + \sigma(X_t, \theta_t) dB_t,$$

and

$$\mathbb{P}(\theta_{t+\Delta t} = j | \theta_t = i, X_t = x) = \lambda_{ij}(x) \Delta t + o(\Delta t), \quad i \neq j.$$

where  $B_t$  is a  $d$ -dimensional standard Brownian motion,  $b(\cdot, \cdot) : \mathbb{R}^n \times \mathbb{M} \rightarrow \mathbb{R}^n$  and  $\sigma(\cdot, \cdot) : \mathbb{R}^n \times \mathbb{M} \rightarrow \mathbb{R}^{n \times d}$ .

Throughout the paper, we assume that  $Z$  is a strong Markov process and we denote by  $\eta_0$  the law of  $Z_0$ . Let  $D \subset \mathbb{R}^d$  be some closed critical region, in which the continuous component of  $Z$  could enter but with a very small probability. Let us introduce an embedded sequence of closed regions

$$[0, \infty) \times D = D_n \subset \dots \subset D_1 \subset D_0 \subset [0, \infty) \times \mathbb{R}^d,$$

which are usually defined by

$$D_k = \{(t, x) \in [0, \infty) \times \mathbb{R}^d : h(t, x) \leq c_k\},$$

where  $h$  is some lower semi-continuous function, called the *importance function*, and the cylinders  $B = D \times \mathbb{M}$  and  $A_k = D_k \times \mathbb{M}$ . Set the corresponding hitting times

$$T_k = \inf\{t \geq 0 : (t, Z_t) \in A_k\} = \inf\{t \geq 0 : (t, X_t) \in D_k\},$$

which satisfy

$$0 = T_0 \leq T_1 \leq \dots \leq T_n = T_B.$$

To capture the precise behaviour of the process  $Z$  in each region, we consider the random excursions  $\mathcal{Z}_k$  of  $Z$  between the successive random times  $T_{k-1}$  and  $T_k$ . More precisely, we introduce the discrete-time Markov chain  $\mathcal{Z} = \{\mathcal{Z}_k, k = 1, \dots, n\}$  with value in the excursion set  $E$ , defined by

$$\mathcal{Z}_k = ((t, X_t, \theta_t), T_{k-1} \wedge T \leq t \leq T_k \wedge T),$$

with  $t \wedge T = \inf\{T, t\}$  and  $T$  a deterministic or stopping time. We observe that these excursions can be decomposed in a string of diffusions, one by discrete mode; each of them having a random length. The non-homogeneous Markov kernels  $\mathcal{M}_k$ , which describe the Markovian transitions of the Markov chain  $\mathcal{Z}$ , are defined for all excursion  $e$  and all functions  $f$  on  $E$  by

$$\mathcal{M}_k f(e) = \mathbb{E}[f(\mathcal{Z}_k) | \mathcal{Z}_{k-1} = e].$$

To check whether or not a given path  $e = ((t, Z_t), t_1 \leq t \leq t_2)$ , starting at  $(t_1, Z_{t_1}) \in A_{k-1}$  at time  $t_1$ , has succeeded to reach the level  $A_k$  at time  $t_2$ , it is convenient to introduce the terminal point  $\pi(e) = (t_2, Z_{t_2})$  of the excursion and the indicator functions  $g_k$  defined by

$$g_k(e) = 1_{\{\pi(e) \in A_k\}},$$

and to capture the discrete component, we introduce the potential functions

$$g_k^j(e) = 1_{\{\pi(e) \in D_k \times \{j\}\}}, \quad j \in \mathbb{M},$$

giving the following decomposition

$$g_k(e) = \sum_{j \in \mathbb{M}} g_k^j(e). \quad (2.1)$$

With these notations, and for each  $k$ , we have  $T_k \leq T$  if and only if  $g_k(\mathcal{Z}_k) = 1$ , and we have  $T_k \leq T$  with  $\theta_{T_k} = j$  if and only if  $g_k^j(\mathcal{Z}_k) = 1$ , hence

$$1_{\{T_k \leq T\}} = g_k(\mathcal{Z}_k), \quad \text{and} \quad 1_{\{T_k \leq T, \theta_{T_k} = j\}} = g_k^j(\mathcal{Z}_k).$$

Now define for  $k = 1, \dots, n$  the so-called unnormalized Feynman-Kac measures,  $\gamma_k$  and  $\widehat{\gamma}_k$  on the path space  $E$  in such a way that the integral of all bounded measurable functions  $f$  relatively to these measures are given by

$$\begin{aligned}\gamma_k(f) &= \mathbb{E}[f(\mathcal{Z}_k)g_{k-1}(\mathcal{Z}_{k-1})] = \mathbb{E}\left[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k \wedge T)1_{\{T_{k-1} \leq T\}}\right], \\ \widehat{\gamma}_k(f) &= \mathbb{E}[f(\mathcal{Z}_k)g_k(\mathcal{Z}_k)] = \mathbb{E}\left[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k)1_{\{T_k \leq T\}}\right].\end{aligned}$$

In particular, when  $f$  is the constant function 1, we obtain

$$\gamma_k(1) = \mathbb{P}[T_{k-1} \leq T], \quad \text{and} \quad \widehat{\gamma}_k(1) = \mathbb{P}[T_k \leq T].$$

The Feynman-Kac distributions  $\eta_k$  and  $\widehat{\eta}_k$  are derived by normalizing these measures,

$$\begin{aligned}\eta_k(f) &= \frac{\gamma_k(f)}{\gamma_k(1)} = \mathbb{E}[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k \wedge T)|T_{k-1} \leq T], \\ \widehat{\eta}_k(f) &= \frac{\widehat{\gamma}_k(f)}{\widehat{\gamma}_k(1)} = \mathbb{E}[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k)|T_k \leq T],\end{aligned}$$

and using the convention  $T_{-1} = 0$  leads to the observation that  $\gamma_0 = \eta_0$ .

We observe that

$$1_{\{T_k \leq T\}} = 1_{\{T_{k-1} \leq T, T_k \leq T\}},$$

or equivalently

$$g_k(\mathcal{Z}_k) = g_{k-1}(\mathcal{Z}_{k-1})g_k(\mathcal{Z}_k),$$

hence

$$\gamma_k(fg_k) = \widehat{\gamma}_k(f), \quad \text{and} \quad \widehat{\eta}_k(f) = \frac{\gamma_k(fg_k)}{\gamma_k(g_k)} = \frac{\eta_k(fg_k)}{\eta_k(g_k)}, \quad (2.2)$$

and more interestingly, how the probabilities of transition from one region to the following are related to the Feynman-Kac distributions,

$$\eta_k(g_k) = \mathbb{P}[T_k \leq T|T_{k-1} \leq T] := P_k, \quad \eta_k(g_k^j) = \mathbb{P}[T_k \leq T, \theta_{T_k} = j|T_{k-1} \leq T] := P_k^j.$$

From now on, we assume that not only  $P_k \neq 0$  for all  $k$ , but also that  $P_k^j$  are non zero all modes  $j \in \mathbb{M}$ .

Furthermore, the ‘‘unnormalized models’’  $(\gamma_k, \widehat{\gamma}_k)$  are related to the Feynman-Kac distribution flow  $(\eta_p)_{p \leq k}$ , by the following key formula

$$\gamma_k(f) = \eta_k(f) \prod_{p=0}^{k-1} \eta_p(g_p) \quad \text{and} \quad \widehat{\gamma}_k(f) = \widehat{\eta}_k(f) \prod_{p=0}^k \eta_p(g_p). \quad (2.3)$$

In order to keep trace of the discrete mode, we construct for any  $j \in \mathbb{M}$  the unnormalized Feynman-Kac measures  $\gamma_k^j$  and  $\widehat{\gamma}_k^j$  defined by

$$\begin{aligned}\gamma_k^j(f) &= \mathbb{E}\left[f(\mathcal{Z}_k)g_{k-1}^j(\mathcal{Z}_{k-1})\right] = \mathbb{E}\left[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k \wedge T)1_{\{T_{k-1} \leq T, \theta_{T_{k-1}} = j\}}\right], \\ \widehat{\gamma}_k^j(f) &= \mathbb{E}\left[f(\mathcal{Z}_k)g_k^j(\mathcal{Z}_k)\right] = \mathbb{E}\left[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k)1_{\{T_k \leq T, \theta_{T_k} = j\}}\right].\end{aligned}$$

The Feynman-Kac distributions  $\eta_k^j$  and  $\widehat{\eta}_k^j$  are derived by normalizing these measures, respectively

$$\begin{aligned}\eta_k^j(f) &= \frac{\gamma_k^j(f)}{\gamma_k^j(1)} = \mathbb{E}[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k \wedge T)|T_{k-1} \leq T, \theta_{T_{k-1}} = j], \\ \widehat{\eta}_k^j(f) &= \frac{\widehat{\gamma}_k^j(f)}{\widehat{\gamma}_k^j(1)} = \mathbb{E}[f((t, X_t, \theta_t), T_{k-1} \leq t \leq T_k)|T_k \leq T, \theta_{T_k} = j].\end{aligned}$$

We observe that

$$1_{\{T_k \leq T, \theta_{T_k} = j\}} = 1_{\{T_{k-1} \leq T, T_k \leq T, \theta_{T_k} = j\}},$$



or equivalently

$$g_k^j(\mathcal{Z}_k) = g_{k-1}(\mathcal{Z}_{k-1})g_k^j(\mathcal{Z}_k),$$

hence

$$\gamma_k(fg_k^j) = \widehat{\gamma}_k^j(f), \quad \text{and} \quad \widehat{\eta}_k^j(f) = \frac{\gamma_k(fg_k^j)}{\gamma_k(g_k^j)} = \frac{\eta_k(fg_k^j)}{\eta_k(g_k^j)}. \quad (2.4)$$

Clearly, we have the decompositions

$$\widehat{\eta}_k = \sum_{j \in \mathbb{M}} \omega_k^j \widehat{\eta}_k^j, \quad \eta_{k+1} = \sum_{j \in \mathbb{M}} \omega_k^j \eta_{k+1}^j. \quad (2.5)$$

where

$$\omega_k^j = \widehat{\eta}_k(g_k^j) = \frac{\eta_k(g_k^j)}{\eta_k(g_k)} = \frac{\widehat{\gamma}_k^j(1)}{\widehat{\gamma}_k(1)} = \mathbb{P}(\theta_{T_k} = j | T_k \leq T). \quad (2.6)$$

## 2.2 Feynman-Kac semigroups

Previously, we have introduced the Feynman-Kac distributions  $\eta_k$ . Now, we will investigate the time evolution of the Feynman-Kac flow ( $\eta_k$ ;  $0 \leq k \leq n$ ). In fact, from the Markov property of the process  $Z$ , we see that

$$\gamma_k(f) = \mathbb{E}[g_{k-1}(\mathcal{Z}_{k-1})\mathbb{E}[f(\mathcal{Z}_k)|\mathcal{Z}_{k-1}]] = \gamma_{k-1}(g_{k-1}\mathcal{M}_k f) = \widehat{\gamma}_{k-1}(\mathcal{M}_k f), \quad (2.7)$$

and in the context of switching jump diffusions, for any  $j \in \mathbb{M}$

$$\gamma_k^j(f) = \mathbb{E}[g_{k-1}^j(\mathcal{Z}_{k-1})\mathbb{E}[f(\mathcal{Z}_k)|\mathcal{Z}_{k-1}]] = \gamma_{k-1}(g_{k-1}^j\mathcal{M}_k f) = \widehat{\gamma}_{k-1}^j(\mathcal{M}_k f). \quad (2.8)$$

These formulas suggest the introduction of the linear operators  $Q_k$  and  $Q_k^j$  defined respectively by

$$Q_k f = g_{k-1}\mathcal{M}_k f \quad \text{and} \quad Q_k^j f = g_{k-1}^j\mathcal{M}_k f. \quad (2.9)$$

An immediate consequence of (2.7) and (2.8) is that  $\gamma_k$  and  $\gamma_k^j$  satisfy linear equations of the form

$$\gamma_k = \gamma_{k-1}Q_k \quad \text{and} \quad \gamma_k^j = \gamma_{k-1}Q_k^j. \quad (2.10)$$

Nevertheless, we seek the evolution of the normalized Feynman-Kac measures  $\eta_k$  and  $\eta_k^j$ , that we suspect to be nonlinear. To establish it, we introduce the mappings  $\Phi_k$  from the set of measures  $\mathcal{P}_k(E) = \{\eta : \eta(g_k) > 0\}$  into the set  $\mathcal{P}(E)$  of measure on  $E$  defined by

$$\Phi_k(\eta)(f) = \Psi_{k-1}(\eta)(\mathcal{M}_k f), \quad \text{with} \quad \Psi_k(\eta)(f) = \frac{\eta(fg_k)}{\eta(g_k)}.$$

It can now be easily verified that the Feynman-Kac flow is the solution of a nonlinear measure-valued dynamical system

$$\eta_k = \Phi_k(\eta_{k-1}), \quad (2.11)$$

Since  $\Psi_k(\eta_k) = \widehat{\eta}_k$ , we see that the recursion (2.11) involves two separate selection/mutation transitions

$$\eta_k \in \mathcal{P}_k(E) \xrightarrow{\text{selection}} \widehat{\eta}_k := \Psi_k(\eta_k) \in \mathcal{P}(E) \xrightarrow{\text{mutation}} \eta_{k+1} = \widehat{\eta}_k \mathcal{M}_{k+1} \in \mathcal{P}(E). \quad (2.12)$$

In the specific case of switching jump diffusions, we introduce for any  $j \in \mathbb{M}$  the following transformations

$$\Phi_k^j(\eta)(f) := \Psi_{k-1}^j(\eta)(\mathcal{M}_k f) \quad \text{with} \quad \Psi_k^j(\eta)(f) := \frac{\eta(g_k^j f)}{\eta(g_k^j)},$$

and from (2.4), we check that

$$\widehat{\eta}_k^j = \Psi_k^j(\eta_k), \quad \text{and} \quad \eta_{k+1}^j = \Phi_{k+1}^j(\eta_k) = \widehat{\eta}_k^j \mathcal{M}_{k+1}. \quad (2.13)$$

Furthermore, the operator  $\Phi_k$  can be written as the weighted sum

$$\Phi_k(\eta) = \sum_{j \in \mathbb{M}} \frac{\eta(g_{k-1}^j)}{\eta(g_{k-1})} \Phi_k^j(\eta), \quad (2.14)$$

and from (2.6), we deduce the particular case

$$\Phi_k(\eta_{k-1}) = \sum_{j \in \mathbb{M}} \omega_{k-1}^j \Phi_k^j(\eta_{k-1}).$$

We have seen how the measure  $\eta_n$  is related to the previous measure  $\eta_{n-1}$ , but we will need to express  $\eta_n$  in terms of  $\eta_p$  for all  $0 \leq p \leq n$ . Nevertheless, the evolution being nonlinear it is easier to start by considering the evolution of  $\gamma_n$  and  $\gamma_n^j$ , since a straightforward iteration of (2.10) leads us to introduce for any  $j \in \mathbb{M}$ , the linear semigroups  $Q_{p,n}$  and  $Q_{p,n}^j$ , defined respectively by

$$Q_{p,n} = Q_{p+1} Q_{p+2} \cdots Q_n, \quad Q_{p,n}^j = Q_{p+1} Q_{p+2} \cdots Q_{n-1} Q_n^j = Q_{p,n-1} Q_n^j, \quad (2.15)$$

with the convention  $Q_{n,n} = \text{Id}$ . We remark firstly that

$$Q_{p,n} 1 = Q_{p,n-1} g_{n-1}, \quad Q_{p,n}^j 1 = Q_{p,n-1} g_{n-1}^j.$$

Secondly, we get

$$\eta_n(f) = \frac{\gamma_p(Q_{p,n} f)}{\gamma_p(Q_{p,n} 1)} = \frac{\eta_p(Q_{p,n} f)}{\eta_p(Q_{p,n} 1)}, \quad (2.16)$$

and, for any  $j \in \mathbb{M}$

$$\eta_n^j(f) = \gamma_p(Q_{p,n}^j f), \quad \eta_n^j(f) = \frac{\eta_p(Q_{p,n}^j f)}{\eta_p(Q_{p,n}^j 1)}. \quad (2.17)$$

Note that an immediate consequence of (2.16) and (2.17) is that

$$\eta_p(Q_{p,n} 1) = \frac{\gamma_n(1)}{\gamma_p(1)} = \prod_{q=p}^{n-1} \eta_q(g_q), \quad \text{and} \quad \eta_p(Q_{p,n}^j 1) = \frac{\gamma_n^j(1)}{\gamma_p(1)}. \quad (2.18)$$

We obtain also

$$\omega_{n-1}^j = \frac{\eta_p(Q_{p,n}^j 1)}{\eta_p(Q_{p,n} 1)}. \quad (2.19)$$

## 2.3 Interacting particle system approximations

### 2.3.1 Introduction

From a pure mathematical point of view, particles methods can be interpreted as a kind of stochastic linearization technique for solving nonlinear equations in measure space. The idea is to associate to the nonlinear dynamical structure (2.11) a sequence  $\xi = (\xi^1, \dots, \xi^N)$ , of  $N$  excursion-valued particles, such that the empirical measure of the configurations converge as  $N \rightarrow \infty$  to the desired distribution  $\eta$ .

More precisely, at time  $k$ , we consider the particles

$$\begin{aligned} \xi_k^i &= ((t, X_t^i, \theta_t^i), ; T_{k-1}^i \leq t \leq T_k^i \wedge T) \in E \cup \{\Delta\}, \\ \widehat{\xi}_k^i &= ((t, \widehat{X}_t^i, \widehat{\theta}_t^i), ; \widehat{T}_{k-1}^i \leq t \leq \widehat{T}_k^i) \in E \cup \{\Delta\}, \end{aligned}$$

where  $\Delta$  stands for a cemetery point that we introduce here to take into account the possible stopping of the algorithm. The random lengths of the corresponding excursions  $\xi_k^i$  and  $\widehat{\xi}_k^i$  are  $T_k^i \wedge T - T_{k-1}^i$  and  $\widehat{T}_k^i - \widehat{T}_{k-1}^i$  respectively.

The interacting particle system (IPS) approach consists in approximating the two step transitions (2.12) of the system (2.11) by the two step transitions

$$\eta_k^N := \frac{1}{N} \sum_{i=1}^N \delta_{\xi_k^i} \xrightarrow{\text{selection}} \widehat{\eta}_k^N := \frac{1}{N} \sum_{i=1}^N \delta_{\widehat{\xi}_k^i} \xrightarrow{\text{mutation}} \eta_{k+1} = \frac{1}{N} \sum_{i=1}^N \delta_{\xi_{k+1}^i}. \quad (2.20)$$

So, starting from an approximation  $\eta_0^N$  to  $\eta_0$ , during the mutation transition  $\widehat{\xi}_k \rightarrow \xi_{k+1}$ , each selected particle  $\widehat{\xi}_k^\kappa$  evolves randomly according to the Markov transition  $\mathcal{M}_{k+1}$ , independently of each other. The selection transition  $\xi_{k+1} \rightarrow \widehat{\xi}_{k+1}$  is defined as follows: only some of the particles  $\xi_{k+1}$  have succeeded in reaching the desired set  $A_{k+1}$ ; unless the amount of these is equal to zero (in this case, the algorithm is stopped), we sample, randomly and independently, the  $N$  particles  $\widehat{\xi}_{k+1}$  distributed according to the distribution  $\Psi_{k+1}(\eta_{k+1}^N)$ .

The IPS is nothing else than a sequence of nonhomogeneous Markov chains on the product space  $E^N$  with transition kernels given by

$$\begin{aligned} \mathbb{P}(\widehat{\xi}_k \in dx | \xi_k) &= \prod_{i=1}^N \Psi_k(\eta_k^N)(dx^i), \\ \mathbb{P}(\xi_k \in dz | \widehat{\xi}_{k-1}) &= \prod_{i=1}^N \mathcal{M}_k(\widehat{\xi}_{k-1}^i, dz^i). \end{aligned}$$

Nevertheless, the classical IPS algorithm is not really suitable for switching jump diffusion, mainly because of the potential existence of discrete modes with very small initial probability. To avoid the particles never been drawn in these “light” modes, a conditional “sampling per mode” algorithm has been proposed in [9]; instead of starting the algorithm with  $N_0$  particles randomly distributed in  $A_0 \setminus A_1$  according to  $\eta_0$ , we draw in each mode  $j$ , a fixed number  $N^j$  particles in  $(D_0 \setminus D_1) \times \{j\}$ , randomly distributed according to the conditional law  $\eta_0^j$ . Resampling per mode allows us to avoid loss of “light” particles in “light modes”, and so helps to maintain fixed number of particles in each mode. We will see hereafter, that the empirical measure  $\eta_k^N$  is a weighted sum of the conditional empirical measure  $\eta_k^{j,N}$  (see (2.5))

$$\eta_k^N = \sum_j \omega_{k-1}^{j,N} \eta_k^{j,N} = \sum_\kappa \beta_k^\kappa \delta_{\xi_k^\kappa}, \quad \text{with} \quad \eta_k^{j,N} = \frac{1}{N^j} \sum_\kappa \delta_{\xi_k^\kappa},$$

where the three sums are taken over, respectively, all discrete modes containing at least one particle, all particles and finally all particles in the discrete mode  $j$ . Nevertheless, the total number of particles can decrease, typically if one or several modes become empty at some time, and it can also increase, typically if one or several of these empty modes become non empty at a later time. That means we have to introduce the total number  $N_k$  of particles, at each step  $k$ .

### 2.3.2 Detailed description

Prior to the detailed description of the IPS, it is convenient to introduce some notations. Let  $J_k$  be the set of non empty discrete modes after the end of the step  $k$ , with the convention  $J_0 = \mathbb{M}$ , and for each  $j \in J_k$ , let

$$\begin{aligned} J_k^j &= \{\kappa : \pi(\xi_k^\kappa) \in D_k \times \{j\}\}, \\ \widehat{J}_k^j &= \{\kappa : \pi(\widehat{\xi}_k^\kappa) \in D_k \times \{j\}\}, \\ \widehat{I}_k^N &= \bigcup_{j \in J_k} \widehat{J}_k^j, \end{aligned}$$

be respectively the set of (labels of) particles  $\xi_k^\kappa$  which have reached before the final time  $T$  the set  $A_k = D_k \times \mathbb{M}$  in the discrete mode  $j$ , the set of (labels of) selected particles  $\widehat{\xi}_k^\kappa$  whose terminal point is in the discrete mode  $j$ , and the set of (labels of) all the selected particles  $\widehat{\xi}_k^\kappa$ .

As the total number of particles can decrease as soon as none of the particles have succeeded to reach the desired level in some mode, we need to introduce the following sequence of integers

$$\widehat{N}_k = |\widehat{I}_k^N| = \sum_{j \in J_k} N^j = \sum_{j \in \mathbb{M}} \widehat{N}_k^j,$$

where  $\widehat{N}_k^j = N^j$  if  $j \in J_k$  and  $\widehat{N}_k^j = 0$  otherwise. Thus, our model is not restricted to a fixed population size but it takes values in the state space

$$\mathbf{E} = \bigcup_{p \in \mathbb{N}} (\{p\} \times [0, \infty)^p \times E^p)$$

with the convention  $E^0 = \{\Delta\}$ ; the parameter  $p$  represents the size of the system. It is advisable to observe that we can have  $\widehat{N}_{k+1}^j = N^j$  while  $\widehat{N}_k^j = 0$ , then the size of the population can increase or decrease by jumps.

We associate with the time evolution of this model

$$(N_k, \beta_k, \xi_k) \xrightarrow{\text{selection}} (\widehat{N}_k, \widehat{\beta}_k, \widehat{\xi}_k) \xrightarrow{\text{mutation}} (N_{k+1}, \beta_{k+1}, \xi_{k+1}) \quad (2.21)$$

the canonical filtrations  $F_k \subset \widehat{F}_k \subset F_{k+1}$ . Then the IPS algorithm is conducted inductively as follows:

**Initialization:** For each  $j \in J_0$ , we sample  $N^j$  particles  $\xi_0^\kappa = \widehat{\xi}_0^\kappa = (0, (X_0^\kappa, j)) \sim \eta_0^j$ . Let  $\omega_0^j = \mathbb{P}(\theta_0 = j)$ , then the empirical measures of particles  $\eta_0^N$  and  $\widehat{\eta}_0^N$  are given by

$$\begin{aligned} \eta_0^N &= \sum_{j \in M} \omega_0^j \eta_0^{j,N} = \sum_{\kappa=1}^{N_0} \beta_0^\kappa \delta_{\xi_0^\kappa}, \quad \text{with} \quad \eta_0^{j,N} = \frac{1}{N^j} \sum_{\kappa \in J_0^j} \delta_{\xi_0^\kappa}, \\ \widehat{\eta}_0^N &= \sum_{j \in M} \omega_0^j \widehat{\eta}_0^{j,N} = \sum_{\kappa=1}^{N_0} \widehat{\beta}_0^\kappa \delta_{\widehat{\xi}_0^\kappa}, \quad \text{with} \quad \widehat{\eta}_0^{j,N} = \frac{1}{N^j} \sum_{\kappa \in \widehat{J}_0^j} \delta_{\widehat{\xi}_0^\kappa}, \end{aligned}$$

where  $\beta_0^\kappa = \widehat{\beta}_0^\kappa = \omega_0^j / N^j$ , for each  $\kappa \in J_0^j = \widehat{J}_0^j$ .

At each step  $k$ , the empirical measure  $\widehat{\eta}_k^N$  will be given by

$$\widehat{\eta}_k^N = \sum_{\kappa \in \widehat{I}_k^N} \widehat{\beta}_k^\kappa \delta_{\widehat{\xi}_k^\kappa} = \sum_{j \in J_k} \omega_k^{j,N} \widehat{\eta}_k^{j,N}, \quad \text{with} \quad \widehat{\eta}_k^{j,N} = \frac{1}{N^j} \sum_{\kappa \in \widehat{J}_k^j} \delta_{\widehat{\xi}_k^\kappa},$$

and the weights  $\omega_k^{j,N}$  are nonnegative and such that  $\sum_{j \in J_k} \omega_k^{j,N} = 1$ . Clearly,  $\widehat{\beta}_k^\kappa = \omega_k^{j,N} / N^j$  for each  $\kappa \in \widehat{J}_k^j$ . We notice that the particles in the same discrete mode have the same weight, or in other words, the weight of a particle depends only on the mode, i.e. on its discrete component.

The **mutation transition**  $\widehat{\xi}_k \rightarrow \xi_{k+1}$  at time  $k+1$  is defined as follows. If  $\widehat{N}_k = 0$ , the particle system dies and we set  $N_{k+1} = 0$ . Otherwise during mutation, independently of each other, each selected particle  $\widehat{\xi}_k^\kappa$  evolves randomly according to the Markov transition  $\mathcal{M}_{k+1}$ ; in other words,

$$\xi_{k+1}^\kappa = ((t, X_t^\kappa, \theta_t^\kappa), T_k^\kappa \leq t \leq T_{k+1}^\kappa \wedge T),$$

is a random variable with distribution  $\mathcal{M}_{k+1}(\widehat{\xi}_k^\kappa, \cdot)$ . More precisely, we set  $T_k^\kappa = \widehat{T}_k^\kappa$  and the path  $((t, X_t^\kappa, \theta_t^\kappa), t \geq T_k^\kappa)$  advances randomly as a copy of the process  $((t, X_t, \theta_t), t \geq T_k^\kappa)$ , i.e. according to the dynamic of the switching diffusion, starting at  $(T_k^\kappa, \widehat{X}_{T_k^\kappa}, \widehat{\theta}_{T_k^\kappa})$  and up to the first time  $T_{k+1}^\kappa$  it visits  $A_{k+1}$ , or up to  $T$ , whichever occurs first. During this transition, the total number of particles does not change, so we set  $N_{k+1} = \widehat{N}_k$ .

The weight of each particle is set as  $\beta_{k+1}^\kappa = \widehat{\beta}_k^\kappa$  and the  $N$ -particle approximation measure is given by

$$\eta_{k+1}^N = \sum_{\kappa \in \widehat{I}_{k+1}^N} \beta_{k+1}^\kappa \delta_{\xi_{k+1}^\kappa} = \sum_{j \in J_k} \omega_k^{j,N} \eta_{k+1}^{j,N}, \quad \text{with} \quad \eta_{k+1}^{j,N} = \frac{1}{N^j} \sum_{\kappa \in \widehat{J}_k^j} \delta_{\xi_{k+1}^\kappa}.$$

We easily verify that

$$\mathbb{E} \left[ \eta_{k+1}^N(f) \middle| \widehat{F}_k \right] = \widehat{\eta}_k^N \mathcal{M}_{k+1} f. \quad (2.22)$$

The **selection transition**  $\xi_{k+1} \rightarrow \widehat{\xi}_{k+1}$  is defined as follows. From the  $N_{k+1}$  particles  $\xi_{k+1}^i$ , only some of them have succeeded to reach the desired set  $A_{k+1}$ ; we recall that

$$J_{k+1}^j = \{\kappa : \pi(\xi_{k+1}^\kappa) \in D_{k+1} \times \{j\}\}$$

and we set

$$I_{k+1}^N = \bigcup_{j \in J_k} J_{k+1}^j.$$

If  $I_{k+1}^N = \emptyset$ , then none of the particles have succeeded to reach the desired region; the algorithm is stopped and  $\widehat{\xi}_{k+1} = \Delta$ . Otherwise, it may happen that there are some  $j$  for which  $J_{k+1}^j = \emptyset$ , but as long as  $I_{k+1}^N$  is not empty, we still continue the algorithm.

For each mode  $j$  such that  $J_{k+1}^j \neq \emptyset$ , we need to sample  $N^j$  particles in this mode. Nevertheless, more or less than  $N^j$  particles may have reached the set  $D_{k+1} \times \{j\}$ , so we need to resample  $N^j$  particles among the particles  $\xi_{k+1}^\kappa$ , with  $\kappa \in J_{k+1}^j$ . If  $|J_{k+1}^j| \leq N^j$ , then not enough particles have managed to reach the set  $D_{k+1} \times \{j\}$  before the final time  $T$ , and these successful particles should be replicated, whereas if  $|J_{k+1}^j| > N^j$ , then too many particles have managed to reach the set  $D_{k+1} \times \{j\}$  before the final time  $T$ , and some of these successful particles should be eliminated.

More precisely, we choose randomly the  $N^j$  particles  $\widehat{\xi}_{k+1}^\kappa$ ,  $\kappa \in \widehat{J}_{k+1}^j$ , identically distributed according to the distribution

$$\Psi_{k+1}^j(\eta_{k+1}^N) = \sum_{\kappa \in J_{k+1}^j} \left( \frac{\beta_{k+1}^\kappa}{\sum_{\kappa \in J_{k+1}^j} \beta_{k+1}^\kappa} \right) \delta_{\xi_{k+1}^\kappa}.$$

By construction  $\widehat{\xi}_{k+1}^\kappa$  is the copy of a successful particle  $\xi_{k+1}^\tau$  with  $\tau \in J_{k+1}^j$ ; necessarily  $\widehat{T}_{k+1}^\kappa = T_{k+1}^\tau \leq T$ . Each particle  $\xi_{k+1}^\kappa$  for  $\kappa \in J_{k+1}^j$  branches into a random number of offsprings  $M_{k+1}^{j,\kappa}$  and the sequence  $(M_{k+1}^{j,\kappa}, \kappa \in J_{k+1}^j)$  is distributed according to a

$$(M_{k+1}^{j,\kappa}, \kappa \in J_{k+1}^j) = \text{Multinomial} \left( N^j, \left( \frac{\beta_{k+1}^\kappa}{\sum_{\kappa \in J_{k+1}^j} \beta_{k+1}^\kappa}, \kappa \in J_{k+1}^j \right) \right).$$

For each  $j \in J_{k+1}$ , we obtain

$$\widehat{\eta}_{k+1}^{j,N} = \frac{1}{N^j} \sum_{\kappa \in \widehat{J}_{k+1}^j} \delta_{\widehat{\xi}_{k+1}^\kappa} = \frac{1}{N^j} \sum_{\kappa \in J_{k+1}^j} M_{k+1}^{j,\kappa} \delta_{\xi_{k+1}^\kappa};$$

namely, we approximate the empirical measure  $\Psi_{k+1}^j(\eta_{k+1}^N)$  by a new probability measure whose atom weights are integer multiples of  $1/N^j$ .

Furthermore, the mechanism is such that for any bounded test function  $f$ , we have

$$\mathbb{E} \left( \widehat{\eta}_{k+1}^{j,N}(f) | F_{k+1} \right) = \Psi_{k+1}^j(\eta_{k+1}^N)(f), \quad (2.23)$$

and

$$\mathbb{E} \left( \left[ \widehat{\eta}_{k+1}^{j,N}(f) - \Psi_{k+1}^j(\eta_{k+1}^N)(f) \right]^2 | F_{k+1} \right) = \frac{1}{N^j} \text{var} \left( f, \Psi_{k+1}^j(\eta_{k+1}^N) \right) \leq \frac{\|f\|^2}{N^j}, \quad (2.24)$$

where  $\text{var}(f, \mu) = \mu[(f - \mu(f))^2]$  and  $\|f\| = \sup_{x \in E} |f(x)|$ .

The total number  $\widehat{N}_{k+1}$  of particles  $\widehat{\xi}_{k+1}^\kappa$  and the weights  $\omega_{k+1}^{j,N}$  are given respectively by

$$\widehat{N}_{k+1} = \sum_{j \in J_{k+1}} N^j, \quad \text{and} \quad \omega_{k+1}^{j,N} = \widehat{\eta}_{k+1}^N(g_{k+1}^j) = \frac{\eta_{k+1}^N(g_{k+1}^j)}{\eta_{k+1}^N(g_{k+1})}.$$

It follows that

$$\omega_{k+1}^{j,N} = \sum_{\kappa \in \widehat{J}_{k+1}^j} \widehat{\beta}_{k+1}^\kappa = \sum_{\kappa \in J_{k+1}^j} \left( \frac{\beta_{k+1}^\kappa}{\sum_{\kappa \in I_{k+1}^N} \beta_{k+1}^\kappa} \right).$$

As the particles  $\widehat{\xi}_{k+1}^\kappa$  in the same mode have the same weight, we deduce that for any  $\kappa \in \widehat{J}_{k+1}^j$

$$\widehat{\beta}_{k+1}^\kappa = \frac{1}{N^j} \sum_{\kappa \in J_{k+1}^j} \left( \frac{\beta_{k+1}^\kappa}{\sum_{\kappa \in I_{k+1}^N} \beta_{k+1}^\kappa} \right).$$

Using (2.3) and (2.6), we also introduce the measures  $\gamma_n^N$  and  $\gamma_n^{j,N}$  defined respectively by

$$\gamma_n^N(f) = \eta_n^N(f) \prod_{p=0}^{n-1} \eta_p^N(g_p) \quad \text{and} \quad \gamma_n^{j,N}(f) = \omega_{n-1}^{j,N} \gamma_n^N(1) \eta_n^{j,N}(f),$$

as approximations of  $\gamma_n$  and  $\gamma_n^j$ . In particular for  $f \equiv 1$ , the IPS algorithm provides

$$\gamma_{n+1}^N(1) = \prod_{p=0}^n \eta_p^N(g_p) = \prod_{p=0}^n \sum_{\kappa \in I_p^N} \beta_p^\kappa,$$

as an estimate of the rare event probability  $\mathbb{P}(T_n \leq T) = \gamma_{n+1}(1)$ . In other words,  $\gamma_{n+1}^N(1)$  is the product of proportions of excursions having entered levels  $A_1, \dots, A_n$ .

### 3 Asymptotic behavior

In this section, our aim is to examine the asymptotic behavior of particle approximation models as the number of particles tends to infinity. We start with the analysis of the unnormalized measure  $\gamma_n^N$  and we show that this approximation has no bias. We follow the approach used in [4] which is based on a martingale decomposition. Finally, we establish a central limit theorem for unnormalized particle approximation measures.

#### 3.1 Law of large numbers

We begin by introducing some useful formulae and inequalities. Firstly, on the event  $\{N_k > 0\}$ , we have

$$\mathbb{E}(\widehat{\eta}_k^N(f)|F_k) = \sum_{j \in J_k} \mathbb{E}(\omega_k^{j,N} \widehat{\eta}_k^{j,N}(f)|F_k) = \sum_{j \in J_k} \omega_k^{j,N} \Psi_k^j(\eta_k^N)(f) = \Psi_k(\eta_k^N)(f), \quad (3.1)$$

since  $\omega_k^{j,N} = \eta_k^N(g_k^j)/\eta_k^N(g_k) \in F_k$ . Introducing the integer  $N_{\text{inf}}$  defined by  $N_{\text{inf}} = \inf\{N^j : j = 1, \dots, M\}$ , we deduce that

$$\mathbb{E} \left( \left[ \widehat{\eta}_k^N(f) - \Psi_k(\eta_k^N)(f) \right]^2 \middle| F_k \right) = \sum_{j \in J_k} \frac{(\omega_k^{j,N})^2}{N^j} \text{var}(f, \Psi_k^j(\eta_k^N)) \leq \|f\|^2 \sum_{j \in J_k} \frac{(\omega_k^{j,N})^2}{N^j} \leq \frac{\|f\|^2}{N_{\text{inf}}}. \quad (3.2)$$

Secondly, when  $J_k \neq \emptyset$  we obtain by (2.22) that

$$\begin{aligned} \mathbb{E} \left( \eta_{k+1}^N(f) \middle| F_k, \widehat{F}_k \right) &= \sum_{j \in J_k} \omega_k^{j,N} \mathbb{E} \left( \widehat{\eta}_k^{j,N}(\mathcal{M}_{k+1} f) \middle| F_k \right) \\ &= \sum_{j \in J_k} \omega_k^{j,N} \Psi_k^j(\eta_k^N)(\mathcal{M}_{k+1} f) \\ &= \sum_{j \in J_k} \omega_k^{j,N} \Phi_{k+1}^j(\eta_k^N)(f) \\ &= \sum_{j \in J_k} \frac{\eta_k^N(g_k^j)}{\eta_k^N(g_k)} \Phi_{k+1}^j(\eta_k^N)(f) \\ &= \Phi_{k+1}(\eta_k^N)(f), \end{aligned}$$

where we used (2.14) for the last line. It follows easily that

$$\mathbb{E}(\eta_{k+1}^N(f)|F_k) = \Phi_{k+1}(\eta_k^N)(f). \quad (3.3)$$

As the particles  $\xi_{k+1}^\kappa$  are independent conditionally to  $\widehat{F}_k$ , we get

$$\begin{aligned} \mathbb{E} \left( \left[ \eta_{k+1}^{j,N}(f) - \Phi_{k+1}^j(\eta_k^N)(f) \right]^2 \middle| \widehat{F}_k \right) &= \frac{1}{N^j} \widehat{\eta}_k^{j,N} \left( \mathcal{M}_{k+1} [f - \Phi_{k+1}^j(\eta_k^N)(f)]^2 \right) \\ \mathbb{E} \left( \left[ \eta_{k+1}^{j,N}(f) - \Phi_{k+1}^j(\eta_k^N)(f) \right]^2 \middle| F_k \right) &= \frac{1}{N^j} \Phi_{k+1}^j(\eta_k^N) \left( [f - \Phi_{k+1}^j(\eta_k^N)(f)]^2 \right). \end{aligned}$$

Hence, we deduce the following

$$\mathbb{E} \left( \left[ \eta_{k+1}^N(f) - \Phi_{k+1}(\eta_k^N)(f) \right]^2 \middle| F_k \right) = \sum_{j \in J_k} \frac{(\omega_k^{j,N})^2}{N^j} \text{var} \left( f, \Phi_{k+1}^j(\eta_k^N) \right) \leq \frac{\|f\|^2}{N_{\text{inf}}}. \quad (3.4)$$

Now, we study the difference between the particle measure  $\gamma_n^N$  and the limiting Feynman-Kac measures  $\gamma_n$ . Following the approach given in [4], we use the decomposition for each bounded function  $f$

$$\begin{aligned} 1_{\{N_n > 0\}} \gamma_n^N(f) - \gamma_n(f) &= \sum_{p=1}^n \left[ 1_{\{N_p > 0\}} \gamma_p^N(Q_{p,n} f) - 1_{\{N_{p-1} > 0\}} \gamma_{p-1}^N(Q_{p-1,n} f) \right] \\ &\quad + [\gamma_0^N(Q_{0,n} f) - \gamma_0(Q_{0,n} f)]. \end{aligned} \quad (3.5)$$

In other respects, since

$$1_{\{N_p > 0\}} = 1_{\{N_{p-1} > 0, N_p > 0\}} = 1_{\{N_{p-1} > 0\}} - 1_{\{N_{p-1} > 0, N_p = 0\}},$$

and

$$1_{\{N_{p-1} > 0, N_p = 0\}} \eta_p^N = 0,$$

we conclude that

$$\begin{aligned} 1_{\{N_{p-1} > 0\}} \gamma_{p-1}^N(Q_{p-1,n} f) &= 1_{\{N_{p-1} > 0\}} \gamma_{p-1}^N(1) \eta_{p-1}^N(Q_p(Q_{p-1,n} f)) \\ &= 1_{\{N_{p-1} > 0\}} \gamma_p^N(1) \Phi_p(\eta_{p-1}^N)(Q_{p,n} f), \end{aligned}$$

and

$$\begin{aligned} 1_{\{N_p > 0\}} \gamma_p^N(Q_{p,n} f) &= 1_{\{N_p > 0\}} \eta_p^N(Q_{p,n} f) \gamma_p^N(1) \\ &= 1_{\{N_{p-1} > 0\}} \gamma_p^N(1) \eta_p^N(Q_{p,n} f). \end{aligned}$$

This yields the formula

$$\begin{aligned} &1_{\{N_p > 0\}} \gamma_p^N(Q_{p,n} f) - 1_{\{N_{p-1} > 0\}} \gamma_{p-1}^N(Q_{p-1,n} f) \\ &= 1_{\{N_{p-1} > 0\}} \gamma_p^N(1) [\eta_p^N(Q_{p,n} f) - \Phi_p(\eta_{p-1}^N)(Q_{p,n} f)]. \end{aligned} \quad (3.6)$$

On taking expectation and using (3.3) we thus deduce that

$$\mathbb{E}[\gamma_p^N(Q_{p,n} f) - \gamma_{p-1}^N(Q_{p-1,n} f) | F_{p-1}] = 0.$$

**Proposition 3.1** For any  $n \geq 0$  and bounded function  $f$ , the  $\mathbb{R}$ -valued process  $\Gamma_{\cdot,n}(f)$  defined by

$$\Gamma_{p,n}(f) := 1_{\{N_p > 0\}} \gamma_p^N(Q_{p,n} f) - \gamma_p(Q_{p,n} f), \quad p \leq n,$$

is an  $F$ -martingale with increasing process given by the formula

$$\begin{aligned} \langle \Gamma_{\cdot,n}(f) \rangle_p &= \sum_{q=1}^p 1_{\{N_{q-1} > 0\}} (\gamma_q^N(1))^2 \mathbb{E} \left( [\eta_q^N(Q_{q,n} f) - \Phi_q(\eta_{q-1}^N)(Q_{q,n} f)]^2 | F_{q-1} \right) \\ &\quad + \mathbb{E} \left( [\eta_0^N(Q_{0,n} f) - \eta_0(Q_{0,n} f)]^2 \right). \end{aligned} \quad (3.7)$$

**PROOF.** For all function  $\phi$ , we deduce from (3.5) and (3.6) that

$$\begin{aligned} 1_{\{N_p > 0\}} \gamma_p^N(\phi) - \gamma_p(\phi) &= \sum_{q=1}^p \gamma_q^N(1) 1_{\{N_{q-1} > 0\}} [\eta_q^N(Q_{q,p} \phi) - \Phi_q(\eta_{q-1}^N)(Q_{q,p} \phi)] \\ &\quad + [\eta_0^N(Q_{0,p} \phi) - \eta_0(Q_{0,p} \phi)]. \end{aligned}$$

Therefore by choosing  $\phi = Q_{p,n} f$ , we get

$$\Gamma_{p,n}(f) = \sum_{q=1}^p 1_{\{N_{q-1} > 0\}} \gamma_q^N(1) [\eta_q^N(Q_{q,n} f) - \Phi_q(\eta_{q-1}^N)(Q_{q,n} f)] + [\eta_0^N(Q_{0,n} f) - \eta_0(Q_{0,n} f)],$$

and (3.7) is a clear consequence of the above decomposition.

**Remark 3.2** We can also express  $\Gamma_{p,n}(f)$  as  $\Gamma_{p,n}(f) = \sum_{j \in \mathbb{M}} \Gamma_{p,n}^j(f)$ , with

$$\Gamma_{p,n}^j(f) = 1_{\{|J_{p-1}^j| > 0\}} \gamma_p^{j,N}(f_p) - \gamma_p^j(f_p),$$

since

$$\gamma_p^N(f) = \sum_{j \in \mathbb{M}} 1_{\{|J_{p-1}^j| > 0\}} \gamma_p^{j,N}(f),$$

and

$$1_{\{N_p > 0; |J_{p-1}^j| > 0\}} = 1_{\{\widehat{N}_{p-1} > 0; |J_{p-1}^j| > 0\}} = 1_{\{|J_{p-1}^j| > 0\}}.$$

Furthermore, from (3.5) we deduce that

$$\Gamma_{p,n}^j(f) = \sum_{q=1}^p 1_{\{|J_{q-1}^j| > 0\}} \gamma_q^{j,N}(1) [\eta_q^{j,N}(Q_{q,n} f) - \Phi_q^j(\eta_{q-1}^N)(Q_{q,n} f)] + \omega_0^j [\eta_0^{j,N}(Q_{0,n} f) - \eta_0^j(Q_{0,n} f)].$$

We also can check that for each  $j \in \mathbb{M}$ , the process  $\Gamma_{\cdot,n}^j(f)$  is an  $F$ -martingale.

**Corollary 3.3** For any  $n \geq 0$  and any bounded function  $f$ , we have

$$\mathbb{E}(\gamma_n^N(f) 1_{\{N_n > 0\}}) = \gamma_n(f),$$

and

$$\sup_{f: \|f\| \leq 1} \mathbb{E} \left( [1_{\{N_n > 0\}} \gamma_n^N(f) - \gamma_n(f)]^2 \right) \leq \frac{b^2(n)}{N_{\text{inf}}}.$$

**PROOF.** The first assertion follows from the martingale property of  $(\Gamma_{p,n}(f))_p$  whereas the second follows from the martingale property of  $(\Gamma_{p,n}^2(f) - \langle \Gamma_{\cdot,n}(f) \rangle_p)_p$  and the observation that in view of (3.4)

$$\mathbb{E}(\langle \Gamma_{\cdot,n}(f) \rangle_n) \leq (n+1) \frac{C}{N_{\text{inf}}} \|f\|^2 := \frac{b^2(n)}{N_{\text{inf}}} \|f\|^2.$$

This corollary shows that the IPS approximation  $\gamma_n^N$  of the unnormalized Feynman-Kac measure  $\gamma_n$  has zero bias and mean-square error (or variance) of order  $1/N_{\text{inf}}$ . The end of this section is devoted to showing that similar results hold for the IPS approximation  $\eta_n^N$  of the normalized Feynman-Kac distribution  $\eta_n$  as well: the bias is of order  $1/N_{\text{inf}}$  and the mean square error is also of order  $1/N_{\text{inf}}$ . The second statement follows readily from the decomposition

$$1_{\{N_n > 0\}} \eta_n^N(f) - \eta_n(f) = \frac{1_{\{N_n > 0\}} \gamma_n^N(f) - \gamma_n(f)}{\gamma_n(1)} - 1_{\{N_n > 0\}} \eta_n^N(f) \frac{1_{\{N_n > 0\}} \gamma_n^N(1) - \gamma_n(1)}{\gamma_n(1)},$$

which implies

$$\begin{aligned} \left[ \mathbb{E} \left| 1_{\{N_n > 0\}} \eta_n^N(f) - \eta_n(f) \right|^2 \right]^{1/2} &\leq \left[ \mathbb{E} \left| \frac{1_{\{N_n > 0\}} \gamma_n^N(f) - \gamma_n(f)}{\gamma_n(1)} \right|^2 \right]^{1/2} \\ &\quad + \|f\| \left[ \mathbb{E} \left| \frac{1_{\{N_n > 0\}} \gamma_n^N(1) - \gamma_n(1)}{\gamma_n(1)} \right|^2 \right]^{1/2}, \end{aligned}$$

using the triangle inequality, hence

$$\sup_{f: \|f\| \leq 1} \left[ \mathbb{E} \left| 1_{\{N_n > 0\}} \eta_n^N(f) - \eta_n(f) \right|^2 \right]^{1/2} \leq \frac{2b^2(n)}{(\gamma_n(1))^2 N_{\text{inf}}}.$$

Now, we address the estimation of the bias. Firstly, using Chebyshev's inequality, we also deduce the following corollary



**Corollary 3.4** *We have*

$$\mathbb{P}\left(\gamma_n^N(1)1_{\{N_n > 0\}} \geq \gamma_n(1)/2\right) \geq 1 - \frac{a^2(n)}{N_{\inf}},$$

for some constant  $a(n)$ .

Now, we use these results in the analysis of the normalized particle model. Arguing as in the proof of Theorem 7.4.3 in [4], we write the following decomposition

$$(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}} = \frac{\gamma_n(1)}{\gamma_n^N(1)} \gamma_n^N(f_n) 1_{\{N_n > 0\}}, \quad (3.8)$$

where  $f_n = \frac{1}{\gamma_n(1)}(f - \eta_n(f))$ . Since  $\gamma_n(f_n) = 0$ , then (3.8) also reads

$$(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}} = \frac{\gamma_n(1)}{\gamma_n^N(1)} \left( \gamma_n^N(f_n) 1_{\{N_n > 0\}} - \gamma_n(f_n) \right) 1_{\{N_n > 0\}}.$$

Now by Corollary 3.3, we have  $\mathbb{E}\left(\gamma_n^N(f_n) 1_{\{N_n > 0\}}\right) = \gamma_n(f_n)$  this implies that

$$\begin{aligned} & \mathbb{E}\left[(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}}\right] \\ &= \mathbb{E}\left[\frac{\gamma_n(1)}{\gamma_n^N(1)} \left(1 - \frac{\gamma_n^N(1)}{\gamma_n(1)}\right) 1_{\{N_n > 0\}} \left(\gamma_n^N(f_n) 1_{\{N_n > 0\}} - \gamma_n(f_n)\right)\right]. \end{aligned}$$

If we set  $h_n = \frac{1}{\gamma_n(1)}$ , we get the formula

$$\begin{aligned} \mathbb{E}\left[(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}}\right] &= -\mathbb{E}\left[\frac{\gamma_n(1)}{\gamma_n^N(1)} \left(\gamma_n^N(h_n) 1_{\{N_n > 0\}} - \gamma_n(h_n)\right)\right. \\ &\quad \left. \times \left(\gamma_n^N(f_n) 1_{\{N_n > 0\}} - \gamma_n(f_n)\right) 1_{\{N_n > 0\}}\right]. \end{aligned} \quad (3.9)$$

Now, let  $\Omega_n^N$  be the set of events defined by

$$\Omega_n^N = \{\gamma_n^N(1)1_{\{N_n > 0\}} \geq \gamma_n(1)/2\} = \left\{\frac{\gamma_n(1)}{\gamma_n^N(1)} \leq 2 \text{ and } N_n > 0\right\} \subset \{N_n > 0\}$$

which satisfies  $\mathbb{P}(\Omega_n^N) \geq 1 - a^2(n)/N_{\inf}$  according to Corollary 3.3. If we combine this estimate with (3.9), we find that for any  $f$  with  $\|f\| \leq 1$

$$\begin{aligned} & \left| \mathbb{E}\left[(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}}\right] \right| \\ & \leq \left| \mathbb{E}\left[(\eta_n^N(f) - \eta_n(f)) 1_{\Omega_n^N}\right] \right| + 2\mathbb{P}((\Omega_n^N)^c) \\ & \leq 2\mathbb{E}\left(\left| \gamma_n^N(h_n) 1_{\{N_n > 0\}} - \gamma_n(h_n) \right| \left| \gamma_n^N(f_n) 1_{\{N_n > 0\}} - \gamma_n(f_n) \right|\right) + \frac{b^2(n)}{N_{\inf}}. \end{aligned}$$

By Corollary 3.3 and the Cauchy-Schwarz inequality, this implies that

$$\left| \mathbb{E}\left[(\eta_n^N(f) - \eta_n(f)) 1_{\{N_n > 0\}}\right] \right| \leq \frac{b^2(n)}{N_{\inf}}.$$

Finally, we get that

$$\left| \mathbb{E}\left[\eta_n^N(f) 1_{\{N_n > 0\}} - \eta_n(f)\right] \right| \leq \frac{b^2(n)}{N_{\inf}} + \mathbb{P}(N_n = 0).$$

Now, we need to estimate the extinction probability of the algorithm. In [4, Theorem 7.4.1], a tricky proof gives the following bound for the extinction time  $\tau^N$  of the general particle algorithm

$$\mathbb{P}(\tau^N \geq n) \leq a(n)e^{-N/c(n)}.$$

Nevertheless, we can apply this bound with  $N_{\text{inf}}$  instead of  $N$ , since the extinction time of our algorithm, defined by the first time  $n$  such that all the set  $J_n^j$  are empty, is greater than the extinction time for a classical IPS algorithm. Actually, we can imagine a classical IPS algorithm with only  $N^j$  particles starting all in the mode  $j$ , whose extinction time is clearly less than the extinction time of our algorithm (remember that a mode could be empty at some time, and non empty again at a later time). Hence, we have obtained the following result

**Theorem 3.5** *For each  $n \in \mathbb{N}$  and for any bounded function  $f$ , we have*

$$\sup_{f: \|f\| \leq 1} \left| \mathbb{E} \left[ \eta_n^N(f) 1_{\{N_n > 0\}} - \eta_n(f) \right] \right| \leq \frac{b^2(n)}{N_{\text{inf}}} + a(n) e^{-N_{\text{inf}}/c(n)},$$

and

$$\sup_{f: \|f\| \leq 1} \left[ \mathbb{E} \left| 1_{\{N_n > 0\}} \eta_n^N(f) - \eta_n(f) \right|^2 \right]^{1/2} \leq \frac{2b^2(n)}{(\gamma_n(1))^2 N_{\text{inf}}}.$$

### 3.2 Central limit theorem

We now tackle the central limit theorem (CLT) whose importance for stochastic algorithms is well known. We will follow the approach based on the CLT for triangular arrays introduced in [4]. Nevertheless, as the number of particles is random, we will need to adapt this approach according to [11].

In order to introduce a triangular array of random variables, we split up the terminal term of the martingale  $(\Gamma_{\cdot, n}(f))$  as a sum over the particles, so we obtain

$$\sqrt{N} \Gamma_{n, n}(f) = \sum_{q=1}^n \sum_{\kappa=1}^{N_q} \sqrt{N} \beta_q^\kappa 1_{\{N_{q-1} > 0\}} \gamma_q^N(1) [f_q(\xi_q^\kappa) - \Phi_q(\eta_{q-1}^N)(f_q)] + \sqrt{N} [\eta_0^N(f_0) - \eta_0(f_0)],$$

with  $f_q := Q_{q, n} f$  for  $0 \leq q \leq n$ . Introducing the random variables

$$\begin{aligned} X_{q, \kappa}^N &:= \sqrt{N} \beta_q^\kappa 1_{\{N_{q-1} > 0\}} \gamma_q^N(1) [f_q(\xi_q^\kappa) - \Phi_q(\eta_{q-1}^N)(f_q)], \quad 1 \leq q \leq n, \\ X_{0, \kappa}^N &:= \sqrt{N} \beta_0^\kappa [f_0(\xi_0^\kappa) - \eta_0(f_0)], \end{aligned}$$

we can rewrite  $\sqrt{N} \Gamma_{n, n}(f)$  in the following form

$$\sqrt{N} \Gamma_{n, n}(f) = \sum_{q=0}^n \sum_{\kappa=1}^{N_q} X_{q, \kappa}^N.$$

Furthermore, we can write also

$$\sqrt{N} \Gamma_{n, n}(f) = \sum_{j \in \mathbb{M}} \sqrt{N/N^j} \sqrt{N^j} \Gamma_{n, n}^j(f),$$

with

$$\begin{aligned} \sqrt{N^j} \Gamma_{n, n}^j(f) &= \frac{1}{\sqrt{N^j}} \sum_{q=1}^n \sum_{\kappa \in \widehat{J}_{q-1}^j} 1_{\{|\widehat{J}_{q-1}^j| > 0\}} \gamma_q^{j, N}(1) [f_q(\xi_q^\kappa) - \Phi_q^j(\eta_{q-1}^N)(f_q)] \\ &\quad + \frac{1}{\sqrt{N^j}} \sum_{\kappa \in J_0^j} \omega_0^j [f_0(\xi_0^\kappa) - \eta_0^j(f_0)]. \end{aligned}$$

We recall that  $|\widehat{J}_{q-1}^j| = N^j$ , and by introducing the random variables

$$\begin{aligned} X_{q, \kappa}^{j, N} &:= \frac{1}{\sqrt{N^j}} 1_{\{|\widehat{J}_{q-1}^j| > 0\}} \gamma_q^{j, N}(1) [f_q(\xi_q^\kappa) - \Phi_q^j(\eta_{q-1}^N)(f_q)], \quad 1 \leq q \leq n, \\ X_{0, \kappa}^{j, N} &:= \frac{1}{\sqrt{N^j}} \omega_0^j [f_0(\xi_0^\kappa) - \eta_0^j(f_0)], \end{aligned}$$

we obtain the following form

$$\sqrt{N^j} \Gamma_{n,n}^j(f) = \sum_{q=1}^n \sum_{\kappa \in \widehat{J}_{q-1}^j} X_{q,\kappa}^{j,N} + \sum_{\kappa \in J_0^j} X_{0,\kappa}^{j,N}.$$

Finally, let us mention that  $X_{q,\kappa}^N = \sqrt{N/N^j} X_{q,\kappa}^{j,N}$  for  $\kappa \in \widehat{J}_{q-1}^j$ .

Henceforth, at the end of each selection/resampling step, the indices of each particle will be ordered according to the order induced by the modes. More precisely, setting  $N_q = \sum_{j \in \mathbb{M}} N_q^j$  with  $N_q^j = N^j$  if  $j \in J_{q-1}$  and  $N_q^j = 0$  otherwise, we label the  $N_q = \widehat{N}_{q-1}$  particles  $(\widehat{\xi}_{q-1}^\kappa)$  in such a way that  $\kappa \in \widehat{J}_{q-1}^j$  if and only if the interval

$$I_j := \left[ \sum_{i=1}^{j-1} N_q^i + 1, \sum_{i=1}^j N_q^i \right] \quad \text{with } N_q^0 := 0$$

is not empty and  $\kappa \in I_j$ . We remind the reader that for  $\kappa \in \widehat{J}_{q-1}^j$  a particle  $\xi_q^\kappa$  is an excursion started in  $D_{q-1} \times \{j\}$  with  $j \in J_{q-1}$ . Now, let us introduce some new notations: firstly, we set

$$K_q^N = N_0 + \dots + N_q, \quad 0 \leq q \leq n,$$

with the convention  $K_{-1}^N = 0$ . Secondly, we subdivide each interval  $[K_{q-1}^N + 1, K_q^N]$  in a the reunion of the mutually disjoint sub-intervals  $[K_{q,j-1}^N + 1, K_{q,j}^N]$  ( $j \in \mathbb{M}$ ), where

$$K_{q,j}^N = K_{q-1}^N + N_q^1 + \dots + N_q^j, \quad K_{q,0}^N = K_q^N \quad \text{and} \quad K_{q,M}^N = K_{q+1}^N.$$

Obviously, for any  $j \notin J_{q-1}$ , such an interval is empty.

We notice that the  $\kappa$ -th particle within the  $q$ -th generation can be associated in a unique way with an integer  $k$  between 1 and  $K_n^N$  and a mode  $j \in J_{q-1}$ : clearly  $k_{q,\kappa} = K_{q-1}^N + \kappa$  and  $j = \inf\{i : K_{q,i}^N \geq k\}$ . Conversely, for a fixed integer  $1 \leq k \leq K_n^N$ , the random integers  $q_k$ ,  $\kappa_k$  and  $j_k$  are defined by

$$q_k = \inf\{q \geq 0 : K_q^N \geq k\}, \quad \kappa_k = k - K_{q_k-1}^N \quad \text{and} \quad j_k = \inf\{j : K_{q_k,j}^N \geq k\},$$

or in other words  $q_k = q$  and  $\kappa_k = \kappa$  if and only if

$$K_{q-1}^N + 1 \leq k = K_{q-1}^N + \kappa \leq K_q^N.$$

We also check that the set of particles  $\xi_q^\kappa$  such that  $\kappa \in \widehat{J}_{q-1}^j$  for any  $j \in J_{q-1}$  is the set of particles whose index  $\kappa$  belongs to the interval  $[K_{q,j-1}^N + 1, K_{q,j}^N]$  which is non empty as soon as  $j \in J_{q-1}$ .

Now, we introduce a filtration  $\mathcal{G}^N = \{\mathcal{G}_k^N, k \geq 1\}$  in such a way that  $K_n^N$  is a stopping time w.r.t.  $\mathcal{G}^N$ . For any  $q = 0, 1, \dots, n$  and any integer  $\kappa \geq 1$ , let  $\mathcal{F}_{q,\kappa}^N = \mathcal{F}_{q,0}^N \vee \sigma(\xi_q^1, \dots, \xi_q^\kappa)$ , where  $\mathcal{F}_{0,0}^N = \{\emptyset, \Omega\}$  and  $\mathcal{F}_{q,0}^N = F_{q-1}$  (for  $q = 1, \dots, n$ ). Since  $\{N_q = p\} \in \mathcal{F}_{q,p}^N$  by definition, the random number  $N_q$  is a stopping time w.r.t. the filtration  $\mathcal{F}_q^N = \{\mathcal{F}_{q,\kappa}^N, \kappa \geq 0\}$ , which allows to define the  $\sigma$ -algebra  $\mathcal{F}_{q,N_q}^N = F_q$ . Therefore, the random variable  $K_q^N$  is measurable w.r.t.  $F_q$ .

For any  $q = 0, 1, \dots, n$  and any integer  $\kappa \geq 1$ ,

$$\{q_k = q, \kappa_k = \kappa\} = \{k = K_{q-1}^N + \kappa, \kappa \leq N_q\} \subset \mathcal{F}_{q,\kappa-1}^N \subset \mathcal{F}_{q,\kappa}^N,$$

since  $\{k = K_{q-1}^N + \kappa\} \in F_{q-1}$  and  $\{\kappa \leq N_q\} \in \mathcal{F}_{q,\kappa-1}^N$ . Then, we can define in the usual way the  $\sigma$ -algebra  $\mathcal{G}_k^N = \mathcal{F}_{q_k,\kappa_k}^N$  by:  $A \in \mathcal{G}_k^N$  if and only if  $A \cap \{q_k = q, \kappa_k = \kappa\} \in \mathcal{F}_{q,\kappa}^N$ , for any  $q = 0, 1, \dots, n$  and any integer  $\kappa \geq 1$ . Using this new labelling of the particle system yields

$$\sqrt{N} \Gamma_{n,n}(f) = \sum_{k=1}^{K_n^N} U_k^N,$$

where  $U_k^N := X_{q_k,\kappa_k}^N$  is measurable w.r.t.  $\mathcal{G}_k^N$ , for any  $k = 1, \dots, K_n^N$ : indeed for any Borel subset  $B$

$$\{U_k^N \in B\} \cap \{q_k = q, \kappa_k = \kappa\} = \{X_{q,\kappa}^N \in B\} \cap \{q_k = q, \kappa_k = \kappa\},$$

hence  $\{U_k^N \in B\} \in \mathcal{G}_k^N$ , since  $\{X_{q,\kappa}^N \in B\} \in \mathcal{F}_{q,\kappa}^N$  and  $\{q_k = q, \kappa_k = \kappa\} \in \mathcal{F}_{q,\kappa-1}^N$ .

Moreover, the random variable  $K_n^N$  is a stopping time w.r.t.  $\mathcal{G}^N$ , since

$$\begin{aligned} \{K_n^N = k\} \cap \{q_k = q, \kappa_k = \kappa\} &= \{K_n^N = k\} \cap \{k = K_{q-1}^N + \kappa, 1 \leq \kappa \leq N_q\}, \\ &= \begin{cases} \emptyset & \text{if } q \neq n, \\ \{k = K_{n-1}^N + \kappa\} \cap \{N_n = \kappa\} & \text{if } q = n, \end{cases} \end{aligned}$$

hence  $\{K_n^N = k\} \in \mathcal{G}_k^N$  since  $\{k = K_{n-1}^N + \kappa\} \in \mathcal{F}_{n-1}$  and  $\{N_n = \kappa\} \in \mathcal{F}_{n,\kappa}^N$ . We also obtain the following expression

$$\begin{aligned} \sqrt{N} \Gamma_{n,n}(f) &= \sum_{q=0}^n \sum_{j \in \mathbb{M}} \sum_{k=K_{q,j-1}^N+1}^{K_{q,j}^N} U_k^N \\ &= \sum_{q=0}^n \sum_{j \in \mathbb{M}} \sum_{\kappa \in \mathcal{J}_{q-1}^j} X_{q,\kappa}^N \\ &= \sum_{q=0}^n \sum_{j \in \mathbb{M}} \sqrt{N/N^j} \sum_{\kappa \in \mathcal{J}_{q-1}^j} X_{q,\kappa}^{j,N}. \end{aligned}$$

Now, since  $K_n^N < \infty$  a.s., to apply the Theorem VIII.3.33 in [8] we need to check the following three conditions

- (i)  $\mathbb{E}(U_k^N | \mathcal{G}_{k-1}^N) = 0$ ,
- (ii)  $\sum_{k=1}^{K_n^N} \mathbb{E}(|U_k^N|^2 | \mathcal{G}_{k-1}^N) \xrightarrow{P} \sigma^2$ ,
- (iii) for all  $\varepsilon > 0$ ,  $\sum_{k=1}^{K_n^N} \mathbb{E}\left(|U_k^N|^2 \mathbf{1}_{\{|U_k^N| > \varepsilon\}} | \mathcal{G}_{k-1}^N\right) \xrightarrow{P} 0$ .

However, since  $U_k^N = X_{q_k, \kappa_k}^N$  is a time changed random variable, we need the following result [11, Lemma 4, Corollaries 1 and 2]

**Lemma 3.6** *If for any  $q = 0, 1, \dots, n$  and any integer  $\kappa \geq 1$*

$$\mathbb{E}[F_{q,\kappa} | \mathcal{F}_{q,\kappa-1}^N] = \widehat{F}_q,$$

where the random variable  $\widehat{F}_q$  is measurable w.r.t.  $\mathcal{F}_{q,0}^N$ , then for any integer  $k \geq 1$ , the time changed random variables  $G_k = F_{q_k, \kappa_k}$  and  $\widehat{G}_k = \widehat{F}_{q_k}$  satisfy

$$\mathbb{E}[G_k | \mathcal{G}_{k-1}^N] = \widehat{G}_k.$$

If for any  $q = 0, 1, \dots, n$  and any integer  $\kappa \geq 1$

$$F_{q,\kappa} \leq F_q^*,$$

where the random variable  $F_q^*$  is measurable w.r.t.  $\mathcal{F}_{q,0}^N$ , then for any integer  $k \geq 1$ , the time changed random variables  $G_k = F_{q_k, \kappa_k}$  and  $G_k^* = F_{q_k}^*$  satisfy

$$\mathbb{E}[G_k | \mathcal{G}_{k-1}^N] \leq G_k^*.$$

Firstly, for any  $\kappa = 1, \dots, N_0$ , the random variable  $X_{0,\kappa}^N$  is measurable w.r.t.  $\mathcal{F}_{0,\kappa}^N$ . Moreover,

$$\mathbb{E}[X_{0,\kappa}^N | \mathcal{F}_{0,\kappa-1}^N] = 0,$$

and for any  $\kappa \in \mathcal{J}_0^j$

$$\mathbb{E}\left[|X_{0,\kappa}^{j,N}|^2 | \mathcal{F}_{0,\kappa-1}^N\right] = \frac{1}{N^j} (\omega_0^j)^2 \text{var}(f_0, \eta_0^j) := V_{0,j}^N.$$

Notice that

$$|X_{0,\kappa}^{j,N}| \leq \frac{1}{\sqrt{N^j}} 2\|f_0\|,$$

hence for any  $\varepsilon > 0$

$$|X_{0,\kappa}^{j,N}|^2 \mathbf{1}_{\{|X_{0,\kappa}^{j,N}| > \varepsilon\}} \leq \frac{1}{N^j} (2\|f_0\|)^2 \mathbf{1}_{\{\frac{1}{\sqrt{N^j}} 2\|f_0\| > \varepsilon\}} := Y_{0,j}^N(\varepsilon).$$

For any  $q = 1, \dots, n$ , and any  $j \in J_{q-1}$ , the random weight  $\gamma_q^{j,N}(1)$  is measurable w.r.t.  $F_{q-1} = \mathcal{F}_{q,0}^N$ . Furthermore, for any  $\kappa \in \widehat{J}_{q-1}^j$  the random variable  $X_{q,\kappa}^{j,N}$  is measurable w.r.t.  $\mathcal{F}_{q,\kappa}^N$ , where conditionally w.r.t. the  $\sigma$ -algebra  $F_{q-1}$ , the particles  $\xi_\kappa^j$ , with  $\kappa \in \widehat{J}_{q-1}^j$ , are i.i.d. with common distribution  $\Phi_q^j(\eta_{q-1}^N)$ . Moreover,

$$\mathbb{E}[X_{q,\kappa}^{j,N} | \mathcal{F}_{q,\kappa-1}^N] = 0,$$

and for  $\kappa \in \widehat{J}_{q-1}^j$

$$\mathbb{E}[|X_{q,\kappa}^{j,N}|^2 | \mathcal{F}_{q,\kappa-1}^N] = \frac{1}{N^j} \mathbf{1}_{\{|J_{q-1}^j| > 0\}} (\gamma_q^{j,N}(1))^2 \text{var}(f_q, \Phi_q^j(\eta_{q-1}^N)) := V_{q,j}^N,$$

where the random variables  $V_{q,j}^N$  are measurable w.r.t.  $\mathcal{F}_{q,0}^N$ . Notice that

$$|X_{q,\kappa}^{j,N}| \leq \frac{\gamma_q^{j,N}(1)}{\sqrt{N^j}} 2\|f_q\|,$$

hence for any  $\varepsilon > 0$ ,

$$|X_{q,\kappa}^{j,N}|^2 \mathbf{1}_{\{|X_{q,\kappa}^{j,N}(f)| > \varepsilon\}} \leq \frac{(\gamma_q^{j,N}(1))^2}{N^j} (2\|f_q\|)^2 \mathbf{1}_{\{\frac{\gamma_q^{j,N}(1)}{\sqrt{N^j}} 2\|f_q\| > \varepsilon\}} := Y_{q,j}^N(\varepsilon),$$

where the random variable  $Y_{q,j}^N(\varepsilon)$  is measurable w.r.t.  $\mathcal{F}_{q,0}^N$ . It follows from Lemma 3.6 that

$$\mathbb{E}(U_k^N | \mathcal{G}_{k-1}^N) = 0$$

$$\mathbb{E}(|U_k^N|^2 | \mathcal{G}_{k-1}^N) = (N/N^{j_k}) V_{q_k, j_k}^N, \quad \text{if } \kappa_k \in J_{q_k-1}^{j_k},$$

and

$$\mathbb{E}\left(|U_k^N|^2 \mathbf{1}_{\{|U_k^N| > \varepsilon\}} | \mathcal{G}_{k-1}^N\right) \leq (N/N^{j_k}) Y_{q_k, j_k}^N(\varepsilon \sqrt{N^{j_k}/N}),$$

hence,

$$\sum_{k=1}^{K_n^N} \mathbb{E}(|U_k^N|^2 | \mathcal{G}_{k-1}^N) = \sum_{j \in \mathbb{M}} N V_{0,j}^N + \sum_{q=1}^n \sum_{j \in J_{q-1}} N V_{q,j}^N$$

and

$$\sum_{k=1}^{K_n^N} \mathbb{E}\left(|U_k^N|^2 \mathbf{1}_{\{|U_k^N| > \varepsilon\}} | \mathcal{G}_{k-1}^N\right) \leq \sum_{q=0}^n \sum_{j \in \mathbb{M}} N Y_{q,j}^N(\varepsilon \sqrt{N^j/N}).$$

We deduce from Corollary 3.3 and Theorem 3.5 that  $\gamma_q^{j,N}(1) \mathbf{1}_{\{|J_{q-1}^j| > 0\}} \rightarrow \gamma_q^j(1)$  and  $\text{var}(f_q, \Phi_q^j(\eta_{q-1}^N)) \rightarrow \text{var}(f_q, \Phi_q^j(\eta_{q-1}))$ . Now, let  $\rho_j := N^j/N$  and let  $N \rightarrow \infty$  in such a way that for any  $j \in \mathbb{M}$ , the ratio  $\rho_j$  is preserved, or at least the ratio  $N^j/N \rightarrow \rho_j$ . Then, we deduce that

$$N V_{0,j}^N \rightarrow \frac{(\omega_0^j)^2}{\rho_j} \text{var}(f_0, \eta_0^j),$$

and

$$NV_{q,j}^N \rightarrow \frac{(\gamma_q^j(1))^2}{\rho_j} \text{var}(f_q, \eta_q^j)$$

in probability. We remark also that  $|J_{q-1}^j| > 0$  if and only if  $\eta_{q-1}^N(g_{q-1}^j) > 0$ , but this probability converges to  $P_{q-1}^j$  as  $N$  tends to infinity, so if we assume  $P_{q-1}^j \neq 0$  (a reasonable assumption), we can conclude that  $\sum_{k=1}^{K_n^N} \mathbb{E}(|U_k^N|^2 | \mathcal{G}_{k-1}^N) \rightarrow W_n(f)$ , with

$$\begin{aligned} W_n(f) &= \sum_{j \in \mathbb{M}} \left( \sum_{q=1}^n \frac{(\gamma_q^j(1))^2}{\rho_j} \text{var}(f_q, \eta_q^j) + \frac{(\omega_0^j)^2}{\rho_j} \text{var}(f_0, \eta_0^j) \right) \\ &= (\gamma_n(1))^2 \sum_{q=0}^n \sum_{j \in \mathbb{M}} \frac{(\omega_{q-1}^j)^2}{\rho_j} \frac{\text{var}(f_q, \eta_q^j)}{\eta_q^2(Q_{q,n}1)}. \end{aligned}$$

To obtain the last line, we used the formula  $\gamma_q^j(1) = \omega_{q-1}^j \gamma_n(1) / \eta_q(Q_{q,n}1)$  valid for  $q \geq 1$ , and the convention  $\omega_{-1}^j = \omega_0^j$ . Moreover, we can check that  $\sum_{q=0}^n \sum_{j \in \mathbb{M}} N Y_{q,j}^N(\varepsilon \sqrt{N/N^j}) \rightarrow 0$  in probability. So, we have proved the CLT.

**Theorem 3.7** *Let  $N$  tends to  $\infty$  in such a way that the ratios  $\rho_j = N^j/N$  are preserved for all  $j \in \mathbb{M}$ . Then, the sequence of random variables*

$$\sqrt{N} \left( \mathbf{1}_{\{N_n > 0\}} \gamma_n^N(g_n) - \mathbb{P}(T_n \leq T) \right)$$

converges in law to a Gaussian random variable with mean 0 and variance  $W_n := W_n(g_n)$ .

**Remark 3.8** The asymptotic variance of the classical algorithm i.e. without sampling per mode is given by [3]

$$V_n(f) = (\gamma_n(1))^2 \sum_{q=0}^n \frac{\text{var}(f_q, \eta_q)}{\eta_q^2(Q_{q,n}1)}.$$

But, we have

$$\text{var}(f_q, \eta_q) = \sum_{j \in \mathbb{M}} \omega_{q-1}^j \text{var}(f_q, \eta_q^j) + \sum_{j \in \mathbb{M}} \omega_{q-1}^j [(\eta_q^j(f_q))^2 - \eta_q^2(f_q)],$$

then  $V_n(f)$  can be written as

$$\frac{V_n(f)}{(\gamma_n(1))^2} = \sum_{q=0}^n \sum_{j \in \mathbb{M}} \omega_{q-1}^j \frac{\text{var}(f_q, \eta_q^j)}{\eta_q^2(Q_{q,n}1)} + \frac{1}{2} \sum_{q=0}^n \sum_{j,l \in \mathbb{M}} \omega_{q-1}^j \omega_{q-1}^l \frac{[\eta_q^j(f_q) - \eta_q^l(f_q)]^2}{\eta_q^2(Q_{q,n}1)}.$$

Consequently, we check that

$$W_n(f) \leq \max_{0 \leq q \leq n-1} \max_{j \in \mathbb{M}} \left( \frac{\omega_q^j}{\rho_j} \right) V_n(f),$$

hence, if we are able to adapt, at each step  $q$ , the number of resampled particles  $N_q^j$  in order that  $\omega_q^j \approx N_q^j/N$ , we will obtain an asymptotic variance less than the asymptotic variance of the classical algorithm.

**Remark 3.9** For any  $q \geq 0$ , the function  $g_{q,n} := Q_{q,n} g_n$  is defined for any excursion  $e = (z(u), s \leq u \leq t)$  by

$$g_{q,n}(e) = g_q(e) \mathbb{P}(T_n \leq T | (T_q, Z_{T_q}) = \pi(e)).$$

Setting

$$\Delta_q^n(t, z) = \mathbb{P}(T_n \leq T | T_q = t, Z_{T_q} = z),$$

and introducing the entrance distribution  $\mu_q := \hat{\eta}_q \circ \pi^{-1}$  give the following identities, since  $g_q^2 = g_q$ ,

$$\begin{aligned} \eta_q(g_{q,n}) &= \eta_q(g_q \Delta_q^n \circ \pi) = \eta_q(g_q) \hat{\eta}_q(\Delta_q^n \circ \pi) = P_q \mu_q(\Delta_q^n), \\ \eta_q[g_{q,n}^2] &= \eta_q(g_q [\Delta_q^n \circ \pi]^2) = \eta_q(g_q) \hat{\eta}_q([\Delta_q^n \circ \pi]^2) = P_q \mu_q([\Delta_q^n]^2). \end{aligned}$$

Moreover, from (2.1), we also get

$$\eta_q(g_{q,n}) = \sum_{j \in \mathbb{M}} P_q^j \mu_q^j(\Delta_q^n), \quad \eta_q(g_{q,n}^2) = \sum_{j \in \mathbb{M}} P_q^j \mu_q^j([\Delta_q^n]^2),$$

with  $\mu_q^j := \widehat{\eta}_q^j \circ \pi^{-1}$ . We also notice that, since  $\Delta_{q-1}^n \circ \pi = \mathcal{M}_q g_{q,n}$ , we obtain

$$\eta_q^j(g_{q,n}) = \widehat{\eta}_{q-1}^j(\mathcal{M}_q g_{q,n}) = \mu_{q-1}^j(\Delta_{q-1}^n).$$

However, we have no so simple expression for  $\eta_q^j(g_{q,n}^2)$ , except perhaps by introducing the Markov kernel  $R_q^{(n)}$  defined by [4, Proposition 2.7.1], for any function  $f_q$  acting on excursions  $e$  such that  $\pi(e) \in A_{q-1}$

$$R_q^{(n)} f_q = \frac{\mathcal{M}_q(g_{q,n} f_q)}{\mathcal{M}_q g_{q,n}},$$

so that we can write

$$\eta_q^j(g_{q,n}^2) = \widehat{\eta}_{q-1}^j(\Delta_{q-1}^n \circ \pi R_q^{(n)} g_{q,n}) = \mu_{q-1}^j(\Delta_{q-1}^n R_q^{(n)} g_{q,n}).$$

As with stratified sampling, a small asymptotic variance can be obtained only if the function  $g_{q,n}$  is relatively homogeneous for each measure  $\eta_q^j$ , i.e. if  $\text{var}(g_{q,n}, \eta_q^j)$  is relatively small for each  $j$  and each  $q$ . This is possible if and only if we have a great variability between the values  $(\eta_q^j(g_{q,n}))_{j \in \mathbb{M}}$  for each  $q$ ; but the sampling per mode algorithm has been introduced precisely in order to improve the classical algorithm in that case.

## 4 Conclusion

In this paper we considered the rare event simulation problem for a switching diffusion, using a multilevel splitting method adapted to the discrete modes: the conditional "sampling per mode" algorithm. Using the Feynman-Kac and interacting particle theory, we established a law of large numbers and a central limit theorem for the estimate of the rare event probability and thus we confirmed that this algorithm has a better asymptotic variance as soon as the probability to hit the target set fluctuates according to the mode. We also observe that an adaptive algorithm which will update the number of resampled particles  $N_q^j$ , at each step  $q$ , in order that  $\omega_q^j \approx N_q^j/N$ , will give a lower asymptotic variance.

It will also be valuable to deduce from the expression of the asymptotic variance some informations about the tuning of the algorithm; for example the optimal number  $N^j$  of particles in each mode. To improve again the quality of our estimate, an importance sampling technique can be added in order to make the rare switches more frequent. This idea has already been explored in [9], and now we think that it will be possible to obtain a central limit theorem also for this kind of algorithm. Let also mention that a possible Rao-Blackwellisation strategy could be investigated, based on a partition of the state space in the continuous component and the discrete modes.

## References

- [1] Henk Blom. Stochastic hybrid processes with hybrid jumps. In *IFAC Conference on Analysis and Design of Hybrid Systems (ADHS03), April 2003, HYBRIDGE deliverable R2.3*, April 2003. <http://www.nlr.nl/public/hostedsites/hybridge/>.
- [2] Henk Blom, Bert Bakker, and Jaroslav Krystul. Rare event estimation for a large-scale stochastic hybrid system with air traffic application. In Gerardo Rubino and Bruno Tuffin, editors, *Rare Event Simulation using Monte Carlo Methods*, pages 194–214. Wiley, 2009.
- [3] Frédéric Cérou, Pierre Del Moral, François Le Gland, and Pascal Lezaud. Genetic genealogical models in rare event analysis. *ALEA, Latin American Journal of Probability and Mathematical Statistics*, 1:181–203, 2006. Paper 01–08.
- [4] Pierre Del Moral. *Feynman-Kac Formulae. Genealogical and Interacting Particle Systems with Applications*. Probability and its Applications. Springer, New York, 2004.

- 
- [5] Pierre Del Moral and Pascal Lezaud. Branching and interacting particle interpretation of rare event probabilities. In Henk Blom and John Lygeros, editors, *Stochastic Hybrid Systems : Theory and Safety Critical Applications*, number 337 in Lecture Notes in Control and Information Sciences, pages 277–323. Springer, Berlin, 2006.
- [6] Marnix J. J. Garvels. *The Splitting Method in Rare Event Simulation*. Ph.D Thesis, Faculty of Mathematical Sciences, University of Twente, Enschede, October 2000.
- [7] Paul Glasserman, Philip Heidelberger, Perwez Shahabuddin, and Tim Zajic. Multilevel splitting for estimating rare event probabilities. *Operations Research*, 47(4):585–600, July–August 1999.
- [8] Jean Jacod and Albert N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer, second edition, 2003.
- [9] Jaroslav Krystul. *Modelling of Stochastic Hybrid Systems with Applications to Accident Risk Assessment*. Ph.D Thesis, Faculty of Mathematical Sciences, University of Twente, Enschede, September 2006.
- [10] Agnès Lagnoux. Rare event simulation. *Probability in the Engineering and Information Sciences*, 20(1):45–66, January 2006.
- [11] François Le Gland and Nadia Oudjane. A sequential algorithm that keeps the particle system alive. In Henk Blom and John Lygeros, editors, *Stochastic Hybrid Systems : Theory and Safety Critical Applications*, number 337 in Lecture Notes in Control and Information Sciences, pages 351–389. Springer, Berlin, 2006.
- [12] Pierre L’Écuyer, François Le Gland, Pascal Lezaud, and Bruno Tuffin. Splitting techniques. In Gerardo Rubino and Bruno Tuffin, editors, *Rare Event Simulation using Monte Carlo Methods*, pages 39–61. Wiley, 2009.
- [13] Pierre L’Écuyer, Michel Mandjes, and Bruno Tuffin. Importance sampling. In Gerardo Rubino and Bruno Tuffin, editors, *Rare Event Simulation using Monte Carlo Methods*, pages 17–38. Wiley, 2009.
- [14] Manuel Villén-Altamirano and José Villén-Altamirano. RESTART : A straightforward method for fast simulation of rare events. In *Proceedings of the 1994 Winter Simulation Conference, Orlando 1994*, pages 282–289, December 1994.
- [15] Manuel Villén-Altamirano and José Villén-Altamirano. Analysis of RESTART simulation : Theoretical basis and sensitivity study. *European Transactions on Telecommunications*, 13(4 (Special issue on Rare Event Simulation)):373–385, July–August 2002.
- [16] Manuel Villén-Altamirano and José Villén-Altamirano. On the efficiency of RESTART for multidimensional state systems. *ACM Transactions on Modeling and Computer Simulation*, 16(3):251–279, July 2006.
- [17] Chao Zhu and George Yin. Asymptotic properties of hybrid diffusion systems. *SIAM J. Control Optim.*, 46(4):1155–1179, 2007.





---

Centre de recherche INRIA Rennes – Bretagne Atlantique  
IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Centre de recherche INRIA Bordeaux – Sud Ouest : Domaine Universitaire - 351, cours de la Libération - 33405 Talence Cedex  
Centre de recherche INRIA Grenoble – Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier  
Centre de recherche INRIA Lille – Nord Europe : Parc Scientifique de la Haute Borne - 40, avenue Halley - 59650 Villeneuve d'Ascq  
Centre de recherche INRIA Nancy – Grand Est : LORIA, Technopôle de Nancy-Brabois - Campus scientifique  
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex  
Centre de recherche INRIA Paris – Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex  
Centre de recherche INRIA Saclay – Île-de-France : Parc Orsay Université - ZAC des Vignes : 4, rue Jacques Monod - 91893 Orsay Cedex  
Centre de recherche INRIA Sophia Antipolis – Méditerranée : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex

---

Éditeur  
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)  
<http://www.inria.fr>  
ISSN 0249-6399