



Steinberg's Conjecture and near-colorings

Gerard Chang, Frédéric Havet, Mickael Montassier, André Raspaud

► **To cite this version:**

Gerard Chang, Frédéric Havet, Mickael Montassier, André Raspaud. Steinberg's Conjecture and near-colorings. [Research Report] RR-7669, INRIA. 2011. <inria-00605810>

HAL Id: inria-00605810

<https://hal.inria.fr/inria-00605810>

Submitted on 4 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Steinberg's Conjecture and near-colorings

Gerard J. Chang

Frédéric Havet Mickael Montassier André Raspaud

N° 7669

July 2011

Thème COM



*R*apport
de recherche

Steinberg's Conjecture and near-colorings*

Gerard J. Chang[†]
Frédéric Havet[‡] Mickael Montassier[§] André Raspaud[§]

Thème COM — Systèmes communicants
Équipe-Projet Mascotte

Rapport de recherche n° 7669 — July 2011 — 13 pages

Abstract: Let \mathcal{F} be the family of planar graphs without cycles of length 4 and 5. Steinberg's Conjecture (1976) that says every graph of \mathcal{F} is 3-colorable remains widely open. Motivées par une relaxation proposée par Erdős (1991), plusieurs études ont montré la conjecture pour des sous-classes de \mathcal{F} . Par exemple, Borodin *et al.* ont prouvé que tout graphe planaire sans cycles de longueur 4 à 7 est 3-colorable. Dans ce rapport, nous relaxons le problème non pas sur la classe de graphes mais sur le type de coloration en considérant des *quasi-colorations*. Un graphe $G = (V, E)$ est dit (i, j, k) -colorable si son ensemble de sommet peut être partitionner en trois ensembles V_1, V_2, V_3 tels que les graphes $G[V_1], G[V_2], G[V_3]$ induits par ces ensembles soit de degré maximum au plus i, j, k respectivement. Avec cette terminologie, la Conjecture de Steinberg dit que tout graphe de \mathcal{F} est $(0, 0, 0)$ -colorable. Un résultat de Xu (2008) implique que tout graphe de \mathcal{F} est $(1, 1, 1)$ -colorable. Nous montrons ici que tout graphe de \mathcal{F} est $(2, 1, 0)$ -colorable et $(4, 0, 0)$ -colorable.

Key-words: graphs, coloring, decomposition, Steinberg's conjecture

* Supported by ANR/NSC projects ANR-09-blanc-0373-01 and NSC99-2923-M-002-007-MY3.

[†] Department of Mathematics and Taida Institute for Mathematical Sciences, National Taiwan University, Taipei 10617, and National Center for Theoretical Sciences, Taipei, Taiwan. E-mail: gjchang@math.ntu.edu.tw.

[‡] Projet MASCOTTE – I3S (CNRS & UNS) and INRIA Sophia-Antipolis, 2004 route des lucioles BP93, 06902 Sophia-Antipolis Cedex, France. E-mail: Frederic.Havet@sophia.inria.fr.

[§] LaBRI – University of Bordeaux, 351 cours de la libération, 33405 Talence Cedex, France.
E-mails: mickael.montassier@labri.fr, andre.raspaud@labri.fr.

Conjecture de Steinberg et quasi-coloration

Résumé : Soit \mathcal{F} la classe des graphes planaires sans cycles de longueur 4 et 5. La Conjecture de Steinberg (1976) affirmant que tout graphe de \mathcal{F} est 3-colorable, reste largement ouverte.

Mots-clés : graphes, coloration, décomposition, conjecture de Steinberg

1 Introduction

In 1976, Appel and Haken proved that every planar graph is 4-colorable [2, 3], and as early as 1959, Grötzsch [20] showed that every planar graph without 3-cycles is 3-colorable. As proved by Garey, Johnson and Stockmeyer [19], the problem of deciding whether a planar graph is 3-colorable is NP-complete. Therefore, some sufficient conditions for planar graphs to be 3-colorable were stated. In 1976, Steinberg [24] raised the following:

Steinberg's Conjecture '76 *Every planar graph without 4- and 5-cycles is 3-colorable.*

There were then no progress in this direction until Erdős (1991) proposed the following relaxation of Steinberg's Conjecture:

Erdős' relaxation '91 Determine the smallest value of k , if it exists, such that every planar graph without cycles of length from 4 to k is 3-colorable.

Abbott and Zhou [1] proved that such a k does exist, with $k \leq 11$. This result was later on improved to $k \leq 10$ by Borodin [4], to $k \leq 9$ by Borodin [5] and Sanders and Zhao [22], to $k \leq 8$ by Salavatipour [21]. The best known bound for such a k is 7 which was proved by Borodin, Glebov, Raspaud and Salavatipour [10].

This approach was at the origin of sufficient conditions of 3-colorability of subfamilies of planar graphs where some families of cycles are forbidden. See for examples [8, 9, 12, 13, 14, 15, 16, 17, 25].

A graph G is called *improperly* (d_1, d_2, \dots, d_k) -colorable, or simply (d_1, d_2, \dots, d_k) -colorable, if the vertex set of G can be partitioned into subsets V_1, V_2, \dots, V_k such that the graph $G[V_i]$ induced by V_i has maximum degree at most d_i for $1 \leq i \leq k$. This notion generalizes those of proper k -coloring (when $d_1 = d_2 = \dots = d_k = 0$) and d -improper k -coloring (when $d_1 = d_2 = \dots = d_k = d \geq 0$). Under this terminology, the Four Color Theorem says that every planar graph is $(0, 0, 0, 0)$ -colorable. Eaton and Hull [18] and independently Škrekovski [23] proved that every planar graph is 2-improperly 3-colorable (in fact, 2-improperly 3-choosable), i.e. $(2, 2, 2)$ -colorable.

In this note we focus on near-colorings and Steinberg's Conjecture. Let \mathcal{F} be the family of planar graphs without cycles of length 4 and 5. We prove:

Theorem 1 *Every graph of \mathcal{F} is $(2, 1, 0)$ -colorable and $(4, 0, 0)$ -colorable.*

The remaining of the paper is dedicated to the proof of this theorem.

2 General setting for (s_1, s_2, s_3) -colorability of \mathcal{F}

The proof of the main theorem is done by reducible configurations and discharging procedure. Suppose the theorem is not true. Let $G = (V, E, F)$ be a counterexample with the minimum order embedded in the plane. We apply a discharging procedure to reach to a contradiction.

We first assign to each vertex v and face f of G a charge ω such that $\omega(v) = 2d(v) - 6$ and $\omega(f) = r(f) - 6$, where $d(v)$ and $r(f)$ denote the degree of the vertex v and the length of the face f respectively. By Euler's Formula $|V| - |E| + |F| = 2$ and formula $\sum_{v \in V} d(v) = 2|E| = \sum_{f \in F} r(f)$, we have:

$$\sum_{v \in V} \omega(v) + \sum_{f \in F} \omega(f) = -12 < 0.$$

We then redistribute the charges according to some discharging rules. During the process, no charges are created or disappear. Let ω^* be the new charge on each vertex and face after the procedure. It follows that:

$$\sum_{v \in V} \omega(v) + \sum_{f \in F} \omega(f) = \sum_{v \in V} \omega^*(v) + \sum_{f \in F} \omega^*(f).$$

However, we will show that under some structural properties of G the new charge on each vertex and face is non-negative. This leads to the following obvious contradiction

$$-12 = \sum_{v \in V} \omega(v) + \sum_{f \in F} \omega(f) = \sum_{v \in V} \omega^*(v) + \sum_{f \in F} \omega^*(f) > 0$$

implying that no counterexample can exist.

Establishing structural properties is essential in the proof of the theorem. Although the properties for $(2, 1, 0)$ -coloring and for $(4, 0, 0)$ -coloring are not the same, they share some common part. In this section, we derive lemmas for a general setting. Suppose $s_1 \geq s_2 \geq s_3 \geq 0$ and $s = s_1 + s_2 + s_3$. In this section we assume that G is a minimum counterexample in \mathcal{F} that is not (s_1, s_2, s_3) -colorable.

A vertex of degree k (resp. at least k , at most k) will be called k -vertex (resp. k^+ -vertex, k^- -vertex). A similar notation will be used for cycles and faces. A k -neighbor (resp. k^+ -neighbor, k^- -neighbor) of some vertex u is a neighbor of u which is a k -vertex. An (a, b, c) -face is a 3-face uvw such that $d(u) = a$, $d(v) = b$ and $d(w) = c$. In addition, a^- (resp. a^+) will mean $d(u) \leq a$ (resp. $d(u) \geq a$) and $*$ will mean any degree. For example, a $(3, 4^-, *)$ -face is a 3-face uvw such that $d(u) = 3$, $d(v) \leq 4$ and w has no restriction on its degree. A *pendent 3-face* of a vertex v is a 3-face not containing v but is incident to a 3-vertex adjacent to v . In the following we will color the vertices of the graphs by partitioning the vertex set into V_1, V_2, V_3 such that each V_i induces a subgraph of maximum degree at most s_i . Coloring a vertex with color i means adding the vertex into V_i . We will say that we *nicely color* a vertex if we color it by i and at most $\max\{0, s_i - 1\}$ of its neighbors are colored by i . We say that we *properly color* a vertex if we color it by a color not used by its neighbors. Properly colored vertices are nicely colored. When the colored neighbors of an uncolored vertex v use at most two colors, in particular when v has at most two colored neighbors, we can always color v properly by using the third color not used by its neighbors. We will use this frequently. As an easy consequence, every vertex of G has degree at least 3.

First, since G has no 4-cycles, we have the following:

Observation 2 *Two 3-faces may not share an edge. If a k -vertex v is incident to α 3-faces and has β pendent 3-faces, then $2\alpha + \beta \leq k$.*

Next, three useful lemmas.

Lemma 3 *Let v be an $(s + 2)^-$ -vertex of G . If $G - v$ has an (s_1, s_2, s_3) -coloring such that all neighbors of v are nicely colored, then G is (s_1, s_2, s_3) -colorable.*

PROOF. For $1 \leq i \leq 3$, if we cannot assign color i to v , then v has at least $s_i + 1$ neighbors colored by i . It follows that v has degree at least $\sum_{i=1}^3 (s_i + 1) = s + 3$, a contradiction. \square

Lemma 4 *Graph G contains no $(s + 2)^-$ -vertex v adjacent only to 4^- -vertices, each 4-neighbor of which is adjacent some 3-neighbor of v .*

PROOF. Suppose to the contrary that G contains such a $(s + 2)^-$ -vertex v . By the minimality of G , the graph G' obtained from G by deleting v and all of its neighbors admits an (s_1, s_2, s_3) -coloring. We first color all 4-neighbors of v properly, and then color all 3-neighbors of v properly. Then all neighbors of v are nicely colored. Thus, by Lemma 3, G is (s_1, s_2, s_3) -colorable, a contradiction. \square

Lemma 5 *The three neighbors x_1, x_2, x_3 of a 3-vertex v of G use different colors in an (s_1, s_2, s_3) -coloring of $G - v$. Moreover, assume x_i is colored by i , we have $d(x_i) \geq s_i + 3$ for $1 \leq i \leq 3$. Furthermore, if $s_i > 0$ and x_i is adjacent to x_j , then either $d(x_i) > s_i + 3$ or $d(x_j) > s_j + 3$.*

PROOF. If x_1, x_2, x_3 do not use three distinct colors, then we can properly color v , a contradiction. Hence w.l.o.g. we can assume that x_i is colored by i for $1 \leq i \leq 3$.

Suppose for a contradiction that some $d(x_i) \leq s_i + 2$ for some i . Then $s_i \geq 1$ as $d(x_i) \geq 3$. If x_i is nicely colored by i , then we color v by i and this extends the coloring to G , a contradiction.

Hence, x_i has at least s_i neighbors colored by i . Since x_i has an uncolored neighbor v , there is at least one color different from i not used by its neighbors. We then color v by i and recolor x_i by the unused color. This extends the coloring to G , a contradiction.

Suppose for a contradiction that x_i is adjacent to x_j , but $d(x_i) = s_i + 3$ and $d(x_j) = s_j + 3$. Let k be the color distinct from i and j . Since G has no 4-cycle, x_k is not adjacent to x_i and x_j . As above, x_i (resp. x_j) has s_i (resp. s_j) neighbors colored by i (resp. j) and another colored neighbor x'_i (resp. x'_j) other than x_j (resp. x_i). If x'_i is colored by j , then we may color v by i and recolor x_i by k to get an (s_1, s_2, s_3) -coloring of G , a contradiction. Hence, x'_i is colored by k . Similarly, x'_j is also colored by k . Then we may color v by i , recolor x_i by j and recolor x_j by i to get an (s_1, s_2, s_3) -coloring of G (notice that $s_i > 0$), again a contradiction. Hence, $d(x_i) > s_i + 3$ or $d(x_j) > s_j + 3$. \square

3 (2, 1, 0)-colorability of \mathcal{F}

In this section we prove that every graph in \mathcal{F} is $(2, 1, 0)$ -colorable, namely we consider the case $(s_1, s_2, s_3) = (2, 1, 0)$ for which $s = s_1 + s_2 + s_3 = 3$.

3.1 Reducible configurations for $(2, 1, 0)$ -coloring

We first establish structural properties of G . More precisely, we prove that some ‘configurations’, i.e. subgraphs, are ‘reducible’, i.e. cannot appear in G because it is a minimum counterexample. Lots of this configurations are depicted in Figure 1.

A *light 5-vertex* is a 5-vertex incident to a $(3, 5, 5)$ -face f and adjacent to three 3-vertices not in f . A *poor $(3, 5, 5)$ -face* is a $(3, 5, 5)$ -face incident to a light 5-vertex. If a 3-vertex is incident to a 3-face, then its neighbor not incident to this 3-face is said to be its *outer neighbor*.

As already mentioned we have the following.

(C1) G contains no 2^- -vertices.

The two following claims come from Lemma 4 with $s = 3$.

(C2) G contains no 5-vertex adjacent to five 3-vertices.

(C3) G does not contain 5-vertices v incident to a $(3, 4, 5)$ -face f and adjacent to three 3-vertices not in f .

(C4) G contains no non-light 5-vertex incident to a poor $(3, 5, 5)$ -face and a $(3, 5^-, 5)$ -face, and adjacent to a 3-vertex not in these faces.

Proof. Suppose to the contrary that G contains such a 5-vertex v . Let uvw be the poor $(3, 5, 5)$ -face, rvs be the $(3, 5^-, 5)$ -face with $d(u) = d(r) = 3$, and x be the neighbor of v not in these faces. Vertex w is light and thus is adjacent to three 3-vertices distinct from u , say w_1, w_2, w_3 . By the minimality of G , the graph $G - \{u, v, w, w_1, w_2, w_3, r, x\}$ admits a $(2, 1, 0)$ -coloring. Now we extend this coloring as follows. We may assume that, if s is colored by 1, then it has at most one neighbor colored by 1, otherwise we can properly recolor it. Then we color r and x properly. If s, r, x use different colors, then we color v with 1; otherwise we color v properly. We then color u, w_1, w_2, w_3 properly. It follows that all neighbors of w are nicely colored. By Lemma 3, G is $(2, 1, 0)$ -colorable, a contradiction. \square

(C5) G does not contain a poor $(3, 5, 5)$ -face incident to two light 5-vertices.

Proof. Suppose to the contrary that G contains a poor $(3, 5, 5)$ -face uvw with light vertices v and w . For $x \in \{v, w\}$, let x_1, x_2, x_3 be the three neighbors of x not in $\{u, v, w\}$. By the minimality of G , the graph $G - \{u, v, w, w_1, w_2, w_3, v_1, v_2, v_3\}$ admits a $(2, 1, 0)$ -coloring. We extend the coloring to $\{v_1, v_2, v_3\}$ by coloring each of them properly. If v_1, v_2, v_3 use three distinct colors, then

we color v with 1, and properly otherwise. After this, we color u, w_1, w_2, w_3 properly. It follows that all neighbors of w are nicely colored. By Lemma 3, G is $(2, 1, 0)$ -colorable, a contradiction. \square

Let v be a 3-vertex adjacent to three vertices y_1, y_2, y_3 . Consider $G - v$. By Lemma 5, the colors 1, 2, and 3 appear on the neighbors of v . Moreover the vertex colored with 1 (resp. 2, 3) has degree at least 5 (resp. 4, 3). Thus (C6) and (C7) follow.

(C6) G does not contain 3-vertices adjacent to two 3-vertices.

(C7) If uvw is a $(3, 4, 4)$ -face with $d(u) = 3$, then the outer neighbor of u has degree at least 5.

Now, if the three vertices y_1, y_2, y_3 satisfy $d(y_1) = 3, d(y_2) \leq 4$ and $d(y_2) \leq d(y_3)$, then y_1 (resp. y_2, y_3) is colored with 3 (resp. 2, 1) and has degree 3 (resp. 4, at least 5). By the last sentence of Lemma 5, the vertices y_1, y_2 are non-adjacent; moreover if $d(y_3) = 5$, then y_3 is not adjacent to y_1 or y_2 . Thus (C8), (C9), and (C10) follow.

(C8) G does not contain $(3, 3, 4^-)$ -faces.

(C9) If uvw is a $(3, 3, 5)$ -face with $d(u) = 3$, then the outer neighbor of u has degree at least 5.

(C10) If uvw is a $(3, 4, 5)$ -face with $d(u) = 3, d(v) = 4$ and $d(w) = 5$, then the outer neighbor of u has degree at least 4.




Figure 1: Reducible configurations (C2)-(C10). Black dots represent vertices all neighbours of which are drawn in the figure; the white dots represent vertices that can have nondepicted neighbours. Dashed lines represent edges that may possibly not exist.

3.2 Discharging procedure for $(2, 1, 0)$ -coloring

We now apply a discharging procedure to reach to a contradiction. The discharging rules are as follows:

- R1.** Every 4-vertex gives $\frac{1}{2}$ to each pendent 3-face.
- R2.** Every 5^+ -vertex gives 1 to each pendent 3-face.
- R3.** Every 4-vertex gives 1 to each incident 3-face.
- R4.** Every non-light 5-vertex gives 2 to each incident poor $(3, 5, 5)$ -face.
- R5.** Every 5-vertex gives $\frac{3}{2}$ to each incident non-poor $(3, 5, 5)$ -face or $(3, 4, 5)$ -face.
- R6.** Every 5-vertex gives 1 to each other incident 3-face.
- R7.** Every 6^+ -vertex gives 2 to each incident 3-face.

Let v be a k -vertex with $k \geq 3$ by (C1).

Case $k = 3$. The discharging procedure does not involves 3-vertices. Hence $\omega^*(v) = \omega(v) = 0$.

Case $k = 4$. Initially $\omega(v) = 2$. Vertex v gives 1 to each of the α incident 3-faces by R3 and $\frac{1}{2}$ to each of the β pendent 3-faces by R1. By Observation 2, $\omega^*(v) \geq 2 - (\alpha + \frac{1}{2}\beta) \geq 2 - \frac{1}{2} \cdot 4 = 0$.

Case $k = 5$. Initially $\omega(v) = 4$. Assume v is not incident to any 3-face. By (C2), v is adjacent to at most four 3-vertices and so has at most four pendent 3-faces. By R2, $\omega^*(v) \geq 4 - 4 \cdot 1 = 0$.

Assume v is incident to exactly one 3-face f . If v is a non-light 5-vertex and f is a poor $(3, 5, 5)$ -face, then v has at most two pendent 3-faces by definition. By R4 and R2, $\omega^*(v) \geq 4 - 2 - 2 \cdot 1 = 0$. If f is a non-poor $(3, 5, 5)$ -face, then v has at most two pendent 3-faces by definition. By R5 and R2, $\omega^*(v) \geq 4 - \frac{3}{2} - 2 \cdot 1 > 0$. If f is a $(3, 4, 5)$ -face, then v has at most two pendent 3-faces by (C3). By R5 and R2, $\omega^*(v) \geq 4 - \frac{3}{2} - 2 \cdot 1 > 0$. If f is a 3-face of other type, then by R6 and R2 $\omega^*(v) \geq 4 - 1 - 3 \cdot 1 = 0$.

Assume v is incident to exactly two 3-faces f_1 and f_2 . If v gives twice at most $\frac{3}{2}$ to the 3-faces, then $\omega^*(v) \geq 4 - 2 \cdot \frac{3}{2} - 1 = 0$. So we may assume that f_1 or f_2 , say f_1 , is a poor $(3, 5, 5)$ -face. If f_2 is a $(3, 5^-, 5)$ -face, then v has no pendent 3-faces by (C4) and $\omega^*(v) \geq 4 - 2 - 2 = 0$. If f_2 is a 3-face of other type, then v may have a pendent 3-face and $\omega^*(v) \geq 4 - 2 - 1 - 1 = 0$ by R6.

Case $k \geq 6$. Initially $\omega(v) = 2k - 6$. Vertex v gives 2 to each of the α incident 3-faces by R7 and 1 to each of the β pendent 3-faces by R2. By Observation 2, $\omega^*(v) \geq 2k - 6 - 2\alpha - \beta \geq 2k - 6 - k = k - 6 \geq 0$.

Let f be a k -face.

Case $k = 3$. Initially $\omega(f) = -3$. By (C8), f is not a $(3, 3, 4^-)$ -face.

Let $f = uvw$ be a $(3, 3, 5)$ -face so that $d(u) = d(v) = 3$ and $d(w) = 5$. By (C9) the outer neighbor of u (resp. v) has degree at least 5 and so gives at least 1 to f by R2. By R6, w gives 1 to f . It follows that $\omega^*(f) = -3 + 2 \cdot 1 + 1 = 0$.

Let $f = uvw$ be a $(3, 3, 6^+)$ -face so that $d(u) = d(v) = 3$ and $d(w) \geq 6$. By (C6), the outer neighbor of u (resp. v) has degree at least 4 and so gives at least $\frac{1}{2}$ to f by R1. By R7, w gives 2 to f . It follows that $\omega^*(f) = -3 + 2 \cdot \frac{1}{2} + 2 = 0$.

Let $f = uvw$ be a $(3, 4, 4)$ -face so that $d(u) = 3$ and $d(v) = d(w) = 4$. By (C7) the outer neighbor of u has degree at least 5 and so gives 1 to f by R2. Vertices v (resp. w) give 1 to f by R3. Hence $\omega^*(f) = -3 + 1 + 2 \cdot 1 = 0$.

Let $f = uvw$ be a $(3, 4, 5)$ -face so that $d(u) = 3$, $d(v) = 4$ and $d(w) = 5$. By (C10), the outer neighbor of u has degree at least 4 and so gives at least $\frac{1}{2}$ to f by R1. Vertices v and w give each 1 and $\frac{3}{2}$ to f respectively by R3 and R5. Hence $\omega^*(f) = -3 + \frac{1}{2} + 1 + \frac{3}{2} = 0$.

Let $f = uvw$ be a $(3, 4, 6^+)$ -face so that $d(u) = 3, d(v) = 4$ and $d(w) \geq 6$. By R3 and R7, vertices v and w give each 1 and 2 to f respectively. Hence $\omega^*(f) = -3 + 1 + 2 = 0$.

Let $f = uvw$ be a $(3, 5, 5)$ -face so that $d(u) = 3, d(v) = d(w) = 5$. Assume f is poor and v is light. By (C5) w cannot be light. Hence $\omega^*(f) = -3 + 1 + 2 = 0$ by R4 and R6. Assume f is not poor. Then $\omega^*(f) = -3 + 2 \cdot \frac{3}{2} = 0$ by R5.

Let $f = uvw$ be a $(3, 5^+, 6^+)$ -face so that $d(u) = 3, d(v) \geq 5, d(w) \geq 6$. Vertices v and w give each at least 1 and 2 respectively by R6-7. Hence $\omega^*(f) \geq -3 + 1 + 2 = 0$.

Let $f = uvw$ be a $(4^+, 4^+, 4^+)$ -face. Each incident vertex gives at least 1 to f by R3-7. Hence $\omega^*(f) \geq -3 + 3 \cdot 1 = 0$.

Case $k \geq 4$. Faces of length 4 and 5 do not exist by hypothesis. Faces of length at least 6 are not involved in the discharging procedure. Hence $\omega^*(f) = \omega(f) = r(f) - 6 \geq 0$.

It follows that every vertex and face has a non-negative charge as required. This completes the proof.

4 $(4, 0, 0)$ -colorability of \mathcal{F}

In this section we prove that every graph of \mathcal{F} is $(4, 0, 0)$ -colorable, namely we consider the case of $(s_1, s_2, s_3) = (4, 0, 0)$ for which $s = s_1 + s_2 + s_3 = 4$.

4.1 Reducible configurations for $(4, 0, 0)$ -coloring

In this section we study structural properties of G and establish a number of reducible configurations. See Figure 3.

A *bad 8-vertex* is a 8-vertex v incident to three $(3, 3, 8)$ -faces and to a $(3, 8, *)$ -face $f = uvw$ with $d(u) = 3, d(v) = 8$, where the vertex w is called the *sponsor* of f and f is a *bad face* of v . See Figure 2.




Figure 2: A bad 8-vertex v whose bad face is uvw with sponsor w . (Drawing conventions are the same as in Figure 1.)

(C1') G contains no 2^- -vertices.

(C2') For $8 \leq k \leq 10$, a k -vertex cannot be incident to exactly $k - 5$ $(3, 3, k)$ -faces and adjacent to k 3-vertices.

Proof. Suppose v is a k -vertex incident to exactly $k - 5$ $(3, 3, k)$ -faces and adjacent to $10 - k$ other 3-vertices not in these $(3, 3, k)$ -faces. By the minimality of G , the graph G' obtained from G by deleting v and all its neighbors admits a $(4, 0, 0)$ -coloring. We color properly and sequentially all neighbors of v . Since each $(3, 3, k)$ -face contains at most one vertex colored by 1, color 1 appears at most $k - 5 + 10 - k = 5$ times on the neighbors of v . If it appears less than 5 times, we can

color v with 1, a contradiction. Hence color 1 appears exactly 5 times, once in each $(3, 3, k)$ -face and on all the $10 - k$ other 3-vertices. For each $(3, 3, k)$ -face vxy with $d(x) = d(y) = 3$, where x is colored by 1, y is colored by 2 or 3. In the case of y is colored by 3, if the outer neighbor of y is colored by 1 (resp. 2), then we can recolor y by 2 (resp. 1). Then we can color v with 3 to obtain a $(4, 0, 0)$ -coloring of G , a contradiction. \square

(C3') Every 3-vertex of G is adjacent to at least one 7^+ -vertex.

Proof. This follows from the fact that the degree sequence for the three neighbors of a 3-vertex is lexicographically at least $(7, 3, 3)$ by Lemma 5. \square

(C4') If uvw is a $(3, 3, 7)$ -face with $d(u) = 3$, then the outer neighbor of u has degree at least 4.

Proof. Suppose to the contrary that G has a $(3, 3, 7)$ -face uvw with $d(u) = d(v) = 3$ and $d(w) = 7$, but the outer vertex x of u has $d(x) = 3$. By Lemma 5, the degree sequence for the three neighbors of u is lex-graphically at least $(7, 3, 3)$. Hence w is colored by 1 and v is colored by 2 or 3. This contradicts the last sentence of Lemma 5 as w is adjacent to v . \square

(C5') The sponsor w of a bad 8-vertex v has degree at least 8 and is not a bad 8-vertex.

Proof. Suppose to the contrary that the bad 8-vertex v is incident to three $(3, 3, 8)$ -faces x_1x_2v , y_1y_2v and z_1z_2v and to a $(3, 8, *)$ -face uvw with $d(u) = 3$ and $3 \leq d(w) \leq 7$ or w a bad 8-vertex. By the minimality of G , the graph $G' = G - \{v, x_1, x_2, y_1, y_2, z_1, z_2, u\}$ admits a $(4, 0, 0)$ -coloring. We can assume that w is nicely colored; otherwise, if $d(w) \leq 7$, then we can recolor it properly, and if w is a bad 8-vertex, then we can recolor properly all its colored neighborhood and then color w nicely. Now we color properly and sequentially $x_1, x_2, y_1, y_2, z_1, z_2, u$, and we assign color 1 to v (color 1 appears at most 4 times on the neighbors of v). This extends the $(4, 0, 0)$ -coloring to G , a contradiction. \square

4.2 Discharging procedure for $(4, 0, 0)$ -coloring

We now apply a discharging procedure to reach a contradiction. The discharging rules are as follows:

- R1'**. For $4 \leq k \leq 6$, every k -vertex gives $\frac{1}{2}$ to each pendent 3-face.
- R2'**. Every 7^+ -vertex gives 1 to each pendent 3-face.
- R3'**. For $4 \leq k \leq 6$, every k -vertex gives 1 to each incident 3-face.
- R4'**. Every 7^+ -vertex gives 1 to each incident $(4^+, 4^+, 4^+)$ -face.
- R5'**. Every non-bad 7^+ -vertex gives 2 to each incident $(3, 4^+, 4^+)$ -face; every bad 8-vertex gives 1 to its bad 3-face.
- R6'**. Every 7-vertex gives 2 to each incident $(3, 3, 7)$ -face.
- R7'**. For $k \geq 8$, every k -vertex gives 3 to each incident $(3, 3, k)$ -face.

Let v be a k -vertex with $k \geq 3$ by (C1'). Initially $\omega(v) = 2k - 6$.

Case $k = 3$. The discharging procedure does not involves 3-vertices. Hence $\omega^*(v) = \omega(v) = 0$.

Case $4 \leq k \leq 6$. Vertex v gives 1 to each of the α incident 3-faces by R3' and $\frac{1}{2}$ to each of the β pendent 3-faces by R1'. By Observation 2, $\omega^*(v) \geq 2k - 6 - (\alpha + \frac{1}{2}\beta) \geq 2k - 6 - \frac{1}{2}k = \frac{3}{2}k - 6 \geq 0$.

Case $k = 7$. Vertex v gives 2 to each of the α' incident $(3, 3^+, 4^+)$ -faces by R5'-6', 1 to each of the α'' incident $(4^+, 4^+, 4^+)$ -faces by R4', and 1 to each of the β pendent 3-faces by R2'. By Observation 2, $\omega^*(v) \geq 2k - 6 - (2\alpha' + \alpha'' + \beta) \geq 2k - 6 - k = k - 6 > 0$.

Case $k \geq 8$. For the case when v is a bad 8-vertex, v gives 3 to each incident $(3, 3, 8)$ -face by R7' and 1 to the bad 3-face by R5'. Hence $\omega^*(v) = 2 \cdot 8 - 6 - 3 \cdot 3 - 1 = 0$.




Figure 3: The reducible configurations (C2')-(C5'). (Drawing conventions are the same as in Figure 1.)

Now assume that v is not a bad 8-vertex. By R7', R5', R4' and R2', v gives 3 to each of the α' incident $(3, 3, k)$ -faces, 2 to each of the α'' incident $(3, 4^+, 4^+)$ -faces, 1 to each of the α''' incident $(4^+, 4^+, 4^+)$ -faces, and 1 to each of the β pendent 3-faces. By Observation 2, $\omega^*(v) = 2k - 6 - (3\alpha' + 2\alpha'' + \alpha''' + \beta) \geq 2k - 6 - \lfloor \frac{3k}{2} \rfloor = \lceil \frac{k}{2} \rceil - 6 \geq 0$ except for the cases (1) $k = 10$ with $\alpha' = 5$, (2) $k = 9$ with $\alpha' = 4$ and $\beta = 1$, (3) $k = 8$ with $\alpha' = 3$ and $\beta = 2$ (note that the bad 8-vertex case, i.e. $\alpha' = 4$ or $\alpha' = 3$ with $\alpha'' = 1$, is excluded). The exceptional cases give a k -vertex, $8 \leq k \leq 10$, with exactly $k - 5$ $(3, 3, k)$ -faces and adjacent only to 3-vertices, a contradiction to (C2').

Let f be a k -face.

Case $k = 3$. Initially $\omega(f) = -3$.

Let $f = uvw$ be a (a_1, a_2, a_3) -face with $3 \leq a_1 \leq 6, 3 \leq a_2 \leq 6$ and $3 \leq a_3 \leq 6$. By (C3'), the outer neighbor of each 3-vertex incident to f has degree at least 7 and gives each at least 1 to f by R2'. By R3', each d -vertex with $4 \leq d \leq 6$ incident to f gives 1 to f . It follows that $\omega^*(f) = -3 + 3 = 0$.

Let $f = uvw$ be a $(3, 3, 7)$ -face so that $d(u) = d(v) = 3$ and $d(w) = 7$. By (C4') the outer neighbor of u (resp. v) has degree at least 4 and so gives at least $\frac{1}{2}$ to f by R1'. By R6', w gives 2 to f . It follows that $\omega^*(f) = -3 + 2 \cdot \frac{1}{2} + 2 = 0$.

Let $f = uvw$ be a $(3, 3, 8^+)$ -face so that $d(u) = d(v) = 3$ and $d(w) \geq 8$. By R7', w gives 3 to f . It follows that $\omega^*(f) = -3 + 3 = 0$.

Let $f = uvw$ be a $(3, 4^+, 7^+)$ -face so that $d(u) \geq 3, d(v) \geq 4$ and $d(w) \geq 7$. By R3'-5', vertices v and w gives at least 3 to f and so $\omega^*(f) = -3 + 3 = 0$, except for the case when f is a bad 3-face with the pair v, w being either two bad 8-vertices or a bad 8-vertex and a 6^- -vertex. But these two exceptional cases are impossible by (C5').

Finally, let $f = uvw$ be a $(4^+, 4^+, 4^+)$ -face. Every incident vertex gives at least 1 to f by R3'-4'. Hence $\omega^*(f) \geq 0$.

It follows that every vertex and face has a non-negative charge as required. This completes the proof.

References

- [1] H. L. Abbott and B. Zhou, On small faces in 4-critical graphs, *Ars Combinatoria* 32:203–207, 1991.
- [2] K. Appel and W. Haken. Every planar map is four colorable. Part I. Discharging. *Illinois J. Math.*, 21:429–490, 1977.
- [3] K. Appel and W. Haken. Every planar map is four colorable. Part II. Reducibility *Illinois J. Math.*, 21:491–567, 1977.
- [4] O. V. Borodin. To the paper of H. L. Abbott and B. Zhou on 4-critical planar graphs. *Ars Combinatoria* 43:191–192, 1996.
- [5] O. V. Borodin. Structural properties of plane graphs without adjacent triangles and an application to 3-colorings. *J. Graph Theory*, 21(2):183–186, 1996.
- [6] O. V. Borodin and A. N. Glebov. A sufficient condition for plane graphs to be 3-colorable. *Diskret. analiz i issled. oper.*, 10(3):3–11, 2004 (in Russian).
- [7] O. V. Borodin and A. N. Glebov. Planar graphs without 5-cycles and with minimum distance between triangles at least two are 3-colorable. Manuscript, 2008. [to update]

-
- [8] O. V. Borodin, A. N. Glebov, M. Montassier, and A. Raspaud. Planar graphs without 5- and 7-cycles without adjacent triangles are 3-colorable. *Journal of Combinatorial Theory, Series B*, 99(4):668–673, 2009.
- [9] O. V. Borodin, A. N. Glebov, and A. R. Raspaud. Planar graphs without triangles adjacent to cycles of length from 4 to 7 are 3-colorable. Manuscript, 2009.[to update]
- [10] O. V. Borodin, A. N. Glebov, A. R. Raspaud, and M. R. Salavatipour. Planar graphs without cycles of length from 4 to 7 are 3-colorable. *Journal of Combinatorial Theory, Series B*, 93:303–311, 2005.
- [11] O. V. Borodin and A. Raspaud. A sufficient condition for planar graph to be 3-colorable. *Journal of Combinatorial Theory, Series B*, 88:17–27, 2003.
- [12] Y. Bu, H. Lu, M. Montassier, A. Raspaud, W. Wang, and Y. Wang. On the 3-colorability of planar graphs without 4-, 7-, 9-cycles. *Discrete Mathematics*, 309(13):4596–4607, 2009.
- [13] M. Chen, X. Luo, and W. Wang. On 3-colorable planar graphs without cycles of four lengths. *Information Processing Letters*, 103(4):150–156, 2007.
- [14] M. Chen, A. Raspaud, and W. Wang. Three-coloring planar graphs without short cycles. *Information Processing Letters* 101:134–138, 2007.
- [15] M. Chen and W. Wang. On 3-colorable planar graphs without prescribed cycles. *Discrete Mathematics*, 307:2820–2825, 2007.
- [16] M. Chen and W. Wang. Planar graphs without 4, 6, 8-cycles are 3-colorable. *Science in China Serie A: Mathematics*, 50:1552–1562, 2007.
- [17] M. Chen and W. Wang. On 3-colorable planar graphs without short cycles. *Applied Mathematics Letters*, 21(9):961–965, 2008.
- [18] N. Eaton and T. Hull. Defective list colorings of planar graphs. *Bull. Inst. Combin. Appl.*, 25:79–87, 1999.
- [19] M. R. Garey, D. S. Johnson, and L.J. Stockmeyer. Some simplified NP-complete graph problems. *Theoretical Computer Science*, 1:237–267, 1976.
- [20] H. Grötzsch. Ein dreifarbensatz für dreiecksfreie netze auf der kugel. *Math.-Nat. Reihe*, 8:109–120, 1959.
- [21] M. R. Salavatipour. The Three Color Problem for planar graphs. *Technical Report CSRG-458*, Department of Computer Science, University of Toronto, 2002.
- [22] D. P. Sanders and Y. Zhao. A note on the three color problem. *Graphs and Combinatorics*, 11:91–94, 1995.
- [23] R. Škrekovski. List improper coloring of planar graphs. *Comb. Prob. Comp.*, 8:293–299, 1999.
- [24] R. Steinberg. The state of the three color problem. *Quo Vadis, Graph Theory?*, Ann. Discrete Math. 55:211–248, 1993.
- [25] L. Zhang and B. Wu. Three-coloring planar graphs without certain small cycles. *Graph Theory Notes of New York*, 46:27–30, 2004.
- [26] B. Xu. A 3-color theorem on plane graph without 5-circuits. *Acta Mathematica Sinica*, 23(6):1059–1062, 2007.



Centre de recherche INRIA Sophia Antipolis – Méditerranée
2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Centre de recherche INRIA Bordeaux – Sud Ouest : Domaine Universitaire - 351, cours de la Libération - 33405 Talence Cedex
Centre de recherche INRIA Grenoble – Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier
Centre de recherche INRIA Lille – Nord Europe : Parc Scientifique de la Haute Borne - 40, avenue Halley - 59650 Villeneuve d'Ascq
Centre de recherche INRIA Nancy – Grand Est : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex
Centre de recherche INRIA Paris – Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex
Centre de recherche INRIA Rennes – Bretagne Atlantique : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex
Centre de recherche INRIA Saclay – Île-de-France : Parc Orsay Université - ZAC des Vignes : 4, rue Jacques Monod - 91893 Orsay Cedex

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399