

Tensor ranks on tangent developable of Segre varieties

Edoardo Ballico, Alessandra Bernardi

► **To cite this version:**

Edoardo Ballico, Alessandra Bernardi. Tensor ranks on tangent developable of Segre varieties. Linear and Multilinear Algebra, Taylor & Francis, 2012, <10.1080/03081087.2012.716430>. <inria-00610362v2>

HAL Id: inria-00610362

<https://hal.inria.fr/inria-00610362v2>

Submitted on 28 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TENSOR RANKS ON TANGENT DEVELOPABLE OF SEGRE VARIETIES

E. BALLICO AND A. BERNARDI

ABSTRACT. We describe the stratification by tensor rank of the points belonging to the tangent developable of any Segre variety. We give algorithms to compute the rank and a decomposition of a tensor belonging to the secant variety of lines of any Segre variety. We prove Comon's conjecture on the rank of symmetric tensors for those tensors belonging to tangential varieties to Veronese varieties.

INTRODUCTION

In this paper we want to address the problem of tensor decomposition over an algebraically closed field K of characteristic 0 for tensors belonging to a tangent space of the projective variety that parameterizes completely decomposable tensors.

Let V_1, \dots, V_d be K -vector spaces of dimensions $n_1 + 1, \dots, n_d + 1$ respectively; the projective variety $X_{n_1, \dots, n_d} \subset \mathbb{P}(V_1 \otimes \dots \otimes V_d)$ that parameterizes projective classes of completely decomposable tensors $v_1 \otimes \dots \otimes v_d \in V_1 \otimes \dots \otimes V_d$ is classically known as a Segre variety (see Definition 1). Given a tensor $T \in V_1 \otimes \dots \otimes V_d$, finding the minimum number of completely decomposable tensors such that T can be written as a linear combination of them (see Definition 2 for the notion of “tensor rank”) is related to the tensor decomposition problem that nowadays seems to be crucial in many applications like Signal Processing (see eg. [1], [20], [13]), Algebraic Statistics ([19], [25]), Neuroscience (eg. [3]). The specific case of tensors belonging to tangential varieties to Segre varieties (Notation 1) is studied in [9] and it turns out to be of certain interest in the context of Computational Biology. In fact in [14] a particular class of statistical models (namely certain context-specific independence model – CSI) is shown to be crucial in machine learning and computational biology. L. Oeding has recently shown in [22] how to interpret the CSI model performed by [14] in terms of tangential variety to Segre variety. In this setting B. Sturmfels and P. Zwiernik in a very recent paper ([23]) show how to derive parametrizations and implicit equations in cumulants for the tangential variety of the Segre variety $X_{1, \dots, 1}$ and for certain CSI models (see [7] for a combinatorial point of view on cumulants).

In this paper, after a preliminary section, we give a complete classification of the tensor rank of an element belonging to the tangent developable of any Segre variety. In particular in Theorem 1 we will prove that if $P \in T_O(X_{n_1, \dots, n_d})$ for certain point $O = (O_1, \dots, O_d) \in X_{n_1, \dots, n_d}$, then the minimum number r of completely decomposable tensors $v_{1,i} \otimes \dots \otimes v_{d,i} \in V_1 \otimes \dots \otimes V_d$ such that $P = \sum_{i=1}^r [v_{1,i} \otimes \dots \otimes v_{d,i}]$ is equal to the minimum number $\eta_{X_{n_1, \dots, n_d}}(P)$

1991 *Mathematics Subject Classification.* 14N05, 14Q05.

Key words and phrases. Secant varieties; tensor rank; tangent developable; Segre Varieties; Comon's conjecture.

The first author was partially supported by MIUR and GNSAGA of INdAM (Italy). The second author was partially supported by CIRM-FBK (TN-Italy), Marie-Curie FP7-PEOPLE-2009-IEF, INRIA Sophia Antipolis Méditerranée Project Galaad (France) and Mittag-Leffler Institut (Sweden).

for which there exist $E \subseteq \{1, \dots, d\}$ such that $\sharp(E) = \eta_{X_{n_1, \dots, n_d}}(P)$ and $T_O(X_{n_1, \dots, n_d}) \subseteq \langle \cup_{i \in E} Y_{O,i} \rangle$ where $Y_{O,i}$ the n_i -dimensional linear subspace obtained by fixing all coordinates $j \in \{1, \dots, d\} \setminus \{i\}$ equal to $O_j \in \mathbb{P}_i^n$ (see Notation 3). Such a result was independently proved by J. Buczyński and J. M. Landsberg (see Theorem 7.1 in the second version of [9]). We propose here a different proof. First of all, the construction that we make in our proof allows to write explicit algorithms for the computation of the rank of a given tensor belonging to the secant variety of lines of any Segre variety (Algorithm 1) and for a decomposition of the same (Algorithm 2). Moreover in the third version of [9], the authors have removed that result.

In Section 3 we give the details for Algorithm 1 and for Algorithm 2.

In the last section we show how to use Theorem 1 in order to prove the so called “Comon’s conjecture” in the particular case in which the points $P \in \tau(X_{n_1, \dots, n_d})$ parameterize symmetric tensors. Let us give more details on that.

Let $V_1 = \dots = V_d = V$ be a vector space of dimension $n + 1$ and consider the subspace $S^d V \subset V^{\otimes d}$ of symmetric tensors. The intersection between the Segre variety $X_{n, \dots, n}$ and $\mathbb{P}(S^d V)$ is a way to interpret the classical Veronese embedding of \mathbb{P}^n via the sections of the sheaf $\mathcal{O}(d)$. Therefore an element of the Veronese variety $\nu_d(\mathbb{P}^n) = X_{n, \dots, n} \cap \mathbb{P}(S^d V)$ is the projective class of a completely decomposable symmetric tensor. Now, given a point $P \in \mathbb{P}(S^d V)$ that parameterizes a projective class of a symmetric tensor, we can look at two different decompositions of it. Let $v_{1,i} \otimes \dots \otimes v_{d,i} \in V^{\otimes d}$ and let $w_j^{\otimes d} \in S^d V$, and ask for the minimum r and the minimum r' such that $P = \sum_{i=1}^r [v_{1,i} \otimes \dots \otimes v_{d,i}] = \sum_{j=1}^{r'} [w_j^{\otimes d}]$. In 2008, at the AIM workshop in Palo Alto, USA (see the report [21]), P. Comon stated the following:

Conjecture 1. [Comon’s Conjecture] The minimum integer r such that a symmetric tensor $T \in S^d V$ can be written as

$$T = \sum_{i=1}^r v_{1,i} \otimes \dots \otimes v_{d,i}$$

for $v_{1,i} \otimes \dots \otimes v_{d,i} \in V^{\otimes d}$, $i = 1, \dots, r$, is equal to the minimum integer r' for which there exist $w_j^{\otimes d} \in S^d V$, $j = 1, \dots, r'$ such that

$$T = \sum_{j=1}^{r'} w_j^{\otimes d}.$$

As far as we know this conjecture is proved if $r \leq \dim(V)$ (for a general d -tensor, d even and large) and if $r = 1, 2$ (see [12]).

In Section 4 we show that our Theorem 1 implies that this conjecture is true also for $[T] \in \tau(X_{n, \dots, n})$ (Corollary 2).

Acknowledgements: We like to thank B. Sturmfels for asking to one of us this question at the Mittag-Leffler Institut during the Spring semester 2011 “Algebraic Geometry with a view towards applications”. We also thank the Mittag-Leffler Institut (Djursholm, Stokholm, Sweden), for its hospitality and opportunities.

1. PRELIMINARIES

Let us start with the classical definition of the Segre varieties.

Definition 1. For all positive integers d and n_i , $1 \leq i \leq d$, let

$$j_{n_1, \dots, n_d} : \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d} \rightarrow \mathbb{P}^{N(n_1, \dots, n_d)},$$

with $N(n_1, \dots, n_d) := (\prod_{i=1}^d (n_i + 1)) - 1$, denote the Segre embedding of $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$ obtained by the section of the sheaf $\mathcal{O}(1, \dots, 1)$. Set $X_{n_1, \dots, n_d} := j_{n_1, \dots, n_d}(\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d})$.

Observe that if we identify each \mathbb{P}^{n_i} with $\mathbb{P}(V_i)$ for certain $(n_i + 1)$ -dimensional vector space V_i for $i = 1, \dots, d$, then an element $[T] \in X_{n_1, \dots, n_d}$ can be interpreted as the projective class of a completely decomposable tensor $T \in V_1 \otimes \dots \otimes V_d$, i.e. there exist $v_i \in V_i$ for $i = 1, \dots, d$ such that $T = v_1 \otimes \dots \otimes v_d$.

We can give now the definition of the rank of an element $P \in \mathbb{P}^{N(n_1, \dots, n_d)} = \mathbb{P}(V_1 \otimes \dots \otimes V_d)$.

Definition 2. For each $P \in \mathbb{P}^{N(n_1, \dots, n_d)}$ the rank (or tensor rank) $r_{X_{n_1, \dots, n_d}}(P)$ of P is the minimal cardinality of a finite set $S \subset X_{n_1, \dots, n_d}$ such that $P \in \langle S \rangle$, where $\langle \cdot \rangle$ denote the linear span.

Notation 1. Let $\tau(X_{n_1, \dots, n_d})$ denote the tangent developable of X_{n_1, \dots, n_d} , i.e. the union of all tangent spaces $T_P X_{n_1, \dots, n_d}$ of X_{n_1, \dots, n_d} . Since $\tau(X_{n_1, \dots, n_d})$ is closed in the Zariski topology, this is equivalent to the usual definition of the tangent developable of a submanifold of a projective space as the closure of the union of all tangent spaces.

Remark 1. Fix any $P \in \tau(X_{n_1, \dots, n_d}) \setminus X_{n_1, \dots, n_d}$. There is a unique $O \in X_{n_1, \dots, n_d}$ and a unique zero-dimensional scheme $Z \subset X_{n_1, \dots, n_d}$ such that $Z_{red} = \{O\}$, $\deg(Z) = 2$ and P is contained in the line $\langle Z \rangle$:

$$(1) \quad P \in T_O X_{n_1, \dots, n_d} = \langle Z \rangle.$$

Notation 2. Let $\tilde{O} = (O_1, \dots, O_d) \in \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$. With an abuse of notation we will write the point $O = j_{n_1, \dots, n_d}(\tilde{O}) \in X_{n_1, \dots, n_d}$ as $O = (O_1, \dots, O_d)$.

Notation 3. Fix $O = (O_1, \dots, O_d) \in X_{n_1, \dots, n_d}$ as above, we indicate with $Y_{O,i} \subset \mathbb{P}^{N(n_1, \dots, n_d)}$ the n_i -dimensional linear subspace obtained by fixing all coordinates $j \in \{1, \dots, d\} \setminus \{i\}$ equal to $O_j \in \mathbb{P}^{n_j}$. To be precise:

$$Y_{O,i} = j_{n_1, \dots, n_d}(O_1, \dots, O_{i-1}, \mathbb{P}^{n_i}, O_{i+1}, \dots, O_d).$$

Remark 2. Let $Y_{O,i} \subset \mathbb{P}^{N(n_1, \dots, n_d)}$ the n_i -dimensional linear subspace just defined. Observe that, as a scheme-theoretic intersection, we have that:

$$(2) \quad T_O X_{n_1, \dots, n_d} \cap X_{n_1, \dots, n_d} = \cup_{i=1}^d Y_{O,i}.$$

Moreover, if $Z \subset X_{n_1, \dots, n_d}$ is the unique degree 2 zero-dimensional scheme such that $\langle Z \rangle = T_O X_{n_1, \dots, n_d}$ as in Remark 1, then there is a unique minimal subset $E \subseteq \{1, \dots, d\}$ such that $\langle Z \rangle \subseteq \langle \cup_{i \in E} Y_{O,i} \rangle$.

The integer $\sharp(E)$ will be called the type $\eta_{X_{n_1, \dots, n_d}}(P)$ of P :

$$(3) \quad \eta_{X_{n_1, \dots, n_d}}(P) := \min\{\sharp(E) \mid E \subseteq \{1, \dots, d\}, \langle Z \rangle \subseteq \langle \cup_{i \in E} Y_{O,i} \rangle\}.$$

Notice that $2 \leq \eta_{X_{n_1, \dots, n_d}}(P) \leq d$. Moreover for a general $Q \in T_O X_{n_1, \dots, n_d}$ we have that $\eta_{X_{n_1, \dots, n_d}}(Q) = d$. Moreover every integer $k \in \{2, \dots, d\}$ is the type of some point of $\tau(X_{n_1, \dots, n_d}) \setminus X_{n_1, \dots, n_d}$. Finally for all $Q \in X_{n_1, \dots, n_d}$ we write $\eta_{X_{n_1, \dots, n_d}}(Q) = 1$ and say that Q has type 1.

In Theorem 1 we will actually prove that if $P \in \tau(X_{n_1, \dots, n_d})$, then the integer $\eta_{X_{n_1, \dots, n_d}}(P)$ just introduced in (3) is actually the rank of P .

Before proving that theorem we need to introduce the notion of secant varieties and other related objects.

Definition 3. For each integer $t \geq 2$ let $\sigma_t(X_{n_1, \dots, n_d})$ denote the Zariski closure in $\mathbb{P}^{N(n_1, \dots, n_d)}$ of the union of all $(t-1)$ -dimensional linear subspaces of $\mathbb{P}^{N(n_1, \dots, n_d)}$ spanned by t points of X_{n_1, \dots, n_d} . This object is classically known as the t -secant variety of X_{n_1, \dots, n_d} .

Notation 4. For each $t \geq 2$ there is a non-empty open subset of $\sigma_t(X_{n_1, \dots, n_d})$, that we indicate with $\sigma_t^0(X_{n_1, \dots, n_d})$, whose elements are points of rank exactly equal to t .

We want to focus our attention on the case $t = 2$ that is very particular. In fact it is classically known that each element of $\sigma_2(X_{n_1, \dots, n_d}) \setminus X_{n_1, \dots, n_d}$ is in the linear span of a unique zero-dimensional scheme $Z \subset X_{n_1, \dots, n_d}$ (this classical assertion uses only that X_{n_1, \dots, n_d} is a smooth submanifold of a projective space; see [5], Proposition 11, and [8], Lemma 2.1.5, for much more). Hence Theorem 1 will give the complete stratification by ranks of points in $\sigma_2(X_{n_1, \dots, n_d})$ (see also 1). Indeed, fix $P \in \sigma_2(X_{n_1, \dots, n_d})$. Obviously $\tau(X_{n_1, \dots, n_d}) \subseteq \sigma_2(X_{n_1, \dots, n_d})$. If $\tau(X_{n_1, \dots, n_d}) \neq \sigma_2(X_{n_1, \dots, n_d})$ and if $P \notin \tau(X_{n_1, \dots, n_d})$, then $r_{X_{n_1, \dots, n_d}}(P) = 2$ (in fact if $P \in \sigma_2(X_{n_1, \dots, n_d}) \setminus \tau(X_{n_1, \dots, n_d})$ there exists, by Definition 3, two distinct points of X_{n_1, \dots, n_d} whose span contains P). If $P \in \tau(X_{n_1, \dots, n_d})$, then in Theorem 1 we will show that $r_{X_{n_1, \dots, n_d}}(P) \in \{1, \dots, d\}$. In particular for each $k \in \{1, \dots, d\}$, Theorem 1 will also imply the existence of $P \in \tau(X_{n_1, \dots, n_d})$ such that $r_{X_{n_1, \dots, n_d}}(P) = k$.

Definition 4. For any $P \in \mathbb{P}^{N(n_1, \dots, n_d)}$ the border rank, or border tensor rank, $b_{X_{n_1, \dots, n_d}}(P)$ is the minimal integer t such that $P \in \sigma_t(X_{n_1, \dots, n_d})$.

Notice that

$$b_{X_{n_1, \dots, n_d}}(P) = 1 \iff r_{X_{n_1, \dots, n_d}}(P) = 1 \iff P \in X_{n_1, \dots, n_d}.$$

Thus Theorem 1 may be considered as the description of the ranks of all points with border rank 2 (Corollary 1).

For the case of Veronese varieties, i.e. the case of symmetric tensors, and symmetric border rank 2 or 3, see [5] and references therein.

2. PROOF OF THEOREM 1.

This section is entirely devoted to the proof of Theorem 1.

Before going into the details of the proof of Theorem 1, we need to remind the following elementary lemma (see e.g. [2], Lemma 1).

Lemma 1. Fix any $P \in \mathbb{P}^{N(n_1, \dots, n_d)}$ and two zero-dimensional subschemes A, B of X_{n_1, \dots, n_d} such that $A \neq B$, $P \in \langle A \rangle$, $P \in \langle B \rangle$, $P \notin \langle A' \rangle$ for any $A' \subsetneq A$ and $P \notin \langle B' \rangle$ for any $B' \subsetneq B$. Then $h^1(\mathbb{P}^{N(n_1, \dots, n_d)}, \mathcal{I}_{A \cup B}(1)) > 0$.

Lemma 2. Fix a zero-dimensional scheme $\tilde{W} \subset \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$. Then $h^1(\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}, \mathcal{I}_{\tilde{W}}(1, \dots, 1)) = h^1(\mathbb{P}^{N(n_1, \dots, n_d)}, \mathcal{I}_{j_{n_1, \dots, n_d}(\tilde{W})}(1))$.

Proof. It is sufficient to observe that j_{n_1, \dots, n_d} is the linearly normal embedding induced by the complete linear system $|\mathcal{O}_{\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}}(1, \dots, 1)|$ and that $h^1(\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}, \mathcal{O}_{\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}}(1, \dots, 1)) = 0$. \square

We are now ready to prove Theorem 1.

Theorem 1. Let $\tau(X_{n_1, \dots, n_d})$ be the tangential variety of the Segre variety X_{n_1, \dots, n_d} . For each $P \in \tau(X_{n_1, \dots, n_d})$ we have that the tensor rank of P is:

$$r_{X_{n_1, \dots, n_d}}(P) = \eta_{X_{n_1, \dots, n_d}}(P)$$

where the integer $\eta_{X_{n_1, \dots, n_d}}(P)$ is the type of P defined in (3).

We write here the strategy of the proof in order to help the reader in following it.

First of all, we observe that if $\eta_{X_{n_1, \dots, n_d}}(P) = 1$ there is nothing to prove. So we assume that there exist a point $O \in X_{n_1, \dots, n_d}$ such that $P \in T_O(X_{n_1, \dots, n_d}) \setminus \{O\}$.

Moreover we point out that the inequality $r_{X_{n_1, \dots, n_d}}(P) \leq \eta_{X_{n_1, \dots, n_d}}(P)$ (see (4)) is obvious, then we need only to prove the reverse inequality.

Then we split the proof in the following cases:

(a) If all the $n_i = 1$ and $\eta_{X_{1, \dots, 1}}(P) = d \Rightarrow r_{X_{1, \dots, 1}}(P) = d$. This is proved by absurd: we assume that $\eta_{X_{1, \dots, 1}}(P) = d$ and that $r_{X_{1, \dots, 1}}(P) < d$ and we show that in each of the following sub-cases we get a contraddiction:

- (a1) $O \notin \langle S \rangle$, where S is the set of points computing the rank of P ;
- (a2) $O \in S$;
- (a3) $O \notin S$ and $O \in \langle S \rangle$.

(b) If all the $n_i = 1$ and $\eta_{X_{1, \dots, 1}}(P) < d \Rightarrow r_{X_{1, \dots, 1}}(P) = \eta_{X_{1, \dots, 1}}(P)$.

(c) We conclude the proof by showing that the theorem is true for all $n_i \geq 2$ (this part may be bypassed quoting [18] where it is shown that secant variety of lines of a Segre variety is contained in the subspace variety).

Proof. Fix $P \in \tau(X_{n_1, \dots, n_d})$ and look for $r_{X_{n_1, \dots, n_d}}(P)$.

Since $\eta_{X_{n_1, \dots, n_d}}(P) = 1 \iff P \in X_{n_1, \dots, n_d} \iff r_{X_{n_1, \dots, n_d}}(P) = 1$, the case $P \in X_{n_1, \dots, n_d}$ is obvious. Hence we may assume $P \notin X_{n_1, \dots, n_d}$. Take $O \in X_{n_1, \dots, n_d}$ and $Z \subset X_{n_1, \dots, n_d}$ with $Z_{red} = \{O\}$, $\deg(Z) = 2$ and $P \in \langle Z \rangle$, hence, as in (1), we have that

$$P \in T_O X_{n_1, \dots, n_d} = \langle Z \rangle.$$

As we have seen above, both the point $O \in X_{n_1, \dots, n_d}$ and the degree 2 zero-dimensional scheme $Z \subset X_{n_1, \dots, n_d}$ that satisfy (1) exist, and moreover they are uniquely determined by P . Moreover we can think of $Z \subset X_{n_1, \dots, n_d}$ as

$$Z = j_{n_1, \dots, n_d}(\tilde{Z})$$

with $\tilde{Z} \subset \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$ and $\tilde{Z} \cong Z$.

Now, as in Remark 2, fix $E \subseteq \{1, \dots, d\}$ such that

$$\sharp(E) = \eta_{X_{n_1, \dots, n_d}}(P)$$

and

$$P \in \langle \cup_{i \in E} Y_{O, i} \rangle$$

(where $Y_{O, i}$ are defined as in Notation 3).

Since each $Y_{O, i} \subset \mathbb{P}^{n_1, \dots, n_d}$ is a linear subspace, then for each $i \in E$ there is $Q_i \in Y_{O, i}$ such that $P \in \langle \cup_{i \in E} Q_i \rangle$. Thus

$$(4) \quad r_{X_{n_1, \dots, n_d}}(P) \leq \eta_{X_{n_1, \dots, n_d}}(P).$$

Therefore we need simply to prove the opposite inequality.

For each $j \in \{1, \dots, d\}$ and each $Q_j \in \mathbb{P}^{n_j}$ (or, with the same abuse of notation as in Notation 2, we can think at a point Q in the Segre variety obtained as $j_{n_1, \dots, n_d}(\tilde{Q})$ with $\tilde{Q} = (Q_1, \dots, Q_d) \in \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$ and then write $Q = (Q_1, \dots, Q_d) \in X_{n_1, \dots, n_d}$), set:

$$(5) \quad X_{n_1, \dots, n_d}(Q_j, j) := \{(A_1, \dots, A_d) \in X_{n_1, \dots, n_d} : A_j = Q_j\}.$$

Hence $X(Q_j, j)$ is an $(n_1 + \dots + n_d - n_j)$ -dimensional product of $d - 1$ projective spaces embedded as a Segre variety in a linear subspace of $\mathbb{P}^{N(n_1, \dots, n_d)}$.

Now our proof splits in two parts: in the first one ((a) together with (b)) we study the case of the Segre product of d copies of \mathbb{P}^1 's (i.e. we prove the theorem for $\tau(X_{1, \dots, 1})$); in part

(c) we generalize the result obtained for $X_{1,\dots,1}$ to the general case X_{n_1,\dots,n_d} with $n_i \geq 1$, $i = 1, \dots, d$.

(a) Here we assume $n_i = 1$ for all i and $\eta_{X_{n_1,\dots,n_d}}(P) = d$.

Assume $r := r_{X_{1,\dots,1}}(P) < d$ and fix a 0-dimensional scheme $S \subset X_{n_1,\dots,n_d}$ that computes the rank r of P , i.e. fix

$$\tilde{S} \subset \mathbb{P}^1 \times \dots \times \mathbb{P}^1$$

such that

$$j_{n_1,\dots,n_d}(\tilde{S}) = S, P \in \langle S \rangle \text{ and } \sharp(j_{n_1,\dots,n_d}(S)) = r.$$

Write

$$S = \{Q_1, \dots, Q_r\}$$

and let $(Q_{i,1}, \dots, Q_{i,d})$ be the components of each $Q_i \in X_{1,\dots,1}$ with $i = 1, \dots, r$, i.e. let $\tilde{Q}_i = (Q_{i,1}, \dots, Q_{i,d}) \in \mathbb{P}^1 \times \dots \times \mathbb{P}^1$ s.t. $j_{n_1,\dots,n_d}(\tilde{Q}_i) = Q_i$ and then, according with Notation 2, write $Q_i = (Q_{i,1}, \dots, Q_{i,d})$.

Now write

$$\tilde{O} = (O_1, \dots, O_d) \in \mathbb{P}^1 \times \dots \times \mathbb{P}^1$$

and

$$O = j_{n_1,\dots,n_d}(\tilde{O}).$$

Choose homogeneous coordinates on \mathbb{P}^1 . Since $X_{1,\dots,1}$ is a homogeneous variety, it is sufficient to prove the case $O_i = [1, 0]$ for all $i = 1, \dots, d$.

Notice that $\deg(Z \cup S) = r + 2$ if $O \notin S$ and $\deg(Z \cup S) = r + 1$ if $O \in S$.

Since S computes $r_{X_{1,\dots,1}}(P)$, we have $P \notin \langle j_{n_1,\dots,n_d}(S') \rangle$ for any $\tilde{S}' \subseteq \tilde{S}$. Since $P \neq O$ and $\{O\}$ is the only proper subscheme of $Z = T_O(X_{n_1,\dots,n_d})$, we have $P \notin \langle Z' \rangle$ for all proper subschemes Z' of Z . Since $P \in \langle Z \rangle \cap \langle S \rangle$, then, by Lemma 1, we have $h^1(\mathcal{I}_{S \cup Z}(1)) > 0$. Thus to get a contradiction and prove Theorem 1 in the case $n_i = 1$ for all $i = 1, \dots, d$ and $\eta_{X_{1,\dots,1}}(P) = d$, it is sufficient to prove $h^1(\mathcal{I}_{S \cup Z}(1)) = 0$, i.e. $h^1(\mathbb{P}^1 \times \dots \times \mathbb{P}^1, \mathcal{I}_{\tilde{Z} \cup \tilde{S}}(1, \dots, 1)) = 0$ where, as above, $\tilde{Z} \subset \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_d}$ s.t. $Z = j_{n_1,\dots,n_d}(\tilde{Z})$.

First assume the existence of an integer $j \in \{1, \dots, d\}$ such that $Q_{i,j} = [1, 0]$ for all $i \in \{1, \dots, r\}$.

We get $S \subset X_{1,\dots,1}([1, 0], j)$, where $X_{n_1,\dots,n_d}(Q_j, j)$ is defined in (5). Hence $P \in \langle X_{1,\dots,1}([1, 0], j) \rangle$.

However $T_O X_{1,\dots,1} \cap X_j = \langle \cup_{i \neq j} Y_{O,i} \rangle$. Hence $\eta(P) \leq d - 1$, but this is a contradiction. Thus:

$$\text{for each } j \in \{1, \dots, d\} \text{ there is } Q_{i,j} \in S \text{ such that } Q_{i,j} \neq [1, 0].$$

(a1) Here we assume $O \notin \langle S \rangle$.

Since S computes $r_{X_{1,\dots,1}}(P)$, it is linearly independent, i.e. (by Lemma 2) $h^1(\mathbb{P}^1 \times \dots \times \mathbb{P}^1, \mathcal{I}_{\tilde{S}}(1, \dots, 1)) = 0$.

Since $O \notin \langle S \rangle$, we get that $\tilde{S} \cup \{\tilde{O}\}$ is linearly independent, i.e. $h^1(\mathbb{P}^1 \times \dots \times \mathbb{P}^1, \mathcal{I}_{\tilde{S} \cup \{\tilde{O}\}}(1, \dots, 1)) = 0$.

We fix $i \in \{1, \dots, r\}$ such that $Q_{i,1} \neq [1, 0]$ (we just saw the existence of such an integer i).

Write $S_1 := S \cap X_{1,\dots,1}(Q_{i,1}, 1)$, where $X_{n_1,\dots,n_d}(Q_j, j)$ is defined in (5). By construction $Q_i \in S_1$ and hence $\sharp(S_1) \geq 1$.

Assume for now that $S_1 \neq S$ and that there exist $j \in S \setminus S_1$ such that $Q_{j,2} \neq [1, 0]$. Set $S_2 := S \cap X_{1,\dots,1}(Q_{j,2})$. And so on constructing subsets S_1, \dots, S_j of S such that:

- $S_j \not\subseteq \cup_{1 \leq i < j} S_i$,
- $Q_{k,i} \neq [1, 0]$ for all $k \in S_i$,
- $S_i = S \cap X_{1,\dots,1}(Q_{h,i}, i)$ for all $h \in S_i$,

until we arrive at one of the following cases:

- (i) $S_1 \cup \cdots \cup S_j = S$;
- (ii) $S_1 \cup \cdots \cup S_j \neq S$ and $Q_{k,j+1} = [1, 0]$ for all $k \in S \setminus (S_1 \cup \cdots \cup S_j)$.

Now fix an index $m_{i+1} \in S_{i+1} \setminus S_i$, $1 \leq i \leq j-1$, and set

$$D_i := X_{1,\dots,1}(Q_{m_i,i}, i), \quad 1 \leq i \leq j,$$

i.e. according with (5), $D_i := \{(A_1, \dots, A_d) \in X_{n_1, \dots, n_d} : A_i = Q_{m_i}$ with $m_i \in S_i \setminus S_{i-1}\}$ for $1 \leq i \leq j$.

First assume that (i) occurs (with j minimal).

Fix $B_i \in \mathbb{P}^1 \setminus \{[1, 0]\}$, $j+1 \leq i \leq d-1$ and set:

- $D_i := X_{1,\dots,1}(B_i, i)$, if $j+1 \leq i \leq d-1$;
- $D_d := X_{1,\dots,1}(O_d, d)$;
- $D := \cup_{i=1}^d D_i$.

Notice that obviously $D \in |\mathcal{O}_{X_{1,\dots,1}}(1)|$ and also that $S \cup \{O\} \subset D$.

Moreover observe that $O \in D_i$ if and only if $i = d$.

Finally, D_d is smooth at O and $T_O D$ is spanned by $\cup_{i=1}^{d-1} X_{1,\dots,1}(O_1, i)$.

Therefore $Z \not\subseteq D$ and $Z \cup S$ imposes one more condition to $|\mathcal{O}_{\mathbb{P}^1 \times \cdots \times \mathbb{P}^1}(1, \dots, 1)|$ than $S \cup \{O\}$.

Since $j_{1,\dots,1}(\tilde{S} \cup \{\tilde{O}\})$ is linearly independent, we get $h^1(\mathbb{P}^1 \times \cdots \times \mathbb{P}^1, \mathcal{I}_{\tilde{Z} \cup \tilde{S}}(1, \dots, 1)) = 0$ that is a contradiction.

Now assume that (ii) occurs and set:

- $M_{j+1} = X_{1,\dots,1}([1, 0], j+1)$;
- $M_h := X_{1,\dots,1}([1, 0], h)$, for all $h \in \{j+2, \dots, d\}$;
- $D' := \cup_{i=1}^j D_i \cup \cup_{h=j+1}^d M_h$.

Notice that $D' \in |\mathcal{O}_{X_{1,\dots,1}}(1)|$ and that $S \cup \{O\} \subset D'$.

The hypersurface M_{j+1} is the unique irreducible component of D' containing O .

Since M_{j+1} is smooth at O and $T_O M_{j+1}$ is spanned by $\cup_{i \neq j+1}^{d-1} X_{1,\dots,1}(O_1, i)$, we get as above that $h^1(\mathbb{P}^1 \times \cdots \times \mathbb{P}^1, \mathcal{I}_{\tilde{Z} \cup \tilde{S}}(1, \dots, 1)) = 0$, and than another contradiction.

(a2) Here we assume $O \in S$.

Hence $S \cup \{O\} = S$ and $j_{1,\dots,1}(\tilde{S} \cup \{\tilde{O}\})$ is linearly independent. Set $S' := S \setminus \{O\}$. We make the construction of step (a1) with S' instead of S , defining the subsets S_i of S' until we get an integer j such that either $S' = S_1 \cup \cdots \cup S_j$ or $S_1 \cup \cdots \cup S_j \neq S'$ and $Q_{j+1,i} = [1, 0]$ for all $i \in S' \setminus (S_1 \cup \cdots \cup S_j)$. In both cases we add the other $d-j$ hypersurfaces, exactly one of them containing O . Since $\deg(Z \cup S) = \deg(S \cup \{O\}) + 1$, we get $h^1(\mathbb{P}^1 \times \cdots \times \mathbb{P}^1, \mathcal{I}_{\tilde{Z} \cup \tilde{S}}(1, \dots, 1)) = 0$ as in step (a1) and hence we get a contradiction.

(a3) Here assume $O \notin S$ and $O \in \langle S \rangle$.

Hence $\langle Z \rangle \subset \langle S \rangle$. Thus there is $S' \subset S$ such that $\sharp(S') = \sharp(S) - 1$ and $\langle S' \cup \{O\} \rangle = \langle S \rangle$. Hence the set $S_1 := S' \cup \{O\}$ computes $r_{X_{1,\dots,1}}(P)$. Apply step (a2) to the set S_1 .

(b) Here we assume $n_i = 1$ for all i and $r := \eta_{X_{1,\dots,1}}(P) < d$.

Let $E \subset \{1, \dots, d\}$ be the minimal subset such that $P \in \langle \cup_{i \in E} Y_{O,i} \rangle$. By the definition of the type $\eta_{X_{1,\dots,1}}(P)$ of P we have $\sharp(E) = \eta_{X_{1,\dots,1}}(P)$. Set $X' := \{(U_1, \dots, U_d) \in X_{1,\dots,1} : U_i = [1, 0] \text{ for all } i \notin E\}$. We identify X' with a Segre product of r copies of \mathbb{P}^1 . Obviously $\eta_{X'}(P) = \eta_{X_{1,\dots,1}}(P)$. By step (a) we have $r_{X'}(P) = \eta_{X'}(P)$. We have $r_{X_{1,\dots,1}}(P) = r_{X'}(P)$ by the concision property of tensors ([9], Corollary 2.2, or [16], Proposition 3.1.4.1).

(c) Here we assume $n_i \geq 2$ for some i .

Since $P \in \langle \cup_{i=1}^d Y_{O,i} \rangle$, there is $U_i \in Y_{O,i}$ such that $P \in \langle \{U_1, \dots, U_d\} \rangle$. Let $U_i^i \in \mathbb{P}^{n_i}$ be the i -th component of U_i . The line $L_i \subseteq \mathbb{P}^{n_i}$ is the line spanned by O_i and U_i^i . We have

$P \in \langle \prod_{i=1}^d L_i \rangle$ and $\eta_{X_{n_1, \dots, n_d}}(P) = \eta_{X_{1, \dots, 1}}(P)$, where we identify $j_{n_1, \dots, n_d}(\prod_{i=1}^d L_i)$ with the Segre variety $X_{1, \dots, 1}$. By parts (a) and (b) we have $r_{X_{1, \dots, 1}}(P) = \eta_{X_{1, \dots, 1}}(P)$. We have $r_{X_{n_1, \dots, n_d}}(P) = r_{X_{1, \dots, 1}}(P)$ by the concision property of tensors ([9], Corollary 2.2, or [16], Proposition 3.1.4.1). \square

Corollary 1. *Let $P \in \sigma_2(X_{n_1, \dots, n_d})$, then:*

- $r_{X_{n_1, \dots, n_d}}(P) = 1$ iff $P \in X_{n_1, \dots, n_d}$;
- $r_{X_{n_1, \dots, n_d}}(P) = 2$ iff either $P \in \sigma_2(X_{n_1, \dots, n_d}) \setminus \tau(X_{n_1, \dots, n_d})$ or there exist $O \in X_{n_1, \dots, n_d}$, $O \neq P$, and $Y_{O,i}, Y_{O,j} \subset \mathbb{P}^{N(n_1, \dots, n_d)}$ as in Notation 3, such that $P \in T_O(X_{n_1, \dots, n_k}) \subset Y_{O,i} \cup Y_{O,j}$ for certain $i \neq j \in \{1, \dots, d\}$;
- $r_{X_{n_1, \dots, n_d}}(P) = k$ with $3 \leq k \leq d$ iff k is the minimum integer s.t. there exist $Y_{O,i_1}, \dots, Y_{O,i_k} \subset \mathbb{P}^{N(n_1, \dots, n_d)}$ as in Notation 3, such that $P \in T_O(X_{n_1, \dots, n_k}) \subset \cup_{j=1, \dots, k} Y_{O,i_j}$ for certain $i_j \in \{1, \dots, d\}$, $j = 1, \dots, k$.

Proof. This corollary follows straightforward from Theorem 1 and the fact that $\sigma_2(X_{n_1, \dots, n_d}) \setminus \tau(X_{n_1, \dots, n_d}) = \sigma_2^0(X_{n_1, \dots, n_d})$ when it is not empty. \square

The three cases of this Corollary actually occur and can be deduced from the proof of Theorem 1.

Example 1. Let us write for convenience $\mathbb{P}^{n_i} = \mathbb{P}(V_i)$ for certain $(n_i + 1)$ -dimensional vector spaces over K .

- The points $P \in \mathbb{P}(V_1 \otimes \dots \otimes V_d)$ for which there exist $v_i \in V_i$, for $i = 1, \dots, d$, such that $P = [v_1 \otimes \dots \otimes v_d]$, have $r_{X_{n_1, \dots, n_d}}(P) = 1$.
- Let $P_1 = [v_{1,1} \otimes \dots \otimes v_{1,d}]$, $P_2 = [v_{2,1} \otimes \dots \otimes v_{2,d}] \in X_{n_1, \dots, n_d}$ with $v_{1,1} \otimes \dots \otimes v_{1,d}, v_{2,1} \otimes \dots \otimes v_{2,d} \in V_1 \otimes \dots \otimes V_d$ linearly independent, then $P = \lambda_1 P_1 + \lambda_2 P_2$, for non-zero coefficients $\lambda_1, \lambda_2 \in K$, has $r_{X_{n_1, \dots, n_d}}(P) = 2$.
- We can observe that, for any $r \leq d$, with an abuse of notation, there is an obvious way to see $V_1 \otimes \dots \otimes V_r$ as a natural subspace of $V_1 \otimes \dots \otimes V_d$. Roughly speaking this is the same to say that the Segre variety of r factors can be seen as a subvariety of the Segre variety of d factors. Let $O = [w_1 \otimes \dots \otimes w_r] \in X_{n_1, \dots, n_r} \subseteq X_{n_1, \dots, n_d}$. Take $v_i \in V_i$, $i = 1, \dots, r$, such that $\{w_1, \dots, v_i, \dots, w_r\}$ are linearly independent, then $P = \lambda_1 [v_1 \otimes w_2 \otimes \dots \otimes w_r] + \dots + \lambda_r [w_1 \otimes \dots \otimes w_{r-1} \otimes v_r]$ has rank r certain non zero $\lambda_1, \dots, \lambda_r \in K$.

3. ALGORITHMS

The proof of Theorem 1 turns out to be useful to produce an algorithm to compute the rank of tensor of border rank 2 (Algorithm 1) and also an Algorithm that gives one of its decompositions (Algorithm 2).

We need to introduce the notion of Flattening and the definition of Hankel operator.

Definition 5. Let V_1, \dots, V_d be vector spaces of dimension $n_1 + 1, \dots, n_d + 1$ respectively. Let (J_1, J_2) be a partition of the set $\{1, \dots, d\}$. If $J_1 = \{h_1, \dots, h_s\}$ and $J_2 = \{1, \dots, d\} \setminus J_1 = \{k_1, \dots, k_{d-s}\}$, the (J_1, J_2) -Flattening of $V_1 \otimes \dots \otimes V_d$ is the following:

$$V_{J_1} \otimes V_{J_2} = (V_{h_1} \otimes \dots \otimes V_{h_s}) \otimes \dots \otimes (V_{k_1} \otimes \dots \otimes V_{k_{d-s}}).$$

Definition 6. Let $n := \sum_{i=1}^d n_i$, set $R := K[x_1, \dots, x_n]$. For any $\Lambda \in R^*$, we define the Hankel operator H_Λ as $H_\Lambda : R \rightarrow R^*$, $p \mapsto p \cdot \Lambda$ where $p \cdot \Lambda$ is the linear operator $p \cdot \Lambda : R \rightarrow K$, $q \mapsto \Lambda(pq)$.

Algorithm 1 (Rank of a border rank 2 tensor).

Input: A tensor $T \in V_1 \otimes \cdots \otimes V_d$, with V_1, \dots, V_d vector spaces of dimension n_1+1, \dots, n_d+1 respectively.

Output: Either $T \notin \sigma_2(X_{n_1, \dots, n_d})$, or the rank of T .

- (1) Write T as an element of $V_{J_1} \otimes V_{J_2}$ for any (J_1, J_2) -Flattening of $V_1 \otimes \cdots \otimes V_d$.
- (2) Compute all the 2×2 minors of $V_{J_1} \otimes V_{J_2}$ for any (J_1, J_2) -Flattening of $V_1 \otimes \cdots \otimes V_d$. If all of them are equal to 0, then $r(T) = 1$ (see eg [15]), otherwise go to Step (3).
- (3) Compute all the 3×3 minors of $V_{J_1} \otimes V_{J_2}$ for any (J_1, J_2) -Flattening of $V_1 \otimes \cdots \otimes V_d$. If at least one of them is different from 0, then $T \notin \sigma_2(X_{n_1, \dots, n_d})$ and this algorithm stops here; otherwise $T \in \sigma_2(X_{n_1, \dots, n_d})$ (see [17]) and go to Step (4).
- (4) Find $\Lambda \in (V_1 \otimes \cdots \otimes V_d)^*$ that extends T^* (for a precise definition of extension see [4]) and such that $rl(H_\Lambda) = 2$ then pass to Step (5).
- (5) Compute the roots of $\ker H_\Lambda$ by generalized eigenvector computation (see [8]) and check if the eigenspaces are simple. If yes then the rank of T is 2 (see [4]), otherwise go to Step (6).
- (6) Write T as a multilinear polynomial t in the ring $K[x_{1,0}, \dots, x_{1,n_1}; \dots; x_{d,0}, \dots, x_{d,n_d}]$, then pass to Step (7).
- (7) Use [10] to write t in the minimum number q of variables. Then the rank of t is equal to $q/2$ (in fact, from the proof of Theorem 1, it is always possible to write T as an element of $\tau(X_{1, \dots, 1})$, then its representative polynomial will be a multilinear form in $K[l_{1,0}, l_{1,1}; \dots; l_{q,0}, l_{q,1}]$ with $l_{i,0}, l_{i,1}$ linear forms in $K[x_{i,0}, \dots, x_{i,n_i}]$ for $i = 1, \dots, q$).

Algorithm 2 (Decomposition of a border rank 2 tensor).

Input: A tensor $T \in V_1 \otimes \cdots \otimes V_d$, with V_1, \dots, V_d vector spaces of dimension n_1+1, \dots, n_d respectively.

Output: Either $T \notin \sigma_2(X_{n_1, \dots, n_d})$, or a decomposition of T .

- (a) Write T as a multilinear polynomial t in the ring $K[x_{1,0}, \dots, x_{1,n_1}; \dots; x_{d,0}, \dots, x_{d,n_d}]$.
- (b) Use [10] to write t in the minimum number of variables. Then, from the proof of Theorem 1, it is always possible to write t as a multilinear form in $K[l_{1,0}, l_{1,1}; \dots; l_{d,0}, l_{d,1}]$ with $l_{i,0}, l_{i,1}$ linear forms in $K[x_{i,0}, \dots, x_{i,n_i}]$ for $i = 1, \dots, d$.
- (c) Run Algorithm 1. If Algorithm 1 stops at Step (2), go to Step (d). If Algorithm 1 stops at Step (3), then $T \notin \sigma_2(X_{n_1, \dots, n_d})$. If Algorithm 1 stops at Step (5), go to Step (e). Otherwise go to Step (f).
- (d) In this case the rank of T is 1, then it is sufficient to find $m_i(l_{i,0}, l_{i,1})$ linear forms in $K[l_{i,0}, l_{i,1}]$, for $i = 1, \dots, d$ such that $t = m_1(l_{1,0}, l_{1,1}) \cdots m_d(l_{d,0}, l_{d,1})$.
- (e) In this case the rank of T is 2, then it is sufficient to find $m_{i,j}(l_{i,0}, l_{i,1})$ linear forms in $K[l_{i,0}, l_{i,1}]$, for $i = 1, \dots, d$ and $j = 1, 2$, such that $t = m_{1,1}(l_{1,0}, l_{1,1}) \cdots m_{d,1}(l_{d,0}, l_{d,1}) + m_{1,2}(l_{1,0}, l_{1,1}) \cdots m_{d,2}(l_{d,0}, l_{d,1})$.
- (f) In this case the rank of T is $q/2$ and $t \in K[l_{1,0}, l_{1,1}; \dots; l_{q,0}, l_{q,1}]$ for certain $q \leq d$ and there exist q 2-dimensional subspaces $W_i \subset V_i$ such that T belongs to the Segre variety $X_{1, \dots, 1} \subset \mathbb{P}(W_1 \otimes \cdots \otimes W_q)$. Let $L_T \subset W_1 \otimes \cdots \otimes W_q$ be a generic space of dimension q passing through T and compute the unique point $O \in X_{1, \dots, 1}$ such that $[T] \in T_O(X_{1, \dots, 1})$ (it is sufficient to impose that $\mathbb{P}(L_T) \cap X_{1, \dots, 1}$ has a double solution). Let $O = n_1(l_{1,0}, l_{1,1}) \cdots n_q(l_{q,0}, l_{q,1})$ and go to Step (g).
- (g) Now it is sufficient to find $m_i(l_{i,0}, l_{i,1})$ linear forms in $K[l_{i,0}, l_{i,1}]$, for $i = 1, \dots, q$ such that $t = m_1(l_{1,0}, l_{1,1}) \cdots n_q(l_{q,0}, l_{q,1}) + \cdots + n_1(l_{1,0}, l_{1,1}) \cdots m_q(l_{q,0}, l_{q,1})$.

4. ON COMON'S CONJECTURE

In this section we want to relate the result obtained in Theorem 1 to the Comon's conjecture stated in the Introduction.

Let $\nu_d(\mathbb{P}^n)$ be the classical Veronese embedding of \mathbb{P}^n into $\mathbb{P}^{\binom{n+d}{d}-1}$ via the sections of the sheaf $\mathcal{O}(d)$. As pointed out in the introduction if $\mathbb{P}^n \simeq \mathbb{P}(V)$ with V an $(n+1)$ -dimensional vector space, then $\nu_d(\mathbb{P}^n) \subset \mathbb{P}(S^d V)$ can be interpreted as the variety that parameterizes projective classes of completely decomposable symmetric tensors $T \in S^d V$. Moreover

$$\nu_d(\mathbb{P}^n) = X_{n,\dots,n} \cap \mathbb{P}(S^d V) \subset \mathbb{P}(V^{\otimes d}).$$

Definition 7. Let $P \in \mathbb{P}(S^d V)$ be a projective class of a symmetric tensor. We define the symmetric rank $r_{\nu_d(\mathbb{P}^n)}(P)$ of P as the minimum number of r of points $P_i \in \nu_d(\mathbb{P}^n)$ whose linear span contains P .

With this definition, **Comon's conjecture** (Conjecture 1) can be rephrased as follows:

$$\text{if } P \in \mathbb{P}(S^d V) \text{ then } r_{\nu_d(\mathbb{P}^n)}(P) = r_{X_{n,\dots,n}}(P).$$

Obviously $r_{X_{n,\dots,n}}(P) \leq r_{\nu_d(\mathbb{P}^n)}(P)$. In [12] the authors prove the reverse inequality for a general d -tensor (d even and large) with rank at most n (Proposition 5.3) and for $r_{X_{n,\dots,n}}(P) = 1, 2$.

With Theorem 1 we can prove that conjecture for all symmetric tensors of border rank 2.

Corollary 2. Let $P \in \sigma_2(\nu_d(\mathbb{P}^n))$. Then $r_{\nu_d(\mathbb{P}^n)}(P) = r_{X_{n,\dots,n}}(P)$.

Proof. For any projective variety X we can observe that $\sigma_2(X) = X \cup \tau(X) \cup \sigma_2^0(X)$. If $P \in \nu_d(\mathbb{P}^n) \subset X_{n,\dots,n}$ then there exist $v \in V$ such that $P = [v^{\otimes d}] \in \nu_d(\mathbb{P}^n) \subset X_{n,\dots,n}$, therefore obviously $r_{X_{n,\dots,n}}(P) = r_{\nu_d(\mathbb{P}^n)}(P) = 1$.

If $P \in \sigma_2^0(\nu_d(\mathbb{P}^n))$ then $r_{\nu_d(\mathbb{P}^n)}(P) = 2$, that implies that $r_{X_{n,\dots,n}}(P) \leq 2$, and therefore by [12], that we have that $r_{X_{n,\dots,n}}(P) = r_{\nu_d(\mathbb{P}^n)}(P) = 2$.

Now assume that $P \in \tau(\nu_d(\mathbb{P}^n)) \setminus \nu_d(\mathbb{P}^n)$ and that $\sigma_2(\nu_d(\mathbb{P}^n)) \neq \tau(\nu_d(\mathbb{P}^n))$. For such a P we know that $r_{\nu_d(\mathbb{P}^n)}(P) = d$ (see [24], [11], [6], [5]). Any point $P \in \tau(\nu_d(\mathbb{P}^n)) \setminus \nu_d(\mathbb{P}^n)$ can be thought as the projective class of a homogeneous degree d polynomial in $n+1$ variables for which there exist two linear forms L, M in $n+1$ variables such that $P = [L^{d-1}M]$; hence d is the minimum integer k such that $P \in \langle \nu_k(\mathbb{P}^n) \rangle$. Therefore $\eta_{X_{n,\dots,n}}(P) = d$. Since obviously $\tau(\nu_d(\mathbb{P}^n)) \subset \tau(X_{n,\dots,n})$ we have that, by Theorem 1, $r_{X_{n,\dots,n}}(P) = \eta_{X_{n,\dots,n}}(P)$. \square

REFERENCES

- [1] A. L. F. de Almeida, G. Favier, and J. C. M. Mota, Parafacbased unified tensor modeling for wireless communication systems with application to blind multiuser equalization. *Signal Processing*, 87 (2007), 337–351.
- [2] E. Ballico and A. Bernardi, Decomposition of homogeneous polynomials with low rank. *Math. Z.*, (2011). DOI: 10.1007/s00209-011-0907-6.
- [3] P. Bartho, C. Curto, A. Luczak, S. Marguet, K.D. Harris, Population coding of tone stimuli in auditory cortex: dynamic rate vector analysis. *European Journal of Neuroscience*, 30 (2009), 1767–1778.
- [4] A. Bernardi, J. Brachat, P. Comon, B. Mourrain, Multihomogeneous Polynomial Decomposition using Moment Matrices. ISAAC 2011, Proceedings of the 36th International Symposium of Symbolic and Algebraic Computation (Held in San Jose, California, USA, June, 8-11, 2011), ACM New York, (2011), 35–42.
- [5] A. Bernardi, A. Gimigliano and M. Idà, On the stratification of secant varieties of Veronese varieties via symmetric rank. *J. Symbolic. Comput.* 46 (2011), 34–55.
- [6] J. Brachat, P. Comon, B. Mourrain, E. P. Tsigaridas, Symmetric tensor decomposition. *Linear Algebra and Applications* 433 (2010) 851–1872.
- [7] W. Bruno, G. Rota, and D. Torney, Probability set functions. *Annals of Combinatorics*, 3 (1999), 13–25.

- [8] J. Buczyński, A. Ginensky and J. M. Landsberg, Determinantal equations for secant varieties and the Eisenbud-Koh-Stillman conjecture. arXiv:1007.0192v3 [math.AG].
- [9] J. Buczyński and J. M. Landsberg, Ranks of tensors and a generalization of secant varieties. arXiv:0909.4262 [math.AG].
- [10] E. Carlini, Reducing the number of variables of a polynomial. Elkadi, M., Mourrain, B., Piene, R. (Eds.), Algebraic Geometry and Geometric Modeling. Springer, (2005), 237–247.
- [11] G. Comas, M. Seiguer, On the rank of a binary form. Found. Comput. Math. 11 (2011), 65–78.
- [12] P. Comon and G. Golub and L-H. Lim and B. Mourrain, Symmetric Tensors and Symmetric Tensor Rank. SIAM Journal on Matrix Analysis Appl. 30 (2008), 1254–1279.
- [13] P. Comon, M. Sorensen, and E. Tsigaridas, Decomposing tensors with structured matrix factors reduces to rank-1 approximations. In ICASSP'2010, Dallas, March (2010), 14–19.
- [14] B. Georgi and A. Schliep, Context-specific independence mixture modeling for positional weight matrices. Bioinformatics, 22 (2006), 166–173.
- [15] H.T.Hà, Box shaped matrices and the defining ideal of certain blowup surface. J. Pure Appl. Algebra 167 (2002), 203– 224.
- [16] J. M. Landsberg, Tensors: Geometry and Applications. book in preparation, version of 8 August 2010.
- [17] J. M. Landsberg, L. Manivel, On the ideals of secant varieties of Segre varieties. Found. Comput. Math. 4 (2004), 397–422.
- [18] J. M. Landsberg, J. Weyman, On the ideals and singularities of secant varieties of Segre varieties. Bull. Lond. Math. Soc. 39 (2007), 685–697.
- [19] P. McCullagh, Tensor Methods in Statistics, Monographs on Statistics and Applied Probability. Chapman and Hall, London, (1987).
- [20] D. Nion and L. De Lathauwer, A block component model based blind DS-CDMA receiver. IEEE Trans. Signal Proc., 56 (2008), 5567–5579.
- [21] L. Oeding, Report on Geometry and representation theory of tensors for computer science, statistics and other areas. arXiv:0810.3940 [math.AG]
- [22] L. Oeding, Set-theoretic defining equations of the tangential variety of the Segre variety. J. Pure and Applied Algebra, 215 (2011), 1516–1527.
- [23] B. Sturmfels and P. Zwiernik, Binary cumulant varieties. arXiv:1103.0153 [math.CO].
- [24] J. J. Sylvester, Sur une extension d'un théorème de Clebsch relatif aux courbes du quatrième degré. Comptes Rendus, Math. Acad. Sci. Paris, 102 (1886), 1532–1534.
- [25] D.C. Torney, Binary cumulants. Advances in Applied Math, 25 (2000), 34–40.

DEPT. OF MATHEMATICS, UNIVERSITY OF TRENTO, 38123 POVO (TN), ITALY
E-mail address: `ballico@science.unitn.it`

GALAAD, INRIA 2004 ROUTE DES LUCIOLES, BP 93, 06902 SOPHIA ANTIPOLIS, FRANCE.
E-mail address: `alessandra.bernardi@inria.fr`