



HAL
open science

Anisotropic Delaunay Mesh Generation

Jean-Daniel Boissonnat, Camille Wormser, Mariette Yvinec

► **To cite this version:**

Jean-Daniel Boissonnat, Camille Wormser, Mariette Yvinec. Anisotropic Delaunay Mesh Generation. [Research Report] RR-7712, Inria Sophia Antipolis; INRIA. 2014, pp.50. inria-00615486v2

HAL Id: inria-00615486

<https://hal.inria.fr/inria-00615486v2>

Submitted on 19 Dec 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Anisotropic Delaunay Mesh Generation

Jean-Daniel Boissonnat, Camille Wormser, Mariette Yvinec

**RESEARCH
REPORT**

N° 7712

Août 2011

Project-Team Geometrica



Anisotropic Delaunay Mesh Generation

Jean-Daniel Boissonnat, Camille Wormser, Mariette Yvinec

Project-Team Geometrica

Research Report n° 7712 — version 2 — initial version Août 2011 —
revised version Décembre 2014 — 50 pages

Abstract: Anisotropic meshes are triangulations of a given domain in the plane or in higher dimensions, with elements elongated along prescribed directions. Anisotropic triangulations are known to be well suited for interpolation of functions or solving PDEs. Assuming that the anisotropic shape requirements for mesh elements are given through a metric field varying over the domain, we propose a new approach to anisotropic mesh generation, relying on the notion of anisotropic Delaunay meshes. An anisotropic Delaunay mesh is defined as a mesh in which the star of each vertex v consists of simplices that are Delaunay for the metric associated to vertex v . This definition works in any dimension and allows to define a simple refinement algorithm. The algorithm takes as input a domain and a metric field and provides, after completion, an anisotropic mesh whose elements are shaped according to the metric field.

Key-words: mesh generation, anisotropic meshes, Delaunay triangulation

This research has been partially supported by the Agence Nationale de la Recherche (project GIGA).

**RESEARCH CENTRE
SOPHIA ANTIPOLIS – MÉDITERRANÉE**

2004 route des Lucioles - BP 93
06902 Sophia Antipolis Cedex

Génération de maillages de Delaunay anisotropes

Résumé : Les maillages anisotropes sont des triangulations d'un domaine donné du plan ou d'un espace de plus grande dimension dont les éléments sont étirés selon des directions prescrites. Les maillages anisotropes sont utiles pour interpoler des fonctions ou résoudre des EDP. Dans cet article, nous supposons que l'anisotropie est prescrite par un champ de métrique défini sur le domaine à mailler. Nous proposons une nouvelle approche de génération de maillages anisotropes qui s'appuie sur la notion de maillage de Delaunay anisotrope. Un tel maillage est défini comme un maillage dont l'étoile de chaque sommet v est formée de simplexes qui sont de Delaunay pour la métrique de v . Cette définition est valide en toutes dimensions et un tel maillage peut être construit par un algorithme simple de raffinement.

Mots-clés : Génération de maillages, maillages anisotropes, triangulation de Delaunay

1 Introduction

Anisotropic meshes are triangulations of a given domain in the plane or in higher dimensions, with elements elongated along prescribed directions. Anisotropic triangulations have been shown to be particularly well suited for interpolation of functions [18, 36] and for solving PDEs [5]. They allow to minimize the number of elements in the mesh while retaining a good accuracy in computations.

The required anisotropy is generally described through a metric field defined over the domain to be meshed. The directions along which the elements should be elongated are usually given, at each point of the domain, as a quadratic form. The eigenvectors and eigenvalues of the quadratic form describe the preferred directions and their anisotropic ratios.

Two main issues arise in this context. The first is to define the metric field. The second one is to generate a mesh whose elements are shaped according to the chosen metric field.

Defining good metric fields and error estimates is still an active research area. Alauzet et al. introduced the notion of continuous metrics and continuous meshes to minimize interpolation error [3, 30, 2]. Loseille et al. [31] applied this notion to a posteriori error estimates in order to minimize the approximation error during the process of solving some PDEs. Chen et al [12] considered anisotropic finite element approximation of functions in the L^p norm. Their result reveals that the accuracy of the approximation is governed by a quantity that depends non-linearly on the hessian of the function. In his thesis, Mirebeau [32] extends this result to finite elements of arbitrary degree and to Sobolev norms, and provides sharp asymptotic error estimates for the approximation of functions of two variables.

Various methods have been proposed to generate anisotropic meshes whose elements are shaped according to a given metric field. In their early work on 2D meshes, Bossen and Heckbert [11] proposed to adapt their *pliant* method for mesh generation to the anisotropic setting. Starting from a constrained Delaunay triangulation, the pliant method performs local optimization operations including centroidal smoothing and retriangulation, and possibly insertion or removal of vertices. Li et al. [29] and Shimada et al. [39] have proposed to place the mesh vertices close to the centers of ellipsoid bubbles optimally packed in the domain. Borouchaki et al. [10] proposed to adapt the standard Delaunay incremental construction to the anisotropic context. This construction is then combined with an anisotropic version of the *unit mesh* approach that aims at producing meshes whose edges have unit length. Lengths, in the anisotropic case, are measured in the Riemannian metric provided by the metric field. The efficiency of the method has been demonstrated in various contexts [24, 20].

Following a different line of research, some attempts have been done recently to define anisotropic Delaunay triangulation and meshes as the duals of some Voronoi diagrams derived from the metric field. Labelle and Shewchuk [25] have defined an anisotropic mesh as the dual of the so-called anisotropic Voronoi diagram. The sites of this diagram are the mesh vertices and the distance to a site is computed with respect to the metric attached to this site. In the 2-dimensional case, Labelle and Shewchuk have proposed a refinement algorithm that can provably produce anisotropic meshes. Their approach has somehow been simplified in [6], leading to a direct computation of the dual mesh, and extended by Cheng et al. [15] to produce anisotropic meshes of surfaces embedded in 3D. Extending Labelle and Shewchuk's approach to higher dimensions seems however difficult due to the presence of flat tetrahedra called *slivers* [35]. Du and Wang [21] have proposed to use a definition of anisotropic Voronoi diagrams which is somewhat symmetric to the definition used by Labelle and Shewchuk. The Voronoi regions are based on distances from points to sites that are computed with respect to the metric of the point. Du and Wang compute centroidal Voronoi diagrams using this definition and show experimentally that the dual structures are generally anisotropic meshes of high quality. However they could

not provide theoretical guarantees or conditions that ensure that the dual structure is a valid triangulation.

In this paper, we introduce a new notion of anisotropic mesh which extends nicely in any dimension. The resulting meshes can be computed using standard Delaunay algorithms. As in the previous approaches, we assume that the anisotropy is prescribed by a metric field that associates to each point p of the domain a symmetric positive definite square matrix M_p , describing the metric at point p . Given a set of points V called *sites*, we consider, for each site $v \in V$, the Delaunay triangulation $\text{Del}_v(V)$ of V , computed for the metric M_v attached to location of v . Each triangulation $\text{Del}_v(V)$ is well defined and can be computed using the standard Euclidean Delaunay triangulation on affinely transformed input points. For each site $v \in V$, we keep the *star* S_v of v in $\text{Del}_v(V)$, i.e. the set of simplices of $\text{Del}_v(V)$ that are incident to v . The collection of stars is called the *star set* of V . In general, there are *inconsistencies* among the stars : a simplex s may appear in the stars of some of its vertices without appearing in the stars of all of them. As a result, the simplices in the star set of V do not form a triangulation of V . However, we show in this paper that, given a compact domain of \mathbb{R}^d and a smooth metric field, one can insert new sites in V at carefully chosen locations so that all inconsistencies are removed. The simplices in the star set then form a d -dimensional triangulation that we call an *anisotropic Delaunay mesh*. [When the domain has smooth boundaries, a faithful representation of those boundaries may be obtained using the method of restricted Delaunay triangulations. The refinement algorithm is then extended to achieve also consistency between *surface stars* which are defined as the restrictions of the stars to the boundary surfaces. The algorithm produces then a mesh whose vertices lie within the input domain and whose boundary is within a controlled Hausdorff distance from the input domain boundary. Sharp features could possibly be handled using protecting balls but this issue is not handled in the present paper.](#)

The idea of maintaining independent stars for each vertex of a mesh has been first proposed by Shewchuk [38] for maintaining triangulations of moving points. The star set has also been used [35] to build the dual of an anisotropic Voronoi diagram as defined by Labelle and Shewchuk [25]. The method we use to ensure consistency among the stars is inspired by the work of Li and Teng [28, 27] for removing slivers in isotropic meshes. In our context, the method is extended so as to take into account the metric distortion between neighboring stars and also to avoid, in addition to slivers, more general quasi-cospherical configurations that may prevent the termination of the algorithm.

In addition to conforming to the given anisotropic metric field, this mesh generation method has several notable advantages.

- It is not limited to the plane and works in any dimension;
- It is easy to implement. Through a stretching transform, the star of each vertex in the mesh can be computed as part of an Euclidean Delaunay triangulation. Therefore the algorithm relies only on the usual Delaunay predicates (applied in some stretched spaces);
- The star of each vertex in the output mesh is formed with simplices that are Delaunay with respect to the metric of the central vertex. This provides a neat characterization of the output mesh from its set of vertices.
- The method provides some theoretical guarantees on the size and shape of the output mesh elements. Each element is guaranteed to be sized and shaped according to the metrics of all its vertices.

[The main downside of this anisotropic Delaunay mesh approach is that no consistent mesh is obtained before reaching the very end of the refinement algorithm. This may leads to over](#)

dense meshes when the metric field is highly distorted. In such cases, the only way out consists in somehow smoothing the input metric field.

This paper is an extension of a preliminary work limited to the 3-dimensional case, that has been presented at the Symposium on Computational Geometry [9].

The paper is organized as follows. Section 2 recalls basic facts about anisotropic metrics, metric fields, metric distortion and sizing fields. Section 3 introduces the main notions underlying our refinement strategy: restricted Delaunay triangulation, star sets, inconsistencies, slivers and quasi-cosphericities. Section 4 presents the anisotropic mesh generation algorithm. For pedagogical reason, we focus in this section on the generation of a mesh covering a given domain and conforming to a varying field of anisotropic metrics defined on this domain. We postpone to section 7 the additional problem to get into the mesh a faithful representation of the domain boundaries and internal subdivisions. Sections 5 and 6 detail the proof that the refinement algorithm terminates. Section 7 explains how to handle domain boundaries and sharp features in the anisotropic setting. At last Section 8 provides concluding remarks and some insights on on-going and future work.

2 Preliminaries

2.1 Anisotropic Metric

An anisotropic metric in \mathbb{R}^d is defined by a symmetric positive definite quadratic form represented, in some vector basis, by a $d \times d$ matrix M . The distance between two points a and b as measured by metric M , is defined as

$$d_M(a, b) = \sqrt{(a - b)^T M (a - b)}.$$

This definition provides a definition for M -lengths and, by integration, for higher dimensional M -volume measures.

In the following, we often use the same notation, M , for a metric and the associated matrix in a given basis. Given the symmetric positive definite matrix M , we denote by F_M any matrix such that $\det(F_M) > 0$ and $F_M^T F_M = M$. Note that F_M is not unique. The Cholesky decomposition provides an upper triangular F_M , while a symmetric F_M can be obtained by diagonalizing the quadratic form M and computing the quadratic form with the same eigenvectors and the square root of each eigenvalue.

Note that

$$d_M(a, b) = \sqrt{(a - b)^T F_M^T F_M (a - b)} = \|F_M(a - b)\| \quad (1)$$

where the notation $\|\cdot\|$ stands for the Euclidean norm. Equation (1) proves that d_M is a distance and, in particular, enjoys the standard triangular inequality. In the following we call such a F_M matrix a *stretching transform* of M .

Given some metric M , an M -sphere $C_M(c, r)$, with center c and radius r , is defined as the set of points p such that $d_M(c, p) = r$, and likewise an M -ball $B_M(c, r)$, is defined as the set of points p such that $d_M(c, p) \leq r$. Note that an M -sphere is an ellipsoid in the Euclidean space, with its axes aligned along the eigenvectors of M .

Given a k -simplex s in \mathbb{R}^d and a metric M , we define the M -circumsphere $C_M(s)$ as the circumscribing M -sphere of s with smallest radius. The M -circumball $B_M(s)$ is the M -ball bounded by $C_M(s)$ and the M -circumradius $r_M(s)$ of a simplex s is the radius of $C_M(s)$. Equation (1) shows that $C_M(s)$ is the reciprocal image $F_M^{-1}(C(F_M(s)))$ of the Euclidean circumscribing sphere of the simplex $F_M(s)$.

Let M be a metric and V be a set of points, called *sites*. The Delaunay triangulation of V for metric M , denoted $\text{Del}_M(V)$, is the triangulation of V such that the interior of the M -circumball of each d -simplex is *empty*, i.e. contains no site of V . Owing to Equation (1), the Delaunay triangulation $\text{Del}_M(V)$ of a finite set of points V for metric M is simply obtained by computing the Euclidean Delaunay triangulation of the stretched image $F(V) = \{F_M v, v \in V\}$, and stretching the result back with F_M^{-1} . The Delaunay triangulation $\text{Del}_M(V)$ is thus viewed as the dual of a stretched Voronoi diagram.

2.2 Metric Field and Distortion

In the rest of the paper, we consider a compact domain $D \subset \mathbb{R}^d$ and assume that we are given a metric field defined over D , i.e. a metric M_x is given at each point $x \in D$.

In the following, to avoid double subscripts, we replace subscript M_x by x and simply write Y_x for Y_{M_x} . Hence, we will write for instance F_x for F_{M_x} and $d_x(a, b)$ for $d_{M_x}(a, b)$.

We recall some definitions due to Labelle and Shewchuk [25].

Given two metrics M and N , with stretching transforms F_M and F_N respectively, the *distortion* $\gamma(M, N)$ between M and N is defined as

$$\gamma(M, N) = \max\{\|F_M^{-1}F_N\|, \|F_N^{-1}F_M\|\},$$

where $\|\cdot\|$ is the matrix norm operator associated with the Euclidean norm, i.e. for a $d \times d$ square matrix A , $\|A\| = \sup_{x \in \mathbb{R}^d} \frac{\|Ax\|}{\|x\|}$. Note that the distortion $\gamma(M, N)$ does not depend the stretching matrices F_M and F_N choosen for the metrics M and N . In the context of a metric field, the relative *distortion* between two points p and q of the domain D is defined as $\gamma(p, q) = \gamma(M_p, M_q)$. Observe that $\gamma \geq 1$ and is equal to 1 iff $M_p = M_q$.

A fundamental property of $\gamma(p, q)$ is that it bounds the ratio between d_p and d_q :

$$\forall x, y, \frac{1}{\gamma(p, q)} d_q(x, y) \leq d_p(x, y) \leq \gamma(p, q) d_q(x, y). \quad (2)$$

Let $s = p_0 p_1 \dots p_d$ be d -simplex, let M_i be the metric attached the vertex p_i , for $i = 0, \dots, d$ and let $B_i(s)$ be the M_i -circumball of s . The *distortion* $\gamma(B_M)$ of a M -ball B_M is defined as the maximal distortion between any pairs of points of $B_M \cap D$. We define the *distortion* $\gamma(s)$ of a simplex s as the maximum of the distortion of its circumballs:

$$\gamma(s) = \max\{\gamma(B_i(s)), i = 0, \dots, d\}.$$

2.3 Sizing field

In this paper, we will assume that the metric field is smooth over the domain D . The distortion $\gamma(p, q)$ is then a continuous function and the maximum distortion over D , $\Gamma = \sup_{x, y \in D} \gamma(x, y)$, is finite since D is compact.

We now consider a local view of the distortion. Given a constant $\gamma_0 > 1$, called the *distortion bound*, we define for each point $p \in D$ the *bounded distortion radius*, $\text{bdr}(p, \gamma_0)$, as the upper bound on distances ℓ such that, for all q and r in D , $\max(d_p(p, q), d_p(p, r)) \leq \ell \Rightarrow \gamma(q, r) \leq \gamma_0$.

Lemma 2.1 (Bounded distortion radius lemma) *The bounded distortion radius $\text{bdr}(p, \gamma_0)$ enjoys the following property for any p, q in D :*

$$\frac{1}{\gamma(p, q)} [\text{bdr}(p, \gamma_0) - d_p(p, q)] \leq \text{bdr}(q, \gamma_0) \leq \gamma(p, q) [\text{bdr}(p, \gamma_0) + d_p(p, q)].$$

Proof Let x, y be any two points in D so that:

$$d_q(q, x) \leq \frac{1}{\gamma(p, q)} (\text{bdr}(p, \gamma_0) - d_p(p, q)), \quad (3)$$

$$d_q(q, y) \leq \frac{1}{\gamma(p, q)} (\text{bdr}(p, \gamma_0) - d_p(p, q)). \quad (4)$$

Then, we have, using the triangular inequality,

$$d_p(p, x) \leq d_p(p, q) + d_p(q, x) \leq d_p(p, q) + \gamma(p, q)d_q(q, x) \leq \text{bdr}(p, \gamma_0)$$

and, similarly,

$$d_p(p, y) \leq \text{bdr}(p, \gamma_0).$$

Then, by definition of the bounded distortion radius, $\gamma(x, y) \leq \gamma_0$. Because the last inequality is true for any pair of points x, y satisfying inequalities (3) and (4), we conclude that

$$\frac{1}{\gamma(p, q)} [\text{bdr}(p, \gamma_0) - d_p(p, q)] \leq \text{bdr}(q, \gamma_0). \quad (5)$$

To prove the second inequality of Lemma 2.1, we simply write inequality (5) for the pair (q, p) , which yields:

$$\frac{1}{\gamma(p, q)} [\text{bdr}(q, \gamma_0) - d_q(p, q)] \leq \text{bdr}(p, \gamma_0)$$

from which we deduce

$$\begin{aligned} \text{bdr}(q, \gamma_0) &\leq \gamma(p, q) \text{bdr}(p, \gamma_0) + d_q(p, q) \\ &\leq \gamma(p, q) [\text{bdr}(p, \gamma_0) + d_p(p, q)]. \end{aligned}$$

□

We will further assume that the bounded distortion radius has a strictly positive lower bound on domain D : $\text{bdr}_0 = \min_{p \in D} \text{bdr}(p, \gamma_0) > 0$.

Our algorithm uses a sizing field to control the size of mesh elements according to the local metric. The most basic sizing field we use is the bounded distortion radius $\text{bdr}(p, \gamma_0)$ which will enforce the mesh density to adapt to the metric distortion. However, our algorithm can take into account additional user defined sizing criteria.

Definition 2.2 (Sizing field) Let $\gamma_0 \geq 1$ be a given distortion bound. We call sizing field and denote by $\text{sf}(p, \gamma_0)$ (or $\text{sf}(p)$ for short if γ_0 is understood), any function defined over the domain D , that satisfies the three following conditions:

$$\text{positiveness} \quad \exists \text{sf}_0 > 0 \text{ such that } \forall x \in D, \text{sf}(x, \gamma_0) \geq \text{sf}_0 \quad (6)$$

$$\text{distortion} \quad \forall x \in D, \text{sf}(x, \gamma_0) \leq \text{bdr}(x, \gamma_0) \quad (7)$$

$$\text{continuity} \quad \forall x, y \in D, \quad (8)$$

$$\frac{1}{\gamma(x, y)} [\text{sf}(x, \gamma_0) - d_x(x, y)] \leq \text{sf}(y, \gamma_0) \leq \gamma(x, y) [\text{sf}(x, \gamma_0) + d_x(x, y)]$$

3 Stars and Refinement

We now define the local structures that are built and refined by our algorithm. These definitions rely on the notion of restricted Delaunay widely used in reconstruction area, see e.g [17, 23, 7, 19].

Let D be as before a domain of \mathbb{R}^d and let V be a finite set of points of D that are called hereafter *sites* or *vertices*.

The *restriction* to D of the Delaunay triangulation $\text{Del}(V)$ of V is the subcomplex of $\text{Del}(V)$ whose maximal faces are the d -simplices of $\text{Del}(V)$ that have their dual Voronoi vertices inside the domain D . The natural extension of this definition in the anisotropic setting is to define the restriction of $\text{Del}_M(V)$ to D as the subcomplex of $\text{Del}_M(V)$ whose maximal faces are the d -simplices s of $\text{Del}_M(V)$ that have their M -circumcenter inside D .

3.1 Stars and Inconsistencies

For each site v in V , we consider the Delaunay triangulation $\text{Del}_v(V)$ of V for the metric M_v . We define the restricted *star* S_v of site v as the set of d -simplices incident to v in the restriction of $\text{Del}_v(V)$ to D .

The collection of all restricted stars $S(V) = \{S_v, v \in V\}$, is called the *restricted star set* of V .

We say that a simplex is *inconsistent* if it appears in the star of at least one of its vertices but does not appear in the stars of all of them. For instance, in Figure 1, edge vw or facet xvw are inconsistent because they appear in the triangulation $\text{Del}_v(V)$ computed with metric M_v but not in the Delaunay triangulations $\text{Del}_w(V)$ computed with metrics M_w .




Figure 1: Example of inconsistent stars in 2D: stars S_v and S_w are inconsistent because edge $[vw]$ belongs to S_v but not to S_w .

Our algorithm incrementally inserts new sites in V and updates the restricted star set $S(V)$ until it contains no more inconsistent simplices. As shown below, when the mesh is dense enough with respect to the variation of the metric field, inconsistencies are related to the occurrence of special configurations of subsets of sites that are called *quasi-cospherical configurations* or *QC-configurations* for short. The algorithm will therefore aim at avoiding those QC-configurations. As it turns out, QC-configurations can be avoided but only when even more special configurations called *slivers* do not occur. Both notions are now defined.

3.2 Slivers

In the following we use the definition of slivers provided by Cheng et al [14] and we extend it in the anisotropic setting.

Let s be a k -simplex. We denote by $\mathcal{C}_M(s)$ the M -circumsphere of s , by $r_M(s)$ the M -circumradius of s , by $e_M(s)$ the M -length of the shortest edge of s for the metric M and by $\text{Vol}_M(s)$ the M -volume of s . We define two quality measures of s for metric M . The M -radius-edge ratio is defined as the ratio $\rho_M(s) = r_M(s)/e_M(s)$. The sliverity ratio $\sigma_M(s)$ is the ratio $(\text{Vol}_M(s)/e_M^k(s))^{\frac{1}{k}}$.

Definition 3.1 (Sliver) Let ρ_0 and σ_0 be two positive constants and let M be a metric. A k -simplex s is said to be

- well-shaped for M , if $\rho_M(s) \leq \rho_0$ and $\sigma_M(s) \geq \sigma_0$
- a sliver for M , if $\rho_M(s) \leq \rho_0$, $\sigma_M(s) < \sigma_0$
- a k -sliver for M , if it is a sliver and all its $(k - 1)$ -dimensional faces are well-shaped.

It is easily shown that any k -dimensional simplex that is a sliver is either a k -sliver or include as a subface a k' -sliver for some $k' < k$.

The following lemma is known for slivers in dimension 3, see e.g. [22]. It has been extended to higher dimensions [27] and extends naturally to anisotropic metrics as proved in the appendix.

Lemma 3.2 (Sliver lemma) Let s be a k -simplex and M a metric. If v is a vertex of s , we denote by $s(v)$ the $(k - 1)$ -face of s opposite to vertex v , by $\text{aff}(s(v))$ the affine hull of $s(v)$, i.e. the $(k - 1)$ -flat spanned by the vertices of $s(v)$, by $\mathcal{C}_M(s(v))$ the M -circumsphere of $s(v)$, and by $r_M(s(v))$ the M -radius of $\mathcal{C}_M(s(v))$.

If s is a k -sliver with respect to M , the M -distance from v to $\text{aff}(s(v))$ is at most $2k\sigma_0 r_M(s(v))$ and the M -distance from v to $\mathcal{C}_M(s(v)) \cap \text{aff}(s(v))$ is at most $4\pi k\rho_0\sigma_0 r_M(s(v))$.

3.3 Quasi-Cosphericity

Let $\gamma_0 > 1$ be a bound on the distortion and M be a metric. We now introduce the notion of (γ_0, M) -cosphericity and show its link with inconsistent simplices.

Definition 3.3 (QC-configuration) A subset U of $d + 2$ sites $\{p_0, p_1, \dots, p_{d+1}\}$ is said to be a (γ_0, M) -cospherical configuration if there exist two metrics N and N' such that :

- $\gamma(M, N) \leq \gamma_0$, $\gamma(M, N') \leq \gamma_0$ and $\gamma(N, N') \leq \gamma_0$;
- the triangulations $\text{Del}_N(U)$ and $\text{Del}_{N'}(U)$ are different.

Metrics N and N' are said to witness the (γ_0, M) -cosphericity of U . If M is clear from the context, we simply say that U is a γ_0 -cospherical configuration and if both M and γ_0 are understood, we say that U is a quasi-cospherical configuration or a QC-configuration for short.

See Figure 2 for an illustration in the plane. Note that the $d + 2$ points in U play symmetric roles in the above definition. In the sequel, U will often consist of the set of vertices of a d -simplex s belonging to the star S_v of some site $v \in V$, together with an additional site p of V . In such a case, we write $U = (s, p)$.

From Radon theorem, there are only two distinct triangulations of $U = (s, p)$ and any d -simplex with vertices in U belongs to exactly one of them [26, 37]. Therefore, we have the following easy lemma.

Lemma 3.4 *A configuration (s, p) is (γ_0, M) -cospherical iff there exist two metrics N and N' such that*

- $\gamma(M, N) \leq \gamma_0$, $\gamma(M, N') \leq \gamma_0$ and $\gamma(N, N') \leq \gamma_0$;
- p belongs to the interior of exactly one of the two circumballs $B_N(s)$ and $B_{N'}(s)$.




Figure 2: $(s = abc, d)$ is a QC-configuration because d is inside $B_N(s)$ but outside $B_{N'}(s)$

The following lemma relates QC-configurations and inconsistencies.

Lemma 3.5 *Let s be an inconsistent simplex of star S_v . If $\gamma(s) < \gamma_0$, then there exists a site $q \in V$ such that the configuration (s, q) is (γ_0, M_v) -cospherical.*

Proof By definition s appears in the Delaunay triangulation $\text{Del}_v(V)$ computed with the metric of vertex v , but not in the triangulation $\text{Del}_w(V)$ computed with the metric of some other vertex w of s . Take $N = M_v$ and $N' = M_w$. Because the distortion of s is less than γ_0 , we have $\gamma(v, w) = \gamma(M_v, M_w) \leq \gamma_0$. Since s is a d -simplex in S_v but not in S_w , it belongs to $\text{Del}_v(V)$ and not to $\text{Del}_w(V)$. Hence, there is a site $q \in V$ such that q is inside $B_w(s)$ and not inside $B_v(s)$. It then follows from Lemma 3.4 that (s, q) is a (γ_0, M_v) -cospherical configuration witnessed by the metrics $N = M_v$ and $N' = M_w$. \square

Given a metric M and a (γ_0, M) -cospherical configuration U , the M -radius $r_M(U)$ of U is defined as the minimum of the M -circumradii of all the d -simplices with vertices in U .

Definition 3.6 (Well-shaped QC-configuration) *A (γ_0, M) -cospherical configuration U is said to be well-shaped if any d -simplex formed with vertices in U is well-shaped with respect to M .*

4 Meshing Algorithm

4.1 Algorithm Outline

To mesh a given compact domain D , the algorithm constructs the set of sites from a small set of initial sites by inserting new sites in a greedy way. The algorithm maintains the star set for the current set of sites and the addition of new sites is steered by this star set : while there remain

bad simplices in the star set, the algorithm selects one bad simplex and kills this simplex by inserting a new site. Bad simplices are d -simplices that have a high distortion or are oversized with respect to a user defined sizing field, those that are badly shaped (high radius-edge ratio or small sliverity ratio), and those that are inconsistent. To kill a bad simplex s appearing in a star S_v , a new site p , called the *refinement point*, is chosen in the M_v -circumscribing ball of s and inserted in the star set. The maintenance of the star set upon the insertion of the refinement point p involves the creation of a new star S_p for p and the insertion of p in the star S_v and in any star S_w where p will appear as a vertex. Note that a new site p has to be inserted in star S_w iff p is in conflict with some d -simplex of S_w , where point p is said to be in conflict with the d -simplex s of S_w if p is included in the M_w -circumball of s . Upon each insertion, the algorithm maintains the star set by calling the following `Insert` procedure:

Algorithm 1 `Insert(p)`

1. insert p in all the stars S_w that contain a simplex in conflict with p ;
 2. create the new star S_p .
-

As noticed in Section 3.3, once the set of vertices is dense enough with respect to the variation of the metric field so that all simplices in the star set have a distortion smaller than γ_0 , inconsistencies arise only from QC-configurations. The refinement algorithm therefore aims at avoiding those configurations. However, as will be clear from the proof of Lemma 5.3, it is not possible to avoid QC-configurations involving slivers. The algorithm thus needs to remove slivers before removing inconsistent simplices.

Recall that for a d -simplex s in some star S_v , we write $B_v(s)$ or $B_v(c_v(s), r_v(s))$ for the M_v -circumball of s with center $c_v(s)$ and radius $r_v(s)$, $\rho_v(s)$ for the M_v -radius-edge ratio of s and $\sigma_v(s)$ for its M_v -sliverity ratio. The refinement algorithm (see Algorithm 2) applies four rules in turn. The rules are applied with a priority order : rule (i) is applied only if no rule (j) with $j < i$ can be applied. The algorithm ends when no rule applies any more. The algorithm relies on two procedures: procedure `Insert` inserts a new site in the data structures, and procedure `Pick_valid` chooses the location of the new site (see the next section).

The refinement algorithm depends on parameters α_0 , ρ_0 , σ_0 , and γ_0 while the `Pick_valid` procedure depends on two more parameters β and δ . The values of constants α_0 , ρ_0 , σ_0 , γ_0 control the quality of the mesh elements and their adaptation to the metric field. Parameters β and δ influence the behaviour of the algorithm and their values are chosen in Section 5 in order to ensure the termination of the refinement algorithm.

Remark. Note that the sizing field $\text{sf}(p)$ used in Rule (1) takes care of the distortion bound γ_0 and also possibly of a user defined sizing field (see Definition 2.2). Parameter α_0 is always chosen less than 1. Therefore, when Rule (1) does not apply anymore, the distortion of any d -simplex in any star is bounded by γ_0 .

Sections 5 and 6 will prove that the algorithm terminates. Before that, Subsection 4.2 describes the procedure `Pick_valid` while Subsection 4.3 analyses the properties of the resulting mesh.

4.2 Picking Region and Hitting Sets

In this section, we describe in more detail procedure `Pick_valid`. The simplest idea to kill a simplex would be to insert a refinement point at its circumcenter. However, with this simple strategy, the algorithm may loop, creating cascading configurations of slivers and QC-configurations, and

Algorithm 2 Refinement algorithm

Rule (1) Size:

If \exists a d -simplex s in star S_v such that $r_v(s) \geq \alpha_0 \text{sf}(c_v(s))$,
 Insert($c_v(s)$);

Rule (2) Radius-edge ratio:

If \exists a d -simplex s in star S_v such that $\rho_v(s) > \rho_0$,
 Insert(Pick_valid(s, M_v));

Rule (3) Sliver removal:

If a d -simplex s in star S_v is a M_v sliver (i.e. $\rho_v(s) \leq \rho_0$ and $\sigma_v(s) < \sigma_0$),
 Insert(Pick_valid(s, M_v));

Rule (4) Inconsistency:

If a d -simplex s in some star S_v is inconsistent,
 Insert(Pick_valid(s, M_v));

is not guaranteed to terminate. To avoid slivers and QC-configurations, the algorithm resorts to a strategy analog to the one used by Li and Teng [28, 27] to avoid slivers in isotropic meshes. The basic idea is to relax the choice of the refinement point of a bad simplex. Instead of using systematically the circumcenter, the refinement point of a bad simplex is picked from a small region around the circumcenter, called the *picking region*. The refinement point is carefully chosen in the picking region so as to avoid the formation of new slivers and new QC-configurations.

Definition 4.1 (Picking region) Let $\delta < 1$ be a constant called the picking ratio. If s is a bad simplex in star S_v , with M_v -circumball $\mathcal{B}_v(c_v(s), r_v(s))$, the M_v -picking region of s , noted $P_v(s)$, is the M_v -ball $\mathcal{B}_v(c_v(s), \delta r_v(s))$.

In fact, it is not possible, when choosing a refinement point in the picking region $P_v(s)$ of a simplex s of S_v to completely avoid the formation of new slivers and new QC-configurations. The Pick_valid procedure will only avoid the creation of *small* slivers and *small* QC-configurations where the meaning of *small*, precisely defined below, is relative to the radius $r_v(s)$ and controlled by a parameter β .

Definition 4.2 (Hitting set) Let p be a point in the M_v -picking region of a simplex s . Let $r_v(s)$ be the M_v -circumradius of s and β be a constant. A subset t of the current set of sites V is said to hit p if one of the two following conditions is satisfied:

- t consists of $k \leq d$ sites and, for some metric M such that $\gamma(M_p, M) \leq \gamma_0$, the k -simplex $s' = (t, p)$ is a k -sliver with M -circumradius $r_M(s') \leq \beta r_v(s)$.
- t consists of $d + 1$ sites and, for some metric M such that $\gamma(M_p, M) \leq \gamma_0$, $U = (t, p)$ is a well-shaped (γ_0, M) -cospherical configuration with M -radius $r_M(U) \leq \beta r_v(s)$.

A point p in $P_v(s)$ is said to be a *valid refinement point* if it is not hit by any subset of V . Each subset t of sites in V induces a *forbidden region* where the refinement point should not lie in order to avoid being hit by t . A subset t of sites in V that hits some point in the picking region $P_v(s)$ is said to be a *hitting set* for $P_v(s)$. A point p in $P_v(s)$ is therefore valid if it avoids the forbidden region of any hitting set of $P_v(s)$.




Figure 3: $\{q, r, u\}$ is a hitting set for the picking region $P_v(s)$. It defines a forbidden region (dashed area) to be avoided by the refinement point p of simplex s .

Note that the definition of valid refinement points depends on the constants δ and β : δ defines the size of the picking regions and β bounds from below the size of acceptable new slivers and new QC-configurations, with respect to the circumradius of the simplex being refined. The definition of valid refinement points also depends on the constants ρ_0 and σ_0 that define well-shaped simplices and slivers, and on the constant γ_0 that defines QC-configurations. We prove in the following sections (Section 5 and Section 6) that it is possible to choose the algorithm parameters β , δ , ρ_0 , σ_0 , γ_0 and α_0 so that valid refinement points do exist in any picking region considered by the refinement algorithm.

To find a valid refinement point in the M_v -picking region $P_v(s)$ of some bad d -simplex s , the insertion algorithm calls the following `Pick_valid` procedure. This procedure randomly chooses a point in the picking region $P_v(s)$ until it finds one that avoids all forbidden regions. This procedure depends on constants ρ_0 , σ_0 , γ_0 , δ and β , to be fixed later in Section 5.

Algorithm 3 `Pick_valid`(s, M_v)

Step 1 Pick randomly a point p in the picking region $P_v(s)$

Step 2 *Avoid small slivers*

For $k = 3$ to d ,

if there exists a subset of k sites in V that hits p ,
then discard p and go back to step 1.

Step 3 *Avoid small QC-configurations*

If there exists a subset of $d + 1$ sites in V that hits p ,
then discard p and go back to step 1.

Step 4 Return p .

4.3 Quality of the final mesh

Upon termination of the algorithm, all stars are consistent. Therefore they can be merged together to form a triangulation \mathcal{T} of the domain. Each simplex s in \mathcal{T} is well-shaped with respect to the metric of all its vertices, i.e. $\rho_p(s) \leq \rho_0$ and $\sigma_p(s) \geq \sigma_0$ for any vertex p of s . Moreover each simplex s in \mathcal{T} complies to the sizing field $\text{sf}(\gamma_0)$ which implies that s has a distortion smaller than γ_0 . If needed, the sizing field $\text{sf}(\gamma_0)$ may also take into account a user defined sizing field.

4.4 Complexity of the meshing algorithm

The complexity of algorithm 2 is roughly linear with respect to the number of vertices of the output mesh. This might appear surprising since the algorithm maintains a distinct 3D triangulation for each vertex in the mesh. In fact, these triangulations are quite small, since they are designed to maintain the star of a single central vertex. In each star, the generated sites form a well spaced set of points with respect to the local metric, and therefore the number of simplices in the star is bounded by a constant depending only on the dimension d . Hence the star set is a data structure whose size is linear with respect to the number of vertices of the output mesh.

Each new vertex is inserted in a constant number of stars and its own stars is initialized by the insertion of a constant number of the current vertices. The insertion of a point in a star takes a constant time. One of the main concerns of our implementation is to efficiently filter out the stars in which a new vertex has to be inserted, and the subset of current vertices that have to be inserted in the star of the new vertex. These problems are handled using additional standard data structures for range queries among bounding boxes.

Finally, the most costly part of the algorithm is the computation of a valid refinement point using the `pick_valid` procedure. This amounts to randomly choose candidates and check if they are valid. The validity check amounts to simulating the insertion of the point and its cost is similar to the cost of an insertion. The proof of the Picking Lemma below, yields that the expected number of performed trials is constant. Therefore, altogether the expected complexity of the algorithm is linear with respect to the number of vertices of the output mesh.

It now remains to prove that the algorithm terminates, which will be done in the two next sections.

5 Termination of the Algorithm

The refinement algorithm (Algorithm 2) depends on parameters $\alpha_0, \gamma_0, \rho_0, \sigma_0$ that respectively control the size, the distortion, the radius-edge ratio and the sliverity ratio of the simplices, and on parameters δ and β that define the picking-region and valid refinement points. In this section and in the following one, we prove that for a suitable choice of those parameters, Algorithm 2 terminates providing as claimed a consistent mesh that is an anisotropic Delaunay mesh.

Let us first notice that our algorithm will never refine a star element that is not part of the restricted Delaunay triangulation of the domain to be meshed. As a consequence, the Steiner vertices inserted by the algorithm are within the domain, or very close to it when they are chosen through a call to the `Pick_valid` procedure. The proof of termination then relies on a volume argument based on a minimum separation distance between any two vertices of the mesh.

For any vertex p in V , we define the *separation distance* and *insertion-radius* as follows.

Definition 5.1 *If p is a vertex in V , the separation distance $\text{sd}(p)$ of p is defined as:*

$$\text{sd}(p) = \min_{q \in V} d_p(p, q)$$

Definition 5.2 If p is a vertex of V and $V(p) \subset V$ the subset of vertices inserted before p , the insertion radius $r(p)$ of p is defined as:

$$r(p) = \min_{q \in V(p)} d_p(p, q).$$

We will mainly show that there is a constant Λ (depending on parameters $\alpha_0, \gamma_0, \rho_0, \sigma_0, \delta$ and β and on properties of the domain and of the metric field) such that the separation bound $\text{sd}(p) \geq \Lambda \text{sf}(p)$ holds for any site in V .

We first need to ensure that valid refinement points will be found in any picking region considered by the algorithm. This is the goal of the next lemma whose proof is deferred to Section 6).

Lemma 5.3 (Picking lemma) For any values of parameters β, δ, α_0 and ρ_0 , it is possible to choose σ_0 small enough and the distortion bound γ_0 close enough to 1, so that, if the separation bound $\text{sd}(p) \geq \Lambda \text{sf}(p)$ holds for any site in the current set V , valid refinement points do exist in the M_v -picking region of any bad simplex s in star S_v .

To prove the separation bound on vertices, we first prove a lower bound on the insertion radii of vertices, considering in turn each of the refinement rules. We begin with a technical lemma relating the circumradius of a simplex with the insertion radius of its refinement point.

Lemma 5.4 (Insertion radius lemma) Let s be a d -simplex of star S_v with the M_v -circumball $B_v(c_v(s), r_v(s))$. Assume that s is a bad simplex. Let p be the refinement point of s and $r(p)$ the insertion radius of p .

- If Rule (1) applies, the refinement point p of s is the circumcenter $c_v(s)$, and

$$r(p) \geq \frac{r_v(s)}{\Gamma}, \quad (9)$$

where Γ is the maximal distortion over D : $\Gamma = \max_{x, y \in D} \gamma(x, y)$.

- If one of Rule (2), (3) or (4) is applied, the refinement point p is taken from the picking region $P_v(s)$ and :

$$r(p) \geq \frac{1 - \delta}{\gamma_0} r_v(s). \quad (10)$$

Proof In the first case, $p = c_v(s)$, and therefore

$$\begin{aligned} \min_{q \in V(p)} d_v(p, q) &\geq r_v(s) \\ r(p) = \min_{q \in V(p)} d_p(p, q) &\geq \frac{r_v(s)}{\Gamma}. \end{aligned}$$

In the second case, p belongs to the picking region $P_v(s)$, and we know that the distortion $\gamma(s)$, hence also the distortion $\gamma(v, p)$, are at most γ_0

$$\begin{aligned} \min_{q \in V(p)} d_v(p, q) &\geq (1 - \delta)r_v(s) \\ r(p) = \min_{q \in V(p)} d_p(p, q) &\geq \frac{1 - \delta}{\gamma_0} r_v(s). \end{aligned}$$

□

Lemma 5.5 (Rule (1) lemma) *When Rule (1) is applied, the insertion radius $r(p)$ of the inserted site p is at least:*

$$r(p) \geq \Lambda_1 \text{sf}(p) \text{ with } \Lambda_1 = \frac{\alpha_0}{\Gamma}. \quad (11)$$

Proof Rule (1) is applied to a simplex s in star S_v when the M_v -circumradius $r_v(s)$ of s is greater than $\alpha_0 \text{sf}(c_v(s))$. The refinement point p is then $c_v(s)$ and we get from Lemma 5.4

$$r(p) \geq \frac{r_v(s)}{\Gamma} \geq \frac{\alpha_0 \text{sf}(c_v(s))}{\Gamma} = \frac{\alpha_0}{\Gamma} \text{sf}(p). \quad (12)$$

□

Lemma 5.5 proves a lower bound on the insertion radius of any vertex p introduced by application of Rule (1). The next lemmas aim at finding a constant Λ_2 , and some conditions on α_0 , ρ_0 , γ_0 , β and δ so that Rules (2)-(4) will maintain the invariant that the insertion radius of any inserted point is at least $\Lambda_2 \text{sf}(p)$.

Lemma 5.6 (Rule (2) lemma) *Let $s \in S_v$ be a simplex to be refined by application of Rule (2) and let p be the refinement point of s . If, for any vertex q inserted before p , $r(q) \geq \Lambda_2 \text{sf}(q)$ then we have $r(p) \geq \Lambda_2 \text{sf}(p)$, provided that the following conditions hold*

$$\frac{(1-\delta)\rho_0}{\gamma_0^3} \geq 2, \quad (13)$$

$$\frac{(1+\delta)\rho_0}{\gamma_0} \Lambda_2 \leq 1. \quad (14)$$

Proof First, observe that $\gamma(s) \leq \gamma_0$ since Rule (1) does not apply. Then, because p is inserted by application of Rule (2), the M_v -circumradius of s , $r_v(s)$, is such that $r_v(s) \geq \rho_0 e_v(s)$, where $e_v(s)$ is the M_v -length of the M_v -shortest edge of s , which is the shortest edge of s according to metric M_v . Let qq' be the M_v -shortest edge of s and q be the last inserted vertex of qq' .

$$\begin{aligned} r_v(s) &\geq \rho_0 e_v(s) = \rho_0 d_v(q, q') \geq \frac{\rho_0}{\gamma_0} d_q(q, q') \\ &\geq \frac{\rho_0}{\gamma_0} r(q) \end{aligned}$$

Using the induction hypothesis and the continuity condition of the sizing field (see Definition 2.2), we have

$$r_v(s) \geq \frac{\rho_0}{\gamma_0} \Lambda_2 \text{sf}(q) \quad (15)$$

$$\begin{aligned} &\geq \frac{\rho_0}{\gamma_0} \Lambda_2 \frac{[\text{sf}(p) - d_p(p, q)]}{\gamma(p, q)} \\ &\geq \frac{\rho_0}{\gamma_0^2} \Lambda_2 [\text{sf}(p) - d_p(p, q)] \\ &\geq \frac{\rho_0}{\gamma_0^2} \Lambda_2 [\text{sf}(p) - \gamma_0 d_v(p, q)] \quad (\text{since } v, p \in B_v(s)). \end{aligned} \quad (16)$$

Now, because q is a vertex of s and p is chosen in the picking region $P_v(s)$, $d_v(p, q) \leq (1+\delta)r_v(s)$ which, together with inequality (16), gives:

$$\begin{aligned} r_v(s) &\geq \frac{\rho_0}{\gamma_0^2} \Lambda_2 [\text{sf}(p) - \gamma_0(1+\delta)r_v(s)] \\ r_v(s) &\geq \frac{\frac{\rho_0}{\gamma_0^2} \Lambda_2 \text{sf}(p)}{1 + \frac{\rho_0}{\gamma_0}(1+\delta) \Lambda_2}. \end{aligned} \quad (17)$$

Then, using the insertion radius lemma (Lemma 5.4), we get:

$$r(p) \geq \frac{\frac{(1-\delta)\rho_0}{\gamma_0^3} \Lambda_2 \text{sf}(p)}{1 + \frac{\rho_0}{\gamma_0}(1+\delta) \Lambda_2}, \quad (18)$$

which proves that $r(p) \geq \Lambda_2 \text{sf}(p)$ when conditions (13) and (14) are fulfilled. \square

Lemma 5.7 (Rule (3)-(4) lemma) *Let $s \in S_v$ be a simplex to be refined by application of Rule (3) or Rule (4), and let p be the refinement point of s . If, for any vertex q inserted before p , $r(q) \geq \Lambda_2 \text{sf}(q)$ then we have $r(p) \geq \Lambda_2 \text{sf}(p)$, provided that the following conditions hold*

$$\frac{\beta(1-\delta)}{\gamma_0^3(1+\delta)} \geq 2 \quad (19)$$

$$\Lambda_2 \leq \frac{(1-\delta)\Lambda_1}{2\gamma_0^3 + \gamma_0^2(1+\delta)\Lambda_1} \quad (20)$$

$$\frac{\beta\Lambda_2}{\gamma_0} \leq 1. \quad (21)$$

Proof Assume first that s was created by application of Rule (1). Then, if q is the last inserted vertex of s , we have $r(q) \geq \Lambda_1 \text{sf}(q)$ by Lemma 5.5. Furthermore, $r_v(s)$ is at least half the M_v -length of any edge of s and, in particular, of any edge of s that is incident to q . Therefore, using the fact that v and q belong to s which has a low distorsion, we get

$$r_v(s) \geq \min_{q' \in s} \frac{d_v(q, q')}{2} \geq \min_{q' \in V} \frac{d_q(q, q')}{2\gamma_0} \geq \frac{r(q)}{2\gamma_0} \geq \frac{\Lambda_1}{2\gamma_0} \text{sf}(q)$$

Now, repeating the calculations performed to deduce inequality (17) from (15), we obtain

$$\begin{aligned} r_v(s) &\geq \frac{\Lambda_1}{2\gamma_0^2} [\text{sf}(p) - \gamma_0(1+\delta)r_v(s)] \\ r_v(s) &\geq \frac{\frac{\Lambda_1}{2\gamma_0^2} \text{sf}(p)}{1 + \frac{\Lambda_1(1+\delta)}{2\gamma_0}} = \frac{\Lambda_1 \text{sf}(p)}{2\gamma_0^2 + \gamma_0(1+\delta)\Lambda_1}. \end{aligned} \quad (22)$$

Then, using the insertion radius lemma (Lemma 5.4), we get:

$$r(p) \geq \frac{(1-\delta)\Lambda_1 \text{sf}(p)}{2\gamma_0^3 + \gamma_0^2(1+\delta)\Lambda_1}. \quad (23)$$

It follows that the bound $r(p) \geq \Lambda_2 \text{sf}(p)$ holds, provided condition (20) is satisfied.

Now consider the case where s was created by application of Rule (2), (3) or (4). Assume that s has been created when inserting the refinement point q of a simplex s' in some star S_w (see Figure 4). The refinement point q was chosen by the procedure `Pick_valid`(s', M_w) and therefore, $r_v(s) \geq \beta r_w(s')$. Let us bound $r_w(s')$ from below. Vertex q is the last inserted vertex of s . It has been chosen in the picking region of s' and therefore the vertices of s' are at M_w -distance at most $(1+\delta)r_w(s')$ from q . Hence, since q and w belong to s' and $\gamma(s') \leq \gamma_0$, $r(q) \leq \gamma_0(1+\delta)r_w(s')$.

Therefore:

$$\begin{aligned} r_v(s) &\geq \beta r_w(s') \geq \frac{\beta}{\gamma_0(1+\delta)} r(q). \\ r_v(s) &\geq \frac{\beta\Lambda_2}{\gamma_0(1+\delta)} \text{sf}(q). \end{aligned}$$




Figure 4: For the proof of Lemma 5.7.

Repeating the calculations performed to deduce inequality (17) from (15), we obtain

$$r_v(s) \geq \frac{\beta\Lambda_2}{\gamma_0^2(1+\delta)} [\text{sf}(p) - \gamma_0(1+\delta)r_v(s)]$$

Hence,

$$r_v(s) \geq \frac{\frac{\beta\Lambda_2}{\gamma_0^2(1+\delta)} \text{sf}(p)}{1 + \frac{\beta\Lambda_2}{\gamma_0}}, \quad (24)$$

and, from the insertion radius lemma (Lemma 5.4),

$$r(p) \geq \frac{\frac{\beta(1-\delta)\Lambda_2}{\gamma_0^3(1+\delta)}}{1 + \frac{\beta\Lambda_2}{\gamma_0}} \text{sf}(p). \quad (25)$$

Conditions (21) and (19) ensure that $r(p) \geq \Lambda_2 \text{sf}(p)$. \square

Lemma 5.8 (Separation bound) *Assume that each vertex p has been inserted in the set V with an insertion radius $r(p)$ such that $r(p) \geq \Lambda_2 \text{sf}(p)$ where Λ_2 is a constant. Then the set V admits the following separation bound :*

$$\text{sd}(p) \geq \Lambda \text{sf}(p)$$

with

$$\Lambda = \frac{\Lambda_2/\Gamma^2}{1 + \Lambda_2/\Gamma^2}.$$

Proof Observe that for any pair of vertices $p, q \in V$, we have

$$\text{either } d_p(p, q) \geq r(p) \text{ or } d_q(p, q) \geq r(q),$$

where the first is true if p has been inserted after q and the second is true otherwise. In the first case, we have

$$d_p(p, q) \geq r(p) \geq \Lambda_2 \text{sf}(p) \geq \Lambda \text{sf}(p)$$

In the second case, we have

$$\begin{aligned} d_p(p, q) &\geq \frac{d_q(p, q)}{\Gamma} \geq \frac{\Lambda_2 \text{sf}(q)}{\Gamma} \\ &\geq \frac{\Lambda_2}{\Gamma^2} (\text{sf}(p) - d_p(p, q)) \quad (\text{using (8)}) \\ d_p(p, q) &\geq \frac{\Lambda_2/\Gamma^2}{1 + \Lambda_2/\Gamma^2} \text{sf}(p) = \Lambda \text{sf}(p) \end{aligned}$$

which proves the separation bound. \square

We can now give the main theorem of this section that proves the separation bound on the set of vertices and ensures that the refinement algorithm terminates.

Theorem 5.9 *Given a compact domain D and a sizing field over D satisfying conditions (6)-(8), assume that the constants ρ_0, γ_0, β and δ are chosen in such a way that:*

$$\frac{(1 - \delta)\rho_0}{\gamma_0^3} \geq 2 \tag{26}$$

$$\frac{(1 - \delta)\beta}{\gamma_0^3(1 + \delta)} \geq 2. \tag{27}$$

Assume furthermore that σ_0 is small enough and γ_0 close enough to 1 so that the picking lemma (Lemma 5.3) holds. Then the refinement algorithm (Algorithm 2) terminates.

Proof Observe that the inequalities 26 and 27 are just conditions (13) and (19) of Lemma 5.6 and 5.7. Assume that these inequalities hold.

We choose the value of Λ_2 small enough so that $\Lambda_2 \leq \Lambda_1 = \frac{\alpha_0}{\Gamma}$ and condition (14) of Lemma 5.6, and conditions (20) and (21) of Lemma 5.7 hold. For further reference, let us notice here that it suffices to choose

$$\Lambda_2 = \min \left\{ \frac{1}{(1 + \delta)\rho_0}, \frac{(1 - \delta)\alpha_0}{16\Gamma + 4(1 + \delta)\alpha_0}, \frac{1}{\beta} \right\} \tag{28}$$

to get a value of Λ_2 , independant of σ_0 and γ_0 , and suitable for any value of σ_0 and any $\gamma_0 \in [1, 2]$.

We first prove by induction that any vertex p is inserted in V with an insertion radius $r(p) \geq \Lambda_2 \text{sf}(p)$. First notice that the induction hypothesis can be enforced on the set of initial points. Then assume that the hypothesis is true up to a given stage. From the separation bound lemma (Lemma 5.8), any vertex in the current set has a separation bound $\text{sd}(p) \geq \Lambda \text{sf}(p)$. From the picking lemma (Lemma 5.3), we know that the algorithm will find a valid refinement point if the next vertex is to be searched in a picking region. Now, from Lemmas 5.5, 5.6, and 5.7, we know that the insertion radius of the next vertex p is still going to be bounded by $r(p) \geq \Lambda_2 \text{sf}(p)$ which achieves the inductive proof.

Finally, we can set up the volume argument for the algorithm termination in the metric M_y of any point $y \in D$. Indeed, for any pair p, q of vertices in V , we have either

$$d_p(p, q) \geq r(p) \geq \Lambda_2 \text{sf}(p) \geq \Lambda_2 \text{sf}_0$$

or

$$d_q(p, q) \geq r(q) \geq \Lambda_2 \text{sf}(q) \geq \Lambda_2 \text{sf}_0.$$

In both cases, we may conclude that

$$d_y(p, q) \geq \frac{\Lambda_2}{\Gamma} \text{sf}_0.$$

Since D is a compact domain and has therefore a bounded M_y -volume, this proves that the algorithm can only insert a finite number of vertices and therefore terminates. \square

6 Proof of Lemma 5.3 (Picking lemma)

To complete the proof of termination of the algorithm, it remains to prove the picking lemma (Lemma 5.3), which is done in this section.

Let us recall briefly the context. Assume that the algorithm needs to refine a simplex s in star S_v , with circumball $B_v(c_v(s), r_v(s))$. The picking lemma states that a valid refinement point can always be found provided that the bound on the sliverity ratio σ_0 is small enough and that the bound on the distortion γ_0 is sufficiently close to 1. The refinement point is searched in the picking region $P_v(s)$, a M_v -ball with radius $\delta r_v(s)$ centered at the circumcenter $c_v(s)$. The refinement point is valid when it does not belong to the so-called forbidden regions. Each forbidden region is associated to a hitting set and consists of the points in the picking region that form with the hitting set either a *small* sliver or a *small* well-shaped QC-configuration. *Small* is here relative to the circumradius $r_v(s)$ and controlled by parameter β (see Definition 4.2).

The proof shows that the union of the forbidden regions does not cover the picking region. In a first step, we show that the volume of each forbidden region is bounded and in fact can be made as small as required with a good choice of the parameters σ_0 and γ_0 (Lemmas 6.5 and 6.7). In a second step, we show that the number of hitting sets, or equivalently of forbidden regions to be avoided, is bounded (Lemma 6.8).

We begin with two technical lemmas. The first one bounds the difference between the two circumspheres of a well-shaped simplex (see Definition 3.1) with respect to two metrics with a bounded distortion.

Lemma 6.1 (Circumsphere lemma) *Let M_v and M_w be two metrics with a distortion $\gamma(M_v, M_w) \leq \gamma_0$ for some $\gamma_0 > 1$. Let s be a k -simplex that is well shaped with respect to metric M_v . We write c_v and r_v for the M_v -circumcenter and M_v -circumradius of s respectively, and we write c_w and r_w for the M_w -circumcenter and M_w -circumradius of s , respectively.*

- The M_v -distance $d_v(c_v, c_w)$ between the circumcenters satisfies

$$d_v(c_v, c_w) \leq f_k(\rho_0, \sigma_0, \gamma_0) r_v \quad (29)$$

where

$$f_k(\rho_0, \sigma_0, \gamma_0) = \left[1 + \frac{2^k}{k} \frac{\gamma_0^2 \rho_0^k}{\sigma_0^k} \right] (\gamma_0^2 - 1).$$

- The circumradius r_w is bounded as follows

$$r_w \in [h_k^-(\rho_0, \sigma_0, \gamma_0) r_v, h_k^+(\rho_0, \sigma_0, \gamma_0) r_v]$$

where

$$\begin{aligned} h_k^-(\rho_0, \sigma_0, \gamma_0) &= \frac{1}{\gamma_0} (1 - f_k(\rho_0, \sigma_0, \gamma_0)), \\ h_k^+(\rho_0, \sigma_0, \gamma_0) &= \gamma_0 (1 + f_k(\rho_0, \sigma_0, \gamma_0)). \end{aligned}$$

Proof The proof is given in the appendix. □

Observe that $f_k(\rho_0, \sigma_0, \gamma_0)$ tends to zero when σ_0 tends to 0 and γ_0 tends to 1 in such a way that $(\gamma_0 - 1)/\sigma_0^k$ tends to 0. We give a name to such functions for further reference.

Definition 6.2 (k -regularly vanishing function) *In the following, a function f of σ_0 and γ_0 is said to be k -regularly vanishing if f tends to zero when σ_0 tends to 0 and γ_0 tends to 1 in such a way that $(\gamma_0 - 1)/\sigma_0^k$ tends to 0.*

The second lemma considers a well-shaped (γ_0, M) -cospherical configuration U and shows that all the d simplices with vertices in U have nearly the same M -circumradii.

Lemma 6.3 (Circumradii in QC-configurations) *Let U be a well shaped (γ_0, M) -cospherical configuration. The M -circumradii of the d -simplices whose vertices belong to U satisfy:*

$$\max_{s \subset U} r_M(s) \leq [1 + \eta(\gamma_0, \rho_0, \sigma_0)] \min_{s \subset U} r_M(s),$$

where $\eta(\gamma_0, \rho_0, \sigma_0)$ is a d -regularly vanishing function.

Note that $\min_{s \subset U} r_M(s)$ has been defined in Subsection 3.3 as the M -radius $r_M(U)$ of the configuration U .

Proof The proof is given in the appendix. □

In the following lemmas, we bound the volume of forbidden regions induced by slivers and QC-configurations by enclosing those regions within spherical shells.

Definition 6.4 (Spherical shells) *The M_v -spherical shell $\mathcal{S}_v(c, r^+, r^-)$ with center c and radii $r^+ > r^-$ is the difference between the two M_v -balls $B_v(c, r^+)$ and $B_v(c, r^-)$.*

The M_v -volume of the spherical shell $\mathcal{S}_v(c, r^+, r^-)$ is upper bounded by:

$$\text{Vol}_v(\mathcal{S}_v(c, r^+, r^-)) \leq \phi_d (r^+)^{d-1} (r^+ - r^-), \tag{30}$$

where ϕ_d is the measure of the unit $(d - 1)$ -sphere and therefore depends only on the dimension d .

Avoiding slivers

Let s be a k -simplex of a star S_v . We again write r_v for its M_v -circumradius. Consider a refinement point p to be taken from the M_v -picking region $P_v(s)$ of s . Point p is required to lie outside all forbidden regions. We first consider the case of the forbidden region $Y_v(t)$ associated to a hitting set t formed by $k \leq d$ sites such that the k -simplex $s' = (t, p)$ is a small k -sliver with respect to a metric M close to M_p . More precisely (see Definition 4.2), by a metric M close to M_p , we mean a metric M such that $\gamma(M, M_p) \leq \gamma_0$, and by a small k -sliver we mean a k -sliver whose M -circumradius is smaller than βr_v . Here, as in the rest of the paper, we use the same notation for a subset of sites and the simplex formed by the convex hull of the subset. Note that t is not required to be a simplex appearing in some current star.

Lemma 6.5 (Forbidden regions due to slivers) *The M_v -volume of the region $Y_v(t)$ forbidden by the hitting set t is bounded from above as follows*

$$\text{Vol}_v(Y_v(t)) \leq \mu_k(\rho_0, \sigma_0, \gamma_0) \beta^d r_v^d,$$

where $\mu_k(\rho_0, \sigma_0, \gamma_0)$ is a k -regularly vanishing function.

Proof By the definition of a hitting set, there is a metric M satisfying $\gamma(M_p, M) \leq \gamma_0$ such that $s' = (t, p)$ is a small k -sliver with respect to M . Now, since p belongs to the M_v -picking region $P_v(s)$ of s , we have $\gamma(M_v, M_p) \leq \gamma_0$. It follows that $\gamma(M_v, M) \leq \gamma_0^2$.

Let $\mathcal{C}(c', r')$ and $\mathcal{C}_v(c'_v, r'_v)$ denotes respectively the M -circumscribing sphere of t and the M_v -circumsphere of t . Since $s' = (t, p)$ is a small k -sliver with respect to M , t is a well-shaped $(k - 1)$ -simplex and its M -circumradius r' smaller than βr_v . From the sliver lemma (Lemma 3.2), we know that p is at M -distance at most $4\pi k \rho_0 \sigma_0 r'$ from $\mathcal{C}(c', r') \cap \text{aff}(t)$, where $\text{aff}(t)$ is the affine

hull of t . Applying the circumsphere lemma (Lemma 6.1) to the well-shaped $(k-1)$ -simplex t , we get

$$\begin{aligned} d_v(c'_v, p) &\leq d_v(c'_v, c') + d_v(c', p) \\ &\leq \gamma_0^2 d_M(c'_v, c') + \gamma_0^2 d_M(c', p) \\ &\leq \gamma_0^2 f_{k-1}(\rho_0, \sigma_0, \gamma_0^2) r' + \gamma_0^2 [1 + 4\pi k \rho_0 \sigma_0] r' \\ &\leq \gamma_0^2 [1 + f_{k-1}(\rho_0, \sigma_0, \gamma_0^2) + 4\pi k \rho_0 \sigma_0] r' \stackrel{\text{def}}{=} \lambda^+ r', \end{aligned}$$

writing $\lambda^+ = \gamma_0^2 [1 + f_{k-1}(\rho_0, \sigma_0, \gamma_0^2) + 4\pi k \rho_0 \sigma_0]$. In the same way, we have:

$$\begin{aligned} d_v(c'_v, p) &\geq d_v(c', p) - d_v(c'_v, c') \\ &\geq \frac{1}{\gamma_0^2} d_M(c', p) - \gamma_0^2 d_M(c'_v, c') \\ &\geq \frac{1}{\gamma_0^2} [1 - 4\pi k \rho_0 \sigma_0] r' - \gamma_0^2 f_{k-1}(\rho_0, \sigma_0, \gamma_0^2) r' \stackrel{\text{def}}{=} \lambda^- r', \end{aligned}$$

writing $\lambda^- = \frac{1}{\gamma_0^2} [1 - 4\pi k \rho_0 \sigma_0] - \gamma_0^2 f_{k-1}(\rho_0, \sigma_0, \gamma_0^2)$.

It follows that the forbidden region $Y_v(s')$ associated to s' is included in the M_v -spherical shell $\mathcal{S}_v(c'_v, r_v^+, r_v^-)$ centered at c'_v and with radii $r_v^+ = \lambda^+ r'$ and $r_v^- = \lambda^- r'$. Then, we get from Equation 30 an upperbound for the volume of $Y_v(s')$. The M_v -volume of the spherical shell $\mathcal{S}_v(c'_v, r_v^+, r_v^-)$ and thus the M_v -volume of $Y_v(s')$ is at most

$$\text{Vol}_v(Y_v(s')) \leq \phi_d(\lambda^+)^{d-1} (\lambda^+ - \lambda^-) r'^d,$$

where

$$\lambda^+ - \lambda^- = \left[\left(\gamma_0^2 - \frac{1}{\gamma_0^2} \right) + 2\gamma_0^2 f_{k-1}(\rho_0, \sigma_0, \gamma_0^2) + \left(\gamma_0^2 + \frac{1}{\gamma_0^2} \right) 4\pi k \rho_0 \sigma_0 \right] \quad (31)$$

Since the M -circumradius r' of t is smaller than βr_v , we get :

$$\text{Vol}_v(Y_v(s')) \leq \mu_k(\rho_0, \sigma_0, \gamma_0) \beta^d r_v^d,$$

with

$$\mu_k(\rho_0, \sigma_0, \gamma_0) = \phi_d(\lambda^+)^{d-1} (\lambda^+ - \lambda^-) \beta^d r_v^d.$$

From the definitions of λ^+ , Equation 31 and Lemma 6.1 (Circumsphere lemma), the function $\mu_k(\rho_0, \sigma_0, \gamma_0)$ is k -regularly vanishing. \square

Avoiding QC-configurations

In Lemma 6.5, we bounded the volume of a forbidden region associated to a sliver. We will now bound the volume of a forbidden region associated to a QC-configuration. We first prove a technical lemma.

Lemma 6.6 (QC-configuration lemma) *Given the following:*

1. a metric M and a distortion bound $\gamma_0 > 1$,
2. a d -simplex s that is well shaped with respect to M . We denote by c and r the M -circumcenter and the M -circumradius of s .

3. a point p such that the configuration (p, s) is a (γ_0, M) -cospherical configuration.

Then p belongs to the M -spherical shell $\mathcal{S}_M(c, g_d^-(\rho_0, \sigma_0, \gamma_0) r, g_d^+(\rho_0, \sigma_0, \gamma_0) r)$ enclosed between two M -spheres centered at c , with respective radii $g_d^-(\rho_0, \sigma_0, \gamma_0)r$ and $g_d^+(\rho_0, \sigma_0, \gamma_0)r$, where:

$$\begin{aligned} g_d^+(\rho_0, \sigma_0, \gamma_0) &= [\gamma_0^2 + (1 + \gamma_0^2)f_d(\rho_0, \sigma_0, \gamma_0)] \\ g_d^-(\rho_0, \sigma_0, \gamma_0) &= \left[\frac{1}{\gamma_0^2} - \left(1 + \frac{1}{\gamma_0^2}\right)f_d(\rho_0, \sigma_0, \gamma_0) \right]. \end{aligned}$$

Proof Let N and N' be two metrics that witness the (γ_0, M) -cospherical configuration (s, p) , such that p belongs to the interior of the N -circumball $B_N(s)$ while p does not belong to the interior of the N' -circumball $B_{N'}(s)$. Let $c_N, c_{N'}$ denote respectively the N and N' -circumcenters of s . Then, using Lemma 6.1,

$$\begin{aligned} d_M(p, c) &\leq d_M(p, c_N) + d_M(c_N, c) \\ &\leq \gamma_0 d_N(p, c_N) + f_d(\rho_0, \sigma_0, \gamma_0)r \\ &\leq \gamma_0 h_d^+(\rho_0, \sigma_0, \gamma_0)r + f_d(\rho_0, \sigma_0, \gamma_0)r \\ &\leq [\gamma_0^2 + (1 + \gamma_0^2)f_d(\rho_0, \sigma_0, \gamma_0)] r \\ &= g_d^+(\rho_0, \sigma_0, \gamma_0) \end{aligned} \tag{32}$$

and

$$\begin{aligned} d_M(p, c) &\geq d_M(p, c_N) - d_M(c_N, c) \\ &\geq \frac{1}{\gamma_0} d_{N'}(p, c_{N'}) - f_d(\rho_0, \sigma_0, \gamma_0)r \\ &\geq \frac{1}{\gamma_0} h_d^-(\rho_0, \sigma_0, \gamma_0)r - f_d(\rho_0, \sigma_0, \gamma_0)r \\ &\geq \left[\frac{1}{\gamma_0^2} - \left(1 + \frac{1}{\gamma_0^2}\right) f_d(\rho_0, \sigma_0, \gamma_0) \right] r \\ &= g_d^-(\rho_0, \sigma_0, \gamma_0) \end{aligned} \tag{33}$$

Inequalities (32) and (33) are just another way to state Lemma 6.6. \square

Let s be a k -simplex of a star S_v , write r_v for its M_v -circumradius, and consider a refinement point p to be taken from the M_v -picking region $P_v(s)$ of s . Point p is required to lie outside all forbidden regions. After considering the case of a forbidden region associated to a sliver in the previous section, we consider now the case of a forbidden region $W_v(t)$ associated to a hitting set t of $d + 1$ sites that form with p a small well-shaped M -cospherical configuration for some metric M close to M_p . Again (see Definition 4.2), by a metric M close to M_p , we mean such that $\gamma(M, M_p) \leq \gamma_0$, and by a small configuration, we mean a configuration whose M -circumradius is smaller than βr_v . For convenience, as before we denote by t either a subset of sites or the simplex formed by the convex hull of these sites. Note that t is not required to be a simplex appearing in some current star.

Lemma 6.7 (Forbidden regions due to QC-configurations) *The M_v -volume of the region $W_v(t)$ forbidden by the hitting set t is upper bounded as follows*

$$\text{Vol}_v(W_v(t)) \leq \omega(\rho_0, \sigma_0, \gamma_0) \beta^d r_v^d,$$

where $\omega(\rho_0, \sigma_0, \gamma_0)$ is a d -regularly vanishing function.

Proof As for the proof of Lemma 6.5, we prove that the forbidden region $W_v(t)$ is included in a M_v -spherical shell $\mathcal{S}_v(c'_v, r_v^+, r_v^-)$ enclosed between two M_v -spheres centered at c'_v , the M_v -circumcenter of t . For the same reason as in the proof of Lemma 6.5, there exists a metric M satisfying $\gamma(M, M_v) \leq \gamma_0^2$ such that t forms with p a (γ_0, M) -cospherical configuration. Let c' , r' be respectively the M -circumcenter and the M -circumradius of t . Applying Lemmas 6.1 and 6.6 to t , we get:

$$\begin{aligned} d_v(p, c'_v) &\leq d_v(p, c') + d_v(c', c'_v) \\ &\leq \gamma_0^2 d_M(p, c') + \gamma_0^2 d_M(c', c'_v) \\ &\leq \gamma_0^2 g_d^+(\rho_0, \sigma_0, \gamma_0) r' + \gamma_0^2 f_d(\rho_0, \sigma_0, \gamma_0^2) r' \stackrel{\text{def}}{=} \lambda^+ r', \end{aligned} \quad (34)$$

where $\lambda^+ = \gamma_0^2 g_d^+(\rho_0, \sigma_0, \gamma_0) + \gamma_0^2 f_d(\rho_0, \sigma_0, \gamma_0^2)$. Similarly,

$$\begin{aligned} d_v(p, c'_v) &\geq d_v(p, c') - d_v(c', c'_v) \\ &\geq \frac{1}{\gamma_0^2} d_M(p, c') - \gamma_0^2 d_M(c', c'_v) \\ &\geq \frac{1}{\gamma_0^2} g_d^-(\rho_0, \sigma_0, \gamma_0) r' - \gamma_0^2 f_d(\rho_0, \sigma_0, \gamma_0^2) r' \stackrel{\text{def}}{=} \lambda^- r', \end{aligned} \quad (35)$$

where $\lambda^- = \frac{1}{\gamma_0^2} g_d^-(\rho_0, \sigma_0, \gamma_0) - \gamma_0^2 f_d(\rho_0, \sigma_0, \gamma_0^2)$.

It follows that the forbidden region $W_v(t)$ associated to t is included in the M_v -spherical shell $\mathcal{S}_v(c'_v, r_v^+, r_v^-)$ enclosed by the two M_v -spheres centered at c'_v of radii $r_v^+ = \lambda^+ r'$ and $r_v^- = \lambda^- r'$. Therefore, using Equation 30, the volume $\text{Vol}_v(W_v(t))$ is upper bounded as follows:

$$\text{Vol}_v(W_v(t)) \leq \phi_d(\lambda^+)^{d-1} (\lambda^+ - \lambda^-) r'^d,$$

with

$$\begin{aligned} \lambda^+ - \lambda^- &= \left[\left(\gamma_0^4 - \frac{1}{\gamma_0^4} \right) \right. \\ &\quad \left. + \left(\gamma_0^2 (1 + \gamma_0^2) + \frac{1}{\gamma_0^2} \left(1 + \frac{1}{\gamma_0^2} \right) \right) f_d(\rho_0, \sigma_0, \gamma_0) + 2\gamma_0^2 f_d(\rho_0, \sigma_0, \gamma_0^2) \right] \end{aligned} \quad (36)$$

By definition of a hitting set, the QC-configuration (t, p) is required to be small. Specifically, (t, p) has a circumradius that is at most βr_v , which, owing to Lemma 6.3, implies that the M -circumradius r' of t is at most $(1 + \eta(\rho_0, \sigma_0, \gamma_0))\beta r_v$.

Hence, we can write

$$\text{Vol}_v(W_v(t)) \leq \omega(\rho_0, \sigma_0, \gamma_0) \beta^d r_v^d$$

with

$$\omega(\rho_0, \sigma_0, \gamma_0) = \phi_d(\lambda^+)^{d-1} (\lambda^+ - \lambda^-) (1 + \eta(\rho_0, \sigma_0, \gamma_0))^d.$$

Owing to the definition of λ^+ (Equation 34), Equation 36, Lemmas 6.1 (Circumsphere lemma) and 6.6 (QC-configuration lemma), function $\omega(\rho_0, \sigma_0, \gamma_0)$ is a d -regularly vanishing function. \square

Bounding the number of forbidden regions

Lemma 6.8 *Assume that the separation bound $\text{sf}(p) \geq \Lambda \text{sf}(p)$ holds for the current set of vertices and that the algorithm parameters $\alpha_0, \beta, \delta, \rho_0, \sigma_0$, and γ_0 satisfy the relation*

$$\gamma_0 \alpha_0 (\delta + 2\gamma_0^2 \beta (1 + \eta(\rho_0, \sigma_0, \gamma_0))) < 1 \quad (37)$$

Assume that a refinement point is searched in the M_v -picking region $P_v(s)$ of the d -simplex s in the star S_v , and write $K_v(s)$ for the set of hitting subsets of $P_v(s)$. The cardinality of $K_v(s)$ is bounded by a constant K that depends on α_0 , β , δ , ρ_0 , σ_0 , and γ_0 and remains bounded when σ_0 tends to 0 and γ_0 tends to 1 in such a way that $\frac{\gamma_0-1}{\sigma_0^d}$ tends to 0.

Proof First observe that the cardinality of each hitting subset t in $K_v(s)$ is at most $d + 1$. To bound the cardinality of $K_v(s)$, we first bound the cardinality of the set $Q_v(s)$ of vertices that may be part of a hitting set t . For this, we use a volume argument based on an upper bound on the distance $d_v(c_v, q)$ for each $q \in Q_v(s)$ and a lower bound on the distance $d_v(q, q')$ for any two sites (q, q') in $Q_v(s)$.

Let q be a vertex of $Q_v(s)$. The slivers or QC-configurations to avoid are required to have respectively M -circumradii and M -radii smaller than βr_v for some metric M such that $\gamma(M, M_v) \leq \gamma_0^2$. If q belongs to a hitting set corresponding to a sliver, the M -distance from q to p is at most $2\beta r_v$. If q belongs to a hitting set corresponding to a QC-configuration Lemma 6.3, implies that the M -distance from q to p is at most $2\beta(1 + \eta(\rho_0, \sigma_0, \gamma_0))r_v$. In any case, the M_v -distance $d_v(p, q)$ is therefore at most $2\gamma_0^2\beta(1 + \eta(\rho_0, \sigma_0, \gamma_0))r_v$. Moreover, if c_v denotes as usual the M_v -circumcenter of the simplex s to be refined,

$$\begin{aligned} d_v(c_v, q) &\leq d_v(c_v, p) + d_v(p, q) \\ &\leq (\delta + 2\gamma_0^2\beta(1 + \eta(\rho_0, \sigma_0, \gamma_0))) r_v \end{aligned}$$

We have $r_v \leq \alpha_0 \text{sf}(c_v)$ since, when a point is searched in the picking region of a simplex, Rule (1) does not apply anymore. Hence, the inequality above becomes

$$\begin{aligned} d_v(c_v, q) &\leq l_1 \text{sf}(c_v), & (38) \\ \text{with } l_1 &= \alpha_0 (\delta + 2\gamma_0^2\beta(1 + \eta(\rho_0, \sigma_0, \gamma_0))). & (39) \end{aligned}$$

We need now to bound the M_v -distance $d_v(q, q')$ between two sites in $Q_v(s)$ as a function of $\text{sf}(c_v)$. Starting from the separation bound hypothesis, we get

$$\text{sd}(q) = \min_{q' \in V} d_q(q, q') \geq \Lambda \text{sf}(q).$$

Then,

$$\begin{aligned} \min_{q' \in V} d_v(q, q') &\geq \frac{\Lambda}{\Gamma} \text{sf}(q) \\ &\geq \frac{\Lambda}{\Gamma\gamma(q, c_v)} (\text{sf}(c_v) - d_{c_v}(c_v, q)) \quad (\text{from (8)}) \\ &\geq \frac{\Lambda}{\Gamma^2} (\text{sf}(c_v) - \gamma_0 d_v(c_v, q)) \quad (\text{from (2)}) \\ &\geq \frac{\Lambda}{\Gamma^2} (1 - \gamma_0 l_1) \text{sf}(c_v) \quad (\text{from (38)}). \end{aligned}$$

Therefore, we have

$$d_v(q, q') \geq l_2 \text{sf}(c_v) \tag{40}$$

$$\text{with } l_2 = \frac{\Lambda}{\Gamma^2} (1 - \gamma_0 l_1) \tag{41}$$

Observe that l_2 is positive when condition (37) is satisfied. Inequality (40) shows that the M_v -balls centered at the vertices of $Q_v(s)$ and with radii $l_2 \text{sf}(c_v)/2$ are disjoint and inequality (38)

shows that those balls are contained in the M_v -ball $B(c_v, (l_1 + l_2/2) \text{sf}(c_v))$. A volume argument then proves that the cardinality of $Q_v(s)$ is bounded by $(1 + 2l_1/l_2)^d$. By considering all possible simplices with vertices in $Q_v(s)$, we get a bound on the number $|K_v(s)|$ of forbidden regions we need to avoid when picking a refinement point in $P_v(s)$

$$|K_v(s)| \leq |Q_v(s)|^{d+1} \leq (1 + 2l_1/l_2)^{d(d+1)}.$$

Therefore,

$$|K_v(s)| \leq K \tag{42}$$

$$\text{with } K = (1 + 2l_1/l_2)^{d(d+1)} \tag{43}$$

$$= \left[1 + \frac{\Gamma^2}{\Lambda} \frac{l_1}{1 - \gamma_0 l_1} \right]^{d(d+1)}. \tag{44}$$

Assume that we choose Λ_2 as in Equation 28. Then Λ is independant of σ_0 and γ_0 . Lemma 6.3 says that $\eta(\rho_0, \sigma_0, \gamma_0)$ tends to 0 when σ_0 tends to 0 and γ_0 tends to 1 in such a way that $\frac{\gamma_0 - 1}{\sigma_0^d}$ tends to 0. Therefore l_1 and K remain bounded in the same conditions, which achieves the proof of Lemma 5.3. \square

Proof of the picking lemma

Proof When a refinement point p has to be picked in the picking region $P_v(s)$ of some d -simplex s in star S_v , the M_v -volume of the picking region $P_v(s)$ is $\delta^d r_v^d(s) u_d$ where u_d is the volume of the unit Euclidean ball of dimension d .

To be valid, the refinement point has to lie outside the forbidden regions. In the previous lemmas, we have bounded the M_v -volume of the forbidden regions. More precisely, in Lemma 6.5, we gave a bound on the volume of the forbidden region associated to a small k -sliver and, in Lemma 6.7, we gave a bound on the volume of the forbidden region associated to a small QC-configuration. Lemma 6.8 bounds the total number of forbidden regions to avoid.

A valid refinement point exists in $P_v(s)$ if the volume of the picking region exceeds the total volume of the forbidden regions which is guaranteed if the two following conditions hold:

$$K \mu_{k'}(\rho_0, \sigma_0, \gamma_0) \beta^d \leq \delta^d u_d, \quad k' = 1, \dots, d \tag{45}$$

$$K \omega(\rho_0, \sigma_0, \gamma_0) \beta^d \leq \delta^d u_d \tag{46}$$

Assume that α_0, β, δ , and ρ_0 have been chosen in such a way that equations (26), (27) and (37) are satisfied for any value of γ_0 in $[1, 2]$. We may for example start with some $\delta \in]0, 1[$, then choose ρ_0 and β such that equations (26) and (27) are satisfied for $\gamma = 2$. We then choose α_0 so that equation (37) is satisfied for $\gamma = 2$. Note that these three inequations will remain satisfied for any value of γ_0 in $[1, 2]$. From Lemma 6.5 and Lemma 6.7, we know that $\mu_{k'}(\rho_0, \sigma_0, \gamma_0)$ and $\omega(\rho_0, \sigma_0, \gamma_0)$ can be made arbitrarily small when σ_0 tends to 0 and γ_0 tends to 1 in such a way that $(\gamma_0 - 1)/\sigma_0^d$ tends to 0, while lemma 6.8 guarantees that K remains bounded under the same circumstances. It is therefore it is possible to choose σ_0 and γ_0 so as to satisfy Equations (45) and (46). \square

7 Boundaries and sharp features

Up to this section, we have focused on generating anisotropic meshes that cover a given d -dimensional domain D and conform to a varying anisotropic metric field defined on D . By

restricting the stars to the domain D , we have ensured to insert Steiner vertices only within the domain D and we got meshes that roughly cover D . Still, no special attention was paid to get into the final mesh a faithful representation of the domain boundary. This is the purpose of this section. Though the following algorithm could be put at work in any dimension, to be more concrete, we assume in the following that we work in \mathbb{R}^3 : the domain is a 3-dimensional domain that is bounded by smooth or piecewise smooth surfaces. The domain may also be subdivided in subdomains by smooth or piecewise smooth surfaces. In the following, we call boundary surface any surface that bounds the domain or one of the subdomains and has to be faithfully represented in the mesh. We denote the domain by D and the set of boundary surfaces by ∂D .

7.1 Domains bounded by smooth surfaces

We handle first the case where boundary surfaces are smooth surfaces. In the isotropic setting, the problem of meshing a 3-dimensional domain bounded by smooth surfaces may be solved by a Delaunay refinement algorithm [33], based on the notion of restricted Delaunay triangulation. The algorithm refines a set of sites V and its Delaunay triangulation $\text{Del}(V)$, the refinement being guided not only by the restriction of $\text{Del}(V)$ to the domain D but also by its restriction to the set of boundary surfaces ∂D . The restriction of the Delaunay triangulation $\text{Del}(V)$ to a surface is the subcomplex of $\text{Del}(V)$ formed by the facets whose dual Voronoi edges intersect the surface. In the isotropic case, the Delaunay refinement algorithm is known to provide a mesh whose boundary is a faithful approximation of the domain surface [33]. The algorithm we propose here combines the Delaunay refinement algorithm with the star set system of an anisotropic Delaunay mesh.

The algorithm is summarized in Algorithm 4 below. For each vertex $v \in V$, it maintains two restrictions of the star of v in the Delaunay triangulation $\text{Del}_v(V)$ computed using the metric M_v of vertex v . The first one is the *restricted star* S_v formed by the tetrahedra of $\text{Del}_v(V)$ incident to v and whose M_v -circumcenter belongs to the domain D . The second one is the *restricted surface star* T_v formed with the facets of $\text{Del}_v(V)$ incident to v and whose M_v -dual edges intersect ∂D .

We note $S(V) = \{S_v, v \in V\}$ and $T(V) = \{T_v, v \in V\}$ those restricted star sets. Facets in $T(V)$ are also sometimes called *surface facets* hereafter. The refinement algorithm will insert new Steiner points in V applying refinement rules that aim to get rid of bad facets in $T(V)$ and bad tetrahedra in $S(V)$.

A facet in $T(V)$ is considered as bad if either some of its vertices do not belong to a surface in ∂D (topological defect) or if it is oversized, overdistrorted, badly shaped or inconsistent. By definition, each facet t in the restricted surface star T_v admits an M_v -circumball $B_v(c_v(t), r_v(t))$ centered on a surface in ∂D and empty of vertices of V . Such a ball is called an *M_v -surface Delaunay ball*. The size condition for t is an upper bound on the radius $r_v(t)$ and in addition the sizing field used to upper bound $r_v(t)$ takes care of the distortion condition. The shape condition for t is an upper bound on the radius-edge ratio $\rho_v(t)$ where $\rho_v(t)$ is the ratio from $r_v(t)$ to the M_v -length of the M_v -shortest edge of t . At last, a facet t in T_v is considered as inconsistent iff it does not belong to the restricted surface star sets of all its vertices.

As in Algorithm 2 above, tetrahedra in $S(V)$ are considered as bad if they are oversized, overdistrorted, badly shaped (radius-edge ratio and sliverity conditions) or inconsistent.

Rules are applied with a priority order: Rule (i) is applied only if no Rule (j) with $j < i$ can be applied. Rules for facets have a higher priority than rules for tetrahedra except the rule for inconsistent facets that we postpone at the before last position.

Algorithm 4 uses the `Pick_valid` procedure to choose refinement points inserted to get rid of bad surface facet or bad tetrahedra. The refinement point computed for a bad tetrahedron s in star S_v is not the M_v circumcenter $c_v(s)$ of s but a point chosen in the picking region $P_v(s)$.

The refinement point inserted to get rid of a bad surface facet t in the surface star T_v is not the center $c_v(t)$ of a M_v -surface Delaunay ball $B_v(c_v(t), r_v(t))$ of t , but a point chosen in the picking region $P_v(t)$ defined as the intersection of the ball $B_v(c_v(t), \delta r_v(t))$ with the surface in ∂D including $c_v(t)$. Note that the refinement point of a surface facet is always a point of the surface.

Let c be the refinement point computed for a tetrahedron s in some restricted star S_v . Point c is said to encroach a facet t in the restricted surface star T_w if it is included in the M_w -surface Delaunay ball of t . When a refinement rule is applied to a tetrahedron, the computed refinement point c is first tested for encroachment against the current surface star set and inserted only if no encroachment occurs. Otherwise, the refinement point c is rejected and one of the facets encroached by c is refined instead of the tetrahedron. Algorithm 5 (`Insert_or_snap_valid`) given below takes care of this behavior. As a result, a refinement point of a tetrahedron is never inserted in the star system if some surface facet is encroached. This ensures that only points in D are inserted as vertices of the star system.

The refinement algorithm uses the constants α_0 , γ_0 , ρ_0 , σ_0 introduced in Algorithm 2, and an additionnal constant α_1 to tune the density of mesh vertices on the domain boundary. The `Pick_valid` procedure still depends on constant β and δ .

At the end of the algorithm, the stars are consistent and the star sets $S(V)$ and $T(V)$ can be merged into a consistent mesh. Each simplex in the resulting mesh is well shaped with respect to the metrics of its vertices. The boundary of the mesh is a two-manifold triangulated surfaces whose Hausdorff distance to the domain boundary is controlled. If the sizing field is dense enough, the mesh includes a faithful approximation of all the domain boundary surfaces.

The proof of termination of Algorithm 4 is still based on a volume argument.

First, we notice that the Picking Lemma (Lemma 5.3) is still valid for a refinement point of a surface facet. Indeed, the number of hitting configurations and the volume of forbidden regions can still be bounded as in section 6.

Then, the following theorem whose proof is given in the appendix, provides a lower bound on the insertion radius of each mesh vertex.

Theorem 7.1 *Assume that the constant δ is chosen in $]0, 0.5[$, that the constants ρ_0 , β , are chosen so as to satisfy:*

$$\rho_0 \geq 5 \frac{(1 + \delta)}{(1 - \delta)^2} \Gamma^2 \gamma_0^8 \quad (47)$$

$$\beta \geq 5 \left(\frac{1 + \delta}{1 - \delta} \right)^2 \Gamma^2 \gamma_0^8, \quad (48)$$

and that α_0 and α_1 are chosen in $]0, 1[$ so as to satisfy

$$\alpha_1 \leq \frac{1}{7\Gamma\gamma_0^7} \frac{\alpha_0}{\left(1 + \frac{\alpha_0}{2\gamma_0}\right)} \quad (49)$$

$$\alpha_1 \leq \frac{1}{8} \frac{1}{(1 - \delta)} \frac{\Gamma^2 \gamma_0}{\rho_0} \quad (50)$$

$$\alpha_1 \leq \frac{1}{8} \frac{(1 + \delta)}{(1 - \delta)} \frac{\Gamma^2 \gamma_0}{\beta}. \quad (51)$$

The constants Λ_2 , Λ_3 and Λ_4 depend on the algorithm parameters α_0 , α_1 , γ_0 , ρ_0 , and β and on the global distortion Γ . Note that conditions (50) and (51) together with conditions (47) and

Algorithm 4 Refinement algorithm for domain with smooth boundary

- Rule (1) Facet size and distortion**
 If there is a facet t in star T_v
 such that $r_v(t) \geq \alpha_1 \text{sf}(c_v(t))$,
 Insert(Pick_valid(t, M_v));
- Rule (2) Facet without topological defect**
 If there is a facet t in star T_v
 with some vertex $\notin \partial D$
 Insert(Pick_valid(t, M_v));
- Rule (3) Facet radius-edge ratio**
 If a facet t in star T_v is such that $\rho_v(t) > \rho_0$,
 Insert(Pick_valid(t, M_v));
- Rule (4) Tet size and distortion**
 If a tetrahedron s in some star S_v ,
 is such that $r_v(s) \geq \alpha_0 \text{sf}(c_v(s))$,
 Insert_or_snap_valid(s, M_v);
- Rule (5) Tet radius-edge ratio:**
 If a tetrahedron s in some star S_v is such that $\rho_v(s) > \rho_0$,
 Insert_or_snap_valid(s, M_v);
- Rule (6) Sliver removal:**
 If a tetrahedron s in star S_v is a M_v -sliver
 Insert_or_snap_valid(s, M_v);
- Rule (7) Facet consistency:**
 If a facet t in some star T_v is inconsistent
 Insert(Pick_valid(t, M_v));
- Rule (8) Tetrahedron consistency**
 If a tetrahedron s in some star S_v is inconsistent
 Insert_or_snap_valid(s, M_v);
-

Algorithm 5 Insert_or_snap_valid(s, M_v)

$c = \text{Pick_valid}(s, M_v)$
 If c encroaches some facet t in some restricted surface star T_w
 insert(Pick_valid(t, M_w))
 else insert(c)

(48) imply that

$$\alpha_1 \leq \frac{1}{40} \left(\frac{1-\delta}{1+\delta} \right) \frac{1}{\gamma_0^7}. \quad (52)$$

Equations (49) and (52) imply a very dense mesh at least on boundary surfaces. Note however that bounds given in Theorem 7.1 are not tight but largely reflect our will to keep the proof relatively simple.

From the lower bound $\Lambda_3 \text{sf}(p)$ on the insertion radius of each mesh vertex p , we establish a separation bound on mesh vertices as in Lemma 5.8 and conclude the proof of termination by a volume argument as in Section 5.

7.2 Domain bounded by piecewise smooth surfaces

In the case of domains bounded by piecewise smooth surfaces, the meshes are also required to include a faithful representation of the sharp edges (creases) of the bounding surfaces. A first idea to handle piecewise smooth boundary surfaces is to generalize to sharp edges the notion of restricted Delaunay triangulation and to add to the Delaunay refinement process a refinement level for sharp edges. This has been attempted for the generation of isotropic meshes [34] and could be generalized to the star set system. The generation of anisotropic meshes for domains bounded by polyhedral input surfaces is handled this way in [9]. However this approach has to cope with the problem of small angles subtended by sharp edges and surface patches incident on sharp edges. Indeed input angles smaller than $\pi/2$ are known to jeopardize the termination of a Delaunay refinement process. The termination of the refinement process is therefore only granted under severe unrealistic restrictions on the angles formed by boundary surface patches and sharp edges. Such restrictions on input angles are even more stringent in the anisotropic setting where the angular condition on input features has to be respected in the local metric of every point around the feature.

A more promising approach is the method of protecting balls proposed by Cheng et al. [16, 13]. In this approach, sharp edges are first covered by a set of protecting balls whose centers belong to the sharp edges and define a subdivision of these edges into smaller edges. Protecting balls are considered as weighted points and included as initial points in a weighted Delaunay triangulation. The Delaunay refinement is then performed using this weighted Delaunay triangulation where every additional Steiner vertex is inserted with a null weight. Such a weighting scheme ensures the preservation in the final mesh of the initial subdivision of sharp edges induced by the centers of the protecting balls. A solution to generate anisotropic meshes respecting sharp edges would be to transpose the protecting balls approach to the star system of anisotropic Delaunay meshes. Obviously protecting balls in the star system should be turned into balls for the local metric. The possible occurrence of metric discontinuities on a sharp edge could be handled by taking for the metric at each point on the sharp edge the intersection of the metrics of both incident patches. The implementation and full study of such an approach will be reported elsewhere.

8 Conclusion

We have proposed a new class of anisotropic meshes, the so-called anisotropic Delaunay meshes. These meshes conform to a given metric field, can be defined in any dimension, and keep locally the nice properties of Delaunay meshes. We also described an algorithm to generate such meshes in any dimension d . Differently from other methods that have been proposed in dimensions

higher than 2, our algorithm produces meshes with a precise characterization and theoretical guarantees.

The algorithm is simple and has been implemented for $d = 2$ and 3 using the CGAL library [1]. We have also implemented a variant of Algorithm 4 using only Rules (1), (3) and (7) to generate anisotropic surface meshes. Results appear in [8]. Figure 8 shows the output of the algorithm on a 3-dimensional ball where the metric is stretched horizontally in the left part and vertically in the right part. The metric field varies slowly on the figure on the left and rapidly on the figure on the right. In this example, we did not enforce any size bound, so that the refinement is only governed by the need to remove inconsistencies. As expected, the mesh density depends on the distortion of the metric. The line where the eigenvectors exchange their eigenvalues is clearly visible on the figure on the right. Further experimental results will be reported elsewhere.

By placing anisotropic meshes in the realm of Delaunay meshes, our framework allows to benefit from recent advances in isotropic mesh generation. In particular, our approach can benefit from local optimization techniques that greatly improve the quality of Delaunay meshes generated by refinement [40]. For example, since generated meshes are locally Delaunay, ODT methods (optimal Delaunay triangulations) [12, 4] can be applied in our anisotropic framework.

At last, since our algorithm computes the stars independently and then look for inconsistencies among neighboring stars, it is naturally amenable to parallel computation.

Acknowledgments

This work has been partially supported by the Agence Nationale de la Recherche (ANR) under project GIGA (Geometric Inference and Geometric Approximation).

The authors would like to thank Tamal Dey for fruitful discussions.




Figure 5: Two examples of anisotropic meshes produced by our algorithm.

References

- [1] CGAL, Computational Geometry Algorithms Library. <http://www.cgal.org>.
- [2] F. Alauzet. Size gradation control of anisotropic meshes. *Finite Elements in Analysis and Design*, 46(1-2):181–202, 2010.

- [3] F. Alauzet, A. Loseille, A. Dervieux, and P. Frey. Multi-dimensional continuous metric for mesh adaptation. In *Proc. 15th Int. Meshing Roundtable*, pages 191–214, 2006.
- [4] P. Alliez, D. Cohen-Steiner, M. Desbrun, P. Schröder, and M. Yvinec. Variational tetrahedral meshing. In *Proceedings SIGGRAPH*, 2005. To appear.
- [5] I. Babuska and W.C. Rheinboldt. Error estimates for adaptive finite element computations. *SIAM Journal on Numerical Analysis*, pages 736–754, 1978.
- [6] J-D. Boissonnat, C. Wormser, and M. Yvinec. Anisotropic diagrams: Labelle and Shewchuk approach revisited. *Theoretical Computer Science*, 408(2-3):163–173, 2008.
- [7] Jean-Daniel Boissonnat and Steve Oudot. Provably good sampling and meshing of surfaces. *Graphical Models*, 67(5):405–451, 2005.
- [8] Jean-Daniel Boissonnat, Kan-Le Shi, Jane Tournois, and Mariette Yvinec. Anisotropic Delaunay meshes of surfaces. *ACM Transactions on Graphics*, to appear.
- [9] Jean-Daniel Boissonnat, Camille Wormser, and Mariette Yvinec. Locally uniform anisotropic meshing. In *Proceedings of the 24th Annual Symposium on Computational Geometry*, pages 270–277, 2008.
- [10] Houman Borouchaki, Paul Louis George, Frédéric Hecht, Patrick Laug, and Eric Saltel. Delaunay mesh generation governed by metric specifications. part I algorithms. *Finite Elem. Anal. Des.*, 25(1-2):61–83, 1997.
- [11] Frank Bossen and Paul Heckbert. A pliant method for anisotropic mesh generation. In *5th International Meshing Roundtable*, October 1996.
- [12] L. Chen, P. Sun, and J. Xu. Optimal anisotropic meshes for minimizing interpolation errors in l^p -norm. *Mathematics of Computation*, 76(257):179, 2007.
- [13] S. W. Cheng, T.K. Dey, and J.A. Levine. A practical Delaunay meshing algorithm for a large class of domains*. In *Proceedings of the 16th International Meshing Roundtable*, pages 477–494. Springer, 2008.
- [14] Siu-Wing Cheng, Tamal K. Dey, Herbert Edelsbrunner, Michael A. Facello, and Shang-Hua Teng. Silver exudation. *J. ACM*, 47(5):883–904, 2000.
- [15] Siu-Wing Cheng, Tamal K. Dey, Edgar A. Ramos, and Rephael Wenger. Anisotropic surface meshing. In *In SODA'06 : Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*, pages 202–211, New York, NY, USA, 2006. ACM.
- [16] S.W. Cheng, T.K. Dey, and J. Levine. Theory of a practical Delaunay meshing algorithm for a large class of domains. *Algorithms, Architecture and Information Systems Security*, B. Bhattacharya, S. Sur-Kolay, S. Nandy, and A. Bagchi, Eds, 3:17–41, 2008.
- [17] L Paul Chew. Guaranteed-quality mesh generation for curved surfaces. In *Proceedings of the ninth annual symposium on Computational geometry*, pages 274–280. ACM, 1993.
- [18] E.F. D’Azevedo. Optimal triangular mesh generation by coordinate transformation. *SIAM Journal on Scientific and Statistical Computing*, 12:755, 1991.
- [19] Tamal K Dey. *Curve and surface reconstruction*. Cambridge University Press, 2007.

- [20] C. Dobrzynski and P. Frey. Anisotropic Delaunay mesh adaptation for unsteady simulations. *Proceedings of the 17th International Meshing Roundtable*, pages 177–194, 2008.
- [21] Q. Du and D. Wang. Anisotropic centroidal voronoi tessellations and their applications. *SIAM Journal on Scientific Computing*, 26(3):737–761, 2005.
- [22] Herbert Edelsbrunner, Xiang-Yang Li, Gary Miller, Andreas Stathopoulos, Dafna Talmor, Shang-Hua Teng, Alper Üngör, and Noel Walkington. Smoothing and cleaning up slivers. In *STOC '00: Proceedings of the thirty-second annual ACM symposium on Theory of computing*, pages 273–277, New York, NY, USA, 2000. ACM.
- [23] Herbert Edelsbrunner and Nimish R Shah. Triangulating topological spaces. In *Proceedings of the tenth annual symposium on Computational geometry*, pages 285–292. ACM, 1994.
- [24] P.J. Frey and F. Alauzet. Anisotropic mesh adaptation for CFD computations. *Computer methods in applied mechanics and engineering*, 194(48-49):5068–5082, 2005.
- [25] Francois Labelle and Jonathan Richard Shewchuk. Anisotropic Voronoi diagrams and guaranteed-quality anisotropic mesh generation. In *SCG' 03 : Proceedings of the nineteenth annual symposium on Computational geometry*, pages 191–200, New York, NY, USA, 2003. ACM Press.
- [26] Charles L Lawson. Properties of n-dimensional triangulations. *Computer Aided Geometric Design*, 3(4):231–246, 1986.
- [27] Xiang-Yang Li. Generating well-shaped d-dimensional Delaunay meshes. *Theor. Comput. Sci.*, 296(1):145–165, 2003.
- [28] Xiang-Yang Li and Shang-Hua Teng. Generating well-shaped Delaunay meshed in 3d. In *SODA '01: Proceedings of the twelfth annual ACM-SIAM symposium on Discrete algorithms*, pages 28–37. Society for Industrial and Applied Mathematics, 2001.
- [29] Xiang-Yang Li, Shang-Hua Teng, and Alper Üngör. Biting ellipses to generate anisotropic mesh. In *8th International Meshing Roundtable*, October 1999.
- [30] A. Loseille and F. Alauzet. Optimal 3d highly anisotropic mesh adaptation based on the continuous mesh framework. *Proceedings of the 18th International Meshing Roundtable*, pages 575–594, 2009.
- [31] A. Loseille, A. Dervieux, and F. Alauzet. Fully anisotropic goal-oriented mesh adaptation for 3d steady Euler equations. *Journal of Computational Physics*, 229(8):2866–2897, 2010.
- [32] J-M. Mirebeau. *Anisotropic finite element approximation. Theory and algorithms*. PhD thesis, Université Pierre et Marie Curie, Paris 6, 2010.
- [33] Steve Oudot, Laurent Rineau, and Mariette Yvinec. Meshing volumes bounded by smooth surfaces. *Engineering with Computers*, 26:265–279, 2010.
- [34] Laurent Rineau and Mariette Yvinec. Meshing 3D domains bounded by piecewise smooth surfaces. In *Meshing Roundtable conference proceedings*, pages 443–460, 2007.
- [35] Jessica Schoen. Robust, guaranteed-quality anisotropic mesh generation. M.S. thesis, University of California ar Berkeley, 2008.

-
- [36] Jonathan Richard Shewchuk. What is a good linear finite element? Interpolation, conditioning, anisotropy, and quality measures. In <http://www.cs.cmu.edu/~jrs/jrspapers.html>, Manuscript 2002.
 - [37] Jonathan Richard Shewchuk. Updating and constructing constrained delaunay and constrained regular triangulations by flips. In *Proceedings of the nineteenth annual symposium on Computational geometry*, pages 181–190. ACM, 2003.
 - [38] Richard Shewchuk. Star splaying: an algorithm for repairing Delaunay triangulations and convex hulls. In *SCG '05: Proceedings of the twenty-first annual symposium on Computational geometry*, pages 237–246, New York, NY, USA, 2005. ACM.
 - [39] K. Shimada, A. Yamada, and T. Itoh. Anisotropic Triangulation of Parametric Surfaces via Close Packing of Ellipsoids. *Int. J. Comput. Geometry Appl*, 10(4):417–440, 2000.
 - [40] J. Tournois, C. Wormser, P. Alliez, and M. Desbrun. Interleaving Delaunay refinement and optimization for practical isotropic tetrahedron mesh generation. *ACM Transactions on Graphics (TOG)*, 28(3):1–9, 2009.

9 Appendix

9.1 Proof of Lemma 3.2 (Sliver lemma)

Proof (Sliver lemma) In this proof, all lengths, volumes and angles are measured with respect to metric M . We denote by r and $r(v)$ the circumradii of s and $s(v)$ respectively, by V and $V(v)$ their respective volumes, and by e and $e(v)$ the lengths of their respective shortest edges. Let a be the distance from v to the affine hull $\text{aff}(s(v))$ of $s(v)$ and let a' be the distance from v to the sphere $\text{aff}(s(v)) \cap \mathcal{C}(v)$.

Using the fact that s is a sliver, we have

$$V = \frac{1}{k} a V(v) < \sigma_0^k e^k,$$

which yields

$$a < \frac{k\sigma_0^k e^k}{V(v)}.$$

As $s(v)$ is a face of s , we have $e \leq e(v)$, and, since $s(v)$ is not a sliver, $V(v) \geq \sigma_0^{k-1} e(v)^{k-1}$. Then,

$$\begin{aligned} a &< \frac{k\sigma_0^k e^k}{\sigma_0^{k-1} e(v)^{k-1}}, \\ &\leq k\sigma_0 e(v) \\ &\leq 2k\sigma_0 r(v), \end{aligned}$$

which proves the first part of the lemma.

To bound the distance a' , we consider the 2-plane through v and the centers c and c' of the circumspheres \mathcal{C} and $\mathcal{C}(v)$ of s and $s(v)$ respectively. See Figure 6. Let p be the projection of v on the affine hull $\text{aff}(s(v))$ and let p' be the projection of v on the sphere $\text{aff}(s(v)) \cap \mathcal{C}(v)$. Thus $a = \|vp\|$ and $a' = \|vp'\|$. Let q be the point where the ray issued from c that passes through c' intersects \mathcal{C} . Let $\varphi = \widehat{pp'v}$ and $\theta = \widehat{qcp'}$. Observe that $\frac{a}{a'} = \sin \varphi$ and $\sin \theta = \frac{r(v)}{r} \geq \frac{r(v)/e(v)}{r/e} \geq \frac{1}{2\rho_0}$, because $r(v) \geq e(v)/2$ and the radius-edge ratio $\frac{r}{e}$ of s is smaller than ρ_0 .

We distinguish two cases depending on the position of c and v with respect to the affine hull $\text{aff}(s(v))$ of $s(v)$.

In the first case (Figure 6, left part), c and v are on different sides of $\text{aff}(s(v))$. We have $\varphi \geq \frac{\theta}{2}$ and therefore

$$a' = \frac{a}{\sin \varphi} \leq \frac{a}{\sin(\frac{\theta}{2})} \leq \frac{2k\sigma_0 r(v)}{\sin(\frac{1}{2} \arcsin \frac{1}{2\rho_0})} \leq \frac{\pi k\sigma_0 r(v)}{\frac{1}{2} \arcsin \frac{1}{2\rho_0}} \leq 4\pi k\rho_0\sigma_0 r(v)$$

where we have made use of the first part of the lemma and of the fact that $\frac{2}{\pi}u \leq \sin u \leq u$ for any $u \in [0, \frac{\pi}{2}]$ and $u \leq \arcsin u$ for $u \in [0, 1]$

In the second case (Figure 6, right part), c and v are on the same side of $\text{aff}(s(v))$. Then, $\varphi \geq \theta$ and

$$a' = \frac{a}{\sin \varphi} \leq \frac{a}{\sin \theta} \leq 4k\rho_0\sigma_0 r(v),$$

which ends the proof. □




Figure 6: For the proof of the sliver lemma.

9.2 Proof of Lemma 6.1 (Circumsphere lemma)

Proof We first prove the circumsphere lemma when s is a d -simplex. The case of a k -simplex, which easily follows, will be considered in subsection 9.2.4.

9.2.1 Computing the circumcenters

Let $s = (p_0, \dots, p_d)$ be a d -simplex. Since the M_v -circumcenter c_v of s is at equal M_v -distance from all the vertices of s , we have $d_v^2(c_v, p_i) = r_v^2$ for $i = 0, \dots, d$. Therefore,

$$(p_i - c_v)^T F_v^T F_v (p_i - c_v) = (p_0 - c_v)^T F_v^T F_v (p_0 - c_v) \quad i = 1, \dots, d.$$

Equivalently, we have for $i = 0, \dots, d$

$$\begin{aligned} ((p_i - p_0) + (p_0 - c_v))^T F_v^T F_v ((p_i - p_0) + (p_0 - c_v)) &= (p_0 - c_v)^T F_v^T F_v (p_0 - c_v) \\ \Leftrightarrow (p_i - p_0)^T F_v^T F_v (p_i - p_0) &= 2(p_i - p_0)^T F_v^T F_v (c_v - p_0) \end{aligned}$$

Writing $P = (p_1 - p_0, \dots, p_d - p_0)$ for the square matrix whose columns are the vectors $p_i - p_0$, $i = 1, \dots, d$, and $\text{Diag}(A)$ for the column matrix whose elements are the elements of the main diagonal of a square matrix A , the last equation becomes

$$\text{Diag}(P^T F_v^T F_v P) = 2 P^T F_v^T F_v (c_v - p_0),$$

from which we get the position of c_v with respect to the position of the vertices of s

$$c_v - p_0 = \frac{1}{2} (F_v^T F_v)^{-1} P^{-T} \text{Diag}(P^T F_v^T F_v P). \quad (53)$$

An equivalent formula gives the M_w -circumcenter c_w of s .

9.2.2 Bounding the distance between c_v and c_w

In the following, we choose a coordinate system in which $M_v = F_v^T F_v$ and F_v are identity matrices. (Equivalently, we could assume without loss of generality that M_v is the Euclidean metric since the distance $d_v(c_v, c_w)$ is only related to the relative distortion between M_v and M_w .) Then, we deduce from (53) :

$$c_v - p_0 = \frac{1}{2} P^{-T} q \quad (54)$$

$$c_v - c_w = \frac{1}{2} [P^{-T} q - (F_w^T F_w)^{-1} P^{-T} q'] \quad (55)$$

where

$$\begin{aligned} q &= \text{Diag}(P^T P), \\ q' &= \text{Diag}(P^T F_w^T F_w P). \end{aligned}$$

We further write

$$c_v - c_w = \frac{1}{2} [I - (F_w^T F_w)^{-1}] P^{-T} q + \frac{1}{2} (F_w^T F_w)^{-1} P^{-T} (q - q'),$$

where I is the identity matrix. By our choice of the coordinate system, the M_v -norm of a vector x is just the Euclidean norm $\|x\|$ of its coordinates in this reference system. Therefore,

$$d_v(c_v, c_w) = \|c_v - c_w\| \leq \frac{1}{2} \| (I - (F_w^T F_w)^{-1}) P^{-T} q \| + \frac{1}{2} \| (F_w^T F_w)^{-1} P^{-T} (q - q') \|. \quad (56)$$

The following claim provides bounds for the two terms on the right hand side of (56).

Claim 9.1

$$\| (I - (F_w^T F_w)^{-1}) P^{-T} q \| \leq 2(\gamma_0^2 - 1) r_v. \quad (57)$$

$$\| (F_w^T F_w)^{-1} P^{-T} (q - q') \| \leq \gamma_0^2 (\gamma_0^2 - 1) \frac{2^{d+1}}{d} \frac{\rho_0^d}{\sigma_0^d} r_v. \quad (58)$$

Proof Writing $\|A\|$ for the Euclidean norm of a matrix A , ($\|A\| = \sup_{\|x\|=1} \|Ax\|$), we have,

$$\| (I - (F_w^T F_w)^{-1}) P^{-T} q \| \leq \| (I - (F_w^T F_w)^{-1}) \| \| P^{-T} q \|.$$

$F_w^T F_w$ is a symmetric square matrix with eigenvalues in the interval $[\frac{1}{\gamma_0^2}, \gamma_0^2]$. The absolute values of the eigenvalues of matrix $I - (F_w^T F_w)^{-1}$ are thus at most $\gamma_0^2 - 1$. Moreover, from (54), $\|P^{-T} q\| = 2d_v(c_v, p_0)$ is just twice the M_v -circumradius of s , which proves inequality (57).

To prove (58), we write

$$\| (F_w^T F_w)^{-1} P^{-T} (q - q') \| \leq \| (F_w^T F_w)^{-1} \| \| P^{-T} \| \| q - q' \|. \quad (59)$$

We will bound the three terms on the right hand side of (59). We first note that

$$\| (F_w^T F_w)^{-1} \| \leq \gamma_0^2. \quad (60)$$

Then, for $\|P^{-T}\|$, we use the fact that $\|P^{-T}\| \leq \|P^{-T}\|_\infty$ where $\|P^{-T}\|_\infty$ is the maximum absolute value of any entry in P^{-T} . Each entry in P^{-T} is a cofactor of matrix P^T divided by

the determinant of P^T . The determinant of P^T is $d!$ times the M_v -volume of s . Each entry in P^T is a coordinate of some $p_i - p_0$ and therefore less than $\|p_i - p_0\| \leq 2r_v$, which implies that each cofactor of P^T is at most $(d-1)!(2r_v)^{d-1}$. Therefore,

$$\begin{aligned} \|P^{-T}\| &\leq \|P^{-T}\|_\infty \\ &\leq \frac{(d-1)!(2r_v)^{d-1}}{d! \operatorname{Vol}_v(s)} \\ &\leq \frac{2^{d-1}}{d} \frac{\rho_0^{d-1}}{\sigma_0^d e_v}, \end{aligned} \tag{61}$$

where e_v is the M_v -length of the shortest (for M_v) edge of s . We now bound $\|q - q'\|$:

$$\begin{aligned} \|q - q'\| &= \|\operatorname{Diag}(P^T P) - \operatorname{Diag}((P^T F_w^T F_w P))\| \\ &\leq \|\operatorname{Diag}(P^T P) - \operatorname{Diag}((P^T F_w^T F_w P))\|_\infty \\ &\leq \max_i \left| d_v(p_i, p_0)^2 - d_w(p_i, p_0)^2 \right| \\ &\leq 4(\gamma_0^2 - 1) r_v^2 \\ &\leq 4(\gamma_0^2 - 1) \rho_0 e_v r_v \end{aligned} \tag{62}$$

Inequalities (59), (60), (61) and (62) yield (58) which achieves to prove claim 9.1 and inequality (56).

We finally get from (56), (57) and (58) :

$$\begin{aligned} d_v(c_v, c_w) &\leq (\gamma_0^2 - 1)r_v + \frac{1}{2}\gamma_0^2 (\gamma_0^2 - 1) \frac{2^{d+1}}{d} \frac{\rho_0^d}{\sigma_0^d} r_v. \\ &\leq \left[1 + \frac{2^d}{d} \frac{\gamma_0^2 \rho_0^d}{\sigma_0^d} \right] (\gamma_0^2 - 1) r_v \end{aligned}$$

This ends the proof of the first part of Lemma 6.1 in the case of a d -simplex. \square

9.2.3 Bounding the circumradius r_w

Let p be a vertex of s . We have $r_v = d_v(c_v, p)$ and $r_w = d_w(c_w, p)$. Since metric M_v satisfies the triangular inequality,

$$d_v(c_w, p) - d_v(c_v, c_w) \leq d_v(c_v, p) \leq d_v(c_w, p) + d_v(c_v, c_w).$$

Then, using the fact $\gamma(M_v, M_w) \leq \gamma_0$ and the first part of Lemma 6.1,

$$\begin{aligned} \frac{d_w(c_w, p)}{\gamma_0} - f_d(\rho_0, \sigma_0, \gamma_0)r_v &\leq r_v \leq \gamma_0 d_w(c_w, p) + f_d(\rho_0, \sigma_0, \gamma_0)r_v \\ \frac{r_w}{\gamma_0} - f_d(\rho_0, \sigma_0, \gamma_0)r_v &\leq r_v \leq \gamma_0 r_w + f_d(\rho_0, \sigma_0, \gamma_0)r_v. \end{aligned}$$

Therefore,

$$\frac{r_v}{\gamma_0} (1 - f_d(\rho_0, \sigma_0, \gamma_0)) \leq r_w \leq r_v \gamma_0 (1 + f_d(\rho_0, \sigma_0, \gamma_0)), \tag{63}$$

which proves the second part of Lemma 6.1 in the case of a d -simplex.

9.2.4 The case of a k -simplex

In the case of a k -simplex s , the circumcenters c_v and c_w belong to the k -dimensional subspace that is the affine hull, $\text{aff}(s)$, of s . If $\mathcal{C}(v)$ and $\mathcal{C}(w)$ are respectively the M_v and M_w circumspheres of s , the above proof applies verbatim to the spheres $\text{aff}(s) \cap \mathcal{C}(v)$ and $\text{aff}(s) \cap \mathcal{C}(w)$ that are the circumspheres of s in the subspace $\text{aff}(s)$. This yields the proof of Lemma 6.1 in the case of a k -simplex. \square

9.3 Proof of Lemma 6.3 (Circumradii in QC-configurations lemma)

Proof Let s_{\min} and s_{\max} be the simplices with vertices in U having respectively the minimum and maximum M -circumradius.

Let N and N' be the two metrics witnessing the quasi-cosphericity of U . We consider the set of metrics with distortion less than γ_0 from M and a continuous path joining N to N' within this set, for instance the linear interpolation between N and N' . Since the Delaunay triangulations $\text{Del}_N(U)$ and $\text{Del}_{N'}(U)$ are different and since the metric evolve continuously along the path, there is at least a metric M' on the path, with $\gamma(M, M') \leq \gamma_0$ and such that U is M' -cospherical which means that all d -simplices with vertices in U have the same M' -circumradius.

Then, applying twice the Circumsphere lemma 6.1 respectively to s_{\min} and s_{\max} , we get:

$$\begin{aligned} r_M(s_{\max}) &\leq \gamma_0 (1 + f_d(\rho_0, \sigma_0, \gamma_0)) r_{M'}(s_{\max}) \\ &= \gamma_0 (1 + f_d(\rho_0, \sigma_0, \gamma_0)) r_{M'}(s_{\min}) \\ &\leq \gamma_0^2 (1 + f_d(\rho_0, \sigma_0, \gamma_0))^2 r_M(s_{\min}), \end{aligned}$$

which proves lemma 6.3, setting

$$\eta(\rho_0, \sigma_0, \gamma_0) = \gamma_0^2 (1 + f_d(\rho_0, \sigma_0, \gamma_0))^2 - 1.$$

\square

9.4 Proof of Theorem 7.1

Proof We begin by a lemma relating the insertion radius of each mesh vertex to the radius of the bad simplex triggering the insertion. This generalizes Lemma 5.4 by taking care of the surface facet refinement rules and of the effect of snapping mesh vertices to the surface when some encroachment occurs (see the `Insert_or_snap_valid` procedure).

Let p be a mesh vertex. The vertex p is inserted by application of one of the refinement rules 1-8 to either a surface facet t of some star T_v whose M_v -surface Delaunay ball radius is denoted by $r_v(t)$, or to a tetrahedron s of some star S_v whose M_v -circumradius is denoted by $r_v(s)$.

Lemma 9.2 (Second Insertion radius lemma) *The insertion radius $r(p)$ of the mesh vertex p is such that:*

- $r(p) \geq \frac{(1-\delta)}{\Gamma} r_v(t)$ if Rule (1) applies,
- $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(t)$ if one of Rule (2), Rule (3) or Rule (7) applies.
- When Rule (4) applies, $r(p) \geq \frac{(1-\delta)}{\Gamma} r_v(t)$ if no encroachment occurs and $r(p) \geq \frac{(1-\delta)^2}{2\Gamma\gamma_0} r_v(t)$ otherwise.

- When one of Rule (5), (6) or (8) applies, $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(s)$ if no encroachment occurs and $r(p) \geq \frac{(1-\delta)^2}{2\gamma_0^3} r_v(t)$ otherwise.

Proof If Rule(1) applies, $p = \text{Pick_valid}(t, M_v)$, and we have

$$\begin{aligned} r(p) = \min_{q \in P} d_p(p, q) &\geq \frac{1}{\gamma(p, v)} \min_{q \in P} d_w(p, q) \\ &\geq \frac{1}{\Gamma} \min_{q \in P} d_v(p, q) \geq \frac{(1-\delta)}{\Gamma} r_v(t) \end{aligned}$$

The cases of Rule (2), (3) and (7) is analogous to the case of Rule (1) except that now the distortion $\gamma(p, v)$ is known to be less than γ_0 .

When no snapping occurs, the cases of Rule (4), (5), (6) and (8) are analogous to the cases of Rule (2), (3) and (7).

Assume that vertex p is inserted by Rule (4) and that snapping occurs. Then the point c output by $\text{Pick_valid}(s, M_v)$ encroaches a facet t in some surface star T_w , and $p = \text{Pick_valid}(t, M_w)$. We have:

$$\begin{aligned} r(p) = \min_{q \in P} d_p(p, q) &\geq \frac{1}{\gamma_0} \min_{q \in P} d_v(p, q) \\ &\geq \frac{(1-\delta)}{\gamma_0} r_w(t) \end{aligned}$$

Furthermore, because c encroaches facet t in surface star T_w , we have, for any q that is a vertex of t ,

$$r_w(t) \geq \frac{1}{2} d_w(c, q) \geq \frac{1}{2\Gamma} d_v(c, q) \geq \frac{(1-\delta)}{2\Gamma} r_v(s), \quad (64)$$

and therefore

$$r(p) \geq \frac{(1-\delta)^2}{2\Gamma\gamma_0} r_v(s).$$

Assume now that one of Rule (5), (6) or (8) applies, and that snapping occurs. As above $c = \text{Pick_valid}(s, M_v)$ encroaches a facet t in some surface star T_w , and $p = \text{Pick_valid}(t, M_w)$. We have:

$$r(p) \geq \frac{(1-\delta)}{\gamma_0} r_w(t). \quad (65)$$

Furthermore, because c encroaches t , we have for any vertex q of t :

$$r_w(t) \geq \frac{1}{2} d_w(c, q) \geq \frac{1}{2\gamma_0} d_c(c, q) \geq \frac{1}{2\gamma_0^2} d_v(c, q) \geq \frac{(1-\delta)}{2\gamma_0^2} r_v(s) \quad (66)$$

and, using Equations (65) and (66),

$$r(p) \geq \frac{(1-\delta)^2}{2\gamma_0^3} r_v(s).$$

□

We complete now the proof of Theorem 7.1 by induction. Let's assume that there exists constants $\Lambda_2 > \Lambda_4 > \Lambda_3$ such that up to a given stage of the algorithm :

- for any mesh vertex q inserted on the boundary surface, the insertion radius $r(q)$ is such that: $r(q) \geq \Lambda_3 \text{sf}(q)$,

- for any mesh vertex q that is not on the boundary surface, the insertion radius $r(q)$ is such that $r(q) \geq \Lambda_2 \text{sf}(q)$ and the M_q -distance $\delta(q) = d_q(q, \partial D)$ from q to the boundary surface satisfies : $\delta(q) \geq \Lambda_4 \text{sf}(q)$.

Performing a case analysis on the rule that triggers the insertion of the next vertex p , we compute a lower bound on the insertion radius $r(p)$ and a lower bound on the distance $\delta(p)$ if p does not belong to the boundary surface.

Lower bound on the insertion radius $r(p)$

Rule (1). Assume p is inserted by Rule (1) applied on facet t of T_v . Then we have $p = \text{Pick_valid}(t, M_v)$ and, using Lemma 9.2, $r(p) \geq \frac{(1-\delta)}{\Gamma} r_v(t)$. Then,

$$\begin{aligned} r_v(t) &\geq \alpha_1 \text{sf}(c_v(t)) \\ &\geq \frac{\alpha_1}{\Gamma} [\text{sf}(p) - d_p(p, c_v(t))] \\ &\geq \frac{\alpha_1}{\Gamma} [\text{sf}(p) - \Gamma d_v(p, c_v(t))] \\ &\geq \frac{\alpha_1}{\Gamma} [\text{sf}(p) - \Gamma \delta r_v(t)] \end{aligned} \quad (67)$$

Hence,

$$r_v(t) \geq \frac{\alpha_1}{\Gamma(1 + \delta\alpha_1)} \text{sf}(p)$$

and

$$r(p) \geq \frac{(1-\delta)}{\Gamma^2} \frac{\alpha_1}{(1 + \delta\alpha_1)} \text{sf}(p). \quad (68)$$

The induction hypothesis is fulfilled for p , if we have:

$$\Lambda_3 \leq \frac{(1-\delta)}{\Gamma^2} \frac{\alpha_1}{(1 + \delta\alpha_1)}. \quad (69)$$

Rule (2). Assume now that Rule 2 is applied. Then $p = \text{Pick_valid}(t, M_v)$ where t is a facet in T_v with a vertex q that does not belong to ∂D . From the induction hypothesis $\delta(q) \geq \Lambda_4 \text{sf}(q)$. We have:

$$d_v(p, q) \leq (1 + \delta)r_v(t)$$

and

$$\begin{aligned} r_v(t) &\geq \frac{1}{(1 + \delta)} d_v(p, q) \geq \frac{1}{\gamma_0(1 + \delta)} d_q(p, q) \geq \frac{1}{\gamma_0(1 + \delta)} \delta(q) \geq \frac{1}{\gamma_0(1 + \delta)} \Lambda_4 \text{sf}(q), \\ &\geq \frac{\Lambda_4}{\gamma_0^2(1 + \delta)} [\text{sf}(p) - d_p(p, q)] \\ &\geq \frac{\Lambda_4}{\gamma_0^2(1 + \delta)} [\text{sf}(p) - \gamma_0(1 + \delta)r_v(t)] \\ r_v(t) &\geq \frac{\frac{\Lambda_4}{\gamma_0^2(1 + \delta)}}{1 + \frac{\Lambda_4}{\gamma_0}} \text{sf}(p) \end{aligned}$$

Then, using Lemma 9.2:

$$r(p) \geq \frac{1-\delta}{1 + \delta} \frac{\Lambda_4}{\gamma_0^2(\gamma_0 + \Lambda_4)} \text{sf}(p).$$

The induction hypothesis is fulfilled for p , if we have:

$$\Lambda_3 \leq \frac{1 - \delta}{1 + \delta} \frac{\Lambda_4}{\gamma_0^2(\gamma_0 + \Lambda_4)}. \quad (70)$$

Rule (3). Assume p is inserted by Rule (3) applied on facet t of T_v . Then, $p = \text{Pick_valid}(t, M_v)$ and, from Lemma 9.2, we have $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(t)$. To get a lower bound for $r_v(t)$, we argue as in the proof of Lemma 5.6 and, replacing Λ_2 by Λ_3 in Equation (17)), we get:

$$r_v(t) \geq \frac{\frac{\rho_0}{\gamma_0^2} \Lambda_3 \text{sf}(p)}{1 + \frac{\rho_0}{\gamma_0}(1 + \delta) \Lambda_3}.$$

Therefore, in this case

$$r(p) \geq \frac{\rho_0(1 - \delta)}{\gamma_0^3} \frac{\Lambda_3}{1 + \frac{\rho_0}{\gamma_0}(1 + \delta) \Lambda_3} \text{sf}(p),$$

and the inductive hypothesis is fulfilled by p , provided

$$\frac{\rho_0(1 - \delta)}{\gamma_0^3} \frac{1}{1 + \frac{\rho_0}{\gamma_0}(1 + \delta) \Lambda_3} \geq 1. \quad (71)$$

which implies

$$\frac{\rho_0(1 - \delta)}{\gamma_0^3} \geq 1 \quad (72)$$

and

$$\Lambda_3 \leq \frac{\gamma_0}{\rho_0(1 + \delta)} \left(\frac{\rho_0(1 - \delta)}{\gamma_0^3} - 1 \right). \quad (73)$$

Rule (4) Assume p is inserted by Rule (4) applied to the tetrahedron s of the star S_v .

No snapping

If no snapping occurs, $p = \text{Pick_valid}(s, M_v)$, and according to lemma 9.2: $r(p) \geq \frac{(1-\delta)}{\Gamma} r_v(s)$. Then,

$$r_v(s) \geq \alpha_0 \text{sf}(c_v(s))$$

and we perform a computation analogous to the computation leading to Equation (67) to get:

$$r_v(s) \geq \frac{\alpha_0}{\Gamma(1 + \delta\alpha_0)} \text{sf}(p).$$

Then

$$r(p) \geq \frac{(1 - \delta)}{\Gamma^2} \frac{\alpha_0}{(1 + \delta\alpha_0)} \text{sf}(p). \quad (74)$$

and the induction hypothesis is fulfilled for p , if we have:

$$\Lambda_2 \leq \frac{(1 - \delta)}{\Gamma^2} \frac{\alpha_0}{(1 + \delta\alpha_0)}. \quad (75)$$

Snapping

If snapping occurs, $p = \text{Pick_valid}(t, M_w)$ where t is a facet of some star S_w that is encroached by the computed refinement point $c = \text{Pick_valid}(s, M_v)$.

From Lemma 9.2, we have $r(p) \geq \frac{(1-\delta)^2}{2\Gamma\gamma_0} r_v(s)$. Then

$$\begin{aligned} r_v(s) &\geq \alpha_0 \text{sf}(c_v(s)) \\ &\geq \frac{\alpha_0}{\Gamma} [\text{sf}(p) - d_p(p, c_v(s))]. \end{aligned} \quad (76)$$

Furthermore,

$$\begin{aligned} d_p(p, c_v(s)) &\leq d_p(p, c) + d_p(c, c_v(s)) \\ &\leq \Gamma d_w(p, c) + \Gamma d_w(c, c_v(s)) \\ &\leq \Gamma(1 + \delta)r_w(t) + \Gamma\delta r_v(s) \\ &\leq \Gamma\gamma_0 \frac{(1 + \delta)}{(1 - \delta)} r(p) + \Gamma\delta r_v(s) \end{aligned}$$

where the last equation makes use of the fact that, since $p = \text{Pick_valid}(t, M_w)$, $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_w(t)$. Then, we get:

$$\begin{aligned} r_v(s) &\geq \frac{\alpha_0}{\Gamma} \left[\text{sf}(p) - \Gamma\gamma_0 \frac{(1 + \delta)}{(1 - \delta)} r(p) - \Gamma\delta r_v(s) \right] \\ r_v(s) &\geq \frac{\alpha_0}{\Gamma(1 + \alpha_0\delta)} \left[\text{sf}(p) - \Gamma\gamma_0 \frac{(1 + \delta)}{(1 - \delta)} r(p) \right] \end{aligned} \quad (77)$$

from which we deduce:

$$\begin{aligned} r(p) &\geq \frac{(1 - \delta)^2}{2\Gamma^2\gamma_0} \frac{\alpha_0}{1 + \alpha_0\delta} \left[\text{sf}(p) - \Gamma\gamma_0 \frac{(1 + \delta)}{(1 - \delta)} r(p) \right] \\ r(p) &\geq \frac{(1 - \delta)^2}{2\Gamma^2\gamma_0} \frac{\alpha_0}{1 + \alpha_0\delta} \frac{1}{1 + \frac{(1-\delta^2)}{2\Gamma\gamma_0} \frac{\alpha_0}{(1+\alpha_0\delta)}} \text{sf}(p). \end{aligned} \quad (78)$$

It follows that the inductive hypothesis on the insertion radii is still fulfilled in this case if

$$\Lambda_3 \leq \frac{(1 - \delta)^2}{2\Gamma^2\gamma_0} \frac{\alpha_0}{(1 + \alpha_0\delta)} \frac{1}{\left(1 + \frac{(1-\delta^2)}{2\Gamma\gamma_0} \frac{\alpha_0}{(1+\alpha_0\delta)}\right)}.$$

Because δ and α_1 belongs to $[0, 1]$ while Γ and γ_0 are greater than 1,

$$\frac{(1 - \delta^2)}{2\Gamma\gamma_0} \frac{\alpha_0}{(1 + \alpha_0\delta)} \leq 1,$$

so that we can simplify the above condition by requiring

$$\Lambda_3 \leq \frac{(1 - \delta)^2}{4\Gamma^2\gamma_0} \frac{\alpha_0}{(1 + \alpha_0\delta)}. \quad (79)$$

Rule 5. Assume p is inserted by Rule (5) applied to the tetrahedron s of S_v .

No snapping If no encroachment occurs, we have $p = \text{Pick_valid}(s, M_v)$ and from Lemma 9.2 $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(s)$. We argue as in the proof of Lemma 5.6 and, replacing Λ_2 by Λ_3 in Equation (17)), we get:

$$r_v(s) \geq \frac{\frac{\rho_0}{\gamma_0^2} \Lambda_3}{1 + \frac{\rho_0}{\gamma_0} (1 + \delta) \Lambda_3} \text{sf}(p),$$

and

$$r(p) \geq \frac{\frac{(1-\delta)\rho_0}{\gamma_0^3} \Lambda_3}{1 + \frac{\rho_0}{\gamma_0} (1 + \delta) \Lambda_3} \text{sf}(p). \quad (80)$$

The induction hypothesis on the insertion radii is satisfied in that case provided that:

$$\Lambda_2 \leq \frac{\frac{(1-\delta)\rho_0}{\gamma_0^3} \Lambda_3}{1 + \frac{\rho_0}{\gamma_0} (1 + \delta) \Lambda_3}. \quad (81)$$

Snapping

If encroachment occurs, $p = \text{Pick_valid}(t, M_w)$ for some facet t in T_w encroached by $c = \text{Pick_valid}(s, M_v)$ and, according to Lemma 9.2, we have $r(p) \geq \frac{(1-\delta)^2}{2\gamma_0^3} r_v(s)$. Then, we have $r_v(s) \geq \rho_0 e_v(s) \geq \frac{\rho_0}{\gamma_0} r(q)$ where $e_v(s)$ is the M_v -length of the M_v -shortest edge of s and q is the last inserted vertex of $e_v(s)$. Then,

$$r_v(s) \geq \frac{\rho_0}{\gamma_0} r(q) \geq \frac{\rho_0}{\gamma_0} \Lambda_3 \text{sf}(q)$$

For further reference, we set

$$r_v(s) \geq A \text{sf}(q) \quad (82)$$

with

$$A = \frac{\rho_0 \Lambda_3}{\gamma_0}. \quad (83)$$

Then we have:

$$r_v(s) \geq A \frac{[\text{sf}(p) - d_p(p, q)]}{\gamma(p, q)}. \quad (84)$$

Furthermore,

$$\gamma(p, q) \leq \gamma(p, c) \gamma(c, q) \leq \gamma_0^2 \quad (85)$$

and

$$\begin{aligned} d_p(p, q) &\leq d_p(p, c) + d_p(c, q) \\ &\leq \gamma_0 d_w(p, c) + \gamma_0 d_c(c, q) \\ &\leq \gamma_0 ((1 + \delta) r_w(t) + \gamma_0^2 d_v(c, q)) \\ &\leq \gamma_0 (1 + \delta) r_w(t) + \gamma_0^2 (1 + \delta) r_v(s) \\ &\leq \gamma_0^2 \frac{(1 + \delta)}{(1 - \delta)} r(p) + \gamma_0^2 (1 + \delta) r_v(s). \end{aligned} \quad (86)$$

Plugging (85) and (86) in (84), we get:

$$\begin{aligned} r_v(s) &\geq \frac{A}{\gamma_0^2} \left[\text{sf}(p) - \gamma_0^2 \frac{(1 + \delta)}{(1 - \delta)} r(p) - \gamma_0^2 (1 + \delta) r_v(s) \right] \\ r_v(s) &\geq \frac{A}{\gamma_0^2 (1 + (1 + \delta) A)} \left[\text{sf}(p) - \gamma_0^2 \frac{(1 + \delta)}{(1 - \delta)} r(p) \right]. \end{aligned}$$

Then,

$$\begin{aligned} r(p) &\geq \frac{(1-\delta)^2}{2\gamma_0^3} \frac{A}{\gamma_0^2(1+(1+\delta)A)} \left[\text{sf}(p) - \gamma_0^2 \frac{(1+\delta)}{(1-\delta)} r(p) \right]. \\ r(p) &\geq \frac{(1-\delta)^2}{2\gamma_0^5} \frac{A}{(1+(1+\delta)A)} \frac{1}{1 + \frac{(1-\delta^2)}{2\gamma_0^3} \frac{A}{(1+(1+\delta)A)}} \text{sf}(p). \end{aligned} \quad (87)$$

The inductive hypothesis on the insertion radii is fulfilled in this case provided that

$$\frac{(1-\delta)^2}{2\gamma_0^6} \frac{\rho_0}{(1+(1+\delta)\frac{\rho_0\Lambda_3}{\gamma_0})} \frac{1}{1 + \frac{(1-\delta^2)}{2\gamma_0^4} \frac{\rho_0\Lambda_3}{(1+(1+\delta)\frac{\rho_0\Lambda_3}{\gamma_0})}} \geq 1. \quad (88)$$

Rule (6) and (8). Assume p is inserted by Rule (6) or (8) applied to a tetrahedron s of S_v .
No snapping

If no encroachment occurs, $p = \text{Pick_valid}(s, M_v)$ and from Lemma 9.2 $r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(s)$.

Let q be the last inserted vertex of simplex s . Vertex q has been inserted as the refinement point of a simplex s' in some star S_w and we have $r_v(s) \geq \beta r_w(s')$. Then we argue as in the proof of Lemma 5.7 and, replacing Λ_2 by Λ_3 in Equation (25))

$$r(p) \geq \frac{\beta(1-\delta)\Lambda_3}{\gamma_0^3(1+\delta) \left(1 + \frac{\beta\Lambda_3}{\gamma_0}\right)} \text{sf}(p), \quad (89)$$

so that the induction hypothesis on the insertion radii is still verified in this case if

$$\Lambda_2 \leq \frac{\beta(1-\delta)\Lambda_3}{\gamma_0^3(1+\delta) \left(1 + \frac{\beta\Lambda_3}{\gamma_0}\right)}. \quad (90)$$

Snapping

If encroachment occurs, $p = \text{Pick_valid}(t, M_w)$ for some facet t in T_w encroached by p and, according to Lemma 9.2, we have $r(p) \geq \frac{(1-\delta)^2}{2\gamma_0^3} r_v(s)$. Let q be the last inserted vertex of q . We argue as in the proof of Lemma 5.7:

$$\begin{aligned} r_v(s) &\geq \beta r_w(s') \geq \frac{\beta}{\gamma_0(1+\delta)} r(q) \\ &\geq \frac{\beta\Lambda_3}{\gamma_0(1+\delta)} \text{sf}(q), \end{aligned}$$

which is

$$r_v(s) \geq A \text{sf}(q) \text{ with } A = \frac{\beta\Lambda_3}{\gamma_0(1+\delta)}.$$

Repeating the calculation leading from Equation (82) to Equation (87), we get:

$$r(p) \geq \frac{(1-\delta)^2}{2\gamma_0^5} \frac{A}{(1+(1+\delta)A)} \frac{1}{1 + \frac{(1-\delta^2)}{2\gamma_0^3} \frac{A}{(1+(1+\delta)A)}} \text{sf}(p). \quad (91)$$

The inductive hypothesis on the insertion radii is fulfilled in this case provided that

$$\frac{(1-\delta)^2}{2\gamma_0^6(1+\delta)} \frac{\beta}{\left(1 + \frac{\beta\Lambda_3}{\gamma_0}\right)} \frac{1}{1 + \frac{(1-\delta)}{2\gamma_0^3} \frac{\frac{\beta\Lambda_3}{\gamma_0}}{\left(1 + \frac{\beta\Lambda_3}{\gamma_0}\right)}} \geq 1. \quad (92)$$

Rule (7) Assume p is inserted by Rule (7) applied to the facet t of T_v . We have $p = \text{Pick_valid}(t, M_v)$ from Lemma 9.2,

$$r(p) \geq \frac{(1-\delta)}{\gamma_0} r_v(t). \quad (93)$$

Let q be the last inserted vertex of t . Vertex q is the refinement point of a simplex t' and we have:

$$\begin{aligned} r_v(t) &\geq \beta r_w(t') \geq \frac{\beta}{\gamma_0(1+\delta)} r(q). \\ r_v(t) &\geq \frac{\beta\Lambda_3}{\gamma_0(1+\delta)} \text{sf}(q). \end{aligned} \quad (94)$$

Then, since $p = \text{Pick_valid}(t, M_v)$, we have

$$d_p(p, q) \leq \gamma_0 d_v(p, q) \leq \gamma_0(1+\delta) r_v(t)$$

and

$$\text{sf}(q) \geq \frac{1}{\gamma_0} [\text{sf}(p) - d_p(p, q)] \geq \frac{1}{\gamma_0} [\text{sf}(p) - \gamma_0(1+\delta) r_v(t)]. \quad (95)$$

Combining Equation (94) and (95) we get:

$$\begin{aligned} r_v(t) &\geq \frac{\beta\Lambda_3}{\gamma_0^2(1+\delta)} [\text{sf}(p) - \gamma_0(1+\delta) r_v(t)] \\ r_v(t) &\geq \frac{\beta\Lambda_3}{\gamma_0^2(1+\delta)} \frac{1}{1 + \frac{\beta\Lambda_3}{\gamma_0}} \text{sf}(p) \end{aligned}$$

and, from Equation (93),

$$r(p) \geq \frac{(1-\delta)}{(1+\delta)} \frac{\beta\Lambda_3}{\gamma_0^3} \frac{1}{1 + \frac{\beta\Lambda_3}{\gamma_0}} \text{sf}(p).$$

Therefore, the inductive hypothesis on the insertion radii is still satisfied in this case if:

$$\frac{(1-\delta)}{(1+\delta)} \frac{\beta}{\gamma_0^3} \frac{1}{(1 + \frac{\beta\Lambda_3}{\gamma_0})} \geq 1. \quad (96)$$

Lower bound on $\delta(p)$.

It remains to establish a lower bound on $\delta(p)$ in case one of Rules (4), (5), (6) or (8) is applied and no snapping on the surface occurs. Let p be the vertex inserted and x be the point of ∂D closest to p according to the metric M_p :

$$\delta(p) = d_p(x, p).$$

At the time p is inserted, surface Delaunay balls of surface facets cover ∂D and point x belongs to the Delaunay surface ball $B_w(c_w(t), r_w(t))$ of some facet t in the star S_w of some vertex w . Therefore,

$$d_w(c_w(t), x) \leq r_w(t). \quad (97)$$

Then,

$$\begin{aligned} r_w(t) &\leq \alpha_1 \text{sf}(c_w(t)) \leq \alpha_1 \gamma_0 (\text{sf}(x) + d_x(x, c_w(t))) \leq \alpha_1 \gamma_0 (\text{sf}(x) + \gamma_0 r_w(t)) \\ r_w(t) &\leq \frac{\alpha_1 \gamma_0}{1 - \alpha_1 \gamma_0^2} \text{sf}(x). \end{aligned} \quad (98)$$

Furthermore, we may assume that $\gamma(x, p) \leq \gamma_0$. Indeed, otherwise $\delta(p) = d_p(p, x) \geq \text{bdr}_0$ and we are done. Therefore $\gamma(p, w) \leq \gamma(p, x)\gamma(x, w) \leq \gamma_0^2$.

Let us now consider $\delta(p) = d_p(p, x)$. We have:

$$\begin{aligned} \delta(p) &= d_p(p, x) \geq d_p(p, w) - d_p(w, x) \\ &\geq r(p) - \gamma_0^2 d_w(w, x) \leq r(p) - 2\gamma_0^2 r_w(t) \\ &\geq r(p) - \frac{2\gamma_0^3 \alpha_1}{1 - \gamma_0^2 \alpha_1} \text{sf}(x). \end{aligned} \quad (99)$$

where Equation (99) makes use of Equation (98).

To get a lower bound for $\delta(p)$, we now have to get an upper bound for $\text{sf}(x)$. We have:

$$\begin{aligned} \text{sf}(x) &\leq \gamma_0^2 [\text{sf}(p) + d_p(p, x)] \\ &\leq \gamma_0^2 [\text{sf}(p) + \delta(p)]. \end{aligned} \quad (100)$$

Plugging Equation (100) into Equation (99), leads to:

$$\delta(p) \geq r(p) - \frac{2\gamma_0^5 \alpha_1}{1 - \gamma_0^2 \alpha_1} (\text{sf}(p) + \delta(p))$$

and using the induction hypothesis for the insertion radii of internal mesh vertices we get:

$$\delta(p) \geq \frac{1}{1 + \frac{2\gamma_0^5 \alpha_1}{1 - \gamma_0^2 \alpha_1}} \left(\Lambda_2 - \frac{2\gamma_0^5 \alpha_1}{1 - \gamma_0^2 \alpha_1} \right) \text{sf}(p) \quad (101)$$

so that the inductive hypothesis on distance to the surface is satisfied if

$$\Lambda_2 \geq \frac{2\gamma_0^5 \alpha_1}{(1 - \gamma_0^2 \alpha_1)} \quad (102)$$

and

$$\Lambda_4 \leq \frac{1}{1 + \frac{2\gamma_0^5 \alpha_1}{1 - \gamma_0^2 \alpha_1}} \left(\Lambda_2 - \frac{2\gamma_0^5 \alpha_1}{1 - \gamma_0^2 \alpha_1} \right) \quad (103)$$

Close up

The inductive hypothesis is fulfilled if we can satisfy Equations 69 , 70, 72, 73, 75, 79, 81, 88, 90, 92, 96, 102 and 103.

Observe that we can drop Equations (72) and (96) because they are implied respectively by Equations (88) and (92). From Equation (69), we know that we will have $\Lambda_3 \leq (1 - \delta) \frac{\alpha_1}{\Gamma^2}$. In fact, for technical reasons, we will choose:

$$\Lambda_3 = 0.8 \frac{(1 - \delta) \alpha_1}{(1 + \delta) \Gamma^2}. \quad (104)$$

Then from hypothesis in Equations (50) and (51) we will have

$$\frac{(1 + \delta)\rho_0\Lambda_3}{\gamma_0} \leq 0.1 \quad (105)$$

$$\frac{\beta\Lambda_3}{\gamma_0} \leq 0.1, \quad (106)$$

which allows to simplify Equations (81), (88), (90), (92) leading to the system:

$$\Lambda_3 \leq \frac{(1 - \delta)}{\Gamma^2} \frac{\alpha_1}{(1 + \delta\alpha_1)} \quad (69)$$

$$\Lambda_3 \leq \frac{1 - \delta}{1 + \delta} \frac{\Lambda_4}{\gamma_0^2(\gamma_0 + \Lambda_4)} \quad (70)$$

$$\Lambda_3 \leq \frac{\gamma_0}{\rho_0(1 + \delta)} \left(\frac{\rho_0(1 - \delta)}{\gamma_0^3} - 1 \right) \quad (73)$$

$$\Lambda_2 \leq \frac{\alpha_0}{2\Gamma\gamma_0^2 \left(1 + \frac{\alpha_0}{2\gamma_0}\right)} \quad (75)$$

$$\Lambda_3 \leq \frac{(1 - \delta)^2}{4\Gamma^2\gamma_0} \frac{\alpha_0}{(1 + \alpha_0\delta)} \quad (79)$$

$$\Lambda_2 \leq 0.9(1 - \delta) \frac{\rho_0\Lambda_3}{\gamma_0^3} \quad (81s)$$

$$0.8 \frac{(1 - \delta)^2\rho_0}{2\gamma_0^6} \geq 1 \quad (88s)$$

$$\Lambda_2 \leq 0.9 \frac{(1 - \delta)}{(1 + \delta)} \frac{\beta\Lambda_3}{\gamma_0^3} \quad (90s)$$

$$\frac{0.8(1 - \delta)^2\beta}{2\gamma_0^6(1 + \delta)} \geq 1 \quad (92s)$$

$$\Lambda_2 \geq \frac{2\gamma_0^5\alpha_1}{(1 - \gamma_0^2\alpha_1)} \quad (102)$$

$$\Lambda_4 \leq \frac{1}{1 + \frac{2\gamma_0^5\alpha_1}{1 - \gamma_0^2\alpha_1}} \left(\Lambda_2 - \frac{2\gamma_0^5\alpha_1}{1 - \gamma_0^2\alpha_1} \right) \quad (103)$$

Let us then choose Λ_2 from Equations (75) and (102s). From Equation (52), we have $\gamma_0^2\alpha_1 \leq \gamma_0^7\alpha_1 \leq 0.1$, so that

$$\frac{2\gamma_0^5\alpha_1}{(1 - \gamma_0^2\alpha_1)} \leq \frac{2\gamma_0^5\alpha_1}{0.9} \leq 2.23\gamma_0^5\alpha_1 \quad (107)$$

Then Equation (102s) is satisfied if we choose:

$$\Lambda_2 = 3.5\gamma_0^5\alpha_1 \quad (108)$$

Condition (49) ensures that this choice satisfies Equation (75). We then choose Λ_4 from Equation (103). Let

$$A = \frac{1}{1 + \frac{2\gamma_0^5\alpha_1}{1 - \gamma_0^2\alpha_1}} \left(\Lambda_2 - \frac{2\gamma_0^5\alpha_1}{1 - \gamma_0^2\alpha_1} \right)$$

Since from Equation (107), $\frac{2\gamma_0^5\alpha_1}{(1-\gamma_0^2\alpha_1)} \leq 2.3\gamma_0^5\alpha_1$ and from Equation (52), $\gamma_0^5\alpha_1 \leq \gamma_0^7\alpha_1 \leq 0.1$, we have:

$$A \geq \frac{1}{1+2.3\gamma_0^5\alpha_1} \left(\Lambda_2 - \frac{2\gamma_0^5\alpha_1}{1-\gamma_0^2\alpha_1} \right) \geq \frac{1}{1.3} (1.2\gamma_0^5\alpha_1) \geq 0.9\gamma_0^5\alpha_1,$$

so that Equation (103) is satisfied if we choose

$$\Lambda_4 = 0.9\gamma_0^5\alpha_1 \tag{109}$$

It remains to check that Equations (70), (73), (79), (81s), (88s), (90s) and (92s) are satisfied. Equation (70) is satisfied since

$$\begin{aligned} \frac{1-\delta}{1+\delta} \frac{\Lambda_4}{\gamma_0^2(\gamma_0+\Lambda_4)} &= \frac{1-\delta}{1+\delta} \frac{0.9\gamma_0^5\alpha_1}{\gamma_0^2(\gamma_0+0.9\gamma_0^5\alpha_1)} \\ &\geq \frac{1-\delta}{1+\delta} \frac{0.9\gamma_0^2\alpha_1}{(1+0.9\gamma_0^4\alpha_1)} \\ &\geq \frac{1-\delta}{1+\delta} 0.8\gamma_0^2\alpha_1 \\ &\geq \frac{1-\delta}{1+\delta} \frac{0.8\alpha_1}{\Gamma^2} = \Lambda_3 \end{aligned}$$

Since Equation (105) is granted, Equation (73) is satisfied if

$$\frac{\rho_0(1-\delta)}{\gamma_0^3} - 1 \geq 0.1 \iff \rho \geq 1.1 \frac{\gamma_0^3}{(1-\delta)},$$

which is implied by Condition (47).

Equation (79) is equivalent to:

$$0.8 \frac{(1-\delta)}{(1+\delta)} \frac{\alpha_1}{\Gamma^2} \leq \frac{(1-\delta)^2}{4\Gamma^2\gamma_0} \frac{\alpha_0}{(1+\alpha_0\delta)} \iff \alpha_1 \leq \frac{(1-\delta^2)}{3.2\gamma_0} \frac{\alpha_0}{(1+\alpha_0\delta)}$$

Since $\frac{(1-\delta^2)}{3.2\gamma_0} \frac{\alpha_0}{(1+\alpha_0\delta)} \geq \frac{0.75}{3.2\gamma_0} \frac{\alpha_0}{1.5} \geq \frac{\alpha_0}{7\gamma_0}$, this is granted by Condition (49).

Equation (81s) is equivalent to:

$$\rho_0 \geq \frac{1}{0.9(1-\delta)} \gamma_0^3 \frac{\Lambda_2}{\Lambda_3}$$

From our choice for Λ_2 and Λ_3 (Equation (104) and (108)),

$$\frac{\Lambda_2}{\Lambda_3} \leq \frac{3.5(1+\delta)}{0.8(1-\delta)} \Gamma^2 \gamma_0^5 \leq 4.4 \frac{(1+\delta)}{(1-\delta)} \Gamma^2 \gamma_0^5. \tag{110}$$

and Equation (81s) is satisfied iff

$$\rho_0 \geq \frac{4.4(1+\delta)}{0.9(1-\delta)^2} \Gamma^2 \gamma_0^8$$

which is granted by Condition (47).

Equation (88s) is equivalent to

$$\rho_0 \geq \frac{2}{0.8} \frac{\gamma_0^6}{(1-\delta)^2} = 2.5 \frac{\gamma_0^6}{(1-\delta)^2}$$

which is granted by Condition (47).

Equation (90s) is equivalent to

$$\beta \geq \frac{1}{0.9} \left(\frac{1+\delta}{1-\delta} \right) \gamma_0^3 \frac{\Lambda_2}{\Lambda_3}.$$

In view of Equation (110), this is granted iff

$$\beta \geq \frac{4.4}{0.9} \left(\frac{1+\delta}{1-\delta} \right)^2 \Gamma^2 \gamma_0^8.$$

which is granted by Condition (48).

At last, Equation (92s) is equivalent to

$$\beta \geq \frac{2}{0.8} \frac{(1+\delta)}{(1-\delta)^2} \gamma_0^6 = 2.5 \frac{(1+\delta)}{(1-\delta)^2} \gamma_0^6$$

which is still granted by Condition (48).

□



**RESEARCH CENTRE
SOPHIA ANTIPOLIS – MÉDITERRANÉE**

2004 route des Lucioles - BP 93
06902 Sophia Antipolis Cedex

Publisher
Inria
Domaine de Voluceau - Rocquencourt
BP 105 - 78153 Le Chesnay Cedex
inria.fr

ISSN 0249-6399