

Consistency, accuracy and entropic behaviour of remeshed particle methods

Lisl Weynans, Adrien Magni

► **To cite this version:**

Lisl Weynans, Adrien Magni. Consistency, accuracy and entropic behaviour of remeshed particle methods. ESAIM: Mathematical Modelling and Numerical Analysis, EDP Sciences, 2011, 47 (1), pp.57-81. <10.1051/m2an/2012019>. <inria-00631076>

HAL Id: inria-00631076

<https://hal.inria.fr/inria-00631076>

Submitted on 11 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

*Consistency, accuracy and entropic behaviour of
remeshed particle methods*

Lisl Weynans — Adrien Magni

N° 7759

Octobre 2011

— Computational models and simulation —

 *Rapport
de recherche*

Consistency, accuracy and entropic behaviour of remeshed particle methods

Lisl Weynans ^{*}, Adrien Magni [†]

Theme : Computational models and simulation
Applied Mathematics, Computation and Simulation
Équipe-Projet MC2

Rapport de recherche n° 7759 — Octobre 2011 — 30 pages

Abstract: In this paper we analyse the consistency, the accuracy and some entropy properties of particle methods with remeshing in the case of a scalar one-dimensional conservation law. As in [7] we re-write particle methods with remeshing in the finite-difference formalism. This allow us to prove the consistency of these methods, and accuracy properties related to the accuracy of interpolation kernels. Cottet and Magni devised recently in [5] and [19] TVD remeshing schemes for particle methods. We extend these results to the non linear case with arbitrary velocity sign. We present numerical results obtained with these new TVD particle methods for the Euler equations in the case of the Sod shock tube. Then we prove that with these new TVD remeshing schemes the particle methods converge toward the entropy solution of the scalar conservation law.

Key-words: particle methods with remeshing, interpolation kernels, consistency, truncation error, entropic inequalities, total variation, limiters, convergence.

^{*} Univ. Bordeaux, IMB, UMR 5251, F-33400 Talence, France. CNRS, IMB, UMR 5251, F-33400 Talence, France. INRIA, F-33400 Talence, France.

[†] Laboratoire Jean Kuntzmann, Université Joseph Fourier, 51 rue des Mathématiques, 38041 Grenoble Cedex 9, France

Consistency, accuracy and entropic behaviour of remeshed particle methods

Résumé : Nous analysons la consistance, la précision et certaines propriétés entropiques des méthodes particulières avec remaillage dans le cas d'une loi de conservation scalaire monodimensionnelle. Comme dans [7] nous ré-écrivons les méthodes de ce type dans le formalisme des différences finies. Cela nous permet de montrer leur consistance ainsi que des propriétés de précision liées à la précision des noyaux d'interpolation utilisés. Cottet et Magni ont introduit récemment dans [5] et [19] des techniques de remaillage TVD pour les méthodes particulières. Nous étendons ces résultats au cas non linéaire avec signe de la vitesse quelconque. Nous présentons ensuite des résultats numériques obtenus avec ces nouveaux schémas pour les équations d'Euler dans le cas du tube à choc de Sod. Puis nous montrons que les schémas particuliers obtenus avec ces techniques de remaillage TVD convergent vers l'unique solution entropique de la loi de conservation scalaire considérée.

Mots-clés : méthodes particulières avec remaillage, noyaux d'interpolation, consistance, erreur de troncature, inégalités entropiques, variation totale, limiteurs, convergence.

1 Introduction

Particle methods are Lagrangian techniques that have been designed for advection-dominated physical problems. In this class of methods, the fluid is discretized on small masses concentrated on points: the particles, which are moved in a lagrangian way. The classical particles methods used in fluids dynamics are Smoothed Particle Hydrodynamics (SPH) [24], [1], [10] introduced by Monaghan and Particle-In-Cell (PIC) methods [11], [9]. If nothing is done, the distribution of particles becomes less and less uniform as time goes on, because they accumulate naturally in certain zones, for instance near strong gradients, and rarely elsewhere. This phenomenon can lead to a loss of accuracy. A common remedy to this problem consists in periodically creating new particles uniformly distributed by an interpolation of the values of the existing particles, what is usually called remeshing the particles. The remeshing step creates new particles in a conservative way, by distributing the quantities carried by the particles at the nodes of an underlying grid. The frequency at which the interpolation must be performed depends on the simulated flow, but it is often chosen to remesh particles at each time step. This choice allows to solve the non convective part of the considered equations (pressure gradient, diffusion for instance) with variables located on a grid, thus more easily as if they were irregularly distributed. The ability of particle methods with remeshing to simulate satisfactorily fluid dynamics has been studied and validated in the past: [15], [4], [27], [26], [6] for instance. More recent works include [14], [29], and [3].

In [7] remeshed particle methods were rewritten as finite-difference methods and analyzed in this formalism. For example the particle scheme corresponding to a second order interpolation kernel named Λ_2 was found to be equivalent to the Lax-Wendroff scheme in the linear case, whereas in the non-linear case it provided a new finite difference scheme. Recently, Cottet and Magni ([5], [19], [20]) devised TVD particle schemes, with flux limiters in the remeshing kernels. In this paper we keep on focusing on the finite difference analysis of remeshed particles methods, and study their properties of consistency, accuracy and convergence toward entropy solution on a monodimensionnal non linear scalar transport equation in a infinite domain:

$$u_t + (g(u)u)_x = 0, \quad t \geq 0, \quad -\infty < x < +\infty \quad (1)$$

A convergence proof in L^p_{loc} has been established in [25] for a weighted particle method belonging to SPH methods which are purely lagrangian methods. In [17] and [16] are studied the convergence in L^2 and L^1 of renormalized SPH methods. To our knowledge, such an analysis for remeshed particle methods has not been performed yet. Moreover, thanks to the flux limiting, we will deal here with higher degree interpolation kernels than in the latter references, where the kernels were assumed to be positive, and thus performed only linear interpolation. In section 2 we recall briefly the principles of remeshed particle methods, and the interpolation kernels that are classically used to perform remeshing. In section 3 we present how a finite difference scheme can be derived from the particle method with remeshing. Then we study the consistency and accuracy of remeshed particle methods under a CFL condition. Cottet and Magni ([5], [19]) introduced recently a way to perform flux limiting on particles schemes and make them TVD. In section 4 we present how TVD remeshing schemes can be built for non-linear conservation laws with arbitrary sign of the

particle velocity, and a numerical application to the Euler equation. In section 5 we study the convergence of these TVD particle methods toward the entropy solution. This study is motivated by a numerical observation: contrary to some finite difference schemes, as the Lax-Wendroff scheme for instance, the particle methods with remeshing seem to converge toward the entropic solution of the Burgers equation, as it is noticed in [5]. Using techniques inspired by [21], we prove that the new TVD remeshing schemes converge in the L^1_{loc} norm toward the entropy solution.

2 Particle methods with remeshing

2.1 Particle discretization

Here we present how a particle method with remeshing can be introduced to solve the model transport equation (1). If we express this equation using the Lagrangian derivative associated to the material velocity $g(u)$:

$$\frac{du}{dt} = \frac{\partial u}{\partial t} + g(u) \frac{\partial u}{\partial x}$$

we get:

$$\frac{du}{dt} + \frac{\partial g(u)}{\partial x} u = 0.$$

Let $V(t)$ be a material volume element moving at the flow velocity. By applying the Reynolds theorem, we get:

$$\frac{d}{dt} \left(\int_{V(t)} u \, dV \right) = \int_{V(t)} \frac{du}{dt} \, dV + \int_{V(t)} u \frac{\partial g(u)}{\partial x} \, dV$$

Thus:

$$\frac{d}{dt} \left(\int_{V(t)} u \, dV \right) = 0.$$

Following this property, the particle discretization consists in cutting the fluid into small masses concentrated on points: the particles. Each particle j has a location x_j , carries the constant quantity $\alpha_j = V_j u_j$, with V_j the volume of the particle, and moves at velocity $g(u_j)$. The variables carried by the particle satisfy the following equations:

$$\begin{aligned} \frac{d\alpha_j}{dt} &= 0 \\ \frac{dx_j}{dt} &= g(u_j). \end{aligned}$$

To solve this system with the particle method, one has to move particles during one time step and then interpolate them on the nodes of the underlying uniform grid. All particles are initially located on the nodes of a uniform grid, with space step dx . The volumes of particles are equal to the cell volumes. We note x_j^n the location of particle j at time $n \, dt$, u_j^n the value of u carried by the particle, and $\tilde{g}(u_i^n)$ the velocity at which the particle is moved. $\tilde{g}(u_i^n)$ may be equal to $g(u_i^n)$

but not necessarily. An example will be given later. It may also be a function of several variables providing a consistent approximation of $g(u)$:

$$\tilde{g}(u)_i^n = F(u_{j-m}, \dots, u_{j+m}) \quad (2)$$

For consistency reasons that will appear in the proof of the consistency in section 3.2 we impose that $F(u, \dots, u) = g(u)$.

2.2 Interpolation kernels

In this section we shortly review interpolation kernels commonly used for the class of particle methods considered here. More details can be found in [4]. We only present monodimensional interpolation kernels, because very often interpolation kernels in higher dimensions are devised by tensorial products of mono-dimensional interpolation formulas.

Let be a distribution of particles indexed with q , located in x_q , carrying quantities α_q . For instance, in the case of Euler equations for gas dynamics, α_q can be the mass, the momentum or the total energy of particle q . For incompressible flows, in the case of Vortex-In-Cell methods, α_q is the vorticity. Let W be an interpolation kernel. The remeshing process creates new particles at the nodes of an uniform underlying grid, with space step dx . The new quantities $\tilde{\alpha}_i$ at grid points \tilde{x}_i are computed from the former values with the formula:

$$\tilde{\alpha}_i = \sum_q \alpha_q W\left(\frac{\tilde{x}_i - x_q}{dx}\right) \quad (3)$$

The usual interpolation kernels are symmetrical, so as not to favour one direction compared to the others. With a Fourier analysis one can prove that the order of the interpolation is equal to the number of momentums preserved by the new particle distribution, i.e.:

$$\begin{aligned} \sum_i \tilde{\alpha}_i &= \sum_q \alpha_q \\ \sum_i \tilde{\alpha}_i (x - \tilde{x}_i) &= \sum_q \alpha_q (x - x_q) \\ \sum_i \tilde{\alpha}_i (x - \tilde{x}_i)^2 &= \sum_q \alpha_q (x - x_q)^2 \\ &\dots \end{aligned}$$

A family of interpolation kernels can be built by imposing the conservation of a given number of momentums M with a minimal number of grid points. It is about to solve the following system:

$$\sum_{i=1}^M W(x - x_i) x_i^a = x^a \quad \text{for } 0 \leq a \leq M - 1.$$

Λ_1 is the first kernel built with this principle, with $M = 2$. It preserves the first two momentums of the particle distribution.

$$\Lambda_1(x) = \begin{cases} 1 - |x| & \text{if } |x| \leq 1 \\ 0 & \text{if } |x| > 1. \end{cases}$$

This kernel is in practise very diffusive and is not very used by itself. The next interpolation kernel, preserving the first three momenta of the particle distribution, is:

$$\Lambda_2(x) = \begin{cases} 1 - x^2 & \text{if } |x| \leq 1/2 \\ (1 - |x|)(2 - |x|)/2 & \text{if } 1/2 < |x| \leq 3/2 \\ 0 & \text{if } |x| \geq 1.5. \end{cases}$$

Although successfully used in the past, this kernel has the drawback of being very dispersive. This is partially explained by the fact that it is not continuous: a small error on the location of a particle can thus result in a big error on the values of the particles created by the interpolation step. The following interpolation kernels of this family, Λ_3 and Λ_4 respectively need four and five grid points and preserve one or two additional momentums.

Other interpolation kernels can be built by successive convolutions of Λ_1 . They are of increasing regularity, but only preserve the first two momentums thus it is only possible to perform linear interpolation with them. The kernel M_3 , which is C^1 , is traditionnally referred in Particle-In-Cell methods as the TSC (Triangular-shaped cloud) interpolation function.

$$M_3(x) = \begin{cases} 1/2(x + 3/2)^2 - 3/2(x + 1/2)^2 & \text{if } |x| \leq 1/2 \\ 1/2(-|x| + 3/2)^2 & \text{if } 1/2 \leq |x| \leq 3/2 \\ 0 & \text{if } |x| > 3/2. \end{cases}$$

M_4 and M_5 are the following kernels of this family, respectively C^2 and C^3 . Monaghan [23] devised the well known M'_4 kernel with a Richardson extrapolation from a linear combination of M_4 and its derivative.

$$M'_4(x) = \begin{cases} 1 - 5x^2/2 + 3|x|^3/2 & \text{if } |x| \leq 1 \\ (2 - |x|)^2(1 - |x|)/2 & \text{if } 1 < |x| \leq 2 \\ 0 & \text{if } |x| \geq 2. \end{cases}$$

This kernel is C^1 and preserves the first three momentums.

3 Consistency and accuracy of particle methods with remeshing

3.1 Results

Following the notations introduced in subsection 2.1, we denote u_j^n the value carried by the particle initially located on grid point j , at time $n\Delta t$. It is possible to express u_j^{n+1} as a function of the u_i^n :

Proposition 1 *If we note $\tilde{g}(u)_i^n$ the velocity used to move the particle i between the times $n dt$ and $(n+1) dt$, and Λ the interpolation kernel used for the remeshing step, we can express the particle method with remeshing as a finite-difference scheme:*

$$u_j^{n+1} = \sum_i u_i^n \Lambda\left(j - i - \frac{dt \tilde{g}(u)_i^n}{dx}\right). \quad (4)$$

Remark: In the latter formula, we use the symbol \sum_i without specifying to which interval belong the i indices. Because the interpolation kernel considered here have all compact supports, but of various sizes, this notation avoids us to specify the support of the considered interpolation kernel.

Proof of Proposition 1:

The location of the particle i after being moved is:

$$\tilde{x}_i = x_i + dt \tilde{g}(u)_i^n.$$

The particles are remeshed on the same uniform grid as the one on which they were initially defined. Thus the locations of the new particles are again the nodes of the grid, and their volumes are all equal to dx . The new particle distribution is computed from the old one with the interpolation formula:

$$\alpha_j^{n+1} = \sum_i \alpha_i^n \Lambda\left(\frac{x_i - \tilde{x}_i}{dx}\right).$$

We can re-write the latter equation as:

$$u_j^{n+1} V_j^{n+1} = \sum_i u_i^n V_i^n \Lambda\left(\frac{x_j - x_i - dt \tilde{g}(u)_i^n}{dx}\right).$$

As $V_j^{n+1} = V_i^n = dx$ this equation simplifies in:

$$u_j^{n+1} = \sum_i u_i^n \Lambda\left(\frac{x_j - x_i - dt \tilde{g}(u)_i^n}{dx}\right).$$

x_i and x_j being the nodes of a uniform grid, with space step dx , we have:

$$\frac{x_j - x_i}{dx} = j - i.$$

We finally get:

$$u_j^{n+1} = \sum_i u_i^n \Lambda\left(j - i - \frac{dt}{dx} \tilde{g}(u)_i^n\right).$$

□

We recognize the form of a finite-difference or finite-volume scheme. For example, if we develop this formula with the kernel Λ_2 in the linear case $g(u) = a > 0$, with the CFL condition $|a \frac{dt}{dx}| < \frac{1}{2}$ we get the Lax-Wendroff scheme. Monaghan ([22]) had already noticed a similarity between particle methods and finite differences methods. In [8], Wee et Ghoniem used an analysis similar to ours to build modified interpolation kernels taking into account diffusion terms. The formula (4) is difficult to interpret by itself, because the weights associated to the values u_i^n are expressed with the kernel Λ . But it is possible to obtain several properties of the particle scheme only with the knowledge of the number of momentums preserved by the interpolation kernel Λ . The first one addresses the consistency of particles methods:

Proposition 2 *Let Λ be an interpolation kernel piecewise polynomial of degree N with compact support, which preserves at least the two first momentums. The scheme (4) can be written in a conservative form consistent with the equation (1).*

To solve a flow where shocks may appear, it is crucial that the numerical scheme can be written in conservative form consistent with the equation to solve. In hyperbolic problems this property ensures the scheme to satisfy the discrete Rankine-Hugoniot conditions across discontinuities. Thus, if the scheme is converging, it converges necessarily toward a weak solution of the considered equation. At the contrary, a non-conservative scheme used to solve conservation laws will have a problematic convergence ([2] and [13]).

One can distinguish two kinds of interpolation kernels: kernels whose support size is an even integer (for example kernels Λ_1 , Λ_3 , M_4 and M'_4), and kernels whose support size is odd (for example Λ_2 and M_3). For the sake of brevity, in the following we will call them respectively kernels with even or odd support. The finite difference stencil provided by kernels with even support varies with the sign of the velocity of the particles. For this reason these kernels suffer from problems of consistency in the finite difference sense (visible through an analysis of the truncation error) if the velocity of the particles changes sign, as it is proved in [20] and [19]. On the contrary, kernels with odd support like Λ_2 are defined on intervals $[k - 1/2, k + 1/2]$, and the formula used to interpolate a particle on a grid point does not depend on the velocity of the particle. Under a CFL condition $|g(u_j) \frac{dt}{dx}| < \frac{1}{2} \forall j$ these kernels do not suffer from consistency problems. In the following we will therefore focus our study on kernels with odd support. The second property that we will prove addresses the accuracy of particle schemes:

Proposition 3 *Let Λ be an interpolation kernel with odd support, piecewise polynomial of degree N , which preserves the M first momentums, and u a solution of equation (1). For a given $n \geq 0$, we denote $u_j^n = u(jdx, ndt) \forall j$. If we suppose that the functions u et \tilde{g} are at least of class C^{M-1} , and the CFL condition $|g(u_j) \frac{dt}{dx}| < \frac{1}{2} \forall j$ is satisfied, then u_j^{n+1} defined by (4) satisfies:*

$$u_j^{n+1} = \sum_{i=0}^{M-1} \frac{dt^i}{i!} (-1)^i \frac{\partial^i (u \tilde{g}^i)}{\partial x^i} (jdx, ndt) + O(dx^M) \quad (5)$$

This property is also satisfied by kernels with even support, if the sign of the velocity of the particles is constant. In this case actually, the stencil used to remesh the particles does not change, and the reasoning is the same as for kernels with odd support. The proofs of propositions 2 et 3 are detailed in section 3.2.

The formula (5) allows us to evaluate the accuracy of the particle method with respect to time and space. We assume that dx and dt are proportional. If

we take $\tilde{g}(u) = g(u)$, the truncation error E_j^{n+1} is:

$$\begin{aligned}
 E_j^{n+1} &= \frac{u(jdx, (n+1)dt) - u_j^{n+1}}{dt} \\
 &= \frac{u(jdx, (n+1)dt) - u(jdx, ndt)}{dt} + (g(u)u)_x(jdx, ndt) \\
 &\quad - \frac{dt}{2}(g(u)^2u)_{xx}(jdx, ndt) + O(dx^2) \\
 &= u_t(jdx, ndt) + \frac{dt}{2}u_{tt}(jdx, ndt) + (g(u)u)_x(jdx, ndt) \\
 &\quad - \frac{dt}{2}(g(u)^2u)_{xx}(jdx, ndt) + O(dx^2) + O(dt^2) \\
 &= O(dt)
 \end{aligned}$$

The scheme is thus first order accurate even if the interpolation kernel preserves a higher number of moments, because of the non-zero term $\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 (g(u)^2 u)}{\partial x^2}$. Actually, the choice $\tilde{g}(u) = g(u)$ consists in moving the particle during the time step dt with its velocity evaluated at the beginning of the time step:

$$\tilde{x}_j = x_j + dt g(u_j).$$

It is an explicit first order Euler scheme, and consequently the particle moving is also order one. Therefore, in order to increase the scheme accuracy, the velocity of particles during time step dt needs to be evaluated with a better accuracy. In [7] was introduced a new Runge-Kutta 2 advancing scheme for the particles, allowing to recover in the non-linear case a second order accuracy. The idea is to use the velocity of the particle at the middle of the time step, at $t + \frac{dt}{2}$. Because this value is not exactly known, it is replaced by a second order approximation: $g(u + \frac{dt}{2} \frac{du}{dt})$. Thus we move the particles at velocity:

$$\tilde{g}(u) = g\left(u + \frac{dt}{2} \frac{du}{dt}\right). \quad (6)$$

That is:

$$\tilde{x}_j = x_j + dt g\left(u + \frac{dt}{2} \frac{du}{dt}\right)_{x=x_j}.$$

Proposition 4 *For smooth enough solutions, the particle scheme (4) computed with the corrected velocity (6) is second order accurate if Λ preserves at least the first three moments.*

Proof of Proposition 4:

The scheme truncation error becomes:

$$\begin{aligned}
 E_j^{n+1} &= \frac{u(jdx, (n+1)dt) - u_j^{n+1}}{dt} \\
 &= u_t(jdx, ndt) + \frac{dt}{2}u_{tt}(jdx, ndt) + (g(u + \frac{dt}{2} \frac{du}{dt})u)_x(jdx, ndt) \\
 &\quad - \frac{dt}{2}(g(u + \frac{dt}{2} \frac{du}{dt})^2u)_{xx}(jdx, ndt) + O(dx^2) \\
 &= u_t(jdx, ndt) + \frac{dt}{2}u_{tt}(jdx, ndt) + (g(u)u)_x(jdx, ndt) \\
 &\quad + \frac{dt}{2}\left(\frac{du}{dt}ug'(u)\right)_x(jdx, ndt) - \frac{dt}{2}(g(u)^2u)_{xx}(jdx, ndt) + O(dx^2).
 \end{aligned}$$

We want to prove that the terms: $u_{tt} + (\frac{du}{dt}ug'(u))_x - (g(u)^2u)_{xx}$ vanish.

$$\begin{aligned} u_{tt} &= -(g(u)u)_{tx} = -(g'(u)uu_t + g(u)u_t)_x = [(g'(u)u + g(u))(g(u)u)_x]_x \\ \frac{du}{dt} &= -(g(u))_x \end{aligned}$$

We check that:

$$[(g'(u)u + g(u))(g(u)u)_x]_x - ((\frac{\partial(g(u))}{\partial x}ug'(u))_x - (g(u)^2u)_{xx}) = 0.$$

Thus we obtain:

$$E_j^{n+1} = O(dx^2).$$

□

3.2 Proof of propositions 2 and 3

We present in this subsection the proofs of propositions 2 and 3 in the case of a kernel with odd support. These proofs are based on the fact that interpolation kernels preserve a certain number of momenta. We begin with the proof of an lemma that will be used in the following.

Lemma 1 *Let Λ be a interpolation kernel with odd support, preserving the M first momentums. $\forall m, \forall i$ such that $0 \leq i \leq M - 1$ we have:*

$$\sum_k k^i \Lambda^{(m)}(k) = (-1)^i i! \delta_i^m \quad (7)$$

Proof of lemma 1

Λ preserves the M first momenta:

$$\sum_k k^i \Lambda(k - x) = x^i \quad \forall 0 \leq i \leq M - 1.$$

We deduce from this formula that:

$$\begin{aligned} \text{if } i = 0 \text{ and } m > 0 & \quad \sum_k k^i \Lambda^{(m)}(k - x) = 0 \\ \text{if } i = m = 0 & \quad \sum_k k^i \Lambda^{(m)}(k - x) = \sum_k \Lambda(k - x) = 1 \\ \text{if } i \neq 0 \text{ and } 0 \leq m \leq i & \quad \sum_k k^i (-1)^m \Lambda^{(m)}(k - x) = i(i-1)\dots(i-m+1)x^{i-m} \\ \text{if } i \neq 0 \text{ and } m > i & \quad \sum_k k^i (-1)^m \Lambda^{(m)}(k - x) = 0. \end{aligned}$$

Thus, for $x = 0$ we have:

$$\begin{aligned} \text{if } i = 0 \text{ and } m > 0 & \quad \sum_k k^i \Lambda^{(m)}(k) = 0 \\ \text{if } i = m = 0 & \quad \sum_k k^i \Lambda(k) = 1 \\ \text{if } i \neq 0 \text{ and } m \neq i & \quad \sum_k k^i \Lambda^{(m)}(k) = 0 \\ \text{if } i \neq 0 \text{ and } m = i & \quad \sum_k k^i \Lambda^{(m)}(k) = (-1)^i i!. \end{aligned}$$

□

Proof of proposition 2 (consistency of the scheme)

Let u be a smooth solution of equation $u_t + (g(u)u)_x = 0$ for a given initial condition. Let us denote:

$$u_j^n = u(jdx, ndt) \quad \forall n \geq 0, \forall j.$$

We consider the formula (4) written in a slightly different form:

$$u_j^{n+1} = \sum_k u_{j+k}^n \Lambda(k + \frac{dt}{dx} \tilde{g}(u)_{j+k}^n). \quad (8)$$

For the sake of clarity we note $\tilde{g}_{j+k} = \tilde{g}(u)_{j+k}^n$, $u_j = u_j^n$ and $\lambda = \frac{dt}{dx}$. Formula (8) becomes:

$$u_j^{n+1} = \sum_k u_{j+k} \Lambda(k + \lambda \tilde{g}_{j+k}).$$

N is the degree of Λ . The kernel Λ being derivable on each interval $[k - 1/2, k + 1/2]$, and because of the CFL condition, we can develop each term in a Taylor expansion:

$$\Lambda(k + \lambda \tilde{g}_{j+k}) = \sum_{i=0}^N \frac{\Lambda^{(i)}(k)}{i!} (\lambda \tilde{g}_{j+k})^i.$$

We deduce from the latter that:

$$u_j^{n+1} = \sum_k \sum_{i=0}^N \frac{\lambda^i}{i!} \Lambda^{(i)}(k) u_{j+k} \tilde{g}_{j+k}^i \quad (9)$$

As we have assumed that the support of Λ is compact, it can be included in $[-d, d]$, with d an integer, and equation (9) can be re-written:

$$\begin{aligned} u_j^{n+1} &= \sum_{k=-d, k \neq 0}^d \left[\sum_{i=0}^N u_{j+k} \frac{\lambda^i}{i!} \Lambda^{(i)}(k^+) \tilde{g}_{j+k}^i \right] \\ &+ \left[\sum_{i=0}^N u_j \frac{\lambda^i}{i!} \Lambda^{(i)}(0^+) \tilde{g}_j^i \right] \end{aligned}$$

As $\sum_k \Lambda(k+x) = 1 \quad \forall x$, we have:

$$\Lambda^{(i)}(0) = \delta_0^i - \sum_{k \neq 0} \Lambda^{(i)}(k)$$

Thus:

$$u_j^{n+1} = u_j + \sum_{k=-d, k \neq 0}^d \sum_{i=0}^N \frac{\lambda^i}{i!} \Lambda^{(i)}(k) [u_{j+k} \tilde{g}_{j+k}^i - u_j \tilde{g}_j^i]$$

$$u_j^{n+1} = u_j + \sum_{k=1}^d \sum_{i=0}^N \frac{\lambda^i}{i!} \left[\Lambda^{(i)}(k) [u_{j+k} \tilde{g}_{j+k}^i - u_j \tilde{g}_j^i] + \Lambda^{(i)}(-k) [u_{j-k} \tilde{g}_{j-k}^i - u_j \tilde{g}_j^i] \right]$$

We notice that:

$$\begin{aligned} u_{j+k}\tilde{g}_{j+k}^i - u_j\tilde{g}_j^i &= \sum_{a=1}^k u_{j+a}\tilde{g}_{j+a}^i - \sum_{a=0}^{k-1} u_{j+a}\tilde{g}_{j+a}^i \\ u_{j-k}\tilde{g}_{j-k}^i - u_j\tilde{g}_j^i &= -\sum_{a=-k+1}^0 u_{j+a}\tilde{g}_{j+a}^i + \sum_{a=-k}^{-1} u_{j+a}\tilde{g}_{j+a}^i \end{aligned}$$

Thus:

$$\begin{aligned} u_j^{n+1} &= u_j + \sum_{k=1}^d \sum_{i=0}^N \frac{\lambda^i}{i!} \left[\Lambda^{(i)}(k) \left[\sum_{a=1}^k u_{j+a}\tilde{g}_{j+a}^i - \sum_{a=0}^{k-1} u_{j+a}\tilde{g}_{j+a}^i \right] \right. \\ &\quad \left. + \Lambda^{(i)}(-k) \left[-\sum_{a=-k+1}^0 u_{j+a}\tilde{g}_{j+a}^i + \sum_{a=-k}^{-1} u_{j+a}\tilde{g}_{j+a}^i \right] \right] \end{aligned}$$

We can then write:

$$u_j^{n+1} = u_j - \lambda \left[G(u_{j+d}, \dots, u_{j-d+1}) - G(u_{j+d-1}, \dots, u_{j-d}) \right]$$

with:

$$G(u_{j+d}, \dots, u_{j-d+1}) = -\sum_{k=1}^d \sum_{i=0}^N \frac{\lambda^{i-1}}{i!} \left[\Lambda^{(i)}(k) \sum_{a=1}^k u_{j+a}\tilde{g}_{j+a}^i - \Lambda^{(i)}(-k) \sum_{a=-k+1}^0 u_{j+a}\tilde{g}_{j+a}^i \right]$$

Consequently the scheme (4) can be written in conservative form. Moreover:

$$G(u, \dots, u) = -\sum_{k=1}^d \sum_{i=0}^N \frac{\lambda^{i-1}}{i!} \left[\Lambda^{(i)}(k) \sum_{a=1}^k u g(u)^i - \Lambda^{(i)}(-k) \sum_{a=-k+1}^0 u g(u)^i \right]$$

$$G(u, \dots, u) = -\sum_{k=1}^d \sum_{i=0}^N u g(u)^i \frac{\lambda^{i-1}}{i!} \left[k\Lambda^{(i)}(k) - k\Lambda^{(i)}(-k) \right]$$

According to lemma 1, we have then:

$$G(u, \dots, u) = u g(u).$$

The scheme (4) is therefore consistent with equation (1).

□

Proof of proposition 3 (accuracy of the scheme)

We start again from formula (8):

$$u_j^{n+1} = \sum_k u_{j+k} \Lambda(k + \lambda\tilde{g}_{j+k}).$$

The kernel Λ being derivable on each interval $[k - 1/2, k + 1/2]$, we have:

$$\Lambda(k + \lambda\tilde{g}_{j+k}) = \sum_{i=0}^N \frac{\Lambda^{(i)}(k)}{i!} (\lambda\tilde{g}_{j+k})^i.$$

Thus:

$$u_j^{n+1} = \sum_k \sum_{i=0}^N \frac{\lambda^i}{i!} \Lambda^{(i)}(k) u_{j+k} \tilde{g}_{j+k}^i.$$

We define $(f_i)_{j+k} = u_{j+k} \tilde{g}_{j+k}^i$ and develop these terms in a Taylor serie:

$$(f_i)_{j+k} = \sum_{a=0}^{M-1} \frac{\partial^a (f_i)}{\partial x^a}(x_j) \frac{k^a dx^a}{a!} + O(dx^M)$$

Thus:

$$u_j^{n+1} = \sum_k \sum_{i=0}^N \frac{\lambda^i}{i!} \sum_{a=0}^{M-1} \frac{k^a dx^a}{a!} \Lambda^{(i)}(k) \frac{\partial^a (f_i)}{\partial x^a}(x_j) + O(dx^M).$$

We exchange the symbols \sum :

$$u_j^{n+1} = \sum_{i=0}^N \frac{\lambda^i}{i!} \sum_{a=0}^{M-1} \frac{dx^a}{a!} \frac{\partial^a (f_i)}{\partial x^a}(x_j) \sum_k k^a \Lambda^{(i)}(k) + O(dx^M).$$

With the results of lemma 1 we simplify the latter expression:

$$\begin{aligned} u_j^{n+1} &= \sum_{i=0}^{M-1} \frac{\lambda^i}{i!} \frac{dx^i}{i!} \left[\frac{\partial^i (f_i)}{\partial x^i}(x_j) (-1)^i i! \right] + O(dx^M) \\ &= \sum_{i=0}^{M-1} \frac{\lambda^i}{i!} (-1)^i dx^i \frac{\partial^i (u \tilde{g}(u)^i)}{\partial x^i}(x_j) + O(dx^M). \end{aligned}$$

□

4 TVD remeshing formulas


Recently Cottet and Magni derived in [5], [19] and [20] TVD remeshing formulas for particle methods. This subsection gives the principles to construct TVD remeshing formulas based on the Λ_2 kernel in the non-linear case and for CFL conditions less than 1/2, avoiding consistency problems evocated in [19]. Exemples are given for Burgers and Euler equations.

4.1 Principle of TVD remeshing schemes

Let the velocity be positive ($0 < \lambda \tilde{g} < 1/2$). The remeshing of a particle with the Λ_2 kernel is equivalent to affect the weights

$$\begin{cases} \alpha(y_j) &= \alpha_j = y_j (y_j - 1) / 2 \\ \beta(y_j) &= \beta_j = 1 - y_j^2 \\ \gamma(y_j) &= \gamma_j = y_j (y_j + 1) / 2 \end{cases} \quad (10)$$


to the amount carried by the particle and redistribute it on the neighbouring grid points. This is sketched on the Figure 1, where $y_j = \lambda \tilde{g}_j$ is the relative


 Figure 1: Λ_2 remeshing formulas, $0 < \lambda\tilde{g} < 1/2$.

distance between the remeshed particle j and the grid point at the left. The scheme resulting of the Λ_2 remeshing (10) at the grid point x_j and the time $t^n = n dt$ is

$$\begin{aligned} u_j^{n+1} &= \sum_k u_{j+k}^n \Lambda_2(k + \lambda\tilde{g}_{j+k}) \\ &= \gamma_{j-1} u_{j-1}^n + \beta_j u_j^n + \alpha_{j+1} u_{j+1}^n. \end{aligned} \quad (11)$$

If now the velocity is negative ($-1/2 < \lambda\tilde{g} < 0$) as shown in the figure 2, the


 Figure 2: Λ_2 remeshing formulas, $-1/2 < \lambda\tilde{g} < 0$.

weights are

$$\begin{cases} \alpha'(y_j) &= \alpha(y_j - 1) \\ \beta'(y_j) &= \beta(y_j - 1) \\ \gamma'(y_j) &= \gamma(y_j - 1) \end{cases} \quad (12)$$

and the scheme is

$$u_j^{n+1} = \gamma'_{j-1} u_{j-1}^n + \beta'_j u_j^n + \alpha'_{j+1} u_{j+1}^n. \quad (13)$$

Since $y_j = \lambda\tilde{g}_j + 1$, (11) is equivalent to (13) and a unique scheme is given at the point x_j independently of the velocity sign. The order of accuracy is two, and the drawback of this scheme is to be not TVD, allowing numerical oscillations. In order to have a TVD scheme, it is possible to introduce some numerical diffusion in (11) by adding a parameter σ to the remeshing formulas (10)-(12). This new formula (14) is called \bar{M}_3 , in reference to the M_3 formula obtained when $\sigma = 1/8$ and used in traditional Particle In Cell methods. The interpolation kernel \bar{M}_3 preserves the two first momenta and thus provides order one accuracy. The value of σ will be evaluated in section 4.2 to ensure the TVD property of the \bar{M}_3 scheme.

$$\begin{cases} \alpha^{M3}(y_j) &= y_j(y_j - 1)/2 + \sigma, & \alpha^{M3'}(y_j) &= \alpha^{M3}(y_j - 1) \\ \beta^{M3}(y_j) &= 1 - y_j^2 - 2\sigma, & \beta^{M3'}(y_j) &= \beta^{M3}(y_j - 1) \\ \gamma^{M3}(y_j) &= y_j(y_j + 1)/2 + \sigma, & \gamma^{M3'}(y_j) &= \gamma^{M3}(y_j - 1). \end{cases} \quad (14)$$

A TVD remeshing formula of order two in smooth regions can be built introducing a limiter ϕ to combine the TVD formula \bar{M}_3 (obtained when $\phi = 0$) and the second order Λ_2 formula ($\phi = 1$). The weights (10) and (12) must be replaced respectively by (15) if $0 < \lambda\tilde{g} < 1/2$ and (16) if $-1/2 < \lambda\tilde{g} < 0$.

$$\begin{cases} \alpha^{TVD}(y_j, \phi) &= y_j(y_j - 1)/2 + \sigma(1 - \phi_{j-1/2}) \\ \beta^{TVD}(y_j, \phi) &= 1 - y_j^2 - \sigma(1 - \phi_{j-1/2}) - \sigma(1 - \phi_{j+1/2}) \\ \gamma^{TVD}(y_j, \phi) &= y_j(y_j + 1)/2 + \sigma(1 - \phi_{j+1/2}), \end{cases} \quad (15)$$

$$\begin{cases} \alpha^{TVD'}(y_j, \bar{\phi}) &= \alpha^{TVD}(y_j - 1, \bar{\phi}) \\ \beta^{TVD'}(y_j, \bar{\phi}) &= \beta^{TVD}(y_j - 1, \bar{\phi}) \\ \gamma^{TVD'}(y_j, \bar{\phi}) &= \gamma^{TVD}(y_j - 1, \bar{\phi}). \end{cases} \quad (16)$$

The limiters $\phi_{j\pm 1/2} = \phi(r_{j\pm 1/2})$ with $r_{j+1/2} = (u_j - u_{j-1}) / (u_{j+1} - u_j)$, $r_{j-1/2} = (u_{j-1} - u_{j-2}) / (u_j - u_{j-1})$ and $\bar{\phi}_{j\pm 1/2} = \phi(\bar{r}_{j\pm 1/2})$ with $\bar{r}_{j+1/2} = (u_{j+2} - u_{j+1}) / (u_{j+1} - u_j)$, $\bar{r}_{j-1/2} = (u_{j+1} - u_j) / (u_j - u_{j-1})$ will be calculated in section 4.3 to ensure the TVD property of the scheme for all grid points x_j .

In the case of a velocity changing sign, we must take care about the limiters. $\phi(r) \neq \phi(\bar{r})$ so the scheme in some grid points is different of (11) and (13). To overcome this difficulty it is possible to replace formulas (15)-(16) to remesh some specific particles. More precisely, if the particle $j-1$ has a positive velocity and j has a negative velocity, this particle must be remeshed by the formulas

$$\begin{cases} \alpha_j^{TVD'} &= (y_j - 1)((y_j - 1) - 1)/2 + \sigma(1 - \phi(\mathbf{r}_{j-1/2})) \\ \beta_j^{TVD'} &= 1 - (y_j - 1)^2 - \sigma(1 - \phi(\mathbf{r}_{j-1/2})) - \sigma(1 - \phi(\bar{\mathbf{r}}_{j+1/2})) \\ \gamma_j^{TVD'} &= (y_j - 1)((y_j - 1) + 1)/2 + \sigma(1 - \phi(\bar{\mathbf{r}}_{j+1/2})). \end{cases} \quad (17)$$

In the same way, if $j-1$ has a negative velocity and j is a particle with positive velocity, it must be remeshed with

$$\begin{cases} \alpha_j^{TVD} &= y_j(y_j - 1)/2 + \sigma(1 - \phi(\bar{\mathbf{r}}_{j-1/2})) \\ \beta_j^{TVD} &= 1 - y_j^2 - \sigma(1 - \phi(\bar{\mathbf{r}}_{j-1/2})) - \sigma(1 - \phi(\mathbf{r}_{j+1/2})) \\ \gamma_j^{TVD} &= y_j(y_j + 1)/2 + \sigma(1 - \phi(\mathbf{r}_{j+1/2})). \end{cases} \quad (18)$$

Remark:

The TVD remeshing formulas (15)-(16)-(17)-(18) are still consistent with an order one accuracy at least, and conservative since the sum of the weights used to remesh any particle is one.

4.2 The TVD scheme \bar{M}_3

We give the proof of the TVD property of the \bar{M}_3 scheme discussing on the value of the parameter σ . This parameter can be viewed as an artificial viscosity since when $\sigma = 0$ the \bar{M}_3 scheme reduced to the Λ_2 scheme, and to the M_3 scheme when $\sigma = 1/8$. Let f and h be the two following functions

$$\begin{aligned} f(u, g(u)) &= \lambda g u (\lambda g + 1) \\ h(u, g(u)) &= \lambda g u (\lambda g - 1). \end{aligned} \quad (19)$$

Let us define $\Delta f_{j+1/2} = f_{j+1} - f_j$, $\Delta f_{j-1/2} = f_j - f_{j-1}$, and the same notation hold replacing f by h or u . If the particles are remeshed with the formulas (15)-(16), the scheme at the grid point x_j is

$$u_j^{n+1} = u_j^n + C_{j+1/2} (u_{j+1}^n - u_j^n) - D_{j-1/2} (u_j^n - u_{j-1}^n), \quad (20)$$

with

$$\begin{cases} C_{j+1/2} = \sigma + \frac{1}{2} \frac{\Delta h_{j+1/2}}{\Delta u_{j+1/2}} \\ D_{j-1/2} = \sigma + \frac{1}{2} \frac{\Delta f_{j-1/2}}{\Delta u_{j-1/2}} + \sigma \left(\frac{\phi_{j+1/2}}{r_{j+1/2}} - \phi_{j-1/2} \right), \end{cases} \quad (21)$$

and following the Taylor-Lagrange formula, there exists $\tilde{u} \in]u_{j-1}, u_j[$ and $\tilde{u} \in]u_{j-1}, u_j[$ such as

$$\begin{cases} C_{j-1/2} = \sigma + \frac{1}{2} h'(\tilde{u}) \\ D_{j-1/2} = \sigma + \frac{1}{2} f'(\tilde{u}) + \sigma \left(\frac{\phi_{j+1/2}}{r_{j+1/2}} - \phi_{j-1/2} \right). \end{cases} \quad (22)$$

The scheme \bar{M}_3 is reached doing $\phi = 0$ in (22). Following the Harten theorem [12], the scheme (20) will be TVD if

$$\begin{cases} C_{j-1/2} \geq 0 \\ 0 \leq D_{j-1/2} \leq 1 - C_{j-1/2}. \end{cases} \quad (23)$$

Or if forall u

$$\begin{cases} \sigma + \frac{1}{2} h'(u) \geq 0 \\ \sigma + \frac{1}{2} f'(u) \geq 0 \\ 2\sigma + \frac{1}{2} f'(u) + \frac{1}{2} h'(u) \leq 1, \end{cases} \quad (24)$$

wich is written again

$$\begin{cases} \sigma + \frac{1}{2} \lambda g (\lambda g - 1) + \frac{1}{2} g'(u) \lambda u (2\lambda g - 1) \geq 0 \\ \sigma + \frac{1}{2} \lambda g (\lambda g + 1) + \frac{1}{2} g'(u) \lambda u (2\lambda g + 1) \geq 0 \\ 2\sigma + \lambda^2 g^2 + 2g'(u) \lambda^2 g u \leq 1. \end{cases} \quad (25)$$

Assuming

$$\begin{cases} |\lambda g| \leq C \\ |\lambda g'| \leq C', \end{cases} \quad (26)$$

$$\begin{aligned} & \frac{1}{2} |f'| = \frac{1}{2} |h'| \\ & \leq \frac{1}{2} |\lambda g|^2 + \frac{1}{2} |\lambda g| + \frac{1}{2} |u| |\lambda g'| + |\lambda g| |\lambda g'| |u| \\ & \leq \frac{1}{2} C (C + 1) + \frac{1}{2} C' \max |u| (1 + 2C) \\ & \leq \sigma \end{aligned} \quad (27)$$

if C and $C' \max |u|$ are small enough. So,

$$\begin{cases} \frac{1}{2} |f'| \leq \sigma \\ \frac{1}{2} |h'| \leq \sigma, \end{cases} \quad (28)$$

and chosing $\sigma = 1/4$,

$$\begin{cases} \frac{1}{2} (|f'| + |h'|) \leq 2\sigma = 1/2 \\ \frac{1}{2} (|f'| + |h'|) \leq 1 - 2\sigma = 1/2, \end{cases} \quad (29)$$

the scheme \bar{M}_3 is TVD. This proof holds in the general case, but requires some constraint on the velocity field and his derivative: (26) with C and $C' \max |u|$ small enough. When the velocity field is given, an optimal value of σ can be determined with the CFL condition corresponding. We study the case of the Burgers and the Euler equations.

4.2.1 Burgers equation

Let us consider the Burgers equation, ie $g(u) = u/2$. Since $g'(u) = 1/2$ there exists two constants C and C' satisfying (26) under a CFL condition. Knowing the velocity g it is possible to find an optimal parameter σ allowing the biggest CFL condition to have the scheme \bar{M}_3 TVD.

Indeed, the conditions (25) becomes

$$\begin{cases} \sigma \geq \frac{1}{2}\lambda u - \frac{3}{8}(\lambda u)^2 \\ \sigma \geq -\frac{1}{2}\lambda u - \frac{3}{8}(\lambda u)^2 \\ \sigma \leq \frac{1}{2} - \frac{3}{8}(\lambda u)^2. \end{cases} \quad (30)$$

Setting $c = \lambda \left(\frac{u}{2}\right)$ and according to the two first inequalities of (30), $\sigma \geq 1/6$ if $|c| \geq \frac{1}{3}$. According to the last condition, CFL condition must be $\max|c| \leq \sqrt{2/3} \simeq 0.47$ to have $\sigma \leq 1/6$, and then we chose $\sigma = 1/6$.

Remark:

If $\sigma = 1/4$ the CFL condition is reduced to $1/\sqrt{6} = 0.41$ and the \bar{M}_3 formula is more diffusive than with $\sigma = 1/6$.

4.2.2 Euler equations

Let be $u_1 = \rho$, $u_2 = \rho u$ and $u_3 = \rho E$. Then, the Euler equations are written

$$\begin{cases} \frac{\partial u_1}{\partial t} + \frac{\partial}{\partial x} \left(u_1 \frac{u_2}{u_1} \right) = 0 \\ \frac{\partial u_2}{\partial t} + \frac{\partial}{\partial x} \left(u_2 \frac{u_2}{u_1} \right) = -\frac{\partial p}{\partial x} \\ \frac{\partial u_3}{\partial t} + \frac{\partial}{\partial x} \left(u_3 \frac{u_2}{u_1} \right) = -\frac{\partial}{\partial x} \left(p \frac{u_2}{u_1} \right) \\ p = (\gamma - 1) \left(u_3 - \frac{1}{2} \frac{u_2^2}{u_1} \right). \end{cases} \quad (31)$$

This equations are solved using a splitting method of two steps. The pressure effects are solved in a Lagrange step and the convective part is solved in an advection step [29]. We are interested in this last step wich consists to advect particles to the velocity $g(u_1, u_2, u_3) = u_2/u_1$ and remesh them on a grid with TVD formulas. In this paragraph, we find optimal parameters σ to construct TVD formulas \bar{M}_3 allowing the biggest CFL condition. The limiters used in the remeshing formulas (15)-(16) will be constructed in the section 4.3.

Since the optimal parameter depends on the amount u carrying by the particles, the calculation must be done for the three equations of the system (31). Let us consider the first equation. $u = u_1$, $g = u_2/u_1$ and $g' = -u_2/u_1^2$, so with $c = \lambda u_2/u_1$ (25) becomes

$$\begin{cases} \sigma - \frac{1}{2}c^2 \geq 0 \\ \sigma - \frac{1}{2}c^2 \geq 0 \\ 2\sigma - c^2 \leq 1. \end{cases} \quad (32)$$

Since to ensure the consistency of the Λ_2 remeshing [19] $-1/2 \leq c \leq 1/2$, the CFL condition is $\mathbf{max}|c| \leq 1/2$ and $\sigma = 1/8$.

In the same way, for the second equation of (31), $u = u_2$, $g = u_2/u_1$, $g' = 1/u_1$ and (25) becomes

$$\begin{cases} \sigma + \frac{3}{2}c^2 - c \geq 0 \\ \sigma + \frac{3}{2}c^2 + c \geq 0 \\ 2\sigma + 3c^2 \leq 1. \end{cases} \quad (33)$$

As for Burgers equation, the optimal value for the parameter σ is $1/6$ and the CFL condition is $\mathbf{max}|c| \leq \sqrt{2}/3$.

Finally, for the energy equation, $u = u_3$, $g = u_2/u_1$ and $g' = 0$. So (25) becomes

$$\begin{cases} \sigma + \frac{1}{2}c^2 - \frac{1}{2}c \geq 0 \\ \sigma + \frac{1}{2}c^2 + \frac{1}{2}c \geq 0 \\ 2\sigma + c^2 \leq 1, \end{cases} \quad (34)$$

and the optimal parameter is $\sigma = 1/8$ allowing a CFL condition of $\sqrt{3}/2$ which is reduced to $1/2$ for the consistency.

We want to use the same parameter σ to remesh ρ , ρu or ρE with the \bar{M}_3 formula. So we chose the smallest value of sigma, ie $\sigma = 1/6$, and the CFL condition is $\mathbf{max}|c| \leq \sqrt{2}/3$.

4.3 Calculation of the limiter $\phi(r)$

Let us assume the velocity $g \geq 0$. The limiters are built from (20)-(22) using again the Harten theorem [12]. Assuming that the parameter σ is chosen to have a \bar{M}_3 scheme TVD as explained in the section 4.2, then the coefficient $C_{j-1/2}$ given in (21) is positive from the conditions (23). So the scheme (20) obtained using the Λ_2 -TVD remeshing formulas (15)-(16) will be TVD if

$$0 \leq D_{j-1/2} \leq 1 - C_{j-1/2} \quad \forall u. \quad (35)$$

This condition is written again

$$0 \leq 1 + \frac{1}{2\sigma} f'(u) - \phi_{j-1/2} + \frac{\phi_{j+1/2}}{r_{j+1/2}} \leq \frac{1}{\sigma} - 1 - \frac{1}{2\sigma} h'(u), \quad (36)$$

which is verified if the limiters satisfy

$$\begin{cases} 0 \leq \phi_{j-1/2} \leq 1 + \frac{1}{2\sigma} f'(u) \\ \phantom{0 \leq \phi_{j-1/2}} = 1 + \frac{1}{2\sigma} \lambda^2 g^2 + \frac{1}{\sigma} \lambda^2 g g' u + \frac{1}{2\sigma} \lambda g + \frac{1}{2\sigma} \lambda g' u \\ 0 \leq \phi_{j+1/2}/r_{j+1/2} \leq \frac{1}{\sigma} - 1 - \frac{1}{2\sigma} h'(u) \\ \phantom{0 \leq \phi_{j+1/2}/r_{j+1/2}} = \frac{1}{\sigma} - 2 - \frac{1}{\sigma} \lambda^2 g^2 - \frac{2}{\sigma} \lambda^2 g g' u. \end{cases} \quad (37)$$

If the velocity $g \leq 0$, the coefficients $C_{j+1/2}$ and $D_{j-1/2}$ can be chosen in order to introduce the slope ratio $\bar{r}_{j-1/2} = \Delta u_{j+1/2}/\Delta u_{j-1/2}$:

$$\begin{cases} C_{j+1/2} = \sigma + \frac{1}{2} \frac{\Delta h_{j+1/2}}{\Delta u_{j+1/2}} + \sigma \left(\frac{\phi_{j-1/2}}{\bar{r}_{j-1/2}} - \phi_{j+1/2} \right) \\ D_{j-1/2} = \sigma + \frac{1}{2} \frac{\Delta f_{j-1/2}}{\Delta u_{j-1/2}}. \end{cases} \quad (38)$$

The limiters are built by imposing $0 \leq C_{j+1/2} \leq 1 - D_{j+1/2}$,

$$\begin{cases} 0 \leq \phi_{j+1/2} \leq 1 + \frac{1}{2\sigma} \lambda^2 g^2 + \frac{1}{\sigma} \lambda^2 g g' u - \frac{1}{2\sigma} \lambda g - \frac{1}{2\sigma} \lambda g' u \\ 0 \leq \phi_{j-1/2}/\bar{r}_{j-1/2} \leq \frac{1}{\sigma} - 2 - \frac{1}{\sigma} \lambda^2 g^2 - \frac{2}{\sigma} \lambda^2 g g' u. \end{cases} \quad (39)$$

In the case of a velocity changing sign, particles are remeshed by the formulas (17) and (18). Then the scheme obtained on the grid point x_j is

$$u_j^{n+1} = \begin{aligned} & u_j^n + \frac{1}{2}(h_{j+1} - h_j) + \sigma(u_{j+1}^n - u_j^n) \\ & - \frac{1}{2}(f_j - f_{j-1}) - \sigma(u_j^n - u_{j-1}^n) \\ & - \sigma[\phi(r_{j+1/2}) (u_{j+1}^n - u_j^n) - \phi(r_{j-1/2}) (u_j^n - u_{j-1}^n)]. \end{aligned} \quad (40)$$

This scheme will be TVD if

$$\begin{cases} \sigma + \frac{1}{2}f' - \sigma\phi_{j-1/2}(r) \geq 0 \\ \sigma + \frac{1}{2}h' - \sigma\phi_{j+1/2}(\bar{r}) \geq 0 \\ 2\sigma + \frac{1}{2}(f' + h') - \sigma(\phi_{j+1/2}(\bar{r}) + \phi_{j+1/2}(r)) \leq 1. \end{cases} \quad (41)$$

The first equation is satisfied since (37) hold and the second also with (39). The last is also satisfied since following the inequality (35),

$$2\sigma + \frac{1}{2}(f' + h') - \sigma\phi(\bar{r}) \leq 1 - \sigma\frac{\phi(\bar{r})}{\bar{r}} \leq 1 + \sigma\phi(r), \quad (42)$$

since

$$-\frac{\phi(\bar{r})}{\bar{r}} \leq 0 \leq \phi(r). \quad (43)$$

So, the scheme obtained in the case of a velocity changing sign is TVD if (37) and (39) hold.

4.3.1 Examples

As in the section 4.2, let us look at the case of the Burgers equation and the Euler equations. For the Burgers equation, $g(u) = u/2$, $g'(u) = 1/2$, and $\sigma = 1/6$, so always with $c = \lambda g$ the equations (37) and (39) are written:

$$\begin{cases} 0 \leq \phi \leq 1 \leq 1 + 9c^2 + 6|c| \\ 0 \leq \phi \leq r(4 - 18c^2). \end{cases} \quad (44)$$

Then the limiter is built in order to be as large as possible:

$$\phi(r) = \max\{0, \min[1 + 9 \max c^2 + 6 \max |c|, (4 - 18 \max c^2) r]\}, \quad (45)$$

or

$$\phi(r) = \max\{0, \min[1, (4 - 18 \max c^2) r]\}, \quad (46)$$

and $r = \bar{r}$ if the velocity g of the remeshed particle is negative.

For the Euler equations, $\sigma = 1/6$. So, for the first equation (mass conservation),(37)-(39) becomes

$$\begin{cases} 0 \leq \phi \leq 1 - \frac{1}{2\sigma}c^2 = 1 - 3c^2 \\ 0 \leq \phi \leq r(\frac{1}{\sigma} - 2) = 4r \leq r(\frac{1}{\sigma}(1 + c^2) - 2). \end{cases} \quad (47)$$

For the second Euler equation (momentum conservation),(37)-(39) becomes

$$\begin{cases} 0 \leq \phi \leq 1 \leq 1 + \frac{3}{2\sigma}c^2 + \frac{1}{\sigma}c = 1 + 9c^2 + 6c, \quad c \geq 0 \\ 0 \leq \phi \leq 1 \leq 1 + \frac{3}{2\sigma}c^2 - \frac{1}{\sigma}c = 1 + 9c^2 - 6c, \quad c \leq 0 \\ 0 \leq \phi \leq r(\frac{1}{\sigma} - 2) - \frac{3}{\sigma}c^2 = r(4 - 18c^2), \end{cases} \quad (48)$$

and for the last equation (energy conservation),(37)-(39) becomes

$$\begin{cases} 0 \leq \phi \leq 1 \leq 1 + \frac{1}{2\sigma}c^2 + \frac{1}{2\sigma}c = 1 + 3c^2 + 3c, & c \geq 0 \\ 0 \leq \phi \leq 1 \leq 1 + \frac{1}{2\sigma}c^2 - \frac{1}{2\sigma}c = 1 + 3c^2 - 3c, & c \leq 0 \\ 0 \leq \phi \leq r \left(\frac{1}{\sigma} (1 - c^2) - 2 \right) = r (4 - 6c^2). \end{cases} \quad (49)$$

We have chosen to remesh the variables ρ , ρu and ρE by the same formula. To do this, the same limiter must be used in (15)-(16), so it must satisfy in the same time inequalities (47), (48) and (49). Setting

$$r_{j+1/2}^{(1)} = \frac{\Delta \rho_{j-1/2}}{\Delta \rho_{j+1/2}}, \quad r_{j+1/2}^{(2)} = \frac{\Delta (\rho u)_{j-1/2}}{\Delta (\rho u)_{j+1/2}}, \quad r_{j+1/2}^{(3)} = \frac{\Delta (\rho E)_{j-1/2}}{\Delta (\rho E)_{j+1/2}}, \quad (50)$$

and

$$\bar{r}_{j-1/2}^{(1)} = \frac{\Delta \rho_{j+1/2}}{\Delta \rho_{j-1/2}}, \quad \bar{r}_{j-1/2}^{(2)} = \frac{\Delta (\rho u)_{j+1/2}}{\Delta (\rho u)_{j-1/2}}, \quad \bar{r}_{j-1/2}^{(3)} = \frac{\Delta (\rho E)_{j+1/2}}{\Delta (\rho E)_{j-1/2}}, \quad (51)$$

the limiter is built like this

$$\phi \left(r^{(1)}, r^{(2)}, r^{(3)} \right) = \max \left\{ 0, \min \left[1 - 3 \max c^2, 4r^{(1)}, (4 - 18 \max c^2) r^{(2)}, (4 - 6 \max c^2) r^{(3)} \right] \right\}, \quad (52)$$

replacing r with \bar{r} for negative velocities.

4.4 Numerical validation: Sod shock tube

In this section we present numerical results for the TVD particle methods applied to the 1D Euler equations. We address the classical test case ([28]) of a shock tube initially made up two compartments, each containing a perfect gas ($\gamma = 1.4$). The initial conditions are the following:

$$\begin{aligned} \vec{U}(x, 0) &= 0 \\ \rho(x, 0) &= 1 \text{ if } 0 \leq x \leq 0.5 \\ &0.125 \text{ if } 0.5 < x \leq 1 \\ P(x, 0) &= 2.5 (\gamma - 1) \text{ if } 0 \leq x \leq 0.5 \\ &0.25 (\gamma - 1) \text{ if } 0.5 < x \leq 1 \end{aligned}$$

We solve the Euler equations using a splitting between the Lagrangian step, where are taken into account pressure effects, and the advection step, during which the particles are moved and remeshed. The Lagrangian step is solved with an approximate Riemann solver (acoustic approximation) with a limited MUSCL reconstruction. On Figure 3 are presented the results at $t = 0.2$ for the density, velocity, pressure and thermal energy. The oscillations near the discontinuities observed in [29] that were created by classical interpolation kernels as Λ_2 , Λ_3 or M'_4 have disappeared.




Figure 3: Sod shock tube, 100 particles, from left to right and from top to bottom: density, velocity, pressure, thermal energy.

5 Convergence of TVD particle methods toward entropy solution

5.1 Results

In this section we address the convergence of TVD particle schemes as defined in previous section toward the unique entropy solution of equation (1). In the remaining of the section we make the following assumptions:

- the initial condition u_0 of equation (1) has its total variation bounded: $TV(u_0) < +\infty$ and is bounded in L^∞ norm.
- the function g is of class $C^1(\mathbb{R})$.

Let us give more details about how the numerical scheme that we study is defined. We denote $x_j = j\Delta x$, $x_{j+1/2} = (j+1/2)\Delta x$ and $t^n = n\Delta t$. We suppose that Δt et Δx are proportional to each other: $\frac{\Delta t}{\Delta x} = \lambda$ with λ a constant. The sequence $(u_j^n)_{n \geq 0, j \in \mathbb{Z}}$ is defined by recurrence by equation (4) and the initial sequence $(u_j^0)_{n \geq 0, j \in \mathbb{Z}}$ is defined by:

$$u_j^0 = \frac{1}{\Delta x} \int_{x_{j-1/2}}^{x_{j+1/2}} u_0(x) dx \quad \forall j \in \mathbb{Z}.$$

Because we have assumed that $TV(u_0) < +\infty$, we have also on the discrete level:

$$\sum_j |u_{j+1}^0 - u_j^0| < +\infty.$$

We define the piecewise constant function:

$$u^{\Delta x}(x, t) = u_j^n \text{ for } (x, t) \in [x_{j+1/2}, x_{j-1/2}] \times [t_n, t_{n+1}).$$

The velocity of the particles is evaluated with second order accuracy, in order that a particle scheme with a second order kernel is really second order. We consider the TVD particle scheme built with the kernels Λ_2 and \bar{M}_3 as presented in section 4. If we name $G_{j+1/2}^2$ the flux of the Λ_2 kernel, and $\bar{G}_{j+1/2}^3$ the flux of the \bar{M}_3 kernel, the TVD particle scheme can be expressed as:

$$u_j^{n+1} = u_j - \lambda [(1 - \phi_{j+1/2}) G_{j+1/2}^2 + \phi_{j+1/2} \bar{G}_{j+1/2}^3 - (1 - \phi_{j-1/2}) G_{j-1/2}^2 - \phi_{j-1/2} \bar{G}_{j-1/2}^3].$$

With the expression of the flux obtained in section 3 this can be re-written:

$$\begin{aligned} u_j^{n+1} = & u_j - \sum_{k=1}^d \sum_{i=0}^N \frac{\lambda^i}{i!} \left[(1 - \phi_{j+1/2}) \left(-\Lambda_2^{(i)}(k) \sum_{a=1}^k u_{j+a} \tilde{g}_{j+a}^i + \Lambda_2^{(i)}(-k) \sum_{a=-k+1}^0 u_{j+a} \tilde{g}_{j+a}^i \right) \right. \\ & + \phi_{j+1/2} \left(-\bar{M}_3^{(i)}(k) \sum_{a=1}^k u_{j+a} \tilde{g}_{j+a}^i + \bar{M}_3^{(i)}(-k) \sum_{a=-k+1}^0 u_{j+a} \tilde{g}_{j+a}^i \right) \\ & - (1 - \phi_{j-1/2}) \left(-\Lambda_2^{(i)}(k) \sum_{a=0}^{k-1} u_{j+a} \tilde{g}_{j+a}^i + \Lambda_2^{(i)}(-k) \sum_{a=-k}^{-1} u_{j+a} \tilde{g}_{j+a}^i \right) \\ & \left. - \phi_{j-1/2} \left(-\bar{M}_3^{(i)}(k) \sum_{a=0}^{k-1} u_{j+a} \tilde{g}_{j+a}^i + \bar{M}_3^{(i)}(-k) \sum_{a=-k}^{-1} u_{j+a} \tilde{g}_{j+a}^i \right) \right]. \end{aligned}$$

Therefore we can write for the expression of such a TVD particle scheme:

$$u_j^{n+1} = \sum_{k=-d}^d u_{j+k}^n \bar{\Lambda}(k, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_{j+k}) \quad (53)$$

The remeshing weights depend of the values of the other neighbouring particles, through the limiting function ϕ . The two interpolation kernels used in this non-linear combination can in fact be any interpolation kernel preserving the two first moments and giving TVD remeshing formulas, and any other more accurate interpolation kernel, as long as the combination of their fluxes is TVD and consistent.

Proposition 5 *The particle scheme defined by (53), built as a TVD combination of the kernels Λ_2 and \bar{M}_3 as described in 4: and satisfying the following CFL condition:*

$$\forall k, \forall n, \lambda |\tilde{g}(u_k^n)| < 1/2 \text{ and } \lambda |g(u_k^n)| < 1/2.$$

converges in L_{loc}^1 norm to the unique entropic solution of (1).

5.2 Proof of Proposition 5

We firstly need to prove that the TVD particle scheme is stable in $L^\infty(\mathbb{R}^+ \times \mathbb{R})$ norm. In that purpose we assume that the particle scheme is not stable in this

norm and show that it leads to a contradiction. Without a loss of generality one can assume for example that, $\forall M > 0, \exists n_0, j_0$ such that $u_{j_0}^{n_0} > M$. Because the particle scheme can be written in conservative form, we have:

$$\Delta x \sum_j u_j^{n+1} = \Delta x \sum_k u_k^0.$$

There exists necessarily an index j_1 such that $u_{j_1}^{n_0} < \frac{M}{2}$. Thus:

$$\sum_j |u_{j+1}^{n_0} - u_j^{n_0}| \geq |u_{j_0}^{n_0} - u_{j_1}^{n_0}| \geq \frac{M}{2}$$

which is not possible if we choose M such that $\frac{M}{2} > \sum_j |u_{j+1}^0 - u_j^0|$. Therefore the TVD particle scheme is bounded in $L^\infty(\mathbb{R}^+ \times \mathbb{R})$ norm, and thus also bounded in $L^1(\Omega)$ for every bounded open set $\Omega \in \mathbb{R}^+ \times \mathbb{R}$.

Now we want to prove that for every bounded open set $\Omega \in \mathbb{R}^+ \times \mathbb{R}$ the total variation of the scheme over Ω is bounded. Cottet and Magni have devised their flux limited particle method in order to have:

$$\sum_j |u_{j+1}^{n+1} - u_j^{n+1}| \leq \sum_j |u_{j+1}^n - u_j^n| \forall n.$$

We use a result in [18], saying that if a conservative scheme with a Lipschitz-continuous flux satisfies the property $\sum_k |u_{j+1}^{n+1} - u_j^{n+1}| \leq \sum_k |u_{j+1}^n - u_j^n|$, then the total variation of the scheme is bounded. The flux $G_{j+1/2}^1 + \phi_{j+1/2}(G_{j+1/2}^2 - G_{j+1/2}^1)$ with $G_{j+1/2}^1$ and $G_{j+1/2}^2$ defined as in subsection 5.1 is locally Lipschitz-continuous, and the scheme is stable in $L^\infty(\mathbb{R}^+ \times \mathbb{R})$ norm. Thus the total variation of the scheme is bounded.

Now we follow the type of reasoning presented in [18]: we consider a sequence of numerical approximations $u^{(\Delta x_i)}$ obtained with the particle scheme with grid parameters Δx_i tending to zero when i tends to infinity. We assume that this sequence does not converge in $L^1(\Omega)$ toward u the entropic solution of (1), and show that this assumption leads to a contradiction. If the particle scheme does not converge toward u , then there exists some $\epsilon > 0$ and a subsequence $u^{(\Delta \tilde{x}_i)}$ such that:

$$\|u^{(\Delta \tilde{x}_i)} - u\|_{L^1(\Omega)} > \epsilon \forall i.$$

Because $u^{(\Delta \tilde{x}_i)}$ is bounded in $L^1(\Omega) \cap TV(\Omega)$, and because of the Helly theorem, one can extract a subsubsequence $u^{(\Delta \tilde{x}_i)}$ that converges in $L^1(\Omega)$. Let us call \bar{u} the limit of $u^{(\Delta \tilde{x}_i)}$. The particle scheme is bounded in $L^\infty(\mathbb{R}^+ \times \mathbb{R})$ norm, can be written in conservative form consistent with the equation (1), thus because of the Lax-Wendroff theorem, $u^{(\Delta \tilde{x}_i)}$ is necessarily a weak solution of (1).

Now we want to prove that \bar{u} satisfies a weak entropic inequality for all Kruzkov entropies, ie: $\forall \phi \in C_0^1(\mathbb{R}^+ \times \mathbb{R}), \phi \geq 0, \forall K \in \mathbb{R}$,

$$\int_0^{+\infty} \int_{-\infty}^{+\infty} \frac{\partial \phi}{\partial t} |\bar{u} - K| + \text{sgn}(\bar{u} - K) \left(g(\bar{u})\bar{u} - g(K)K \right) \frac{\partial \phi}{\partial x} dx dt + \int_{-\infty}^{+\infty} |\bar{u}(0, x) - K| \phi(0, x) dx \geq 0.$$

This will be possible thanks to the form of the particle scheme: the new particle values are expressed as a weighted sum of the old values, unlike finite difference schemes like the Lax-Wendroff scheme. We define:

$$(I) = \sum_{n=0}^{\infty} \sum_j \frac{1}{\Delta t} \left[\phi_j^n |u_j^{n+1} - K| - \phi_j^n \sum_k \operatorname{sgn}(u_k^n - K) [u_k^n \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - K \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K))] \right] \Delta x \Delta t.$$

with $\phi \in C_0^1(\mathbb{R}^+ \times \mathbb{R})$ a positive function whose support is included in $[-D, D] \times [0, T]$. We firstly want to prove that:

$$\lim_{\Delta t \rightarrow 0} (I) = - \int_0^{+\infty} \int_{-\infty}^{+\infty} \phi_t |\bar{u} - K| + \operatorname{sgn}(\bar{u} - K) (g(\bar{u})\bar{u} - g(K)K) \phi_x \, dx dt - \int_{-\infty}^{+\infty} |\bar{u}(0, x) - K| \phi(0, x) \, dx$$

and secondly that:

$$\lim_{\Delta t \rightarrow 0} (I) \leq 0.$$

By exchanging the indices j and k and making a change of variable on k we get:

$$(I) = \sum_{n=0}^{\infty} \sum_j \frac{1}{\Delta t} \left[\phi_j^n |u_j^{n+1} - K| - \operatorname{sgn}(u_j^n - K) u_j^n \sum_k \phi_{j-k}^n \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}_j^n) + \operatorname{sgn}(u_j^n - K) K \sum_k \phi_{j-k}^n \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}(K)) \right] \Delta x \Delta t.$$

With a change of variables for indices n the equation becomes:

$$(I) = \sum_{n=1}^{\infty} \sum_j \frac{1}{\Delta t} \left[\phi_j^{n-1} |u_j^n - K| - \operatorname{sgn}(u_j^n - K) u_j^n \sum_k \phi_{j-k}^n \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}_j^n) + \operatorname{sgn}(u_j^n - K) K \sum_k \phi_{j-k}^n \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}(K)) \right] \Delta x \Delta t - \frac{1}{\Delta t} \sum_j \left[\operatorname{sgn}(u_j^0 - K) u_j^0 \sum_k \phi_{j-k}^0 \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}_j^0) - \operatorname{sgn}(u_j^0 - K) K \sum_k \phi_{j-k}^0 \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}(K)) \right] \Delta x \Delta t.$$

The idea is to recognize in the terms $\sum_k \phi_{j-k}^n \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}_j^n)$ a discrete approximation of $\phi(x_j, t^n) - \Delta t g(u_j^n) \frac{\partial \phi}{\partial x}(x_j, \Delta t)$. The TVD remeshing scheme is obtained by combining the fluxes of a linear interpolation kernel, which gives a first order scheme, and another kernel which provides at least second order interpolation. The resulting scheme is thus at least first order accurate on every grid point. Therefore, we can write as in the proof of Proposition 3 in section 3:

$$\sum_k \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}_j^n) = \phi(x_j, t^n) - \Delta t g(u_j^n) \frac{\partial \phi}{\partial x}(x_j, t^n) + O(\Delta t^2)$$

$$\sum_k \bar{\Lambda}(k, \phi_{j-k-1/2}, \phi_{j-k+1/2}, \lambda \tilde{g}(K)) = \phi(x_j, t^n) - \Delta t g(K) \frac{\partial \phi}{\partial x}(x_j, t^n) + O(\Delta t^2).$$

Therefore when Δt tends to zero:

$$(I) \rightarrow - \int_0^{+\infty} \int_{-\infty}^{+\infty} \phi_t |u - K| + \operatorname{sgn}(u - K)(g(u)u - g(K)K) \phi_x \, dx dt \\ - \int_{-\infty}^{+\infty} |u(0, x) - K| \phi_x(0, x) \, dx.$$

Now we want to prove that $\lim_{\Delta t \rightarrow 0}(I) \leq 0$. By noting Φ the maximum of $|\phi|$ on $[-D, D] \times [0, T]$ we have:

$$(I) \leq \Phi \sum_{n=0}^{T/\Delta t} \Delta t \sum_{j, j\Delta x \in [-D, D]} \left| |u_j^{n+1} - K| - \sum_k \operatorname{sgn}(u_k^n - K) [u_k^n \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \right. \\ \left. - K \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K))] \right| \Delta x.$$

We define:

$$(B^n) = \sum_{j, j\Delta x \in [-D, D]} \left| |u_j^{n+1} - K| - \sum_k \operatorname{sgn}(u_k^n - K) [u_k^n \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \right. \\ \left. - K \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K))] \right| \Delta x.$$

We want to prove that (B^n) tends to zero when Δx tends to zero. Because of the definition of the particle scheme (53):

$$(B^n) = \sum_{j, j\Delta x \in [-D, D]} \left| \left| \sum_k u_k^n \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - K \right| \right. \\ \left. - \sum_k \operatorname{sgn}(u_k^n - K) [u_k^n \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - K \bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K))] \right| \Delta x.$$

The kernels involved in the formula of Λ have a compact support, and we have assumed that a CFL condition was satisfied. So there exists a real S such that:

$$\bar{\Lambda}(k - j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k) = 0$$

if $k - j > S$. Let $\eta > 0$ be a real such that $S \Delta x < \eta \forall \Delta x$. Because u is locally bounded, for all $\epsilon > 0$ there exists $u_\epsilon \in C^1([-D - \eta, D + \eta] \times [0, T])$ such that

$$\|u(t, \cdot) - u_\epsilon(t, \cdot)\|_{L^1([-D - \eta, D + \eta])} \leq \epsilon$$

. We split (B^n) into several terms:

$$\begin{aligned}
 (B^n) &\leq \underbrace{\sum_{j,j\Delta x \in [-D,D]} \left| \sum_k (u_k^n - u_\epsilon(x_k, t^n)) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \right| \Delta x}_{(a)} \\
 &+ \underbrace{\sum_{j,j\Delta x \in [-D,D]} \left| \sum_k u_\epsilon(x_k, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - u_\epsilon(x_j, t^n) \right| \Delta x}_{(b)} \\
 &+ \underbrace{\sum_{j,j\Delta x \in [-D,D]} \left| \text{sgn}(u_\epsilon(x_j, t^n) - K)(u_\epsilon(x_j, t^n) - K) \Delta x \right.}_{(c)} \\
 &\quad \left. - \sum_k \text{sgn}(u_\epsilon(x_k, t^n) - K) \left(u_\epsilon(x_k, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \right. \right. \\
 &\quad \left. \left. - K \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K)) \right) \right|}_{(c)} \\
 &+ \underbrace{\sum_{j,j\Delta x \in [-D,D]} \left| \sum_k \text{sgn}(u_\epsilon(x_k, t^n) - K) \left[u_\epsilon(x_k, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \right. \right.}_{(d)} \\
 &\quad \left. \left. - K \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K)) \right] \right|}_{(d)} \\
 &\quad \left. - \text{sgn}(u_k^n - K) \left[u_k^n \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - K \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(K)) \right] \right| \Delta x}_{(d)}.
 \end{aligned}$$

$$\begin{aligned}
 (a) &\leq \sum_{j,j\Delta x \in [-D,D]} \sum_k |u_k^n - u_\epsilon(x_k, t^n)| |\bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n)| \Delta x \\
 &\leq \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} |u_k^n - u_\epsilon(x_k, t^n)| \sum_{j, |j-k| \leq S} |\bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n)| \Delta x \\
 &\leq 2S \|\bar{\Lambda}\|_\infty \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} |u_k^n - u_\epsilon(x_k, t^n)| \Delta x \\
 &\leq 2S \|\bar{\Lambda}\|_\infty \int_{-D-(S+1/2)\Delta x}^{D+(S+1/2)\Delta x} |u^{\Delta x}(x, t) - u_\epsilon(x, t)| dx \\
 &\quad + 2S \|\bar{\Lambda}\|_\infty \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \left| \int_{(k-1/2)\Delta x}^{(k+1/2)\Delta x} u_\epsilon(x, t) - u_\epsilon(x_k, t^n) dx \right|.
 \end{aligned}$$

Thus (a) tends to zero when Δx tend to zero. One can prove similarly that (d) tends to zero. Now the term (b):

$$\begin{aligned}
 (b) &= \sum_{j,j\Delta x \in [-D,D]} \left| \sum_k u_\epsilon(x_k, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - \sum_k u_\epsilon(x_j, t^n) \Lambda_{j,k}(k-j + \lambda \tilde{g}_k) \right. \\
 &\quad \left. + \sum_k u_\epsilon(x_j, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - u_\epsilon(x_j, t^n) \underbrace{\sum_k \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_j^n)}_{=1} \right| \Delta x
 \end{aligned}$$

The term $\sum_k \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_j^n)$ is equal to 1 because if we consider the TVD remeshing formula (53) with $u_k^n = u \forall k$:

$$u_j^{n+1} = \sum_k u \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u))$$

we have, due to the property of consistency of the fluxes: $u_j^{n+1} = u$. Thus

$$\sum_k \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u)) = 1.$$

$$\begin{aligned}
 (b) &\leq \underbrace{\sum_{j,j\Delta x \in [-D,D]} \sum_k \left| u_\epsilon(x_k, t^n) - u_\epsilon(x_j, t^n) \right| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n)}_{(1)} \Delta x \\
 &\quad + \underbrace{\sum_{j,j\Delta x \in [-D,D]} \left| \sum_k u_\epsilon(x_j, t^n) \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_j^n) \right|}_{(2)} \Delta x.
 \end{aligned}$$

$$(1) \leq \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \sum_{j,|j-k| \leq S} \left| u_\epsilon(x_k, t^n) - u_\epsilon(x_j, t^n) \right| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) \Delta x.$$

The function u_ϵ belongs to $C^1([-D-\eta, D+\eta] \times [0, T])$, thus there exists a real K such that:

$$\forall x, y \in [-D-\eta, D+\eta] \quad |u_\epsilon(x, t) - u_\epsilon(y, t)| \leq K|x-y|.$$

$$\begin{aligned}
 (1) &\leq \|\bar{\Lambda}\|_\infty \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \sum_{j,|j-k| \leq S} K|k-j| \Delta x^2 \\
 &\leq 2S^2 \|\bar{\Lambda}\|_\infty K \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \Delta x^2 \leq 4S^2 \|\bar{\Lambda}\|_\infty K \left(\frac{D}{\Delta x} + S \right) \Delta x^2.
 \end{aligned}$$

Therefore (1) tends to zero when Δt tends to zero.

$$\begin{aligned}
 (2) &\leq \|u_\epsilon\|_\infty \sum_{j,j\Delta x \in [-D,D]} \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \\
 &\quad \left| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k^n) - \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_j^n) \right| \Delta x \\
 &\leq \|u_\epsilon\|_\infty \sum_{k,k\Delta x \in [-D-S\Delta x, D+S\Delta x]} \sum_{j,|j-k| \leq S} \\
 &\quad \left| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_k) - \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u_\epsilon(x_k, t^n))) \right| \Delta x \\
 &\quad + \left| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u_\epsilon(x_k, t^n))) - \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u_\epsilon(x_j, t^n))) \right| \Delta x \\
 &\quad + \left| \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}(u_\epsilon(x_j, t^n))) - \bar{\Lambda}(k-j, \phi_{j-1/2}, \phi_{j+1/2}, \lambda \tilde{g}_j) \right| \Delta x.
 \end{aligned}$$

We have assumed that:

$$\forall k, \forall n, \lambda |\tilde{g}(u_k^n)| < 1/2 \text{ and } \lambda |g(u_k^n)| < 1/2.$$

$\bar{\Lambda}$ is of class C^1 . The function g is also of class $C^1(\mathbb{R})$. We have assumed that the scheme converges in L^1_{loc} when Δx and Δt tend to zero. We conclude that (2), thus (b), tend to zero when Δx and Δt tend to zero. With the same kind of reasoning we could also prove that (c) tends to zero.

We conclude that the limit \bar{u} satisfies $\forall \phi \in C_0^1(\mathbb{R}^+ \times \mathbb{R}), \forall K \in \mathbb{R}$:

$$- \int_0^{+\infty} \int_{-\infty}^{+\infty} \phi_t |u - K| + \text{sgn}(u - K)(g(u)u - g(K)K) \phi_x \, dx dt - \int_{-\infty}^{+\infty} |u(0, x) - K| \phi_x(0, x) \, dx \geq 0.$$

Therefore \bar{u} is the unique entropic solution of (1), which contradicts the initial assumption. We conclude that the TVD particle scheme (53) converges in $L^1(\Omega)$ to the unique entropic solution of (1).

6 Conclusion

We have studied the consistency and accuracy properties of remeshed particle methods in the case of a scalar one-dimensional conservation law. The accuracy of the particle scheme depends on the accuracy of the interpolation kernel used. In the linear case, if the interpolation kernel preserves the first M moments then the particle scheme is of order $M - 1$. In the non-linear case, the particle scheme is a priori only of order one, because of the first order evaluation of the particle moving, unless a correction of the evaluation of the particle velocities is used. Cottet and Magni [5] introduced recently TVD remeshing schemes for particle-grid methods. We have extended the construction of these new TVD particle schemes to non-linear conservation laws with a possible change of velocity sign, with application to Burgers and Euler equations. Numerical results obtained in the case of the Sod shock tube for the Euler equations have been presented. Then we have proved that with these new TVD remeshing schemes the particle schemes converge toward the entropy solution. The perspectives of this work are the application of the TVD particle schemes to systems of conservation laws, for instance more complex 2D and 3D compressible flows like hydrodynamic instabilities.

References

- [1] W. Benz. The Numerical Modelling of Nonlinear Stellar Pulsations, Problems and Prospects, chapter Smooth Particle Hydrodynamics: A Review, pages 269–287. NATO ASIS Series, 1989.
- [2] C. Berthon. Contribution à l’analyse numérique des équations de Navier-Stokes compressibles à deux entropies spécifiques. Application à la turbulence compressible. PhD thesis, Université Paris VI, 1998.
- [3] M. Coquerelle and G.-H. Cottet. A vortex level set method for the two-way coupling of an incompressible fluid with colliding rigid bodies. J. Comput. Phys., 227:9121–9137, 2008.
- [4] G.-H. Cottet and P. D. Koumoutsakos. Vortex methods. Cambridge University Press, 2000.
- [5] G.-H. Cottet and A. Magni. Tvd remeshing schemes for particle methods. C. R. Acad. Sci. Paris, Ser. I, 347:1367–1372, 2009.
- [6] G.-H. Cottet, B. Michaux, S. Ossia, and G. Vanderlinden. A comparison of spectral and vortex methods in three-dimensional incompressible flow. J. Comput. Phys., 175:702–712, 2002.
- [7] G.-H. Cottet and L. Weynans. Particle methods revisited: a class of high-order finite-difference schemes. C. R. Acad. Sci. Paris, Ser. I, 343:51–56, 2006.
- [8] A. Ghoniem D. Wee. Modified interpolation kernels for treating diffusion and remeshing in vortex methods. J. Comput. Phys., 213:239–263, 2006.
- [9] M. W. Evans and F. H. Harlow. The particle-in-cell method for hydrodynamics calculations. Technical report, Los Alamos Scientific Laboratory, 1956.
- [10] R. A. Gingold and J. J. Monaghan. Smoothed particle hydrodynamics: theory and application to non-spherical stars. Mon. Not. R. astr. Soc., 181:375–389, 1977.
- [11] F. H. Harlow. Hydrodynamic problems involving large fluid distorsion. J. Assoc. Comp. Mach., 4:137–..., 1957.
- [12] A. Harten. High resolution schemes for hyperbolic conservation laws. J. Comput. Phys., 49:357–393, 1983.
- [13] T. Hou and P. G. Lefloch. Why non-conservative schemes converge to wrong solutions: error analysis. Mathematics of Computation, 62:497–530, 1994.
- [14] P. Koumoutsakos and S. Hieber. A lagrangian particle level set method. J. Comp. Phys., 210:342–367, 2005.
- [15] P. Koumoutsakos and A. Leonard. High resolution simulations of the flow around an impulsively started cylinder using vortex methods. J. Fluid Mech., 296:1–38, 1995.

- [16] N. Larson, , and J.P. Vila. Renormalized meshfree schemes ii: convergence for scalar conservation laws. SIAM J. Numerical. Analysis., 46(4).
- [17] N. Larson and J.P. Vila. Convergence des methodes particulaires renormalisees pour les systemes de friedrichs. C. R. Acad. Sci. Paris, Ser I, 349:465 – 470, 2005.
- [18] R. J. Leveque. Finite-volume methods for hyperbolic problems. Cambridge University Press, 2002.
- [19] A. Magni and G.-H. Cottet. Accurate, non-oscillatory, remeshing schemes for particle methods, to appear in. J. Comput. Phys.
- [20] Adrien Magni. Méthodes particulaires avec remaillage : analyse numérique nouveaux schémas et applications pour la simulation d'équations de transport. PhD thesis, Université de Grenoble, <http://tel.archives-ouvertes.fr/tel-00623128/fr/>, 2011.
- [21] A. Majda and S. Osher. Numerical viscosity and the entropy condition. Communications on Pure and Applied Mathematics, XXXII:797–838, 1979.
- [22] J. J. Monaghan. Why particle methods work. SIAM J. Sci. Stat. Comput, 3(4):422–433, 1982.
- [23] J. J. Monaghan. Extrapolating b-splines for interpolation. J. Comput. Phys., 60(2):253–262, 1985.
- [24] J. J. Monaghan. Smoothed particle hydrodynamics. Annu. Rev. Astron. Astrophys., 30:543–..., 1992.
- [25] B. Ben Moussa and J.P. Vila. Convergence of sph methods for scalar non linear conservation laws. SIAM J. Numerical. Analysis., 37(3).
- [26] P. Ploumhans, G. S. Winckelmans, J. K. Salmon, A. Leonard, and M. S. Warren. Vortex methods for direct numerical simulation of three-dimensional bluff body flows: application to the sphere at $re = 300, 500,$ and 1000 . J. Comput. Phys., 178(2):427–463, 2002.
- [27] P. Poncet. Topological aspects of the three-dimensional wake behind rotary oscillating circular cylinder. J. Fluid Mech., 517:27–53, 2004.
- [28] G. A. Sod. A survey of several finite difference methods for systems of non-linear hyperbolic conservation laws. J. Comput. Phys., 27:1–131, 1978.
- [29] L. Weynans. Méthode particulière multi-niveaux pour la dynamique des gaz, application au calcul d'écoulements multifluides. PhD thesis, Université Joseph Fourier, <http://tel.archives-ouvertes.fr/tel-00121346/en/>, 2006.



Centre de recherche INRIA Bordeaux – Sud Ouest
Domaine Universitaire - 351, cours de la Libération - 33405 Talence Cedex (France)

Centre de recherche INRIA Grenoble – Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier
Centre de recherche INRIA Lille – Nord Europe : Parc Scientifique de la Haute Borne - 40, avenue Halley - 59650 Villeneuve d'Ascq
Centre de recherche INRIA Nancy – Grand Est : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex
Centre de recherche INRIA Paris – Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex
Centre de recherche INRIA Rennes – Bretagne Atlantique : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex
Centre de recherche INRIA Saclay – Île-de-France : Parc Orsay Université - ZAC des Vignes : 4, rue Jacques Monod - 91893 Orsay Cedex
Centre de recherche INRIA Sophia Antipolis – Méditerranée : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399