



HAL
open science

Characterization of local quadratic growth for strong minima in the optimal control of semi-linear elliptic equations

Térence Bayen, J. Frederic Bonnans, Francisco J. Silva

► **To cite this version:**

Térence Bayen, J. Frederic Bonnans, Francisco J. Silva. Characterization of local quadratic growth for strong minima in the optimal control of semi-linear elliptic equations. Transactions of the American Mathematical Society, 2014, 366 (4), pp.2063–2087. 10.1090/S0002-9947-2013-05961-2 . inria-00632308v2

HAL Id: inria-00632308

<https://hal.inria.fr/inria-00632308v2>

Submitted on 26 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CHARACTERIZATION OF LOCAL QUADRATIC GROWTH FOR STRONG MINIMA IN THE OPTIMAL CONTROL OF SEMI-LINEAR ELLIPTIC EQUATIONS

T. BAYEN, J.F. BONNANS, AND F.J. SILVA

ABSTRACT. In this article we consider an optimal control problem of a semi-linear elliptic equation, with bound constraints on the control. Our aim is to characterize local quadratic growth for the cost function J in the sense of strong solutions. This means that the function J grows quadratically over all feasible controls whose associated state is close enough to the nominal one, in the uniform topology. The study of strong solutions, classical in the Calculus of Variations, seems to be new in the context of PDE optimization. Our analysis, based on a decomposition result for the variation of the cost, combines Pontryagin's principle and second order conditions. While these two ingredients are known, we use them in such a way that we do not need to assume that the Hessian of Lagrangian of the problem is a Legendre form, or that it is uniformly positive on an extended set of critical directions.

INTRODUCTION

Over the last decades important progress has been done in the field of optimal control of Partial Differential Equations (PDEs). In the case of a semi-linear elliptic equations, we have particularly in mind (i) the extensions of Pontryagin's minimum principle [28], the first papers being due to [29], and then [4, 6, 7], and (ii) the theory of second order optimality conditions for weak minima [3, 12, 13, 14, 15, 16, 18, 19, 31]. By weak minimum we mean that optimality is ensured in a L^∞ - neighborhood in the *control space* (sometimes local optimality can be established in an L^2 space, see [17]).

To be more precise, regarding second order optimality conditions, the authors of this article are aware of two sufficient conditions that imply local quadratic growth for the cost function in the *weak sense*. This means that the cost grows quadratically, with respect to the L^2 -norm, in a feasible L^∞ - neighborhood of the nominal control. Both aforementioned conditions ask that a *weak* form of the Pontryagin's principle is satisfied, while they differ on the second order condition.

Date: October 16, 2011 and, in revised form, July 19, 2012.

2010 Mathematics Subject Classification. Primary 49J20, 49K20; Secondary 35J61.

Key words and phrases. Optimal control of PDE, strong minima, Pontryagin's minimum principle, second order optimality conditions, control constraints, local quadratic growth.

REVISED VERSION, TO APPEAR IN TRANSACTIONS AMER. MATH. SOC. The support of the Chair "Modélisation Mathématique et Simulation Numérique", EADS-Polytechnique-INRIA and of the European Union under the "7th Framework Program FP7-PEOPLE-2010-ITN Grant agreement number 264735-SADCO" are gratefully acknowledged. The first author thanks the CNRS for providing him a research opportunity of one year at the Centro de Modelamiento Matemático (Chile).

In references [3, 12] it is supposed that the Hessian of the Lagrangian of the problem is a Legendre form and that it is uniformly positive over a cone of *critical directions*, which is exactly the cone provided by second order necessary conditions. On the other hand, in references [13, 14, 15, 16, 18, 19, 31] the Legendre form assumption is not needed, but it is required a uniform positivity condition of the Hessian of the Lagrangian over a slightly larger cone than the critical one.

This paper is devoted to the study of *strong solutions* for the optimal control problem of a semi-linear elliptic equation with Dirichlet boundary conditions, under bound constraints on the controls. By strong solutions we mean, as in the classical Calculus of Variations, optimality in a L^∞ neighborhood in the *state space* only. Thus, the neighborhoods are considered in the state space rather than in the control space, which is the setting of *weak solutions* (see Definition 1.6).

The motivation for studying strong solutions is quite the same as in the classical Calculus of Variations. By the definition, a strong solution remains optimal over a much larger set than a weak one. The standard *needle perturbation argument* can be applied for these types of solutions, which implies that the Pontryagin's principle holds true (see [4, 6, 7, 29] and Section 3.1). Moreover, since a strong solution is in particular a weak solution, it satisfies the classical second order necessary optimality condition (see e.g. [18, 3] and Section 3.2). One of the goals of this article is to prove that a sort of converse assertion holds true: If a *strict form* of the Pontryagin principle and the standard second order sufficient condition are verified at a control \bar{u} , then this control is a strong solution.

Let us also mention that Pontryagin's principle, which is a consequence of strong optimality, is used in some numerical algorithms for solving optimal control problems. We refer the reader to [5] in the framework of Ordinary Differential Equations (ODEs) and to [2] for the infinite dimensional case.

In this article, we are particularly interested in providing a characterization of local quadratic growth for the cost function in the *strong sense*. This means that quadratic growth for the cost, with respect to the L^2 -norm, holds over all feasible controls whose associated states are uniformly close to the nominal one. Our main result is Theorem 3.24 which states that local quadratic growth for the cost holds in the strong sense at \bar{u} if and only if the Hessian of Lagrangian of the problem is uniformly positive over the cone of critical directions, and the Hamiltonian satisfies a global quadratic growth property at \bar{u} . Since this is the first result in this direction for the PDE framework, we decided to present it in the rather simple framework of bounds constraints. Important extensions like non-local constraints over the control and state constraints will be addressed in future works.

The proof of Theorem 3.24 relies on the combination of Pontryagin's minimum principle and second order conditions, which is possible thanks to the extension of a decomposition result in [8, Theorem 2.14] to the elliptic framework. Roughly speaking, the decomposition result obtained in Theorem 2.5 and its Corollary 2.6 says that the variation of the cost function under a perturbation of the control can be expressed as the sum of two terms: the first one is the variation due to a *large* perturbation in the L^∞ -norm but with support over a set of small measure, while the second one corresponds to the variation due to a perturbation *small* in the L^∞ -norm. The key elements in the proof of the decomposition result are the well known regularity estimates from the L^s -theory of linear elliptic equations (see e.g. [22]).

The article is organized as follows: in Section 1 we set some useful notations, recall some basics facts about linear and semi-linear elliptic equations, define the optimal control problem (\mathcal{CP}) and set some standard assumptions over the data. Estimates of Section 1 are used in Section 2 to provide first and second order expansions for the state and cost functions. The latter are expressed in terms of an associated Hamiltonian. The novelty of this section is Theorem 2.5, where the decomposition result is proved. Section 3 begins with the statement and proof of an extension of the standard Pontryagin's minimum principle. This result allows us to show that weak local solutions satisfy a *local Pontryagin's minimum principle*. Next, after recalling some well-known facts about necessary conditions for weak solutions, we prove in Theorem 3.17 that if local quadratic growth for the cost holds at \bar{u} in the strong sense, then \bar{u} satisfies a strict Pontryagin inequality and the associated quadratic form, i.e. the Hessian of the Lagrangian of (\mathcal{CP}), is uniformly positive over the cone of critical directions. Regarding sufficient conditions, the decomposition result allows us in Theorems 3.20 and 3.23 to extend to the strong sense, respectively, Theorems [3, Theorem 2.9] and [18, Theorem 2]. Finally, in Theorem 3.24 we characterize the local quadratic growth in the strong sense, i.e. we provide the converse implication of Theorem 3.17. In order to prove the result, we adapt the technique of projection on the pointwise critical cone due to [8, Theorem 5.5]. The article concludes with an appendix containing the proofs of some technical lemmas of Sections 2 and 3.

1. PROBLEM STATEMENT AND PRELIMINARY RESULTS.

We first fix some useful notations. For a function $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ ($d \in \mathbb{N}$), a nominal $\bar{x} \in \mathbb{R}^d$ and a perturbation $z \in \mathbb{R}$, we set $\psi_{x_i}(\bar{x})z := D_{x_i}\psi(\bar{x})z$ for $i = 1, \dots, d$. Analogously, for $z_1, z_2 \in \mathbb{R}$ we use the following convention

$$(1.1) \quad \begin{aligned} \psi_{x_i x_i}(\bar{x})z_1^2 &:= D_{x_i x_i}^2 \psi(\bar{x})(z_1, z_1), & \psi_{x_i x_j}(\bar{x})z_1 z_2 &:= D_{x_i x_j}^2 \psi(\bar{x})(z_1, z_2), \\ \psi_{(x_i, x_j)^2}(\bar{x})(z_1, z_2)^2 &:= \psi_{x_i x_i}(\bar{x})z_1^2 + 2\psi_{x_i x_j}(\bar{x})z_1 z_2 + \psi_{x_j x_j}(\bar{x})z_2^2. \end{aligned}$$

From now on, we fix a non-empty bounded open set $\Omega \subseteq \mathbb{R}^n$ ($n \in \mathbb{N}$) with a $C^{1,1}$ boundary and for $s \in [1, \infty]$, $k \in \mathbb{N}$, we denote by $\|\cdot\|_s$ and $\|\cdot\|_{k,s}$ the standard norms in $L^s(\Omega)$ and $W^{k,s}(\Omega)$, respectively. For any Borel set $A \subseteq \mathbb{R}^n$ we denote by $|A|$ its Lebesgue measure (not to be confused with the same notation for the absolute value in \mathbb{R}) and when a property holds for almost all $x \in \Omega$, with respect to the Lebesgue measure, we use the abbreviation “for a.a. $x \in \Omega$ ”. Given $g_1, g_2 : L^\infty(\Omega) \rightarrow [0, \infty)$, we use the notation $g_1(\cdot) = O(g_2(\cdot))$ to indicate the existence of a constant $c > 0$ such that $g_1(\cdot) \leq c g_2(\cdot)$. Given two sequences $a_k, b_k \in [0, \infty]$ we say that $b_k = o(c_k)$ if there exists a sequence $c_k \in [0, \infty)$, with $c_k \rightarrow 0$ as $k \uparrow \infty$, such that $b_k = c_k a_k$.

For future reference we recall the following Sobolev embeddings (cf. [1], [20], [22]):

$$(1.2) \quad W^{m,s}(\Omega) \subseteq \begin{cases} L^{q_1}(\Omega) & \text{with } \frac{1}{q_1} = \frac{1}{s} - \frac{m}{n} \text{ if } s < \frac{n}{m}, \\ L^q(\Omega) & \text{with } q \in [1, +\infty) \text{ if } s = \frac{n}{m}, \\ C^{m - \lfloor \frac{n}{s} \rfloor - 1, \gamma(n,s)}(\bar{\Omega}) & \text{if } s > \frac{n}{m}, \end{cases}$$

where the injections are continuous and $\gamma(n, s)$ is defined as

$$(1.3) \quad \gamma(n, s) = \begin{cases} \lfloor \frac{n}{s} \rfloor - \frac{n}{s} + 1, & \text{if } \frac{n}{s} \notin \mathbb{Z}, \\ \text{any positive number} < 1 & \text{if } \frac{n}{s} \in \mathbb{Z}. \end{cases}$$

The next well known regularity result for *linear elliptic equations* will be very useful.

Theorem 1.1. *Let $R > 0$ be given. Consider the following Dirichlet problem*

$$(1.4) \quad \begin{cases} -\Delta z + \alpha(x)z = f(x) & \text{in } \Omega, \\ z = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\alpha \in L^\infty(\Omega)$ satisfies $\alpha(x) \geq 0$ for a.a. $x \in \Omega$, and $\|\alpha\|_\infty \leq R$. Then, for every $s \in (1, \infty)$ and $f \in L^s(\Omega)$, equation (1.4) admits a unique strong solution $z \in W^{2,s}(\Omega) \cap W_0^{1,s}(\Omega)$ and there exists $c_s = c_s(R) > 0$ such that for every $f \in L^s(\Omega)$,

$$(1.5) \quad \|z\|_{2,s} \leq c_s \|f\|_s.$$

Moreover, there exists $c_1 = c_1(R) > 0$ such that the following L^1 -estimate holds true

$$(1.6) \quad \|z\|_1 \leq c_1 \|f\|_1.$$

Remark 1.2. The proof of (1.5) can be found in [22, Theorem 9.15 and Lemma 9.17] while (1.6) is a corollary of Stampacchia's results in [32] (see also [11, Lemma 2.11] for a simple proof). By the Sobolev embeddings (1.2), inequality (1.5) implies that if $s > n/2$ ($s = 2$ if $n \leq 3$), then $z \in C(\overline{\Omega})$ and $\|z\|_\infty \leq c_s \|f\|_s$.

In this work, we are concerned with the following controlled semi-linear elliptic equation:

$$(1.7) \quad \begin{cases} -\Delta y + \varphi(x, y, u) = 0 & \text{in } \Omega, \\ y = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\varphi : \overline{\Omega} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$. For $a, b \in C(\overline{\Omega})$ with $a \leq b$ in $\overline{\Omega}$, we suppose that the controls u take values in the set \mathcal{K} of *admissible controls*:

$$(1.8) \quad \mathcal{K} = \{u \in L^\infty(\Omega) \mid a(x) \leq u(x) \leq b(x), \text{ for a.a. } x \in \Omega\}.$$

Let us denote by $M := \max\{\|a\|_\infty, \|b\|_\infty\}$. We will assume the following assumption:

(H1) The function φ is continuous and satisfies:

(i) For all $x \in \overline{\Omega}$ we have that $\varphi(x, \cdot, \cdot)$ is C^1 . Moreover, uniformly on $x \in \overline{\Omega}$ we have that

- (i.1) $D_{(y,u)}\varphi(x, 0, 0)$ is bounded;
- (i.2) $D_{(y,u)}\varphi(x, \cdot, \cdot)$ is Lipschitz on bounded sets.

(ii) For all $(x, y) \in \overline{\Omega} \times \mathbb{R}$ and $|u| \leq M$, we have $\varphi_y(x, y, u) \geq 0$.

Example 1.3. A typical example of φ satisfying **(H1)** is $\varphi(x, y, u) = g(y) + u + f(x)$, where $f \in C(\overline{\Omega})$, and $g \in C^2(\mathbb{R})$ and satisfies $g_y \geq 0$.

The following result is well known (see e.g. [25, Chapter 5, Proposition 1.1]).

Proposition 1.4. *Under **(H1)**, for every $u \in \mathcal{K}$ and $s \in (n/2, \infty)$, equation (1.7) has a unique strong solution $y_u \in W_0^{1,s}(\Omega) \cap W^{2,s}(\Omega)$. In particular, we have that y_u is continuous. Moreover, there exists a constant C_s depending only on s , such that*

$$(1.9) \quad \|y_u\|_\infty + \|y_u\|_{2,s} \leq C_s, \quad \text{for all } u \in \mathcal{K}.$$

Consider a function $\ell : \bar{\Omega} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and assume that:

(H2) The function ℓ satisfies the same assumptions for φ in **(H1)** except for (ii).

Let us define the *cost function* $J : L^\infty(\Omega) \rightarrow \mathbb{R}$ by

$$(1.10) \quad J(u) := \int_{\Omega} \ell(x, y_u(x), u(x)) dx.$$

In this work we are concerned with the following *optimal control problem*:

$$\min J(u) \quad \text{subject to } u \in \mathcal{K}. \quad (\mathcal{CP})$$

Remark 1.5. The existence of a global solution $\bar{u} \in \mathcal{K}$ (i.e. $J(\bar{u}) \leq J(u)$ for all $u \in \mathcal{K}$) of (\mathcal{CP}) can be proved only for specific structures (see e.g. [21, Chapter 1] and [33, Chapter 4]). In particular, if $v \in \mathbb{R} \rightarrow \phi(x, y, v) \in \mathbb{R}$ is affine, $v \in \mathbb{R} \rightarrow \ell(x, y, v) \in \mathbb{R}$ is convex and continuous and **(H1)**-**(H2)** are satisfied, then there exists at least one global solution \bar{u} of (\mathcal{CP}) . In what follows, we assume the existence of (*weak or strong*) local solutions (see Definition 1.6).

As we will see in the next section, assumptions **(H1)**-**(H2)** will allow us to obtain well known first order expansions for the state and the cost functions. However, in order to provide second order expansions we will need the following assumption:

(H3) For all $x \in \bar{\Omega}$ and $\psi = \varphi, \ell$, we have that $\psi(x, \cdot, \cdot)$ is C^2 . Moreover, uniformly on $x \in \bar{\Omega}$ we have that

$$(i) \ D_{(y,u)^2}^2 \psi(x, 0, 0) \text{ is bounded;} \quad (ii) \ D_{(y,u)^2}^2 \psi(x, \cdot, \cdot) \text{ is locally Lipschitz.}$$

See Example 3.21 for a typical problem satisfying **(H1)**-**(H3)**. We end this section by recalling the notion of weak and strong local solutions and the so-called local quadratic growth conditions for J .

Definition 1.6. For a fixed $\bar{u} \in \mathcal{K}$ and $s \in [1, \infty)$, we say that:

(i) \bar{u} is a *L^s -weak local minimum* (*weak local minimum if $s = \infty$*) of J on \mathcal{K} if there exists $\varepsilon > 0$ such that

$$(1.11) \quad J(u) \geq J(\bar{u}) \quad \text{for all } u \in \mathcal{K} \text{ with } \|u - \bar{u}\|_s \leq \varepsilon.$$

In this case we also speak of *L^s -weak local solution* (*weak local solution if $s = \infty$*) of (\mathcal{CP}) .

(ii) \bar{u} is a *strong local minimum* of J on \mathcal{K} if there exists $\varepsilon > 0$ such that

$$(1.12) \quad J(u) \geq J(\bar{u}) \quad \text{for all } u \in \mathcal{K} \text{ with } \|y_u - y_{\bar{u}}\|_\infty \leq \varepsilon.$$

In this case we also speak of *strong local solution* of (\mathcal{CP}) .

The following remark exploits the fact that the set \mathcal{K} is bounded in $L^\infty(\Omega)$.

Remark 1.7. (i) Let $1 \leq p < q < \infty$. We claim that $L^p(\Omega)$ and $L^q(\Omega)$ have the same topology (open sets) on \mathcal{K} . Indeed, since Ω is bounded it is known that $L^q(\Omega) \subset L^p(\Omega)$ with continuous injection, proving that each subset of \mathcal{K} that is (relatively) open in $L^p(\Omega)$ is open in $L^q(\Omega)$. On the other hand, for $M = \max(\|a\|_\infty, \|b\|_\infty)$ and $u \in \mathcal{K}$, we have $\|u\|_q^q \leq M^{q-p} \|u\|_p^p$, proving that each open set of $L^q(\Omega) \cap \mathcal{K}$ is open in $L^p(\Omega) \cap \mathcal{K}$.

(ii) It follows that the notion of L^s -weak local solution is equivalent for all $s \in [1, \infty)$. In this case we simply speak of L^1 -weak local solution. Obviously, any L^1 -weak local solution is a weak local solution.

(iii) In Lemma 2.1, stated in next section, we check that $\|y_u - y_{\bar{u}}\|_\infty = O(\|u - \bar{u}\|_s)$ for all $\bar{u}, u \in \mathcal{K}$ and $s \in (n/2, \infty)$. Therefore, every strong local solution is a L^s -weak local solution and, in view of point (ii), is also a L^1 -weak local solution.

We now define the corresponding types of local quadratic growth for J at $\bar{u} \in \mathcal{K}$.

Definition 1.8. Given $\bar{u} \in \mathcal{K}$ and $s \in [1, \infty)$, we say that:

(i) J has local quadratic growth on \mathcal{K} at \bar{u} in the L^s -weak sense (in the weak sense if $s = \infty$) if there exists $\alpha, \varepsilon > 0$ such that

$$(1.13) \quad J(u) \geq J(\bar{u}) + \alpha \|u - \bar{u}\|_2^2 \quad \text{for all } u \in \mathcal{K} \text{ with } \|u - \bar{u}\|_s \leq \varepsilon.$$

(ii) J has local quadratic growth on \mathcal{K} at \bar{u} in the strong sense if there exists $\alpha, \varepsilon > 0$ such that

$$(1.14) \quad J(u) \geq J(\bar{u}) + \alpha \|u - \bar{u}\|_2^2 \quad \text{for all } u \in \mathcal{K} \text{ with } \|y_u - y_{\bar{u}}\|_\infty \leq \varepsilon.$$

Remark 1.9. In view of Remark 1.7, local quadratic growth in the L^s -weak sense is equivalent to the local quadratic growth in the L^1 -weak sense. The latter implies local quadratic growth in the weak sense, and it is implied by local quadratic growth in the strong sense.

2. EXPANSIONS FOR THE STATE AND THE COST FUNCTIONS

In this section, we establish first and second order expansions for the state and the cost functions. The novelty is Theorem 2.5 where a decomposition result for the variation of the cost is provided. In the entire section, we fix some $\bar{u} \in \mathcal{K}$ and we set $\bar{y} := y_{\bar{u}}$ for its associated state. For notational convenience, we often omit the dependence on x of certain functions such as $y_u(\cdot)$ and $u(\cdot)$.

2.1. First order expansions. We define the Hamiltonian $H : \Omega \times \mathbb{R}^3 \rightarrow \mathbb{R}$ associated to $(\mathcal{C}\mathcal{P})$ by

$$(2.1) \quad H(x, y, p, u) = \ell(x, y, u) - p\varphi(x, y, u).$$

The *adjoint state* \bar{p} , associated to \bar{u} , is defined as the unique solution in $H_0^1(\Omega) \cap C(\bar{\Omega})$ of

$$(2.2) \quad \begin{cases} -\Delta \bar{p} &= H_y(x, \bar{y}, \bar{p}, \bar{u}) \text{ in } \Omega, \\ \bar{p} &= 0 \quad \text{on } \partial\Omega. \end{cases}$$

Let us now fix some useful notation. For $\psi = \ell, \varphi$, when there is no ambiguity, we write $\psi(x), \psi_y(x), \psi_u(x), \psi_{yy}(x), \psi_{yu}(x)$ for the value of $\psi, \psi_y, \psi_u, \psi_{yy}, \psi_{yu}$

on $(x, \bar{y}(x), \bar{u}(x))$, respectively. Similar notations are used for H and its derivatives evaluated at $(x, \bar{y}(x), \bar{p}(x), \bar{u}(x))$, for example $H(x) := H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x))$. Moreover, for a fixed $u \in \mathcal{K}$ we set

$$(2.3) \quad \begin{cases} \delta\psi(x) = \psi(x, \bar{y}(x), u(x)) - \psi(x), & \delta\psi_y(x) = \psi_y(x, \bar{y}(x), u(x)) - \psi_y(x), \\ \delta H(x) = H(x, \bar{y}(x), \bar{p}(x), u(x)) - H(x), & \delta H_y(x) = H_y(x, \bar{y}(x), \bar{p}(x), u(x)) - H_y(x). \end{cases}$$

The *first order Pontryagin linearization* $z_1[u]$ of $u \in \mathcal{K} \rightarrow y_u \in H_0^1(\Omega) \cap C(\bar{\Omega})$ in the direction $u - \bar{u}$ is defined as the unique solution in $H_0^1(\Omega) \cap C(\bar{\Omega})$ of

$$(2.4) \quad \begin{cases} -\Delta z_1 + \varphi_y(x)z_1 + \delta\varphi(x) & = 0 \quad \text{on } \Omega, \\ z_1 & = 0 \quad \text{in } \partial\Omega. \end{cases}$$

For a fixed $u \in \mathcal{K}$, set

$$(2.5) \quad \delta u := u - \bar{u}, \quad \delta y[u] = y_u - \bar{y}, \quad d_1[u] = \delta y - z_1[u].$$

When the context is clear, we will write $z_1 = z_1[u]$, $\delta y = \delta y[u]$, $d_1 = d_1[u]$. The next lemma is proved in the appendix.

Lemma 2.1. *Under (H1)-(H2), for every $s \in (n/2, \infty)$ ($s = 2$ if $n \leq 3$) we have:*

$$(2.6) \quad \begin{cases} \|\delta y\|_1 = O(\|\delta u\|_1), \quad \|\delta y\|_2 = O(\|\delta u\|_2), \quad \|\delta y\|_\infty = O(\|\delta u\|_s), \\ \|z_1\|_1 = O(\|\delta u\|_1), \quad \|z_1\|_2 = O(\|\delta u\|_2), \quad \|z_1\|_\infty = O(\|\delta u\|_s), \\ \|d_1\|_1 = O(\|\delta u\|_1 \|\delta u\|_s), \quad \|d_1\|_2 = O(\|\delta y\|_\infty \|\delta u\|_2). \end{cases}$$

We now provide a first order expansion of the cost function. For future reference, we note that multiplying (2.4) by \bar{p} , integrating by parts and using the adjoint equation, we get (recalling (2.3))

$$(2.7) \quad \int_{\Omega} \ell_y(x, \bar{y}, \bar{u}) z_1 dx + \int_{\Omega} \delta\varphi(x) \bar{p} dx = 0.$$

Lemma 2.2. *Under the assumptions of Lemma 2.1 we have, recalling (2.3):*

$$(2.8) \quad J(u) - J(\bar{u}) = \int_{\Omega} \delta H(x) dx + O(\|\delta y\|_\infty \|\delta u\|_2),$$

$$(2.9) \quad J(u) - J(\bar{u}) = \int_{\Omega} \delta H(x) dx + O(\|\delta u\|_1 \|\delta u\|_s).$$

Proof. By doing a Taylor expansion for ℓ , we get the following equalities:

$$\begin{aligned} J(u) - J(\bar{u}) &= \int_{\Omega} [\ell(x, y, u) - \ell(x, \bar{y}, \bar{u})] dx = \int_{\Omega} [\ell(x, y, u) - \ell(x, \bar{y}, u) + \delta\ell(x)] dx, \\ &= \int_{\Omega} [\ell_y(x, \bar{y}, u) \delta y + \delta\ell(x)] dx + O\left(\int_{\Omega} |\delta y|^2 dx\right), \\ &= \int_{\Omega} [\ell_y(x, \bar{y}, \bar{u}) \delta y + \delta\ell_y(x) \delta y + \delta\ell(x)] dx + O\left(\int_{\Omega} |\delta y|^2 dx\right). \end{aligned}$$

Since ℓ_y is uniformly Lipschitz, we have $|\delta\ell_y(x)| = O(|\delta u(x)|)$, hence, introducing z_1 leads to:

$$J(u) - J(\bar{u}) = \int_{\Omega} [\ell_y(x, \bar{y}, \bar{u}) z_1 + \delta\ell(x)] dx + O\left(\int_{\Omega} [|\delta y|^2 + |\delta u \delta y| + |d_1|] dx\right).$$

Using (2.7), we get:

$$(2.10) \quad J(u) - J(\bar{u}) = \int_{\Omega} \delta H(x) dx + O \left(\int_{\Omega} [|\delta y|^2 + |\delta u \delta y| + |d_1|] dx \right).$$

Using that $\|d_1\|_1 \leq |\Omega|^{1/2} \|d_1\|_2$, Lemma 2.1 implies that

$$\int_{\Omega} (|\delta y|^2 + |\delta u \delta y| + |d_1|) dx = O(\|\delta u\|_1 \|\delta y\|_{\infty} + \|\delta u\|_2 \|\delta y\|_{\infty}) = O(\|\delta u\|_2 \|\delta y\|_{\infty}),$$

which proves (2.8). Similarly, combining (2.10) and Lemma 2.1 yields to (2.9). \square

2.2. Second order expansions for the state and the cost function. The second order Pontryagin linearization $z_2[u]$ of $u \in \mathcal{K} \rightarrow y_u \in H_0^1(\Omega) \cap C(\bar{\Omega})$ in the direction $u - \bar{u}$ is defined as the unique solution in $H_0^1(\Omega) \cap C(\bar{\Omega})$ of

$$(2.11) \quad \begin{aligned} -\Delta z_2 + \varphi_y(x) z_2 + \frac{1}{2} \varphi_{yy}(x) (z_1[u])^2 + \delta \varphi_y(x) z_1[u] &= 0 \text{ in } \Omega, \\ z_2 &= 0 \text{ on } \partial\Omega, \end{aligned}$$

where $z_1[u]$ is defined by (2.4) and we recall the notations (1.1) and (2.3). Let us define (recall (2.5))

$$(2.12) \quad d_2[u] := \delta y[u] - (z_1[u] + z_2[u]) = d_1[u] - z_2[u].$$

When the context is clear, we will write $z_2 = z_2[u]$ and $d_2 = d_2[u]$. The next lemma is proved in the appendix.

Lemma 2.3. *Under (H1)-(H3) we have $\|d_2\|_1 = O(\|\delta y\|_{\infty} \|\delta u\|_2^2)$.*

Given sequence $u_k \in \mathcal{K}$, let us set (recall (2.4))

$$(2.13) \quad \delta_k u := u_k - \bar{u}, \quad y_k := y_{u_k}, \quad \delta_k y = y_k - \bar{y} \quad \text{and} \quad z_1^k := z_1[u_k],$$

with a similar convention for z_2^k , d_1^k and d_2^k . We also set

$$(2.14) \quad \delta H^k(x) = H(x, \bar{y}(x), \bar{p}(x), u_k(x)) - H(x), \quad \delta H_y^k(x) = H_y(x, \bar{y}(x), \bar{p}(x), u_k(x)) - H_y(x)$$

with an analogous notation for $\delta \ell^k$, $\delta \varphi^k$ and their derivatives. We now prove the following second order expansion of the cost.

Proposition 2.4. *Suppose that (H1)-(H3) hold true and that $\|\delta_k u\|_2 \rightarrow 0$. Then, we have that*

$$(2.15) \quad J(u_k) - J(\bar{u}) = \int_{\Omega} [\delta H^k(x) + \delta H_y^k(x) z_1^k + \frac{1}{2} H_{yy}(x) (z_1^k)^2] dx + o(\|\delta_k u\|_2^2).$$

Proof. Writing $\ell(x, y_k, u_k) - \ell(x, \bar{y}, \bar{u}) = \ell(x, y_k, u_k) - \ell(x, \bar{y}, u_k) + \delta \ell^k(x)$, a Taylor expansion gives

$$J(u_k) - J(\bar{u}) = \int_{\Omega} [\delta \ell^k(x) + \ell_y(x, \bar{y}, u_k) \delta_k y + \frac{1}{2} \ell_{yy}(x, \bar{y}, u_k) (\delta_k y)^2] dx + O \left(\int_{\Omega} |\delta_k y|^3 dx \right).$$

Using that $\ell_{yy}(x, \cdot, \cdot)$ is locally Lipschitz, we get

$$\begin{aligned} J(u_k) - J(\bar{u}) &= \int_{\Omega} [\delta \ell^k(x) + \delta \ell_y^k(x) \delta_k y + \ell_y(x, \bar{y}, \bar{u}) \delta_k y + \frac{1}{2} \ell_{yy}(x, \bar{y}, \bar{u}) (\delta_k y)^2] dx \\ &\quad + O \left(\int_{\Omega} [|\delta_k y|^2 |\delta_k u| + |\delta_k y|^3] dx \right). \end{aligned}$$

Introducing z_1^k , z_2^k and using that $\ell_y(x, \cdot, \cdot)$ is locally Lipschitz gives:

$$\begin{cases} \int_{\Omega} \delta \ell_y^k(x) \delta_k y dx &= \int_{\Omega} \delta \ell_y^k(x) z_1^k dx + O \left(\int_{\Omega} |\delta_k u| |d_1^k| dx \right), \\ \int_{\Omega} \ell_y(x, \bar{y}, \bar{u}) \delta_k y dx &= \int_{\Omega} \ell_y(x, \bar{y}, \bar{u}) (z_1^k + z_2^k) dx + O \left(\int_{\Omega} |d_2^k| dx \right), \\ \int_{\Omega} \ell_{yy}(x, \bar{y}, \bar{u}) (\delta_k y)^2 dx &= \int_{\Omega} \ell_{yy}(x, \bar{y}, \bar{u}) (z_1^k)^2 dx + O \left(\int_{\Omega} |d_1^k| (\delta_k y + z_1^k) dx \right). \end{cases}$$

Lemmas 2.1 and 2.3 yield:

$$\int_{\Omega} \left(|\delta_k u| |d_1^k| + |d_2^k| + |d_1^k| |\delta_k y| + |d_1^k| |z_1^k| + |\delta_k y|^2 |\delta_k u| + |\delta_k y|^3 \right) dx = O(\|\delta_k u\|_{\infty} \|\delta_k u\|_2^2).$$

On the other hand, by Remark 1.7, $\|\delta_k y\|_{\infty} \rightarrow 0$ when $\|\delta_k u\|_2^2 \rightarrow 0$, so that

$$(2.16) \quad J(u_k) - J(\bar{u}) = \int_{\Omega} [\delta \ell^k(x) + \delta \ell_y^k(x) z_1^k + \ell_y(x, \bar{y}, \bar{u})(z_1^k + z_2^k) + \frac{1}{2} \ell_{yy}(x, \bar{y}, \bar{u})(z_1^k)^2] dx + o(\|\delta_k u\|_2^2).$$

Multiplying (2.11) by \bar{p} and integrating by parts gives

$$(2.17) \quad \int_{\Omega} \ell_y(x, \bar{y}, \bar{u}) z_2^k dx + \int_{\Omega} \frac{1}{2} \varphi_{yy}(x, \bar{y}, \bar{u})(z_1^k)^2 \bar{p} dx + \int_{\Omega} \delta \varphi_y^k(x) z_1^k \bar{p} dx = 0.$$

We conclude by combining (2.7) and (2.17) with (2.16). \square

2.3. A Decomposition result. Our aim is to prove in this section Theorem 2.5 and Corollary 2.6 which roughly say that the effect of small perturbations on the control in the L^2 -norm can be decomposed as the effect of a *small* perturbation in the L^{∞} -norm and the effect of perturbations which can be *large* in the L^{∞} norm but are supported on sets of small measure.

For a sequence u_k in \mathcal{K} , we use the notation introduced in (2.13). Let us suppose that $\|\delta_k u\|_2 \rightarrow 0$ as $k \uparrow \infty$ and consider a sequence of measurable sets A_k and B_k such that

$$(2.18) \quad |A_k \cup B_k| = |\Omega|, \quad |A_k \cap B_k| = 0 \quad \text{and} \quad |B_k| \rightarrow 0.$$

We decompose the sequence u_k into u_{A_k} and u_{B_k} defined by :

$$\begin{cases} u_{A_k} = u_k & \text{on } A_k, & u_{A_k} = \bar{u} & \text{on } B_k, \\ u_{B_k} = \bar{u} & \text{on } A_k, & u_{B_k} = u_k & \text{on } B_k. \end{cases}$$

We set

$$(2.19) \quad \delta_{A_k} u := u_{A_k} - \bar{u}, \quad \delta_{B_k} u := u_{B_k} - \bar{u} \quad \text{and hence} \quad \delta_k u = \delta_{A_k} u + \delta_{B_k} u.$$

Let us set $z_{A_k} := z_1[u_{A_k}]$ and $z_{B_k} := z_1[u_{B_k}]$. Since $|A_k \cap B_k| = 0$ we have, by uniqueness of the Dirichlet problem, that $z_1^k = z_{A_k} + z_{B_k}$. From (1.5), we obtain

$$(2.20) \quad \|z_{A_k}\|_{2,s} \leq c_s \|\delta_{A_k} u\|_s, \quad \|z_{B_k}\|_{2,s} \leq c_s \|\delta_{B_k} u\|_s \quad \text{for all } s \in (1, \infty).$$

We have the following *decomposition result* (recall the notations (2.13) and (2.14)):

Theorem 2.5. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Let $u_k \in \mathcal{K}$ be such that $\|\delta_k u\|_2 \rightarrow 0$, and A_k , B_k and $\delta_{A_k} u$, $\delta_{B_k} u$ be as in (2.18) and (2.19), respectively. If $\|\delta_{A_k} u\|_{\infty} \rightarrow 0$, then (recalling (2.14))*

$$(2.21) \quad J(u_k) - J(\bar{u}) = \int_{B_k} \delta H^k(x) dx + \int_{A_k} H_u(x) \delta_{A_k} u dx + \frac{1}{2} \int_{\Omega} [H_{uu}(x)(\delta_{A_k} u)^2 + 2H_{yu}(x) z_{A_k} \delta_{A_k} u + H_{yy}(x)(z_{A_k})^2] dx + o(\|\delta_k u\|_2^2).$$

Proof. Given $p \in (1, \infty)$ we will denote by p^* its conjugate, i.e. $p^* = p/(p-1)$. Proposition 2.4, with u_k in place of u , yields

$$(2.22) \quad J(u_k) - J(\bar{u}) = \int_{\Omega} [\delta H^k(x) + \delta H_y^k(x) z_1^k + \frac{1}{2} H_{yy}(x)(z_1^k)^2] dx + o(\|\delta_k u\|_2^2).$$

a) We first prove that

$$(2.23) \quad \int_{\Omega} \left[\delta H_y^k(x) z_1^k + \frac{1}{2} H_{yy}(x) (z_1^k)^2 \right] dx = \int_{A_k} \left[\delta H_y^k(x) z_{A_k} + \frac{1}{2} H_{yy}(x) (z_{A_k})^2 \right] dx + o(\|\delta_k u\|_2^2).$$

By (2.20) and (1.2), there exists $q_1 \in (1, 2)$ such that $\|z_{B_k}\|_2 = O(\|\delta_{B_k} u\|_{q_1})$. Thus, by Hölder inequality and the fact that $|B_k| \rightarrow 0$, we get

$$(2.24) \quad \|z_{B_k}\|_2 = O(\|\delta_{B_k} u\|_{q_1}) = o(\|\delta_k u\|_2).$$

Henceforth, an easy computation, using the Cauchy-Schwarz inequality and (2.24), implies that

$$(2.25) \quad \int_{\Omega} \left[\delta H_y^k(x) z_1^k + \frac{1}{2} H_{yy}(x) (z_1^k)^2 \right] dx = \int_{\Omega} \left[\delta H_y^k(x) z_{A_k} + \frac{1}{2} H_{yy}(x) (z_{A_k})^2 \right] dx + o(\|\delta_k u\|_2^2).$$

On the other hand, using (2.20) and (1.2), there exists $q_2 \in (2, \infty)$ such that

$$(2.26) \quad \|z_{A_k}\|_{q_2} = O(\|\delta_{A_k} u\|_2).$$

Since $q_2^* \in (1, 2)$, as in estimate (2.24) we get $\|\delta_{B_k} u\|_{q_2^*} = o(\|\delta_k u\|_2)$. Therefore, (2.26) yields

$$(2.27) \quad \int_{B_k} |z_{A_k}| |\delta_k u| dx \leq \|z_{A_k}\|_{q_2} \|\delta_{B_k} u\|_{q_2^*} = o(\|\delta_k u\|_2^2),$$

showing that $\int_{B_k} \delta H_y^k(x) z_{A_k} dx = o(\|\delta_k u\|_2^2)$. Moreover, using (2.26) and Hölder inequality we obtain

$$\int_{B_k} |z_{A_k}|^2 dx \leq |B_k|^{\frac{1}{(q_2/2)^*}} \left(\int_{B_k} |z_{A_k}|^{q_2} dx \right)^{\frac{2}{q_2}} = o(\|\delta_k u\|_2^2),$$

which, together with (2.25), implies expression (2.23).

b) We now prove that

$$(2.28) \quad \int_{A_k} \delta H_y^k(x) z_{A_k} dx = \int_{A_k} H_{yu}(x) z_{A_k} \delta_{A_k} u dx + o(\|\delta_k u\|_2^2).$$

By a Taylor expansion, we have:

$$(2.29) \quad \int_{A_k} \delta H_y^k(x) z_{A_k} dx = \int_{A_k} H_{yu}(x) z_{A_k} \delta_{A_k} u dx + O\left(\int_{A_k} |z_{A_k}| |\delta_k u|^2 dx \right).$$

Hölder inequality, estimates (2.26) and $\|\delta_k u\|_{q_2^*} = O(\|\delta_k u\|_2)$ and the fact that $\|\delta_{A_k} u\|_{\infty} \rightarrow 0$, give

$$(2.30) \quad \int_{A_k} |z_{A_k}| |\delta_k u|^2 dx \leq \|z_{A_k}\|_{q_2} \left(\int_{A_k} |\delta_k u|^{2q_2^*} dx \right)^{\frac{1}{q_2^*}} = O(\|\delta_{A_k} u\|_{\infty} \|\delta_k u\|_2^2) = o(\|\delta_k u\|_2^2),$$

from which (2.28) follows.

c) Finally, a Taylor expansion and the fact that $\|\delta_{A_k} u\|_{\infty} \rightarrow 0$ imply that

$$(2.31) \quad \int_{A_k} \delta H^k(x) dx = \int_{A_k} \left[H_u(x) \delta_{A_k} u + \frac{1}{2} H_{uu}(x) (\delta_{A_k} u)^2 \right] dx + o(\|\delta_k u\|_2^2).$$

The conclusion follows from (2.22) and expressions (2.23), (2.28) and (2.31). \square

The following corollary is a straightforward consequence of Theorem 2.5.

Corollary 2.6. *Under the assumptions of Theorem 2.5 we have that*

$$(2.32) \quad J(u_k) - J(\bar{u}) = J(u_{B_k}) - J(\bar{u}) + J(u_{A_k}) - J(\bar{u}) + o(\|\delta_k u\|_2^2).$$

We now set some useful notation. Let us first define the linear map $Q_1[\bar{u}] : L^2(\Omega) \rightarrow \mathbb{R}$ as

$$(2.33) \quad Q_1[\bar{u}]v := \int_{\Omega} H_u(x)v(x)dx.$$

Secondly, we define the quadratic form $Q_2[\bar{u}] : L^2(\Omega) \rightarrow \mathbb{R}$ as

$$(2.34) \quad Q_2[\bar{u}](v) = \int_{\Omega} [H_{yy}(x)(\zeta[v])^2 + 2H_{yu}(x)\zeta[v]v + H_{uu}(x)v^2] dx,$$

where $\zeta[v]$ is defined as the unique solution in $H^2(\Omega) \cap H_0^1(\Omega)$ of the linearized state equation

$$(2.35) \quad \begin{cases} -\Delta\zeta + \varphi_y(x)\zeta + \varphi_u(x)v &= 0, & \text{in } \Omega, \\ \zeta &= 0, & \text{on } \partial\Omega. \end{cases}$$

It is easy to check, using Theorem 1.1, that $\zeta[v]$ satisfies the same estimates as $z_1[v]$ in Lemma 2.1. Therefore, we have that $Q_2[\bar{u}](v) = O(\|v\|_2^2)$, and thus $Q_2[\bar{u}]$ is a continuous quadratic form on $L^2(\Omega)$. For future reference, we denote by $\hat{Q}_2[\bar{u}] : L^2(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$ for the associated continuous symmetric bilinear form, i.e.

$$(2.36) \quad \hat{Q}_2[\bar{u}](v_1, v_2) := \frac{1}{2} [Q_2[\bar{u}](v_1 + v_2) - Q_2[\bar{u}](v_1) - Q_2[\bar{u}](v_2)].$$

Using the previous definitions we have the following result:

Theorem 2.7. *Under the assumptions of Theorem 2.5 we have:*

$$(2.37) \quad J(u_k) - J(\bar{u}) = \int_{B_k} \delta H^k(x)dx + Q_1[\bar{u}]\delta_{A_k}u + \frac{1}{2}Q_2[\bar{u}](\delta_{A_k}u) + o(\|\delta_k u\|_2^2).$$

Proof. Let ζ_k be the solution of (2.35) with $\delta_{A_k}u$ in place of v . By a Taylor expansion, the function $w_k := z_{A_k} - \zeta_k$ satisfies on Ω :

$$-\Delta w_k + \varphi_y(x)w_k = O(|\delta_{A_k}u|^2),$$

with Dirichlet boundary condition. Hence, we have

$$\|w_k\|_2 = O(\|\delta_{A_k}u\|_2 \|\delta_{A_k}u\|_{\infty}) = o(\|\delta_{A_k}u\|_2).$$

Using this estimate it is straightforward to prove that we can replace z_{A_k} by ζ_k in (2.21) up to an error $o(\|\delta_{A_k}u\|_2^2)$, from which the result follows. \square

3. OPTIMALITY CONDITIONS

The purpose in this section is to provide some new results concerning optimality conditions for (\mathcal{CP}) . More precisely, we first provide a general first order result that yields to the well-known Pontryagin's minimum principle for L^1 -weak local solutions. Moreover, it also implies that weak local solutions satisfy a *local* Pontryagin's minimum principle. Next, we study second order conditions and we extend to the strong sense two second order sufficient conditions for local quadratic growth (see [18, 3]). Finally, we characterize local quadratic growth in the strong sense.

3.1. Pontryagin's minimum principle for semi-linear elliptic equations.

While Pontryagin's minimum principle for semi-linear elliptic equations is well-known for L^s -weak local solutions ($s \in [1, \infty)$), we will obtain it as a particular case of a more general statement (see Proposition 3.5); the latter will allow us to obtain a version of Pontryagin's minimum principle for weak local solutions. Let us set

$$(3.1) \quad \mathcal{K}(x) := [a(x), b(x)] \quad \text{for any } x \in \Omega.$$

Definition 3.1. We say that the set-valued map $U : \Omega \rightarrow 2^{\mathbb{R}}$ (denoted by $U : \Omega \rightrightarrows \mathbb{R}$) is a measurable multifunction if, for any closed set $C \subset \mathbb{R}$, we have that $U^{-1}(C)$ is measurable. We say that it is feasible if $U(x) \subseteq \mathcal{K}(x)$ for a.a. $x \in \Omega$, and closed-valued if $U(x)$ is closed for a.a. $x \in \Omega$. If $u : \Omega \rightarrow \mathbb{R}$ is a measurable function, with $u(x) \in U(x)$ for a.a. $x \in \Omega$, we say that u is a *measurable selection* of U . We denote by $\text{select}(U)$ the set of measurable selections.

Example 3.2. Given $\bar{u} \in \mathcal{K}$, consider the two following examples:

(i) For $\varepsilon > 0$, set $U^\varepsilon(x) = [a(x), b(x)] \cap [\bar{u}(x) - \varepsilon, \bar{u}(x) + \varepsilon]$ for a.a. $x \in \Omega$. We can interpret a weak solution as a solution of the problem obtained by adding the constraint $u(x) \in U^\varepsilon(x)$ a.e., for $\varepsilon > 0$ small enough.

(ii) For $\bar{x} \in \Omega$ and $\varepsilon > 0$, let $\bar{B}(\bar{x}, \varepsilon)$ the closed ball in \mathbb{R}^n of center \bar{x} and radius ε . The following measurable multifunction will be used in the proof of theorem 3.6:

$$(3.2) \quad U^{\bar{x}, \varepsilon}(x) := \begin{cases} \mathcal{K}(x) & \text{if } x \in \bar{B}(\bar{x}, \varepsilon) \cap \Omega, \\ \{\bar{u}(x)\} & \text{if } x \in \Omega \setminus \bar{B}(\bar{x}, \varepsilon). \end{cases}$$

Definition 3.3. Let $U : \Omega \rightrightarrows \mathbb{R}$ be a feasible, closed valued measurable multifunction. We say that $\bar{u} \in \mathcal{K}$ is a *Pontryagin extremal in integral form with respect to U* if

$$(3.3) \quad \int_{\Omega} H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) dx \leq \int_{\Omega} H(x, \bar{y}(x), \bar{p}(x), v(x)) dx, \quad \text{for all } v \in \text{select}(U),$$

and that it is a *Pontryagin extremal with respect to U* if for a.a. $x \in \Omega$ we have

$$(3.4) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) \leq H(x, \bar{y}(x), \bar{p}(x), v) \quad \text{for all } v \in U(x).$$

If $U(x) = \mathcal{K}(x)$ for a.a. $x \in \Omega$, a Pontryagin extremal with respect to U will be simply called *Pontryagin extremal*. By analogy with weak solutions, if \bar{u} is a Pontryagin extremal with respect to U^ε (defined in Example 3.2(i)) for some $\varepsilon > 0$, we say that \bar{u} is a *weak Pontryagin extremal*.

Obviously, a Pontryagin extremal is a Pontryagin extremal in integral form. Setting $F(x, u) := H(x, \bar{y}(x), \bar{p}(x), u)$ in the proposition below, whose proof can be found in [30, Theorem 3A], we have that the converse is also true.

Proposition 3.4. *Let $U : \Omega \rightrightarrows \mathbb{R}$ be a feasible, closed valued measurable multifunction, $\bar{u} \in \text{select}(U)$, and $F : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, such that \bar{u} is solution of the problem*

$$(3.5) \quad \text{Min}_u \int_{\Omega} F(x, u(x)) dx \quad u \in \text{select}(U).$$

Then, for a.a. $x \in \Omega$,

$$(3.6) \quad F(x, \bar{u}(x)) \leq F(x, v), \quad \text{for all } v \in U(x).$$

As a consequence we have:

Proposition 3.5. *Let $U : \Omega \rightrightarrows \mathbb{R}$ be a feasible, closed valued measurable multifunction. Suppose that $\bar{u} \in \mathcal{K}$ satisfies*

$$(3.7) \quad J(\bar{u}) \leq J(u), \quad \text{for all } u \in \text{select}(U).$$

Then \bar{u} is a Pontryagin extremal with respect to U .

Proof. In view of Proposition 3.4 it is enough to show that \bar{u} is a Pontryagin extremal in integral form with respect to U . Set $F(x, u) := H(x, \bar{y}(x), \bar{p}(x), u)$ and $\mathcal{F}(u) := \int_{\Omega} F(x, u(x)) dx$. By contradiction, suppose that there exists $\bar{v} \in \text{select}(U)$ such that $(\bar{v}) < (\bar{u})$. Consequently, for some $\varepsilon > 0$, there exists a measurable subset Ω_{ε} of Ω such that

$$(3.8) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{v}(x)) + \varepsilon \leq H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)), \quad \text{for a.a. } x \in \Omega_{\varepsilon}.$$

Let Ω_k be a sequence of measurable subsets of Ω_{ε} such that $|\Omega_k| = 1/k$ and define $v_k : \Omega \rightarrow \mathbb{R}$ as

$$(3.9) \quad v_k(x) = \bar{v}(x) \text{ if } x \in \Omega_k, \quad v_k(x) = \bar{u}(x) \text{ otherwise.}$$

Then $v_k \in \text{select}(U)$ and $\|v_k - \bar{u}\|_1 = O(1/k)$. Thus, $\|v_k - \bar{u}\|_1 \|v_k - \bar{u}\|_s = o(1/k)$ for $s > n/2$. Therefore, expression (2.9) in Lemma 2.2, inequality (3.8) and the optimality condition (3.7) imply

$$0 \leq J(v_k) - J(\bar{u}) \leq -\varepsilon/k + o(1/k) \quad \text{as } k \uparrow \infty,$$

which gives the desired contradiction. \square

As a corollary we obtain the well known Pontryagin's minimum principle for semi-linear elliptic problems, e.g. [4, 6, 7, 29], as well as a version for weak local solutions.

Theorem 3.6. *Let \bar{u} be a L^1 -weak (respectively weak) local solution of (\mathcal{CP}) . Then \bar{u} is a Pontryagin extremal (respectively weak Pontryagin extremal).*

Proof. The result is a straightforward consequence of Proposition 3.5 applied to the multifunctions of example 3.2, in case (i) (respectively (ii)) for dealing with weak (respectively, L^1 -weak) extremals. \square

Definition 3.7. Let \bar{u} be a Pontryagin extremal with respect to the feasible, closed valued measurable multifunction U . We can change U and \bar{u} over a negligible set, so that $U(x)$ is compact for all x , and $\bar{u}(x)$ minimizes the Hamiltonian for all $x \in \bar{\Omega}$. In that case we say that \bar{u} is a *Pontryagin representative* (of the equivalence class of functions a.e. equal to \bar{u}). Note that a Pontryagin representative is also defined on $\partial\Omega$. In the sequel we will identify Pontryagin extremals with one of their Pontryagin representatives.

Corollary 3.8. *Let \bar{u} be a weak local solution of (\mathcal{CP}) . Let $x \in \Omega$ be such that $a(x) < b(x)$. Then*

$$(3.10) \quad \begin{aligned} H_u(x) \geq 0 \quad \text{if } \bar{u}(x) = a(x), \quad H_u(x) \leq 0 \quad \text{if } \bar{u}(x) = b(x), \\ \text{and } H_u(x) = 0 \text{ otherwise.} \end{aligned}$$

Moreover, we have that

$$(3.11) \quad H_{uu}(x) \geq 0 \quad \text{if } H_u(x) = 0.$$

Proof. This is a straightforward consequence of Theorem 3.6. \square

Definition 3.9. Let \bar{u} be a Pontryagin extremal. We say that the *strict Pontryagin inequality* holds at $\bar{u} \in \mathcal{K}$ if for all $x \in \bar{\Omega}$,

$$(3.12) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) < H(x, \bar{y}(x), \bar{p}(x), v), \quad \text{when } v \neq \bar{u}(x), \quad \text{for all } v \in \mathcal{K}(x).$$

Lemma 3.10. *A Pontryagin extremal that satisfies the strict Pontryagin inequality belongs to $C(\bar{\Omega})$.*

Proof. Let \bar{u} satisfy the hypothesis of the lemma, and let $x_k \rightarrow \bar{x}$ in $\bar{\Omega}$. Denote by

$$\hat{u}_k := \max(a(x_k), \min(\bar{u}(\bar{x}), b(x_k))),$$

the projection of $\bar{u}(\bar{x})$ onto $[a(x_k), b(x_k)]$. Since a and b are continuous, $\hat{u}_k \rightarrow \bar{u}(\bar{x})$. Extracting if necessary a subsequence, we may assume that $\bar{u}(x_k) \rightarrow \tilde{u} \in \mathcal{K}(\bar{x})$, and so

$$(3.13) \quad \begin{aligned} H(\bar{x}, \bar{y}(\bar{x}), \bar{p}(\bar{x}), \tilde{u}) &= \lim_k H(x_k, \bar{y}(x_k), \bar{p}(x_k), \bar{u}(x_k)) \\ &\leq \lim_k H(x_k, \bar{y}(x_k), \bar{p}(x_k), \hat{u}_k) = H(\bar{x}, \bar{y}(\bar{x}), \bar{p}(\bar{x}), \bar{u}(\bar{x})). \end{aligned}$$

By (3.12), $\tilde{u} = \bar{u}(\bar{x})$. The conclusion follows. \square

3.2. Second order necessary conditions. Now, we establish second order necessary conditions. The novelty in this subsection is a second order necessary condition for local quadratic growth in L^1 -sense (so in particular in the strong sense). Let us start with some standard definitions and results.

Consider a Banach space $(X, \|\cdot\|_X)$ and a non-empty closed convex set $K \subseteq X$. For $x, x' \in X$ define the *segment* $[x, x'] := \{x + \lambda(x' - x) \mid \lambda \in [0, 1]\}$ and for $A \subseteq X$ set $\text{clo}_X(A)$ for its closure. In order to simplify the notation, when $X = L^p(\Omega)$ for some $p \in [1, \infty]$ we will write $\text{clo}_p(A) := \text{clo}_{L^p(\Omega)}(A)$ for $A \subseteq L^\infty(\Omega)$. The radial, the tangent and the normal cone to K at \bar{x} are defined respectively by

$$(3.14) \quad \begin{aligned} R_K(\bar{x}) &:= \{h \in X \mid \exists \sigma > 0 \text{ such that } [\bar{x}, \bar{x} + \sigma h] \subseteq K\}, \\ T_K(\bar{x}) &:= \{h \in X \mid \exists x(\sigma) = \bar{x} + \sigma h + o(\sigma) \in K, \sigma > 0, \|o(\sigma)/\sigma\|_X \rightarrow 0, \text{ as } \sigma \downarrow 0\}, \\ N_K(\bar{x}) &:= \{x^* \in X^* \mid \langle x^*, x - \bar{x} \rangle_{X^*, X} \leq 0, \text{ for all } x \in K\}, \end{aligned}$$

where X^* is the dual topological space of X and $\langle \cdot, \cdot \rangle_{X^*, X}$ is the duality product. Recall that, since K is a closed convex set, we have $T_K(\bar{x}) = \text{clo}_X(R_K(\bar{x}))$ and $N_K(\bar{x})$ is the polar cone of $T_K(\bar{x})$, i.e.,

$$(3.15) \quad N_K(\bar{x}) = \{x^* \in X^* \mid \langle x^*, h \rangle_{X^*, X} \leq 0, \text{ for all } h \in T_K(\bar{x})\}.$$

In what follows we will consider the set \mathcal{K} , defined in (1.8), as a subset of $L^2(\Omega)$ rather than a subset of $L^\infty(\Omega)$. This will allow us to give explicit expressions for the tangent and normal cone. Note that, since $a, b \in L^\infty(\Omega)$, for all $\bar{u} \in \mathcal{K}$ we have that $R_{\mathcal{K}}(\bar{u})$ is a subset of $L^\infty(\Omega)$. The next lemma is standard (at least when $\min(b - a) > 0$) and gives the relation between the tangent cone $T_{\mathcal{K}}(\bar{u})$ (normal cone $N_{\mathcal{K}}(\bar{u})$) and the *pointwise* tangent cone $T_{\mathcal{K}(x)}(\bar{u}(x))$ (*pointwise* normal cone $N_{\mathcal{K}(x)}(\bar{u}(x))$) for $x \in \Omega$. The proof of the lemma can be found in the appendix for the sake of completeness of the paper.

Lemma 3.11. *Let \mathcal{K} be defined by (1.8) and $\bar{u} \in \mathcal{K}$. Then the following assertions hold true:*

(i) *The tangent cone to \mathcal{K} at \bar{u} is given by*

$$(3.16) \quad \begin{aligned} T_{\mathcal{K}}(\bar{u}) &= \{v \in L^2(\Omega) \mid v(x) \in T_{\mathcal{K}(x)}(\bar{u}(x)) \text{ for a.a. } x \in \Omega\}, \\ &= \{v \in L^2(\Omega) \mid v(x) \geq 0 \text{ if } \bar{u}(x) = a(x), v(x) \leq 0 \text{ if } \bar{u}(x) = b(x) \\ &\quad \text{a.e. in } \Omega\}. \end{aligned}$$

(ii) *The normal cone to \mathcal{K} at \bar{u} is given by*

$$(3.17) \quad N_{\mathcal{K}}(\bar{u}) = \{v \in L^2(\Omega) \mid v(x) \in N_{\mathcal{K}(x)}(\bar{u}(x)) \text{ for a.a. } x \in \Omega\}.$$

(iii) *For every $q^* \in N_{\mathcal{K}}(\bar{u})$ we have that*

$$(3.18) \quad T_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp = \{v \in T_{\mathcal{K}}(\bar{u}) \mid v(x)q^*(x) = 0 \text{ for a.a. } x \in \Omega\},$$

where $(q^*)^\perp$ is the subspace of $L^2(\Omega)$ consisting of the orthogonal vectors to q^* .

The following classical technical notion is essential in the study of second order necessary conditions in abstract optimization theory, see e.g. [23, 26, 10].

Definition 3.12. The set \mathcal{K} is said to be polyhedral at $\bar{u} \in \mathcal{K}$ if for every $q^* \in N_{\mathcal{K}}(\bar{u})$ we have that

$$(3.19) \quad \text{clo}_2(R_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp) = T_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp.$$

The following lemma is a particular instance of the more general result [10, Theorem 3.58] which holds true in general Banach lattices. For the reader's convenience we provide the simple proof for our case.

Lemma 3.13. *The set \mathcal{K} is polyhedral at \bar{u} .*

Proof. The inclusion $\text{clo}_2(R_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp) \subseteq T_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp$ being trivial, we prove the other one. Now, fix $q^* \in N_{\mathcal{K}}(\bar{u})$ and let $v \in T_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp$. For $\varepsilon > 0$ define

$$(3.20) \quad \widehat{v}_\varepsilon(x) := \begin{cases} v(x) & \text{if } \bar{u}(x) + \varepsilon v(x) \in \mathcal{K}(x), \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, $\widehat{v}_\varepsilon \in R_{\mathcal{K}}(\bar{u}) \cap (q^*)^\perp$ and, as $\varepsilon \downarrow 0$, we have $\widehat{v}_\varepsilon(x) \rightarrow v(x)$ for a.a. $x \in \Omega$. The Dominated Convergence Theorem gives that $\widehat{v}_\varepsilon \rightarrow v$ in $L^2(\Omega)$ and the result follows. \square

We now introduce the set of directions that, while being tangent to the feasible set, do not increase the cost function up to the first order. The *critical cone* to \mathcal{K} at a local Pontryagin extremal \bar{u} is defined as

$$(3.21) \quad C_{\mathcal{K}}(\bar{u}) := T_{\mathcal{K}}(\bar{u}) \cap (Q_1[\bar{u}])^\perp,$$

while the *pointwise* critical cone is defined by

$$(3.22) \quad C_x := \{v \in T_{\mathcal{K}(x)}(\bar{u}(x)) \mid H_u(x)v = 0\} \text{ for a.a. } x \in \Omega.$$

Since \bar{u} is a local Pontryagin extremal \bar{u} , we have that $-Q_1[\bar{u}] \in N_{\mathcal{K}}(\bar{u})$. Thus, Lemma 3.11(iii) implies that

$$(3.23) \quad C_{\mathcal{K}}(\bar{u}) = \{v \in L^2(\Omega) \mid v(x) \in C_x \text{ for a.a. } x \in \Omega\}.$$

We have the following second order necessary condition (see e.g. [3, 14, 15] for more general settings).

Theorem 3.14. *Let \bar{u} be a weak local solution of (CP). Then, recalling (2.34), we have*

$$(3.24) \quad Q_2[\bar{u}](v) \geq 0 \quad \text{for all } v \in C_{\mathcal{K}}(\bar{u}),$$

Proof. Let $v \in R_{\mathcal{K}}(\bar{u}) \cap (Q_1[\bar{u}])^\perp$ and set $u_k := \bar{u} + \frac{1}{k}v \in \mathcal{K}$, for k large. The local optimality of \bar{u} and Theorem 2.7 with $A_k \equiv \Omega$, imply that

$$0 \leq J(u_k) - J(\bar{u}) = \frac{1}{k^2}Q_2[\bar{u}](v) + o(1/k^2).$$

Multiplying by k^2 and letting $k \uparrow \infty$ yields (3.24) for all $v \in R_{\mathcal{K}}(\bar{u}) \cap (Q_1[\bar{u}])^\perp$. Using that $Q_1[\bar{u}] \in N_{\mathcal{K}}(\bar{u})$, relation (3.24) follows from the continuity of Q_2 in $L^2(\Omega)$ and the first equality in Lemma 3.11(iii). \square

Now we begin the study of necessary conditions for local quadratic growth on \mathcal{K} .

Definition 3.15. For $\bar{u} \in \mathcal{K}$, we say that

(i) The Hamiltonian satisfies the *a.e. local quadratic growth property* at \bar{u} if and only if there exist $\alpha, \varepsilon > 0$ such that *a.e.* in Ω we have

$$(3.25) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) + \alpha|v - \bar{u}(x)|^2 \leq H(x, \bar{y}(x), \bar{p}(x), v) \quad \forall v \in \mathcal{K}(x) \quad \text{with } |v - \bar{u}(x)| \leq \varepsilon.$$

(ii) The Hamiltonian satisfies the *global quadratic growth property* at Pontryagin representative of \bar{u} if and only if there exists $\alpha > 0$ such that *for all* $x \in \bar{\Omega}$ we have

$$(3.26) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) + \alpha|v - \bar{u}(x)|^2 \leq H(x, \bar{y}(x), \bar{p}(x), v) \quad \text{for all } v \in \mathcal{K}(x).$$

Adapting the techniques of [9] to our framework we can characterize the global quadratic growth for the Hamiltonian.

Lemma 3.16. *Let $\bar{u} \in \mathcal{K}$. Then \bar{u} is a Pontryagin extremal that satisfies the global quadratic growth property for the Hamiltonian iff both the strict Pontryagin inequality (3.12) and the a.e. local quadratic growth property for the Hamiltonian hold.*

Proof. It is enough to show that if (3.12) holds everywhere and (3.25) holds a.e. then (3.26) holds everywhere. First note that by Lemma 3.10 we have that \bar{u} is continuous. It follows that (3.25) holds everywhere. Now, let

$$\beta := \min_{x,v} \{H(x, \bar{y}(x), \bar{p}(x), v) - H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) \mid x \in \bar{\Omega}, v \in \mathcal{K}(x), |v - \bar{u}(x)| \geq \varepsilon\}.$$

By (3.12) and since \bar{u} is continuous, we get that $\beta > 0$. For $v \in \mathcal{K}(x)$ with $|v - \bar{u}(x)| \geq \varepsilon$, we have

$$H(x, \bar{y}(x), \bar{p}(x), v) - H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) \geq \beta \geq \frac{\beta}{4M^2}|v - \bar{u}(x)|^2,$$

where we recall that $M = \max\{\|a\|_\infty, \|b\|_\infty\}$. Hence, by (3.25) we get that (3.26) is satisfied with $\min\left(\alpha, \frac{\beta}{4M^2}\right)$, which concludes the proof. \square

We now provide second order necessary conditions for local quadratic growth on \mathcal{K} .

Theorem 3.17. *The following assertions hold true:*

(i) *If J has local quadratic growth on \mathcal{K} at \bar{u} in the weak sense, then the Hamiltonian satisfies the a.e. local quadratic growth property, and we have*

$$(3.27) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}(\bar{u}).$$

(ii) *If J has local quadratic growth on \mathcal{K} at \bar{u} in the L^1 -weak sense, then the Hamiltonian satisfies the global quadratic growth property and (3.27) holds true. In particular, \bar{u} is continuous.*

Proof. For $\alpha > 0$, let $J_\alpha : L^\infty(\Omega) \rightarrow \mathbb{R}$ be defined by $J_\alpha(u) := J(u) - \alpha \|u - \bar{u}\|_2^2$. Consider the problem

$$\min J_\alpha(u) \quad \text{subject to } u \in \mathcal{K}. \quad (\mathcal{CP}_\alpha)$$

If J has local quadratic growth on \mathcal{K} at \bar{u} in the weak sense, then there exists $\alpha > 0$, $\varepsilon > 0$ such that $J_\alpha(u) \geq J_\alpha(\bar{u})$, whenever $\|u - \bar{u}\|_\infty \leq \varepsilon$, that is \bar{u} is a weak local solution of (\mathcal{CP}_α) . By Theorem 3.14, (3.27) holds, and by Theorem 3.6, \bar{u} is a local Pontryagin extremal for (\mathcal{CP}_α) , i.e.,

$$(3.28) \quad H(x, \bar{y}(x), \bar{p}(x), \bar{u}(x)) \leq H(x, \bar{y}(x), \bar{p}(x), v) - \alpha |v - \bar{u}|^2,$$

for all $v \in \mathcal{K}(x)$ with $|v - \bar{u}(x)| \leq \varepsilon$. This means that Hamiltonian satisfies the a.e. local quadratic growth property (3.25). On the other hand, if J has local quadratic growth on \mathcal{K} at \bar{u} in the L^1 -weak sense, we have that \bar{u} is a L^1 -weak local solution of (\mathcal{CP}_α) . By Theorem 3.6, \bar{u} is a Pontryagin extremal for (\mathcal{CP}_α) , i.e. for a.a. $x \in \Omega$, inequality (3.28) holds for all $v \in \mathcal{K}(x)$. This means that (3.26) holds for all $x \in \Omega' \subset \Omega$ with $\text{mes}(\Omega \setminus \Omega') = 0$. Let $\bar{x} \in \bar{\Omega}$, and $x_k \rightarrow \bar{x}$, $x_k \in \Omega'$. Extracting if necessary a subsequence, we may assume that $u(x_k) \rightarrow \tilde{u}$, and so for $v \in \mathcal{K}(\bar{x})$, denoting by v_k the projection of v onto $[a(x_k), b(x_k)]$, we have:

$$\begin{aligned} & H(\bar{x}, \bar{y}(\bar{x}), \bar{p}(\bar{x}), \tilde{u}) - H(\bar{x}, \bar{y}(\bar{x}), \bar{p}(\bar{x}), v) + \alpha |v - \tilde{u}|^2 = \\ & \lim_{k \rightarrow \infty} \{H(x_k, \bar{y}(x_k), \bar{p}(x_k), \bar{u}(x_k)) - H(x_k, \bar{y}(x_k), \bar{p}(x_k), v_k) + \alpha |v_k - \bar{u}(x_k)|^2\} \leq 0. \end{aligned}$$

By the definition 3.7, we have that $\tilde{u} = u(\bar{x})$ and the above inequality yields that (3.26) holds for all $x \in \Omega$. In particular, Lemma 3.10 implies that \bar{u} is continuous. \square

3.3. Extension of standard second order sufficient conditions. In this subsection we extend to the strong sense two well known second order sufficient conditions for the local quadratic growth of J on \mathcal{K} in the weak sense. The main tool for proving such extensions is the decomposition result in Theorem 2.5.

We first consider the case studied in [3] which supposes that $Q_2[\bar{u}]$ is a Legendre form.

Definition 3.18. Given a Hilbert space X , a quadratic form $Q : X \rightarrow \mathbb{R}$ is said to be a Legendre form if it is sequentially weakly lower semicontinuous and that if h_k converges weakly to h in X and $Q(h_k) \rightarrow Q(h)$ then h_k converges strongly to h in X .

For the reader's convenience, let us reproduce the sufficiency part of [3, Theorem 2.9].

Theorem 3.19. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Assume that $H_u(x)v \geq 0$ for all $v \in T_{\mathcal{K}(x)}(\bar{u}(x))$ a.e. in Ω , that $Q_2[\bar{u}]$ is a Legendre form and that there exists $\alpha > 0$ such that the following second order condition holds true:*

$$(3.29) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}(\bar{u}).$$

Then J has local quadratic growth on \mathcal{K} at \bar{u} in the weak sense.

We recall that J has local quadratic growth on \mathcal{K} at \bar{u} in the strong sense if there exists $\alpha, \varepsilon > 0$ such that

$$(3.30) \quad J(u) \geq J(\bar{u}) + \alpha \|u - \bar{u}\|_2^2 \quad \text{for all } u \in \mathcal{K} \text{ with } \|y_u - y_{\bar{u}}\|_{\infty} \leq \varepsilon.$$

We have the following extension of Theorem 3.19.

Theorem 3.20. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Assume that the strict Pontryagin inequality holds at \bar{u} , that $Q_2[\bar{u}]$ is a Legendre form and that there exists $\alpha > 0$ such that the following second order condition holds true:*

$$(3.31) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}(\bar{u}).$$

Then J has local quadratic growth on \mathcal{K} at \bar{u} in the strong sense.

Proof. a) Let assume that (3.30) does not hold. Then, there exists a sequence $u_k \in \mathcal{K}$ such that $\|y_k - \bar{y}\|_{\infty} \rightarrow 0$ as $k \uparrow \infty$ (we have denoted $y_k := y_{u_k}$) and

$$(3.32) \quad J(u_k) - J(\bar{u}) \leq o(\|\delta_k u\|_2^2),$$

where $\delta_k u := u_k - \bar{u}$. Theorem 3.19 implies that J has local quadratic growth on \mathcal{K} at \bar{u} in the weak sense. Therefore, Theorem 3.17(i) yields that the Hamiltonian satisfies the a.e. local quadratic growth property. Using Lemma 3.16, we obtain that the Hamiltonian satisfies the global quadratic growth property. Consequently, by expression (2.8) in Lemma 2.2, inequality (3.32) implies that $\|\delta_k u\|_2 \rightarrow 0$. Let us define the measurable sets

$$(3.33) \quad A_k := \left\{ x \in \Omega \mid |u_k(x) - \bar{u}(x)| \leq \sqrt{\|u_k - \bar{u}\|_1} \right\} \quad \text{and} \quad B_k := \Omega \setminus A_k.$$

Chebyshev's inequality implies that $|B_k| \leq \sqrt{\|u_k - \bar{u}\|_1}$, hence $|B_k|$ goes to zero. Thus, introducing the notations of subsection 2.3 we clearly have that $\|\delta_{A_k} u\|_{\infty} \rightarrow 0$. Therefore, Theorem 2.5 gives

$$(3.34) \quad \int_{B_k} \delta H^k(x) dx + \int_{A_k} H_u(x) \delta_{A_k} u(x) dx + \frac{1}{2} Q_2[\bar{u}](\delta_{A_k} u) \leq o(\|\delta_k u\|_2^2).$$

Now, set

$$(3.35) \quad \sigma_{A_k} := \|\delta_{A_k} u\|_2, \quad \sigma_{B_k} := \|\delta_{B_k} u\|_2 \quad \text{and hence} \quad \|\delta_k u\|_2^2 = \sigma_{A_k}^2 + \sigma_{B_k}^2.$$

If $\sigma_{A_k} = o(\sigma_{B_k})$, using that $H_u(x) \delta_{A_k} u(x) \geq 0$ and that $Q_2[\bar{u}](\delta_{A_k} u) = O(\sigma_{A_k}^2) = o(\sigma_{B_k}^2)$, inequality (3.34) and (3.26) imply that $\sigma_{B_k}^2 \leq o(\sigma_{B_k}^2)$, which is impossible. Thus, let us assume, up to a subsequence, that $\sigma_{B_k} = O(\sigma_{A_k})$ and define $h_k := \delta_{A_k} u / \sigma_{A_k}$.

By (3.12) the first integral in (3.34) is nonnegative, and therefore, after minorizing it by 0,

$$(3.36) \quad \int_{A_k} H_u(x) h_k(x) dx + \frac{1}{2} \sigma_{A_k} Q_2[\bar{u}](h_k) \leq o(\sigma_{A_k}).$$

It follows that

$$(3.37) \quad \int_{A_k} H_u(x)h_k(x)dx \leq O(\sigma_{A_k}).$$

Also, minorizing the first integral in (3.36) by 0, we obtain that

$$(3.38) \quad Q_2[\bar{u}](h_k) \leq o(1),$$

where we recall that $o(1)$ denotes a sequence α_k that tends to zero as $k \uparrow \infty$.

b) Since $h_k \in T_{\mathcal{K}}(\bar{u})$ and $\|h_k\|_2 = 1$, up to a subsequence, it converges weakly in $L^2(\Omega)$ to some \bar{h} . Recalling that $T_{\mathcal{K}}(\bar{u})$ is weakly closed we get that $\bar{h} \in T_{\mathcal{K}}(\bar{u})$. Noting that $Q_2[\bar{u}](\delta_{A_k}u)/\sigma_k = o(1)$, condition (3.26) and equation (3.34) imply that

$$0 \leq \int_{A_k} H_u(x)h_k(x)dx \leq o(1).$$

By passing to the limit in the above inequality, we get that $\bar{h} \in C_{\mathcal{K}}(\bar{u})$. On the other hand, since equation (3.34) implies that $Q_2[\bar{u}](h_k) \leq o(1)$, the lower semicontinuity of $Q_2[\bar{u}]$ and its positivity over $C_{\mathcal{K}}(\bar{u})$ give

$$0 \leq Q_2[\bar{u}](\bar{h}) \leq \liminf_{k \rightarrow \infty} Q_2[\bar{u}](h_k) \leq \limsup_{k \rightarrow \infty} Q_2[\bar{u}](h_k) \leq 0.$$

The above inequality and (3.31) imply that $\bar{h} = 0$ and that $Q_2[\bar{u}](h_k) \rightarrow Q_2[\bar{u}](\bar{h})$. Thus, since $Q_2[\bar{u}]$ is a Legendre form, we have that $h_k \rightarrow 0$ strongly in $L^2(\Omega)$ which contradicts that $\|h_k\|_2 = 1$. \square

Example 3.21. Let us consider a slight variation of the problem treated in [3]. Let $f, y_d \in C(\bar{\Omega})$, $g \in C^2(\mathbb{R})$ with $g_y \geq 0$ and g_{yy} locally Lipschitz. Consider the following data for (\mathcal{CP}) ,

$$(3.39) \quad \ell(x, y, u) = \frac{1}{2}|u|^2 + \frac{1}{2}(y - y_d(x))^2, \quad \varphi(x, y, u) = g(y) + u + f.$$

By Remark 1.5, the above problem admits at least one solution. Moreover, it is easy to see (see e.g. [3, 10]) that for $\bar{u} \in \mathcal{K}$ the associated quadratic form $Q_2[\bar{u}]$ is a Legendre form. Therefore, since the Hamiltonian for this problem is strictly convex with respect to the control variable, we have that $H_u(x)v \geq 0$ for all $v \in T_{\mathcal{K}(x)}(\bar{u}(x))$ a.e. in Ω together with (3.31) are a sufficient condition for the local quadratic growth on \mathcal{K} in the strong sense. Note that even in this particular convex case, our sufficient condition for strong optimality is still of interest.

Our aim now is to extend to the strong sense the second order sufficient condition in [18], which is stated in terms of a larger cone than $C_{\mathcal{K}}(\bar{u})$ but the assumption for $Q_2[\bar{u}]$ of being a Legendre form is not needed. For $\tau > 0$ define the *strongly active set*

$$(3.40) \quad A^\tau(\bar{u}) := \{x \in \Omega \mid |H_u(x)| > \tau\},$$

and the τ -critical cone

$$(3.41) \quad C_{\mathcal{K}}^\tau(\bar{u}) := \{v \in T_{\mathcal{K}}(\bar{u}) \mid v(x) = 0 \text{ for } x \in A^\tau(\bar{u})\}.$$

For the reader's convenience, we reproduce [18, Theorem 2] adapted to our setting. Let us remark that the result holds true under more general assumptions (more precisely, measurability conditions over the data, instead of our continuity assumptions in **(H1)**-**(H3)**). Moreover, the result in [18] is stated for Neumann boundary controls, but the technique is quite similar for distributed controls and Dirichlet boundary conditions.

Theorem 3.22. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Assume that $H_u(x)v \geq 0$ for all $v \in T_{\mathcal{K}(x)}(\bar{u}(x))$ a.e. in Ω and that there exist $\tau, \alpha > 0$ such that the following second order condition holds true:*

$$(3.42) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}^\tau(\bar{u}).$$

Then J has local quadratic growth on \mathcal{K} at \bar{u} in the weak sense.

We have the following extension of Theorem 3.22 to the strong sense.

Theorem 3.23. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Assume that the strict Pontryagin inequality holds at \bar{u} and that there exist $\tau, \alpha > 0$ such that the following second order condition holds true:*

$$(3.43) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}^\tau(\bar{u}).$$

Then J has local quadratic growth on \mathcal{K} at \bar{u} in the strong sense.

Proof. The beginning of the proof is exactly as in Theorem 3.20. The rest of the proof is a slight modification of the proof in [33] for the weak case. Thus, by (3.34), we may assume that the following inequality holds true

$$(3.44) \quad \int_{A_k} H_u(x) \delta_{A_k} u(x) dx + \frac{1}{2} Q_2[\bar{u}](\delta_{A_k} u) \leq o(\|\delta_{A_k} u\|_2^2).$$

Define $v_k^0(x) := \mathbf{1}_{\{A_k \setminus A^\tau(\bar{u})\}}(x) \delta_{A_k} u(x)$ (where $\mathbf{1}_A$ denotes the indicator function of A) and $v_k^1(x) := \delta_{A_k} u(x) - v_k^0(x)$. Obviously $v_k^0 \in C_{\mathcal{K}}^\tau(\bar{u})$. Using (2.36) and (3.43), we get

$$(3.45) \quad \tau \int_{A_k \cap A^\tau(\bar{u})} |v_k^1| dx + \frac{1}{2} \alpha \|v_k^0\|_2^2 + \frac{1}{2} Q_2[\bar{u}](v_k^1) + \hat{Q}_2[\bar{u}](v_k^0, v_k^1) \leq o(\|\delta_{A_k} u\|_2^2).$$

There exists $c_1 > 0$ such that $\hat{Q}_2[\bar{u}](v_k^0, v_k^1) \geq -c_1 \|v_k^0\|_2 \|v_k^1\|_2$. With Young's inequality, we get

$$(3.46) \quad -Q_2[\bar{u}](v_k^0, v_k^1) \leq \frac{\alpha}{4} \|v_k^0\|_2^2 + c_2 \|v_k^1\|_2^2 \leq \frac{\alpha}{4} \|v_k^0\|_2^2 + c_2 \|v_k^1\|_\infty \|v_k^1\|_1,$$

for some $c_2 > 0$. On the other hand, there exists $c_3 > 0$ such that

$$(3.47) \quad \frac{1}{2} Q_2[\bar{u}](v_k^1) \geq -c_3 \|v_k^1\|_2^2 \geq -c_3 \|v_k^1\|_\infty \|v_k^1\|_1.$$

Recall that $\|\delta_{A_k} u\|_\infty \rightarrow 0$, so we can choose k large such that $\|v_k^1\|_\infty (c_2 + c_3) \leq \tau/2$. Therefore, combining inequalities (3.46)-(3.47) with (3.45) we easily get that

$$\min \left\{ \frac{\tau}{2}, \frac{\alpha}{4} \right\} \|\delta_{A_k} u\|_2^2 \leq \frac{\tau}{2} \int_{A_k \cap A^\tau(\bar{u})} |v_k^1|^2 dx + \frac{\alpha}{4} \int_{A_k \setminus A^\tau(\bar{u})} |v_k^0|^2 dx \leq o(\|\delta_{A_k} u\|_2^2),$$

which gives the desired contradiction. \square

3.4. Characterization of local quadratic growth in the strong sense. We now state the main result of the article, which characterizes local quadratic growth in the strong sense. Note that in particular, the sufficient condition does not need the assumption that $Q_2[\bar{u}]$ is a Legendre form (as in Theorem 3.20) or that it is uniformly positive on $C_{\mathcal{K}}^\tau(\bar{u})$ (as in Theorem 3.23).

Theorem 3.24. *Suppose that (H1)-(H3) hold true and let $\bar{u} \in \mathcal{K}$. Then J has local quadratic growth on \mathcal{K} at \bar{u} in the strong sense if and only if the the Hamiltonian satisfies the global quadratic growth property at \bar{u} and there exists $\alpha > 0$ such that the following second order condition holds true:*

$$(3.48) \quad Q_2[\bar{u}](v) \geq \alpha \|v\|_2^2, \quad \text{for all } v \in C_{\mathcal{K}}(\bar{u}).$$

The proof needs some preparation. Again, the main ingredient of the proof is the decomposition result in Theorem 2.5. However, the choice of the sets A_k and B_k is slightly different from the one used in the preceding results. It takes into account the degeneracy of the so-called Hoffman constants associated with the pointwise critical cone C_x , defined in (3.22). By Hoffman's Lemma [24] for each $x \in \Omega$ there exists a smallest possible (finite) nonnegative number κ_x , called the Hoffman constant, such that

$$(3.49) \quad \text{dist}(v, C_x) \leq \kappa_x \left[|H_u(x)v| + \mathbf{1}_{\{\bar{u}(x)=a(x)\}}(-v)_+ + \mathbf{1}_{\{\bar{u}(x)=b(x)\}}v_+ \right],$$

where for $x \in \mathbb{R}$ we set $(x)_+ := \max\{x, 0\}$. Since these Hoffman constants play an important role in the analysis, we compute them. It is easily checked that

$$(3.50) \quad \begin{cases} C_x = T_{\mathcal{K}(x)}, & \kappa_x = 1 & \text{if } H_u(x) = 0, \\ C_x = \{0\}, & \kappa_x \leq \max(1, |H_u(x)|^{-1}) & \text{otherwise.} \end{cases}$$

Proof of Theorem 3.24. Since by Theorem 3.17(ii) the condition is necessary, we only need to prove that is also sufficient. In order to do this, we proceed as follows: a) We first essentially repeat step (a) of the proof of Theorem 3.20, with a slightly different choice of the sets A_k and B_k . If the conclusion does not hold, let $u_k \in \mathcal{K}$ be such that $\|y_k - \bar{y}\|_{\infty} \rightarrow 0$ and (3.32) holds. Setting $\delta_k u := u_k - \bar{u}$ we have that $\|\delta_k u\|_2 \rightarrow 0$. Remind that κ_x denotes the Hoffman constants of the pointwise cones C_x . For some sequence $\varepsilon_k \downarrow 0$ to be specified later, consider the measurable sets

$$(3.51) \quad \begin{aligned} B_k^1 &:= \left\{ x \in \Omega : |u_k(x) - \bar{u}(x)| \geq \sqrt{\|\delta_k u\|_2} \right\}, & B_k^2 &:= \{x \in \Omega : \kappa_x \geq 1/\varepsilon_k\}, \\ B_k &:= B_k^1 \cup B_k^2, \end{aligned}$$

and $A_k := \Omega \setminus B_k$. Since $|B_k^i| \rightarrow 0$ for $i = 1, 2$, we have that $|B_k| \rightarrow 0$, and therefore, (3.34) holds. Fix σ_{A_k} and σ_{B_k} as in (3.35) and define $h_k := \delta_{A_k} u / \sigma_{A_k}$. If (for a subsequence) $\sigma_{A_k} = o(\sigma_{B_k})$, we obtain a contradiction in the same way. So we may assume that $\sigma_{B_k} = O(\sigma_{A_k})$, and we obtain that (3.37)-(3.38) hold.

b) We make the decomposition

$$(3.52) \quad h_k = \hat{h}_k + \tilde{h}_k, \quad \text{where } \hat{h}_k(x) := P_{C_x}(h_k(x)), \quad \text{for a.a. } x \in \Omega,$$

where $P_{C_x}(\cdot)$ denotes the projection on C_x . Since the multifunction $x \in \Omega \rightarrow C_x$ is convex and measurable, classical results (see e.g. [30]) imply that \hat{h}_k is measurable. By the definition of A_k , and since u_k is feasible, we have that the contribution of the control constraints to the estimate of the distance to C_x in (3.49) is zero, and $H_u(x)h_k(x) \geq 0$, proving that

$$(3.53) \quad |\tilde{h}_k(x)| = \text{dist}(h_k(x), C_x) \leq \frac{1}{\varepsilon_k} H_u(x)h_k(x).$$

With (3.37) we deduce that

$$(3.54) \quad \|\tilde{h}_k\|_1 = O\left(\frac{\sigma_{A_k}}{\varepsilon_k}\right).$$

Since a projection is non expansive, we also have $|\tilde{h}_k(x)| \leq |h_k(x)|$ for a.a. $x \in \Omega$. Therefore

$$(3.55) \quad \sigma_{A_k} \|\tilde{h}_k\|_\infty \leq \sigma_{A_k} \|h_k\|_\infty = \|\delta_{A_k} u\|_\infty.$$

We let $\varepsilon_k := \|\delta_k u\|_2^{1/4}$, that converges to 0 as required. Using that $\|\delta_{A_k} u\|_\infty \leq \sqrt{\|\delta_k u\|_2}$, we get

$$(3.56) \quad \|\tilde{h}_k\|_2^2 \leq \|\tilde{h}_k\|_\infty \|\tilde{h}_k\|_1 = O\left(\frac{\|\delta_{A_k} u\|_\infty \sigma_{A_k}}{\varepsilon_k}\right) = O(\varepsilon_k).$$

By (3.52) and (3.56) we obtain that $\|h_k - \hat{h}_k\|_2 \rightarrow 0$ and so $\|\hat{h}_k\|_2 \rightarrow 1$. Combining with (3.38) and using (3.48) we deduce that $\alpha \|\hat{h}_k\|_2 \leq Q_2[\bar{u}](\hat{h}_k)^2 \leq o(1)$, which is impossible. \square

Remark 3.25. (i) If $\min(b - a) > 0$, an extension of the analysis in [9] to our framework (based on similar arguments), gives that (3.48) implies the a.e. local quadratic growth for the Hamiltonian. Thus, by lemma 3.16, we have that the strict Pontryagin inequality at \bar{u} together with (3.48) imply the global quadratic growth for the Hamiltonian.

(ii) Our analysis excludes the case when the Hamiltonian can have multiple minima, which happens when the optimal control is discontinuous. However, we note that, even in the ODE setting, the second order analysis in this case is quite involved [27], and extending this analysis to the elliptic setting seems to be rather difficult.

4. APPENDIX

In this section we prove the technical lemmas in section 2.

Proof of Lemma 2.1. For convenience, we will omit the dependence of x in some parts of the proof. The equation satisfied by δy , with Dirichlet boundary condition, can be written as

$$-\Delta \delta y(x) + \varphi(x, y_u, u) - \varphi(x, \bar{y}, u) + \delta \varphi(x) = 0, \quad \text{for } x \in \Omega.$$

Equivalently,

$$-\Delta \delta y + \left[\int_0^1 \varphi_y(x, \bar{y} + \theta \delta y, u) d\theta \right] \delta y = O(|\delta u|), \quad \text{for } x \in \Omega.$$

Thus, the estimates for δy , as well as those for z_1 , follow directly from Theorem 1.1. The equation satisfied by d_1 , with Dirichlet boundary condition, becomes:

$$(4.1) \quad -\Delta d_1 + \varphi_y(x) d_1 + \varphi(x, y_u, u) - \varphi(x, \bar{y}, u) - \varphi_y(x) \delta y = 0.$$

Introducing $\varphi_y(x, \bar{y}, u)$ in $\varphi(x, y, u) - \varphi(x, \bar{y}, u) - \varphi_y(x) \delta y$, easily yields

$$\varphi(x, y, u) - \varphi(x, \bar{y}, u) - \varphi_y(x) \delta y = O(|\delta y|^2 + |\delta y \delta u|),$$

hence the equation satisfied by d_1 rewrites:

$$-\Delta d_1 + \varphi_y(x) d_1 = O(|\delta y|^2 + |\delta y \delta u|).$$

Theorem 1.1 implies that for $s \in (n/2, \infty)$

$$\begin{aligned} \|d_1\|_1 &\leq O(\|(\delta y)^2\|_1 + \|\delta y \delta u\|_1) \\ &= O(\|\delta y\|_1 \|\delta y\|_\infty + \|\delta y\|_\infty \|\delta u\|_1) \\ &= O(\|\delta u\|_1 \|\delta u\|_s). \end{aligned}$$

Finally, taking $s = 2$ in Theorem 1.1 gives

$$\|d_1\|_2 = O(\|\delta y \delta u\|_2 + \|(\delta y)^2\|_2) = O(\|\delta y\|_\infty \|\delta u\|_2).$$

□

Proof of Lemma 2.3. Since $d_2 = d_1 - z_2$, combining (4.1) with (2.11) yields

$$-\Delta d_2 + \varphi_y(x) d_2 + \varphi(x, y, u) - \varphi(x, \bar{y}, u) - \varphi_y(x, \bar{y}, u) \delta y + \delta \varphi_y(x) (\delta y - z_1) - \frac{1}{2} \varphi_{yy}(x) z_1^2 = 0.$$

Now, by a Taylor expansion, we have:

$$\varphi(x, y, u) - \varphi(x, \bar{y}, u) - \varphi_y(x, \bar{y}, u) \delta y = \frac{1}{2} \varphi_{yy}(x, \bar{y}, u) (\delta y)^2 + O(|\delta y|^3),$$

and as φ_y is locally Lipschitz, $\delta \varphi_y(\delta y - z_1) = O(|\delta u| |\delta y - z_1|) = O(|\delta u| |d_1|)$. Therefore,

$$-\Delta d_2 + \varphi_y(x) d_2 + \frac{1}{2} \varphi_{yy}(x, \bar{y}, u) (\delta y)^2 - \frac{1}{2} \varphi_{yy}(x) z_1^2 = O(|\delta y|^3 + |\delta u| |d_1|).$$

Now, we have $\varphi_{yy}(x, \bar{y}, u) (\delta y)^2 = \varphi_{yy}(x) (\delta y)^2 + O(|\delta u| |\delta y|^2)$. Therefore, we obtain:

$$-\Delta d_2 + \varphi_y(x) d_2 + \frac{1}{2} \varphi_{yy}(x) [(\delta y)^2 - z_1^2] = O(|\delta y|^3 + |\delta u| |d_1| + |\delta u| \delta y^2).$$

As $d_1 = \delta y - z_1$, we get:

$$-\Delta d_2 + \varphi_y(x) d_2 = O(|\delta y|^3 + |\delta u| |d_1| + |\delta u| |\delta y|^2 + |d_1| |\delta y + z_1|).$$

From the Theorem 1.1 we get the inequality:

$$\|d_2\|_1 = O(\|(\delta y)^3\|_1 + \|\delta u d_1\|_1 + \|\delta u (\delta y)^2\|_1 + \|d_1 \delta y\|_1 + \|d_1 z_1\|_1).$$

Using the previous estimates, we easily obtain the result. □

Proof of Lemma 3.11 (i). Set

$$\hat{T}(\bar{u}) := \{v \in L^2(\Omega) \mid v(x) \in T_{\mathcal{K}(x)}(\bar{u}(x)) \text{ for a.a. } x \in \Omega\},$$

and let $v \in T_{\mathcal{K}}(\bar{u})$. By definition, there exists $r : \mathbb{R} \rightarrow L^2(\Omega)$, with $\|r(\sigma)\|_2/\sigma \rightarrow 0$ as $\sigma \downarrow 0$, such that for small σ

$$(4.2) \quad \bar{u}(x) + \sigma v(x) + r(\sigma)(x) \in \mathcal{K}(x), \quad \text{for a.a. } x \in \Omega.$$

Since, up to subsequence, $|r(\sigma)(x)|/\sigma \rightarrow 0$ for a.a. $x \in \Omega$, relation (4.2) implies that $v \in \hat{T}(\bar{u})$. Conversely, let $v \in \hat{T}(\bar{u})$ and for $\varepsilon > 0$ set

$$(4.3) \quad v_\varepsilon := \varepsilon^{-1} [P_{\mathcal{K}}(\bar{u} + \varepsilon v) - \bar{u}],$$

where $P_{\mathcal{K}}(\cdot)$ denotes the orthogonal projection in $L^2(\Omega)$ onto \mathcal{K} . Clearly, $v_\varepsilon \in R_{\mathcal{K}}(\bar{u})$ and classical results (see e.g. [30]) yield that v_ε is measurable and given by

$$(4.4) \quad v_\varepsilon(x) := \varepsilon^{-1} [P_{\mathcal{K}(x)}(\bar{u}(x) + \varepsilon v(x)) - \bar{u}(x)] \quad \text{for a.a. } x \in \Omega.$$

Clearly, $v_\varepsilon(x) \in R_{\mathcal{K}(x)}(\bar{u}(x))$ and using that $v(x) \in T_{\mathcal{K}(x)}(\bar{u}(x))$ we get $v_\varepsilon(x) \rightarrow v(x)$ for a.a. $x \in \Omega$. Finally, using that $|v_\varepsilon(x)| \leq |v(x)|$, we obtain the convergence in $L^2(\Omega)$ and so $v \in T_{\mathcal{K}}(\bar{u})$.

Proof of (ii): Set $\hat{N}(\bar{u}) := \{v \in L^2(\Omega) \mid v(x) \in N_{\mathcal{K}(x)}(\bar{u}(x)) \text{ for a.a. } x \in \Omega\}$ and let $v^* \in \hat{N}(\bar{u})$. Using that $N_{\mathcal{K}(x)}(\bar{u}(x))$ is the polar cone of $T_{\mathcal{K}(x)}(\bar{u}(x))$, assertion (i) yields $v^* \in N_{\mathcal{K}}(\bar{u})$. Conversely, let $v^* \in N_{\mathcal{K}}(\bar{u})$ and let v_1, v_2 be such that

$$v^*(x) = v_1(x) + v_2(x), \quad v_1(x) v_2(x) = 0, \quad v_1(x) \in T_{\mathcal{K}(x)}(\bar{u}(x)), \quad v_2(x) \in N_{\mathcal{K}(x)}(\bar{u}(x)),$$

for a.a. $x \in \Omega$. Assertion (i) implies that $v_1 \in T_{\mathcal{K}}(\bar{u})$ and so

$$\|v_1\|_2^2 = \int_{\Omega} v^*(x)v_1(x)dx \leq 0,$$

which gives that $v^* = v_2$. □

REFERENCES

1. R.A. Adams, *Sobolev spaces*, Academic Press, New York, 1975.
2. H. Benker and M. Handschug, *An algorithm for abstract optimal control problems using maximum principles and applications to a class of distributed parameter systems*, Optimal control (Freiburg, 1991), Internat. Ser. Numer. Math., vol. 111, Birkhäuser, Basel, 1993, pp. 31–42. MR 1297996 (95g:49050)
3. J. F. Bonnans, *Second order analysis for control constrained optimal control problems of semilinear elliptic systems*, Appl. Math. Optim. **38-3** (1998), 303–325.
4. ———, *Pontryagin's principle for the optimal control of semilinear elliptic systems with state constraints*, Proceedings of the 30th Conference on Decision and Control, Brighthon, England **2** (December 1991), 1976–1979.
5. J.F. Bonnans, *On an algorithm for optimal control using Pontryagin's maximum principle*, SIAM J. Control Optim. **24** (1986), no. 3, 579–588. MR 838058 (87f:49040)
6. J.F. Bonnans and E. Casas, *Un principe de Pontryagin pour le contrôle des systèmes semilinéaires elliptiques*, Journal of Differential Equations **90** (1991), 288–303.
7. ———, *Maximum principles in the optimal control of semilinear elliptic systems*, 5th IFAC Conference on Distributed Parameter Systems, A. El Jai ed. (Perpignan), Pergamon Press **33-1** (26–29 June 1989), 274–298.
8. J.F. Bonnans and N.P. Osmolovskii, *Second-order analysis of optimal control problems with control and initial-final state constraints*, Journal of Convex Analysis **17** (2010), no. 3, 885–913.
9. ———, *Characterization of a local quadratic growth of the hamiltonian for control constrained optimal control problems*, Dynamics of Continuous, Discrete and Impulsive Systems **19** (2012), 1–16.
10. J.F. Bonnans and A. Shapiro, *Perturbation analysis of optimization problems*, Springer-Verlag, New York, 2000.
11. J.F. Bonnans and F.J. Silva., *Asymptotic expansion for the solutions of control constrained semilinear elliptic problems with interior penalties*, SIAM J. on Control and Optimization **49** (2011), 2494–2517.
12. J.F. Bonnans and H. Zidani, *Optimal control problems with partially polyhedral constraints*, SIAM J. Control Optim. **37** (1999), no. 6, 1726–1741 (electronic).
13. E. Casas, J.C. de Los Reyes, and F. Tröeltzsch, *Sufficient second-order optimality conditions for semilinear control problems with pointwise state constraints*, SIAM J. Optim. **19-2** (2008), 616–643.
14. E. Casas and F. Tröeltzsch, *Second order necessary optimality conditions for some state-constrained control problems of semilinear elliptic equations*, Appl. Math. Optim. **39** (1999), 211–228.
15. ———, *Second order necessary and sufficient optimality conditions for optimization problems and applications to control theory*, SIAM J. Optim. **13** (2002), 406–431.
16. ———, *Recent advances in the analysis of pointwise state-constrained elliptic optimal control problems*, ESAIM Cont. Optim. Calc. Var. **16-3** (2010), 581–600.
17. ———, *Second order analysis for optimal control problems: Improving results expected from abstract theory*, SIAM J. on Optimization **22** (2012), no. 1, 261–279.
18. E. Casas, F. Tröeltzsch, and A. Unger, *Second order sufficient optimality conditions for a nonlinear elliptic control problem*, Journal for Analysis and its Applications **15** (1996), 687–707.
19. ———, *Second order sufficient optimality conditions for some state-constrained control problems of semilinear elliptic equations*, SIAM J. Control Optim. **38** (2000), 369–1391.
20. L.C. Evans, *Partial differential equations*, Amer. Math Soc., Providence, RI, 1998, Graduate Studies in Mathematics 19.

21. A. V. Fursikov, *Optimal control of distributed systems. theory and applications*, Translations of Mathematical Monographs, Vol.187, AMS, Providence, Rhode Island, 1999.
22. D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order, 2nd edition*, Springer Verlag, Berlin, 1983.
23. A. Haraux, *How to differentiate the projection on a convex set in Hilbert space. some applications to variational inequalities*, J. Mathematical Society of Japan **29** (1977), 615–631.
24. A. Hoffman, *On approximate solutions of systems of linear inequalities*, Journal of Research of the National Bureau of Standards, Section B, Mathematical Sciences **49** (1952), 263–265.
25. X. Li and J. Yong, *Optimal control theory for infinite dimensional systems*, Birkhäuser, 1994, Systems and Control: Foundations and Applications.
26. F. Mignot, *Contrôle dans les inéquations variationnelles*, Journal of Functional Analysis **22** (1976), 25–39.
27. A. Milyutin and N. Osmolovskii, *Calculus of variations and optimal control*, AMS., 1998, Systems and Control: Foundations and Applications.
28. L. Pontryagin, V. Boltianski, R. Gamkrelidze, and E. Michtchenko, *The mathematical theory of optimal processes*, Wiley Interscience, New York, 1962. MR 0358482 (50 #10943)
29. U. E. Raitums, *The maximum principle in optimal control problems for an elliptic equation*, Z. Anal. Anwendungen **5** (1986), no. 4, 291–306. MR 855633 (87j:49048)
30. R. T. Rockafellar, *Integral functionals, normal integrands and measurable selections*, Nonlinear operators and the calculus of variations (Summer School, Univ. Libre Bruxelles, Brussels, 1975), Springer, Berlin, 1976, pp. 157–207. Lecture Notes in Math., Vol. 543.
31. A. Rösch and F. Tröltzsch, *Sufficient second-order optimality conditions for an elliptic optimal control problem with pointwise control-state constraints.*, SIAM J. Optim. **17-3** (2006), 776–794.
32. G. Stampacchia, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier (Grenoble) **15** (1965), 189–258.
33. F. Tröltzsch, *Optimal Control of Partial Differential Equations - Theory, Methods and Applications*, Graduate Studies in Mathematics, Vol. 112. American Mathematical Society, Providence, Rhode Island, 2010.

UNIVERSITÉ MONTPELLIER 2, CASE COURRIER 051, 34095 MONTPELLIER CEDEX 5, FRANCE.

E-mail address: `tbayen@math.univ-montp2.fr`

INRIA-SACLAY AND CMAP, ÉCOLE POLYTECHNIQUE, 91128 PALAISEAU, FRANCE.

E-mail address: `Frederic.Bonnans@inria.fr`

DIPARTIMENTO DI MATEMATICA GUIDO CASTELNUOVO, 00185 ROME, ITALY.

Current address: XLIM - UMR CNRS 7252, Université de Limoges, 87060 Limoges, France.

E-mail address: `francisco.silva@unilim.fr`