



**HAL**  
open science

# A New Closed-Form Information Metric for Shape Analysis

Adrian Peter, Anand Rangarajan

► **To cite this version:**

Adrian Peter, Anand Rangarajan. A New Closed-Form Information Metric for Shape Analysis. 1st MICCAI Workshop on Mathematical Foundations of Computational Anatomy: Geometrical, Statistical and Registration Methods for Modeling Biological Shape Variability, Oct 2006, Copenhagen, Denmark. pp.100-101. inria-00635898

**HAL Id: inria-00635898**

**<https://inria.hal.science/inria-00635898>**

Submitted on 26 Oct 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A New Closed-Form Information Metric for Shape Analysis

Adrian Peter<sup>1</sup> and Anand Rangarajan<sup>2</sup>

<sup>1</sup>Dept. of ECE, <sup>2</sup>Dept. of CISE, University of Florida, Gainesville, FL

**Abstract.** Recently, a unifying framework was introduced for shape matching that uses mixture-models and the Fisher-Rao metric to couple both the shape representation and deformation. A fundamental drawback of the Fisher-Rao metric is that it is NOT available in closed-form for the mixture models making shape comparisons computationally very expensive. Here, we propose a new Riemannian metric based on generalized  $\phi$ -entropy measures. In sharp contrast to the Fisher-Rao metric, our new metric is available in closed-form.

## 1 Introduction

Shape analysis is a key ingredient to many medical imaging applications that seek to study the intimate relationship between the form and function of medical and biological structures. In recent work [1], it was shown that shape representation and deformation can be unified within a probabilistic framework – using a mixture of densities to represent landmark shapes and then directly using the Fisher information matrix to establish an intrinsic, Riemannian metric and subsequently a shape distance measure.

To address many of the computational inefficiencies that arise when using the standard Fisher-Rao information metric, the current work introduces a new Riemannian metric based on the generalized notion of a  $\phi$ -entropy functional.

## 2 Differential Metrics for Parametric Distributions

The parametric, GMM representation for 2-D shapes is given by

$$p(\mathbf{x}|\theta) = \frac{1}{2\pi\sigma^2 K} \sum_{a=1}^K \exp\left\{-\frac{\|\mathbf{x} - \phi_a\|^2}{2\sigma^2}\right\} \quad (1)$$

where  $\theta$  is our set of landmarks,  $\phi_a = [\theta^{(2a-1)}, \theta^{(2a)}]^T$ ,  $\mathbf{x} = [x^{(1)}, x^{(2)}]^T \in \mathbb{R}^2$  and equal weight priors are assigned to all components, i.e.  $\frac{1}{K}$ . Basically, a shape with  $K$  landmarks is represented as a  $K$ -component GMM where the landmarks are the means of each component.

Burbea and Rao demonstrated that the notion of distances between parametric models can be extended to a large class of generalized metrics [2] using the  $\phi$ -entropy functional

$$H_\phi(p) = - \int_{\chi} \phi(p) dx \quad (2)$$

where  $\chi$  is the measurable space (for our purposes  $\mathbb{R}^2$ ), and  $\phi$  is a  $C^2$ -convex function defined on  $\mathbb{R}_+ \equiv [0, \infty)$ . The metric on the parameter space is obtained by finding the Hessian of (2) along a direction in its tangent space. This directly leads to the following differential metric satisfying Riemannian metric properties

$$ds_\phi^2(\theta) = -\Delta_\theta H_\phi(p) = \sum_{i,j=1}^n g_{i,j}^\phi d\theta^i d\theta^j, \quad (3)$$

where

$$g_{i,j}^{\phi} = \int_{\mathcal{X}} \phi''(p) \left( \frac{\partial p}{\partial \theta^i} \right) \left( \frac{\partial p}{\partial \theta^j} \right) d\mathbf{x}. \quad (4)$$

The metric tensor in (4) is called the  $\phi$ -entropy matrix. We can define distances between probability densities by finding a geodesic between their parameter values as determined by (3).

The new metric can be obtained by selecting a  $\phi$  such that  $\phi''$  becomes a constant in (4). In [3], Havrda and Charvát introduced the notion of a  $\alpha$ -order entropy using the convex function

$$\phi(p) = (\alpha - 1)^{-1} (p^{\alpha} - p), \alpha \neq 1. \quad (5)$$

We set  $\alpha = 2$  which results in  $\frac{1}{2}\phi'' = 1$ . The one-half scaling factor does not impact the metric properties. The new metric is defined as

$$g_{i,j}^{\alpha} = \int_{\mathcal{X}} \left( \frac{\partial p}{\partial \theta^i} \right) \left( \frac{\partial p}{\partial \theta^j} \right) d\mathbf{x} \quad (6)$$

and we refer to it as the  $\alpha$ -order entropy metric tensor.

The new metric can be used to find the distance between two shapes represented as  $p(\mathbf{x}|\theta_1)$  and  $p(\mathbf{x}|\theta_2)$ . The geodesic can be obtained by solving

$$g_{ki}^{\alpha} \ddot{\theta}^i + \Gamma_{k,i,j} \dot{\theta}^i \dot{\theta}^j = 0. \quad (7)$$

Using this new metric, we now have closed-form solutions to  $g_{i,j}^{\alpha}$  and  $\frac{\partial g_{k,j}^{\alpha}}{\partial \theta^i}$ .

### 3 Experimental Results and Analysis

For applications in medical imaging, we have evaluated both metrics and compared to the standard thin-plate spline (TPS) bending energy on real data consisting of nine corpora callosa with 63-landmarks each. (Please see the conference paper for test results and analysis.)

The new  $\phi$ -entropy metric avoids an  $O(N^2)$  computational hit on the metric tensor computation and gains another  $O(N^2)$  savings when computing the derivative of the metric tensor.

### 4 Conclusion

In this paper, we introduced a new Riemannian metric for landmark-based shape matching that combines both shape representation and deformation. This improved on the previous work of [1] where the Fisher-Rao metric was used to establish an intrinsic metric between landmark shapes. The new  $\alpha$ -order entropy metric enables us to obtain closed-form solutions to the metric tensor and its derivatives. Our new approach also illustrated the possibilities of using information metrics besides Fisher-Rao and their benefits.

### References

1. Peter, A., Rangarajan, A.: Shape matching using the Fisher-Rao Riemannian metric: Unifying shape representation and deformation. ISBI 2006 (2006)
2. Burbea, J., Rao, R.: Entropy differential metric, distance and divergence measures in probability spaces: A unified approach. Journal of Multivariate Analysis **12** (1982) 575–596
3. Havrda, M.E., Charvát, F.: Quantification method of classification processes: Concept of structural  $\alpha$ -entropy. Kybernetika **3** (1967) 30–35