

Fractional colorings of cubic graphs with large girth

František Kardoš, Daniel Král, Jan Volec

► **To cite this version:**

František Kardoš, Daniel Král, Jan Volec. Fractional colorings of cubic graphs with large girth. Siam Journal on Discrete Mathematics, Society for Industrial and Applied Mathematics, 2011, 25 (3), pp.1454-1476. <http://epubs.siam.org/sidma/resource/1/sjdmec/v25/i3/p1454_s1?isAuthorized=no>. <inria-00635910>

HAL Id: inria-00635910

<https://hal.inria.fr/inria-00635910>

Submitted on 9 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fractional colorings of cubic graphs with large girth

František Kardoš* Daniel Král'† Jan Volec‡

Abstract

We show that every (sub)cubic n -vertex graph with sufficiently large girth has fractional chromatic number at most 2.2978 which implies that it contains an independent set of size at least $0.4352n$. Our bound on the independence number is valid for random cubic graphs as well as it improves existing lower bounds on the maximum cut in cubic graphs with large girth.

1 Introduction

An *independent set* is a subset of vertices such that no two of them are adjacent. The *independence number* $\alpha(G)$ of a graph G is the size of the largest independent set in G . In this paper, we study independent sets in cubic graphs with large girth. Recall that a graph G is *cubic* if every vertex

*Institute of Mathematics, Faculty of Science, University of Pavol Jozef Šafárik, Jessenná 5, 041 54 Košice, Slovakia. E-mail: frantisek.kardos@upjs.sk. This author was supported by Slovak Research and Development Agency under the contract no. APVV-0007-07.

†Department of Applied Mathematics and Institute for Theoretical Computer Science (ITI), Faculty of Mathematics and Physics, Charles University, Malostranské náměstí 25, 118 00 Prague 1, Czech Republic. E-mail: kral@kam.mff.cuni.cz. Institute for Theoretical computer science is supported as project 1M0545 by Czech Ministry of Education. This author was supported by the grant GACR 201/09/0197 and in part was also supported by the grant GAUK 60310.

‡Department of Applied Mathematics, Faculty of Mathematics and Physics, Charles University, Malostranské náměstí 25, 118 00 Prague 1, Czech Republic. E-mail: volec@kam.mff.cuni.cz. This author was supported by the grant GACR 201/09/0197.

of G has degree three, G is *subcubic* if every vertex has degree at most three, and the *girth* of G is the length of the shortest cycle of G .

A *fractional coloring* of a graph G is an assignment of weights to independent sets in G such that for each vertex v of G , the sum of weights of the sets containing v is at least one. The *fractional chromatic number* $c_f(G)$ of G is the minimum sum of weights of independent sets forming a fractional coloring. The main result of this paper asserts that if G is a cubic graph with sufficiently large girth, then there exists a probability distribution on its independent sets such that each vertex is contained in an independent set drawn based on this distribution with probability at least 0.4352. This implies that every n -vertex cubic graph G with large girth contains an independent set with at least $0.4352n$ vertices and its fractional chromatic number is at most 2.2978 (to see the latter, consider the probability distribution and assign each independent set I in G a weight equal to $cp(I)$ where $p(I)$ is the probability of I and $c = 0.4352^{-1}$). In addition, our lower bound on the independence number also translates to random cubic graphs.

Let us now survey previous results in this area. Inspired by Nešetřil's Pentagon Conjecture [9], Hatami and Zhu [2] showed that every cubic graph with sufficiently large girth has fractional chromatic number at most $8/3$. For the independence number, Hoppen [5] showed that every n -vertex cubic graph with sufficiently large girth has independence number at least $0.4328n$; this bound matches an earlier bound of Wormald [11], also independently proven by Frieze and Suen [3], for random cubic graphs. The bound for random cubic graphs was further improved by Duckworth and Zito [1] to $0.4347n$, providing an improvement of the bound unchallenged for almost 15 years.

1.1 Random cubic graphs

The independence numbers of random cubic graphs and cubic graphs with large girth are closely related. Here, we consider the model where a random cubic graph is randomly uniformly chosen among all n -vertex cubic graphs.

Any lower bound on the independence number for the class of cubic graphs with large girth is also a lower bound for random cubic graphs. First observe that a lower bound on the independence number for cubic graphs with large girth translates to subcubic graphs with large girth: assume that H is a subcubic graph with m_1 vertices of degree one and m_2 vertices of degree two. Now consider an $(2m_1 + m_2)$ -regular graph with high girth and replace each vertex of it with a copy of H in such a way that its edges are incident with

vertices of degree one and two in the copies. The obtained graph G is cubic and has large girth; an application of the lower bound on the independence number for cubic graphs yields that one of the copies of H contains a large independent set, too.

Since a random cubic graph contains asymptotically almost surely only $o(n)$ cycles shorter than a fixed integer g [10], any lower bound for cubic graphs with large girth also applies to random cubic graphs. Conversely, Hoppen and Wormald [12] have recently developed a technique for translating lower bounds (obtained in a specific but quite general way) for the independence number of a random cubic graph to cubic graphs with large girth.

In the other direction, any upper bound on the independence number for a random cubic graph also applies to cubic graphs with large girth (just remove few short cycles from a random graph). The currently best upper bound for random cubic graphs with n vertices and thus for cubic graphs with large girth is $0.455n$ derived by McKay [7]. In [7], McKay mentions that experimental evidence suggests that almost all n -vertex cubic graphs contain independent sets of size $0.439n$; his more recent experiments suggest that the best possible lower bound might be even as large as $0.447n$.

1.2 Maximum cut

Our results also improve known bounds on the size of the maximum cut in cubic graphs with large girth. For a coloring of vertices of a graph G with two colors, the set of edges with end-vertices of different colors is called a *cut*. The *maximum cut* is a cut with the largest number of edges. It is known that if G is an m -edge cubic graph with sufficiently large girth, then its maximum cut has size at least $6m/7 - o(m)$ [13]. On the other hand, there exist m -edge cubic graphs with large girth with the maximum cut of size at most $0.9351m$. This result was first announced by McKay [6]; the proof can be found in [4].

If G has an independent set of size α , then it also has a cut of size 3α (put the independent set on one side and all the other vertices on the other side). So, our result shows that if G is a cubic graph with n vertices and $m = 3n/2$ edges and the girth of G is sufficiently large, then it has a cut of size at least $3 \times 0.4352n = 0.8704m$. This bound can further be improved to $0.8834m$ by considering a randomized procedure tuned up for producing a cut of large size. Since we focus on independent sets in cubic graphs with large girth, we omit details how the improved bound can be obtained.

2 Structure of the proof

Our proof is inspired by the proof of Hoppen from [5]. We develop a procedure for obtaining a random independent set in a cubic graph with large girth. Our main result is the following.

Theorem 1. *There exists $g > 0$ such that for every cubic graph G with girth at least g , there is a probability distribution such that each vertex is contained in an independent set drawn according to this distribution with probability at least 0.4352.*

We describe our procedure for obtaining a random independent set in a cubic graph in Section 3. To analyze its performance, we first focus on its behavior for infinite cubic trees in Section 4. The core of our analysis is the independence lemma (Lemma 6) which is used to simplify the recurrence relation appearing in the analysis. We then show that the procedure can be modified for cubic graphs with sufficiently large girth while keeping its performance. The actual performance of our randomized procedure is based on solving the derived recurrences numerically.

As we have explained in Section 1, Theorem 1 has the following three corollaries.

Corollary 2. *There exists $g > 0$ such that every n -vertex subcubic graph with girth at least g contains an independent set of size at least $0.4352n$.*

Corollary 3. *There exists $g > 0$ such that every cubic graph with girth at least g has the fractional chromatic number at most 2.2978.*

Corollary 4. *There exists $g > 0$ such that every cubic graph G with girth at least g has a cut containing at least $0.8704m$ edges where m is the total number of edges of G .*

3 Description of the randomized procedure

The procedure for creating a random independent set is parametrized by three numbers: the number K of its rounds and probabilities p_1 and p_2 .

Throughout the procedure, vertices of the input graph have one of the three colors: white, blue and red. At the beginning, all vertices are white. In each round, some of the white vertices are recolored in such a way that

red vertices always form an independent set and all vertices adjacent to red vertices as well as some of other vertices (see details below) are *blue*. All other vertices of the input graph are *white*. Red and blue vertices are never again recolored during the procedure. When we talk of a *degree* of a vertex, we mean the number of its white neighbors. Observe that the neighbors of a white vertex that are not white must be blue.

The first round of the procedure is special and differs from the other rounds. As we have already said, at the very beginning, all vertices of the input graph are white. In the first round, we randomly and independently with probability p_1 mark some vertices as active. Active vertices with no active neighbor become red and vertices with at least one active neighbor become blue. In particular, if two adjacent vertices are active, they both become blue (as well as their neighbors). At this point, we should note that the probability p_1 will be very small.

In the second and the remaining rounds, we first color all white vertices of degree zero by red. We then consider all paths formed by white vertices with end vertices of degree one or three and with all inner vertices of degree two. Based on the degrees of their end vertices, we refer to these paths as paths of type $1\leftrightarrow 1$, $1\leftrightarrow 3$ or $3\leftrightarrow 3$. Edges joining two vertices of degree one or three are considered as such paths of length one. Finally, each vertex of degree two is now activated with probability p_2 independently of all other vertices.

For each path of type $1\leftrightarrow 3$, we color the end vertex of degree one with red and we then color all the inner vertices with red and blue in the alternating way. If the neighbor of the end vertex of degree three becomes red, we also color the end vertex of degree three with blue. In other words, we color the end vertex of degree three by blue if the path has an odd length. For a path of type $1\leftrightarrow 1$, we choose randomly one of its end vertices, color it red and color all the remaining vertices of the path with red and blue in the alternating way. We refer to the chosen end vertex as the beginning of this path. Note that whether and which vertices of degree two on the paths of type $1\leftrightarrow 1$ or $1\leftrightarrow 3$ are activated do not influence the above procedure.

Paths of type $3\leftrightarrow 3$ are processed as follows. A path of type $3\leftrightarrow 3$ becomes active if at least one of its inner vertices is activated, i.e., a path of length ℓ is active with probability $1 - (1 - p_2)^{\ell-1}$. Note that paths with no inner vertices, i.e., edges between two vertices of degree three, are never active. For each active path, flip the fair coin to select one of its end vertices of degree three, color this vertex by blue and its neighbor on the path by red. The remaining

inner vertices of the path are colored with red and blue in the alternating way. Again we refer to the chosen end vertex as the beginning of this path. The other end vertex of degree three is colored blue if its neighbor on the path becomes red, i.e., if the path has an even length.

Note that a vertex that has degree three at the beginning of the round cannot become red during the round but it can become blue because of several different paths ending at it.

4 Analysis for infinite cubic trees

Let us start with introducing some notation used in the analysis. For any edge uv of a cubic graph G , $T_{u,v}$ is the component of $G - v$ containing the vertex u ; we sometimes refer to u as to the root of $T_{u,v}$. If G is the infinite cubic tree, then it is a union of $T_{u,v}, T_{v,u}$ and the edge uv . The subgraph of $T_{u,v}$ induced by vertices at distance at most d from u is denoted by $T_{u,v}^d$. Observe that if the girth of G is larger than $2d + 1$, all the subgraphs $T_{u,v}^d$ are isomorphic to the same rooted tree \mathcal{T}^d of depth d . The infinite rooted tree with the root of degree two and all inner vertices of degree three will be denoted as T^∞ .

Since any automorphism of the infinite cubic tree yields an automorphism of the probability space of the vertex colorings constructed by our procedure, the probability that a vertex u has a fixed color after k rounds does not depend on the choice of u . Hence, we can use w_k, b_k and r_k to denote a probability that a fixed vertex of the infinite cubic tree is white, blue and red, respectively, after k rounds. Similarly, $w_k^{(i)}$ denotes the probability that a fixed vertex has degree i after k rounds conditioned by the event that it is white after k rounds, i.e., the probability that a fixed vertex is white and has degree two after k rounds is $w_k \cdot w_k^{(2)}$.

The sets of white, blue and red vertices after k rounds of the randomized procedure described in Section 3 will be denoted by W_k, B_k and R_k , respectively. Similarly, $W_k^{(i)}$ denotes the set of white vertices with degree i after k rounds, i.e., $W_k = W_k^{(0)} \cup W_k^{(1)} \cup W_k^{(2)} \cup W_k^{(3)}$. Finally, $c_k(T_{u,v})$ denotes the coloring of $T_{u,v}$ and $c_k(T_{u,v}^d)$ the coloring of $T_{u,v}^d$ after k rounds. The set \mathcal{C}_k^d will consist of all possible colorings γ of \mathcal{T}^d such that the probability of $c_k(T_{u,v}^d) = \gamma$ is non-zero in the infinite cubic tree (note that this probability does not depend on the edge uv) and such that the root of \mathcal{T}^d is colored white. We extend this notation and use \mathcal{C}_k^∞ to denote all such colorings

of \mathcal{T}^∞ .

Since the infinite cubic tree is strongly edge-transitive, we conclude that the probability that $T_{u,v}$ has a given coloring from \mathcal{C}_k^∞ after k rounds does not depend on the choice of uv . Similarly, the probability that both u and v are white after k rounds does not depend on the choice of the edge uv . To simplify our notation, this event is denoted by $uv \subseteq W_k$. Finally, the probability that u has degree $i \in \{1, 2, 3\}$ after k rounds conditioned by $uv \subseteq W_k$ does also not depend on the choice of the edge uv . This probability is denoted by $q_k^{(i)}$.

4.1 Independence lemma

We now show that the probability that both end-vertices of an edge uv are white after k rounds can be computed as a product of two probabilities. This will be crucial in the proof of the Independence Lemma. To be able to state the next lemma, we need to introduce additional notation. For $\gamma \in \mathcal{C}_{k-1}^\infty$, define

$$P_k(u, v, \gamma) := \mathbf{P}[u \text{ stays white regarding } T_{u,v} \mid c_{k-1}(T_{u,v}) = \gamma \wedge v \in W_{k-1}]$$

where the phrase “ u stays white regarding $T_{u,v}$ ” means

- if $k = 1$, that neither u nor a neighbor of it in $T_{u,v}$ is active, and
- if $k > 1$, that neither
 - u has degree one after $k - 1$ rounds,
 - u has degree two after $k - 1$ rounds and the path from u formed by vertices of degree two in $T_{u,v}$ ends at a vertex of degree one,
 - u has degree two after $k - 1$ rounds, the path from u formed by vertices of degree two in $T_{u,v}$ ends at a vertex of degree three and at least one of the vertices of degree two on this path is activated, nor
 - u has degree three after $k - 1$ rounds and is colored blue because of a path of type $1 \leftrightarrow 3$ or $3 \leftrightarrow 3$ fully contained in $T_{u,v}$.

Informally, this phrase represents that there is no reason for u not to stay white based on the coloring of $T_{u,v}$ and vertices in $T_{u,v}$ activated in the k -th round.

Lemma 5. *Consider the randomized procedure for the infinite cubic tree. Let k be an integer, uv an edge of the tree and γ_u and γ_v two colorings from \mathcal{C}_{k-1}^∞ . The probability*

$$\mathbf{P}[uv \subseteq W_k \mid c_{k-1}(T_{u,v}) = \gamma_u \wedge c_{k-1}(T_{v,u}) = \gamma_v], \quad (1)$$

i.e., the probability that both u and v are white after k rounds conditioned by $c_{k-1}(T_{u,v}) = \gamma_u$ and $c_{k-1}(T_{v,u}) = \gamma_v$, is equal to

$$P_k(u, v, \gamma_u) \cdot P_k(v, u, \gamma_v).$$

Proof. We distinguish the cases $k = 1$ and $k > 1$. If $k = 1$, \mathcal{C}_0^∞ contains a single coloring γ_0 where all vertices are white. Hence, the probability (1) is $\mathbf{P}[uv \subseteq W_1 \mid c_{k-1}(T_{u,v}) = \gamma_0 \wedge c_{k-1}(T_{v,u}) = \gamma_0]$ and it is equal to $\mathbf{P}[uv \subseteq W_1] = (1 - p_1)^6$. On the other hand, $P_1(u, v, \gamma_0) = P_1(v, u, \gamma_0) = (1 - p_1)^3$. The assertion of the lemma follows.

Suppose that $k > 1$. Note that there is a unique coloring of the infinite cubic tree that coincides with γ_u on $T_{u,v}$ and with γ_v on $T_{v,u}$. If u has degree one after $k - 1$ rounds, then (1) is zero as well as $P_k(u, v, \gamma_u) = 0$. If u has degree two and lie on a path of type $1 \leftrightarrow 1$ or $1 \leftrightarrow 3$, then (1) is zero and $P_k(u, v, \gamma_u) = 0$ or $P_k(v, u, \gamma_v) = 0$ depending which of the trees $T_{u,v}$ and $T_{v,u}$ contains the vertex of degree one. If u has degree two and lie on a path of type $3 \leftrightarrow 3$ of length ℓ , then (1) is equal to $(1 - p_2)^{\ell-1}$. Let ℓ_1 and ℓ_2 be the number of vertices of degree two on this path in $T_{u,v}$ and $T_{v,u}$, respectively. Observe that $\ell_1 + \ell_2 = \ell - 1$. Since $P_k(u, v, \gamma_u) = (1 - p_2)^{\ell_1}$ and $P_k(v, u, \gamma_v) = (1 - p_2)^{\ell_2}$, the claimed equality holds.

Hence, we can now assume that the degree of u is three. By symmetry, the degree of v is also three. Note that u can only become blue and only because of an active path of type $3 \leftrightarrow 3$ ending at u . This happens with probability $1 - P_k(u, v, \gamma_u)$. Similarly, v becomes blue with probability $1 - P_k(v, u, \gamma_v)$. Since the event that u becomes blue and v becomes blue conditioned by $c_{k-1}(T_{u,v}) = \gamma_u$ and $c_{k-1}(T_{v,u}) = \gamma_v$ are independent, it follows that (1) is also equal to $P_k(u, v, \gamma_u) \cdot P_k(v, u, \gamma_v)$ in this case. \square

Lemma 5 plays a crucial role in the Independence Lemma, which we now prove. Its proof enlightens how we designed our randomized procedure.

Lemma 6 (Independence Lemma). *Consider the randomized procedure for the infinite cubic tree. Let k be an integer, uv an edge of the tree and Γ_u and*

Γ_v two measurable subsets of \mathcal{C}_{k-1}^∞ . Conditioned by the event $uv \subseteq W_k$, the events that $c_k(T_{u,v}) \in \Gamma_u$ and $c_k(T_{v,u}) \in \Gamma_v$ are independent. In other words,

$$\mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k] = \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_k(T_{v,u}) \in \Gamma_v].$$

Proof. The proof proceeds by induction on k . If $k = 1$, the event $uv \subseteq W_1$ implies that neither u, v nor their neighbors is active during the first round. Conditioned by this, the other vertices of the infinite cubic tree are marked active with probability p_1 randomly and independently. The result of the marking in $T_{u,v}$ fully determine the coloring of the vertices of $T_{u,v}$ and is independent of the marking and coloring of $T_{v,u}$. Hence, the claim follows.

Assume that $k > 1$. Fix subsets Γ_u and Γ_v . We aim at showing that the probabilities

$$\mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k] \tag{2}$$

and

$$\mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid u \in W_k \wedge c_k(T_{v,u}) \in \Gamma_v] \tag{3}$$

are equal.

The definition of the conditional probability and the fact that $\mathbf{P}[uv \subseteq W_{k-1} \mid uv \subseteq W_k] = 1$ yield that (2) is equal to

$$\frac{\mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \wedge v \in W_k \mid uv \subseteq W_{k-1}]}{\mathbf{P}[uv \subseteq W_k \mid uv \subseteq W_{k-1}]} . \tag{4}$$

By the induction hypothesis, for any two subsets Γ'_u and Γ'_v of \mathcal{C}_{k-1}^∞ , the probabilities $\mathbf{P}[c_{k-1}(T_{v,u}) \in \Gamma'_v \mid u \in W_{k-1} \wedge c_{k-1}(T_{u,v}) \in \Gamma'_u]$ and $\mathbf{P}[c_{k-1}(T_{v,u}) \in \Gamma'_v \mid uv \subseteq W_{k-1}]$ are the same. Hence, the numerator of (4) is equal to the following:

$$\int_{\gamma'_u, \gamma'_v \in \mathcal{C}_{k-1}^\infty} \mathbf{P}[uv \subseteq W_k \mid c_{k-1}(T_{u,v}) = \gamma'_u \wedge c_{k-1}(T_{v,u}) = \gamma'_v] \times \\ \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u \wedge c_{k-1}(T_{v,u}) = \gamma'_v] \\ \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}].$$

Observe that when conditioning by $uv \subseteq W_k$, the event $c_k(T_{u,v}) \in \Gamma_u$ is independent of $c_{k-1}(T_{v,u}) = \gamma'_v$. Hence, the double integral can be rewritten

to

$$\int_{\gamma'_u, \gamma'_v \in C_{k-1}^\infty} \mathbf{P}[uv \subseteq W_k \mid c_{k-1}(T_{u,v}) = \gamma'_u \wedge c_{k-1}(T_{v,u}) = \gamma'_v] \times \\ \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u] \\ \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}].$$

An application of Lemma 5 then yields that the double sum is equal to

$$\int_{\gamma'_u, \gamma'_v \in C_{k-1}^\infty} P_k(u, v, \gamma'_u) \cdot P_k(v, u, \gamma'_v) \cdot \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u] \\ \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}].$$

Regrouping the terms containing γ'_u only and γ'_v only, we obtain that the numerator of (4) is equal to

$$\left(\int_{\gamma'_u \in C_{k-1}^\infty} P_k(u, v, \gamma'_u) \times \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u] \\ \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \right) \times \\ \left(\int_{\gamma'_v \in C_{k-1}^\infty} P_k(v, u, \gamma'_v) \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}] \right).$$

Along the same lines, the denominator of (4) can be expressed as

$$\left(\int_{\gamma'_u \in C_{k-1}^\infty} P_k(u, v, \gamma'_u) \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \right) \times \\ \left(\int_{\gamma'_v \in C_{k-1}^\infty} P_k(v, u, \gamma'_v) \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}] \right).$$

Cancelling out the integral over γ'_v which is the same in the numerator and the denominator of (4), we obtain that (2) is equal to

$$\frac{\int_{\gamma'_u \in C_{k-1}^\infty} P_k(u, v, \gamma'_u) \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u] \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}]}{\int_{\gamma'_u \in C_{k-1}^\infty} P_k(u, v, \gamma'_u) \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}]} \quad (5)$$

The same trimming is applied to (3). First, the probability (3) is expressed as

$$\frac{\mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \wedge c_k(T_{v,u}) \in \Gamma_v \mid uv \subseteq W_{k-1}]}{\mathbf{P}[u \in W_k \wedge c_k(T_{v,u}) \in \Gamma_v \mid uv \subseteq W_{k-1}]} \quad (6)$$

The numerator of (6) is then expanded to

$$\begin{aligned} & \left(\int_{\gamma'_u \in \mathcal{C}_{k-1}^\infty} P_k(u, v, \gamma'_u) \mathbf{P}[c_k(T_{u,v}) \in \Gamma_u \mid uv \subseteq W_k \wedge c_{k-1}(T_{u,v}) = \gamma'_u] \right. \\ & \quad \left. \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \right) \times \\ & \left(\int_{\gamma'_v \in \mathcal{C}_{k-1}^\infty} \mathbf{P}[c_k(T_{v,u}) \in \Gamma_v \mid uv \subseteq W_k \wedge c_{k-1}(T_{v,u}) = \gamma'_v \cdot P_k(v, u, \gamma'_v)] \right. \\ & \quad \left. \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}] \right) \end{aligned}$$

and the denominator of (3) is expanded to

$$\begin{aligned} & \left(\int_{\gamma'_u \in \mathcal{C}_{k-1}^\infty} P_k(u, v, \gamma'_u) \mathbf{dP}[c_{k-1}(T_{u,v}) = \gamma'_u \mid uv \subseteq W_{k-1}] \right) \times \\ & \left(\int_{\gamma'_v \in \mathcal{C}_{k-1}^\infty} \mathbf{P}[c_k(T_{v,u}) \in \Gamma_v \mid uv \subseteq W_k \wedge c_{k-1}(T_{v,u}) = \gamma'_v \cdot P_k(v, u, \gamma'_v)] \right. \\ & \quad \left. \mathbf{dP}[c_{k-1}(T_{v,u}) = \gamma'_v \mid uv \subseteq W_{k-1}] \right). \end{aligned}$$

We obtain (5) by cancelling out the integrals over γ'_v . The proof is now finished. \square

4.2 Recurrence relations

We now derive recurrence relations for the probabilities describing the behavior of the randomized procedure. We show how to compute the probabilities after $(k + 1)$ rounds from the probabilities after k rounds only.

Recall that $w_k^{(i)}$ is the probability that a fixed vertex u has degree i after k rounds conditioned by the event that u is white after k rounds. Also recall that $q_k^{(i)}$ is the probability that a fixed vertex u with a fixed neighbor v has degree i after k rounds conditioned by the event that both u and v are white after k rounds, i.e., $uv \subseteq W_k$. Finally, w_k, r_k and b_k are the probabilities that a fixed vertex is white, red and blue, respectively, after k rounds.

If u is white, the *white subtree* of $T_{u,v}$ is the maximal subtree containing u and white vertices only. We claim that the probability that the white subtree of $T_{u,v}$ is isomorphic to a tree in a given subset \mathcal{T}_0 after k rounds, conditioned

by the event that both u and v are white after k rounds, can be computed from the values of $q_k^{(i)}$. Indeed, if $T_0 \in \mathcal{T}_0$, the probability that u has degree i as in T_0 is $q_k^{(i)}$. Now, if the degree of u is i as in T_0 and z is a neighbor of u , the values of $q_k^{(i)}$ again determine the probability that the degree of z is as in T_0 . By Lemma 6, the probabilities that u and z have certain degrees, conditioned by the event that they are both white, are independent. Inductively, we can proceed with other vertices of T_0 . Applying standard probability arguments, we see that the values of $q_k^{(i)}$ fully determine the probability that the white subtree of $T_{u,v}$ after k rounds is isomorphic to a tree in \mathcal{T}_0 .

After the first round, the probabilities w_k, r_k, b_k and $q_k^{(i)}$, $i \in \{0, 1, 2, 3\}$, are the following.

$$\begin{aligned}
b_1 &= 1 - (1 - p_1)^3 & q_1^{(1)} &= (1 - (1 - p_1)^2)^2 \\
r_1 &= p_1(1 - b_1) = p_1(1 - p_1)^3 & w_1^{(3)} &= (1 - p_1)^6 \\
w_1 &= 1 - b_1 - r_1 & w_1^{(2)} &= 3 \cdot (1 - p_1)^4 (1 - (1 - p_1)^2) \\
q_1^{(3)} &= (1 - p_1)^4 & w_1^{(1)} &= 3 \cdot (1 - p_1)^2 (1 - (1 - p_1)^2)^2 \\
q_1^{(2)} &= 2 \cdot (1 - p_1)^2 (1 - (1 - p_1)^2) & w_1^{(0)} &= (1 - (1 - p_1)^2)^3
\end{aligned}$$

A vertex becomes blue if at least one of its neighbors is active and it becomes red if it is active and none of its neighbors is also active. Otherwise, a vertex stays white. This leads to the formulas above.

To derive formulas for the probabilities w_k, r_k, b_k and $q_k^{(i)}$ for $k \geq 2$, we introduce additional notation. The recurrence relations can be expressed using $q_k^{(i)}$ only, but additional notation will help us to simplify expressions appearing in our analysis. For a given edge uv of the infinite tree, let $P_k^{\rightarrow 1}$ be the probability that the white subtree of $T_{u,v}$ contains a path from u to a vertex of degree one with all inner vertices of degree two after k rounds conditioned by the event $uv \subseteq W_k$. Note that such a path may have length zero ending at u (this happens if the degree of u is one). Quantities $P_k^{E \rightarrow 1}$ and $P_k^{O \rightarrow 1}$ are probabilities that the length of such a path is even or odd, respectively. Analogously, $P_k^{\rightarrow 3}$, $P_k^{E \rightarrow 3}$ and $P_k^{O \rightarrow 3}$ are probabilities that the white subtree of $T_{u,v}$ contains a path, an even path and an odd path, respectively, from u to a vertex of degree three with all inner vertices of degree two after k rounds conditioned by the event $uv \subseteq W_k$.

Using Lemma 6, we conclude that the values of the just defined proba-

bilities can be computed as follows.

$$\begin{aligned}
P_k^{\rightarrow 1} &= q_k^{(1)} \cdot \sum_{\ell \geq 0} \left(q_k^{(2)}\right)^\ell = \frac{q_k^{(1)}}{1 - q_k^{(2)}} & P_k^{\rightarrow 3} &= q_k^{(3)} \cdot \sum_{\ell \geq 0} \left(q_k^{(2)}\right)^\ell = \frac{q_k^{(3)}}{1 - q_k^{(2)}} \\
P_k^{O \rightarrow 1} &= q_k^{(1)} \cdot \sum_{\ell \geq 0} \left(q_k^{(2)}\right)^{2\ell} = \frac{q_k^{(1)}}{1 - \left(q_k^{(2)}\right)^2} & P_k^{O \rightarrow 3} &= q_k^{(3)} \cdot \sum_{\ell \geq 0} \left(q_k^{(2)}\right)^{2\ell} = \frac{q_k^{(3)}}{1 - \left(q_k^{(2)}\right)^2} \\
P_k^{E \rightarrow 1} &= P_k^{\rightarrow 1} - P_k^{O \rightarrow 1} = q_k^{(2)} \cdot P_k^{O \rightarrow 1} & P_k^{E \rightarrow 3} &= P_k^{\rightarrow 3} - P_k^{O \rightarrow 3} = q_k^{(2)} \cdot P_k^{O \rightarrow 3}
\end{aligned}$$

Observe that $P_k^{\rightarrow 1} + P_k^{\rightarrow 3} = 1$.

The formulas for the above probabilities can be easily altered to express the probabilities that a path exists and one of its inner vertices is active; simply, instead of multiplying by $q_k^{(2)}$, we multiply by $p_2 q_k^{(2)}$. Now, $\widehat{P}_k^{\rightarrow 3}$ is the probability that the white subtree of T_{uv} contains a path from u to a vertex of degree three with all inner vertices of degree two after k rounds and none of them become active, conditioned by $uv \subseteq W_k$. Analogously to the previous paragraph, we use $\widehat{P}_k^{O \rightarrow 3}$ and $\widehat{P}_k^{E \rightarrow 3}$. The probabilities $\widehat{P}_k^{\rightarrow 3}$, $\widehat{P}_k^{O \rightarrow 3}$ and $\widehat{P}_k^{E \rightarrow 3}$ can be computed in the following way.

$$\begin{aligned}
\widehat{P}_k^{\rightarrow 3} &= q_k^{(3)} \cdot \sum_{\ell \geq 0} \left(q_k^{(2)} \cdot (1 - p_2)\right)^\ell = \frac{q_k^{(3)}}{1 - q_k^{(2)} \cdot (1 - p_2)} \\
\widehat{P}_k^{O \rightarrow 3} &= q_k^{(3)} \cdot \sum_{\ell \geq 1} \left(q_k^{(2)} \cdot (1 - p_2)\right)^{2\ell} = \frac{q_k^{(3)}}{1 - \left(q_k^{(2)}\right)^2 \cdot (1 - p_2)^2} \\
\widehat{P}_k^{E \rightarrow 3} &= \widehat{P}_k^{\rightarrow 1} - \widehat{P}_k^{O \rightarrow 3} = q_k^{(2)} \cdot (1 - p_2) \cdot \widehat{P}_k^{O \rightarrow 3}
\end{aligned}$$

The probabilities $\widetilde{P}_k^{O \rightarrow 3}$ and $\widetilde{P}_k^{E \rightarrow 3}$ are the probabilities that such an odd/even path exists and at least one of its inner vertices become active. Note that $P_k^{O \rightarrow 3} = \widehat{P}_k^{O \rightarrow 3} + \widetilde{P}_k^{O \rightarrow 3}$. The value of $\widetilde{P}_k^{O \rightarrow 3}$ is given by the equation

$$\widetilde{P}_k^{O \rightarrow 3} = \left(q_k^{(2)}\right)^2 \cdot \left((1 - p_2)^2 \cdot \widetilde{P}_k^{O \rightarrow 3} + (1 - (1 - p_2)^2) \cdot P_k^{O \rightarrow 3} \right),$$

which can be manipulated to

$$\widetilde{P}_k^{O \rightarrow 3} = \frac{\left(q_k^{(2)}\right)^2 \cdot (1 - (1 - p_2)^2) \cdot P_k^{O \rightarrow 3}}{1 - \left(q_k^{(2)}\right)^2 \cdot (1 - p_2)^2}.$$

Using the expression for $\widetilde{P}_k^{O \rightarrow 3}$, we derive that $\widetilde{P}_k^{E \rightarrow 3}$ is equal to the following.

$$\widetilde{P}_k^{E \rightarrow 3} = q_k^{(2)} \cdot \left(p_2 \cdot P_k^{O \rightarrow 3} + (1 - p_2) \cdot \widetilde{P}_k^{O \rightarrow 3} \right)$$

We now show how to compute the probabilities w_{k+1} , b_{k+1} and r_{k+1} . Since blue and red vertices keep their colors once assigned, we have to focus on the probability that a white vertex changes a color. We distinguish vertices based on their degrees.

A vertex of degree zero. Such a vertex is always recolored to red.

A vertex of degree one. Such a vertex is always recolored. Its new color is blue only if lies on an odd path to another vertex of degree one and the other end is chosen to be the beginning of the path. This leads to the following equalities.

$$\mathbf{P}[u \in R_{k+1} \mid u \in W_k^{(1)}] = \frac{1}{2}P_k^{O \rightarrow 1} + P_k^{E \rightarrow 1} + P_k^{\rightarrow 3},$$

and

$$\mathbf{P}[u \in B_{k+1} \mid u \in W_k^{(1)}] = \frac{1}{2}P_k^{O \rightarrow 1}.$$

A vertex of degree two. Since we have already computed the probabilities that the paths of white vertices with degree two leading in the two directions from the vertex end at a vertex of degree one/three, have odd/even length and contain an active vertex, we can easily determine the probability that the vertex stays white or becomes red or blue. Note that for odd paths with type $1 \leftrightarrow 1$ and $3 \leftrightarrow 3$, in addition, the random choice of the start of the path comes in the play. It is then straightforward to derive the following.

$$\begin{aligned} \mathbf{P}[u \in R_{k+1} \mid u \in W_k^{(2)}] &= (P_k^{E \rightarrow 1})^2 + \frac{2}{2}P_k^{O \rightarrow 1}P_k^{E \rightarrow 1} + 2P_k^{E \rightarrow 1}P_k^{\rightarrow 3} \\ &+ (1 - p_2) \cdot \left((\tilde{P}_k^{O \rightarrow 3})^2 + 2\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{O \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} \right) \\ &+ p_2 \cdot \left((P_k^{O \rightarrow 3})^2 + \frac{2}{2}P_k^{E \rightarrow 3}P_k^{O \rightarrow 3} \right) \\ \mathbf{P}[u \in B_{k+1} \mid u \in W_k^{(2)}] &= (P_k^{O \rightarrow 1})^2 + \frac{2}{2}P_k^{O \rightarrow 1}P_k^{E \rightarrow 1} + 2P_k^{O \rightarrow 1}P_k^{\rightarrow 3} \\ &+ (1 - p_2) \cdot \left((\tilde{P}_k^{E \rightarrow 3})^2 + 2\tilde{P}_k^{E \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} + \frac{2}{2}\tilde{P}_k^{O \rightarrow 3}\tilde{P}_k^{E \rightarrow 3} \right) \\ &+ p_2 \cdot \left((P_k^{E \rightarrow 3})^2 + \frac{2}{2}P_k^{E \rightarrow 3}P_k^{O \rightarrow 3} \right) \end{aligned}$$

A vertex of degree three. The vertex either stays white or is recolored to blue. It stays white if (and only if) all the three paths with inner

vertices being white and with degree two are among the following: even paths to a vertex of degree one, non-activated paths to a vertex of degree three, or activated odd paths to a vertex of degree three that was chosen as the beginning (and thus recolored with blue). Hence we obtain that

$$\mathbf{P}[u \in B_{k+1} \mid u \in W_k^{(3)}] = 1 - \left(P_k^{E \rightarrow 1} + \widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{O \rightarrow 3} \right)^3. \quad (7)$$

Plugging all the probabilities from the above analysis together yields that

$$\begin{aligned} r_{k+1} &= r_k + w_k \cdot \left(\sum_{i=0}^2 w_k^{(i)} \cdot \mathbf{P}[u \in R_{k+1} \mid u \in W_k^{(i)}] \right) \\ b_{k+1} &= b_k + w_k \cdot \left(\sum_{i=1}^3 w_k^{(i)} \cdot \mathbf{P}[u \in B_{k+1} \mid u \in W_k^{(i)}] \right) \\ w_{k+1} &= 1 - r_{k+1} - b_{k+1}. \end{aligned}$$

The crucial for the whole analysis is computing the values of $w_{k+1}^{(i)}$. Suppose that u is a white vertex after k rounds. The values of $w_k^{(i)}$ determine the probability that u has degree i and the values of $q_k^{(i)}$ determine the probabilities that white neighbors of u have certain degrees. In particular, the probability that u has degree i and its neighbors have degrees j_1, \dots, j_i after k rounds conditioned by u being white after k rounds is equal to

$$w_k^{(i)} \cdot \prod_{j \in \{j_1, \dots, j_i\}} q_k^{(j)}.$$

In what follows, the vector of degrees j_1, \dots, j_i will be denoted by \vec{j} .

Let $R_{k+1}^{(i)}(\vec{j})$ be the probability that u is white after $(k+1)$ rounds conditioned by the event that u is white, has degree i and its white neighbors have degrees \vec{j} after k rounds. Note that the value of $R_{k+1}^{(i)}(\vec{j})$ is the same for all permutation of entries/degrees of the vector \vec{j} .

If u has degree three and all its neighbors also have degree three after k rounds, the probability $R_{k+1}^{(3)}(3, 3, 3)$ is equal to one: no vertex of degree three can be colored by red and thus the color of u stays white. On the other hand, if u or any of its neighbor has degree one, u does definitely not stay white and the corresponding probability $R_{k+1}^{(i)}(\vec{j})$ is equal to zero.

We now analyze the value of $R_{k+1}^{(i)}(\vec{J})$ for the remaining combinations of i and \vec{J} . If $i = 2$, then u stays white only if it lies on a non-active path of degree-two vertices between two vertices of degree three. Consequently, it holds that

$$\begin{aligned} R_{k+1}^{(2)}(2, 2) &= (1 - p_2)^3 \cdot \left(\widehat{P}_k^{\rightarrow 3}\right)^2 \\ R_{k+1}^{(2)}(3, 2) &= (1 - p_2)^2 \cdot \widehat{P}_k^{\rightarrow 3} \\ R_{k+1}^{(2)}(3, 3) &= (1 - p_2) \end{aligned}$$

If $i = 3$, then u stays white if and only if for every neighbor v of degree two of u , the path of degree-two vertices from v to u is

- an odd path to a vertex of degree one,
- a non-activated path to a vertex of degree three,
- an activated even path to vertex of degree three and u is not chosen as the beginning.

Based on this, we obtain that the values of $R_{k+1}^{(3)}(\vec{J})$ for the remaining choices of \vec{J} are the as follows.

$$\begin{aligned} R_{k+1}^{(3)}(2, 2, 2) &= \left(P_k^{O \rightarrow 1} + (1 - p_2) \cdot \left(\widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{E \rightarrow 3} \right) + p_2 \cdot \frac{1}{2} P_k^{E \rightarrow 3} \right)^3 \\ R_{k+1}^{(3)}(2, 2, 3) &= \left(P_k^{O \rightarrow 1} + (1 - p_2) \cdot \left(\widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{E \rightarrow 3} \right) + p_2 \cdot \frac{1}{2} P_k^{E \rightarrow 3} \right)^2 \\ R_{k+1}^{(3)}(2, 3, 3) &= P_k^{O \rightarrow 1} + (1 - p_2) \cdot \left(\widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{E \rightarrow 3} \right) + p_2 \cdot \frac{1}{2} P_k^{E \rightarrow 3} \end{aligned}$$

We now focus on computing the probabilities $R_{k+1}^{(i \rightarrow i')}(\vec{J})$ that a vertex u is a white vertex of degree i' after $(k + 1)$ rounds conditioned by the event that u is a white vertex with degree i with neighbors of degrees in \vec{J} after k rounds and u is also white after $(k + 1)$ rounds. For example, $R_{k+1}^{(2 \rightarrow i')}(2, 2)$ is equal to one for $i' = 2$ and to zero for $i' \neq 2$. To derive formulas for the probabilities $R_{k+1}^{(i \rightarrow i')}$, we have to introduce some additional notation: $S_{k+1}^{(i, j)}$ for $(i, j) \in \{(2, 2), (2, 3), (3, 2), (3, 3)\}$ will denote the probability that a vertex v

is white after $(k+1)$ rounds conditioned by the event that v is a white vertex of degree j after k rounds and a fixed (white) neighbor u of v has degree i after k rounds and u is white after $(k+1)$ rounds. It is easy to see that $S_{k+1}^{(2,2)} = 1$. If $j = 3$, the event we condition by guarantees that one of the neighbors of v is white after $(k+1)$ rounds. Hence, we derive that

$$S_{k+1}^{(2,3)} = S_{k+1}^{(3,3)} = \left(P_k^{E \rightarrow 1} + \widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{O \rightarrow 3} \right)^2.$$

Using the probabilities $S_{k+1}^{(i,j)}$, we can easily express some of the probabilities $R_{k+1}^{(i \rightarrow i')}(\vec{J})$.

$$\begin{aligned} R_{k+1}^{(2 \rightarrow 2)}(2, 3) &= S_{k+1}^{(2,3)} & R_{k+1}^{(2 \rightarrow 2)}(3, 3) &= \left(S_{k+1}^{(2,3)} \right)^2 \\ R_{k+1}^{(2 \rightarrow 1)}(2, 3) &= 1 - S_{k+1}^{(2,3)} & R_{k+1}^{(2 \rightarrow 1)}(3, 3) &= 2 \cdot S_{k+1}^{(2,3)} \left(1 - S_{k+1}^{(2,3)} \right) \\ R_{k+1}^{(2 \rightarrow 0)}(2, 3) &= 0 & R_{k+1}^{(2 \rightarrow 0)}(3, 3) &= \left(1 - S_{k+1}^{(2,3)} \right)^2 \\ R_{k+1}^{(3 \rightarrow 3)}(3, 3, 3) &= \left(S_{k+1}^{(3,3)} \right)^3 & R_{k+1}^{(3 \rightarrow 1)}(3, 3, 3) &= 3 \cdot S_{k+1}^{(3,3)} \left(1 - S_{k+1}^{(3,3)} \right)^2 \\ R_{k+1}^{(3 \rightarrow 2)}(3, 3, 3) &= 3 \cdot \left(S_{k+1}^{(3,3)} \right)^2 \left(1 - S_{k+1}^{(3,3)} \right) & R_{k+1}^{(3 \rightarrow 0)}(3, 3, 3) &= \left(1 - S_{k+1}^{(3,3)} \right)^3 \end{aligned}$$

We now determine $S_{k+1}^{(3,2)}$, i.e., the probability that a vertex v is white after $(k+1)$ rounds conditioned by the event that v has degree two after k rounds and a fixed white neighbor u of v that has degree three after k rounds is white after $(k+1)$ rounds. Observe that v is white after $(k+1)$ rounds only if v is contained in a non-active $3 \leftrightarrow 3$ path. Since we condition by the event that u is white after $(k+1)$ rounds, v cannot be contained in an active $3 \leftrightarrow 3$ path of even length or an active $3 \leftrightarrow 3$ odd path with u being chosen as the beginning of this path. Hence, the value of $S_{k+1}^{(3,2)}$ is the following.

$$S_{k+1}^{(3,2)} = \frac{(1 - p_2) \cdot \widehat{P}_k^{\rightarrow 3}}{P_k^{O \rightarrow 1} + (1 - p_2) \cdot \left(\widehat{P}_k^{\rightarrow 3} + \frac{1}{2} \widetilde{P}_k^{E \rightarrow 3} \right) + p_2 \cdot \frac{1}{2} P_k^{E \rightarrow 3}}.$$

Using $S_{k+1}^{(3,2)}$, the remaining values of $R_{k+1}^{(i \rightarrow i')}(\vec{\mathcal{J}})$ can be expressed as follows.

$$\begin{aligned}
R_{k+1}^{(3 \rightarrow 3)}(3, 3, 2) &= \left(S_{k+1}^{(3,3)}\right)^2 \cdot S_{k+1}^{(3,2)} \\
R_{k+1}^{(3 \rightarrow 2)}(3, 3, 2) &= \left(S_{k+1}^{(3,3)}\right)^2 \left(1 - S_{k+1}^{(3,2)}\right) + 2 \cdot \left(1 - S_{k+1}^{(3,3)}\right) S_{k+1}^{(3,3)} \cdot S_{k+1}^{(3,2)} \\
R_{k+1}^{(3 \rightarrow 1)}(3, 3, 2) &= \left(1 - S_{k+1}^{(3,3)}\right)^2 S_{k+1}^{(3,2)} + 2 \cdot S_{k+1}^{(3,3)} \left(1 - S_{k+1}^{(3,3)}\right) \left(1 - S_{k+1}^{(3,2)}\right) \\
R_{k+1}^{(3 \rightarrow 0)}(3, 3, 2) &= \left(1 - S_{k+1}^{(3,3)}\right)^2 \left(1 - S_{k+1}^{(3,2)}\right) \\
R_{k+1}^{(3 \rightarrow 3)}(2, 2, 3) &= \left(S_{k+1}^{(3,2)}\right)^2 \cdot S_{k+1}^{(3,3)} \\
R_{k+1}^{(3 \rightarrow 2)}(2, 2, 3) &= \left(S_{k+1}^{(3,2)}\right)^2 \left(1 - S_{k+1}^{(3,3)}\right) + 2 \cdot \left(1 - S_{k+1}^{(3,2)}\right) S_{k+1}^{(3,2)} \cdot S_{k+1}^{(3,3)} \\
R_{k+1}^{(3 \rightarrow 1)}(2, 2, 3) &= \left(1 - S_{k+1}^{(3,2)}\right)^2 S_{k+1}^{(3,3)} + 2 \cdot S_{k+1}^{(3,2)} \left(1 - S_{k+1}^{(3,2)}\right) \left(1 - S_{k+1}^{(3,3)}\right) \\
R_{k+1}^{(3 \rightarrow 0)}(2, 2, 3) &= \left(1 - S_{k+1}^{(3,2)}\right)^2 \left(1 - S_{k+1}^{(3,3)}\right) \\
R_{k+1}^{(3 \rightarrow 3)}(2, 2, 2) &= \left(S_{k+1}^{(3,2)}\right)^3 \\
R_{k+1}^{(3 \rightarrow 2)}(2, 2, 2) &= 3 \cdot \left(S_{k+1}^{(3,2)}\right)^2 \left(1 - S_{k+1}^{(3,2)}\right) \\
R_{k+1}^{(3 \rightarrow 1)}(2, 2, 2) &= 3 \cdot S_{k+1}^{(3,2)} \left(1 - S_{k+1}^{(3,2)}\right)^2 \\
R_{k+1}^{(3 \rightarrow 0)}(2, 2, 2) &= \left(1 - S_{k+1}^{(3,2)}\right)^3
\end{aligned}$$

Using $R_{k+1}^{(i \rightarrow i')}(\vec{\mathcal{J}})$, we can compute $w_{k+1}^{(i')}$, $i' \in \{0, 1, 2, 3\}$. In the formula below, \mathcal{J}_i denotes the set of all possible vectors $\vec{\mathcal{J}}$ with i entries and all entries either two or three. The denominator of the formula is the probability that the vertex u is white after $k+1$ rounds conditioned by the event that u is white after k rounds; the nominator is the probability that u is white and has degree i' after $k+1$ rounds conditioned by the event that u is white after k rounds.

$$w_{k+1}^{(i')} = \frac{\sum_{\substack{i \geq 2 \\ i \geq i'}} \sum_{\vec{\mathcal{J}} \in \mathcal{J}_i} w_k^{(i)} \cdot \prod_{j \in \vec{\mathcal{J}}} q_k^{(j)} \cdot R_{k+1}^{(i)}(\vec{\mathcal{J}}) \cdot R_{k+1}^{(i \rightarrow i')}(\vec{\mathcal{J}})}{\sum_{i \geq 2} \sum_{\vec{\mathcal{J}} \in \mathcal{J}_i} w_k^{(i)} \cdot \prod_{j \in \vec{\mathcal{J}}} q_k^{(j)} \cdot R_{k+1}^{(i)}(\vec{\mathcal{J}})}$$

It remains to exhibit the recurrence relations for the values of $q_{k+1}^{(i)}$. Let uv be an edge of the tree, $i \geq 1$ and $\vec{\mathcal{J}} \in \mathcal{J}_i$. Observe that the probability that

u has degree i and its neighbors have degrees \vec{J} after k rounds conditioned by the event $uv \subseteq W_k$ is exactly

$$q_k^{(i)} \cdot \prod_{j \in \vec{J}} q_k^{(j)} .$$

In what follows, we will assume that the first coordinate j_1 of \vec{J} corresponds to the vector v .

The probabilities $Q_{k+1}^{(i)}$ are defined analogically to $R_{k+1}^{(i)}$ with an additional requirement that v is also white after $k+1$ rounds. Formally, $Q_{k+1}^{(i)}(\vec{J})$ is the probability that a vertex u and its fixed neighbor v are both white after $(k+1)$ rounds conditioned by the event that $uv \subseteq W_k$, u has degree i and its white neighbors have degrees \vec{J} after k rounds. Observe that the following holds.

$$\begin{aligned} Q_{k+1}^{(2)}(2, 2) &= R_{k+1}^{(2)}(2, 2) & Q_{k+1}^{(2)}(3, 2) &= R_{k+1}^{(2)}(3, 2) \cdot S_{k+1}^{(2,3)} \\ Q_{k+1}^{(2)}(2, 3) &= R_{k+1}^{(2)}(2, 3) = R_{k+1}^{(2)}(3, 2) & Q_{k+1}^{(2)}(3, 3) &= R_{k+1}^{(2)}(3, 3) \cdot S_{k+1}^{(2,3)} \\ Q_{k+1}^{(3)}(j_1, j_2, j_3) &= R_{k+1}^{(3)}(j_1, j_2, j_3) \cdot S_{k+1}^{(3, j_1)} \end{aligned}$$

Similarly, $Q_{k+1}^{(i \rightarrow i')}$ is the probability that a vertex u is a white vertex of degree $i' \geq 1$ and its fixed neighbor v is white after $(k+1)$ rounds conditioned by the event that u is a white vertex with degree i with neighbors of degrees in \vec{J} after k rounds and $uv \subseteq W_{k+1}$. Using the arguments analogous to those to derive the formulas for $R_{k+1}^{(i \rightarrow i')}(\vec{J})$, we obtain the following formulas for $Q_{k+1}^{(i \rightarrow i')}$. We provide the list of recurrences to compute the values of $Q_{k+1}^{(i \rightarrow i')}$ and leave the actual derivation to the reader.

$$\begin{aligned} Q_{k+1}^{(2 \rightarrow 2)}(2, 2) &= Q_{k+1}^{(2 \rightarrow 2)}(3, 2) = 1 & Q_{k+1}^{(2 \rightarrow 2)}(2, 3) &= Q_{k+1}^{(2 \rightarrow 2)}(3, 3) = S_{k+1}^{(2,3)} \\ Q_{k+1}^{(2 \rightarrow 1)}(2, 2) &= Q_{k+1}^{(2 \rightarrow 1)}(3, 2) = 0 & Q_{k+1}^{(2 \rightarrow 1)}(2, 3) &= Q_{k+1}^{(2 \rightarrow 1)}(3, 3) = 1 - S_{k+1}^{(2,3)} \\ \\ Q_{k+1}^{(3 \rightarrow 3)}(3, 3, 3) &= \left(S_{k+1}^{(3,3)}\right)^2 & Q_{k+1}^{(3 \rightarrow 3)}(3, 2, 2) &= \left(S_{k+1}^{(3,2)}\right)^2 \\ Q_{k+1}^{(3 \rightarrow 2)}(3, 3, 3) &= 2 \cdot S_{k+1}^{(3,3)} \left(1 - S_{k+1}^{(3,3)}\right) & Q_{k+1}^{(3 \rightarrow 2)}(3, 2, 2) &= 2 \cdot \left(1 - S_{k+1}^{(3,2)}\right) S_{k+1}^{(3,2)} \\ Q_{k+1}^{(3 \rightarrow 1)}(3, 3, 3) &= \left(1 - S_{k+1}^{(3,3)}\right)^2 & Q_{k+1}^{(3 \rightarrow 1)}(3, 2, 2) &= \left(1 - S_{k+1}^{(3,2)}\right)^2 \\ Q_{k+1}^{(3 \rightarrow 3)}(3, 3, 2) &= S_{k+1}^{(3,3)} \cdot S_{k+1}^{(3,2)} & Q_{k+1}^{(3 \rightarrow 3)}(2, 3, 3) &= \left(S_{k+1}^{(3,3)}\right)^2 \\ Q_{k+1}^{(3 \rightarrow 2)}(3, 3, 2) &= S_{k+1}^{(3,3)} \left(1 - S_{k+1}^{(3,2)}\right) + \left(1 - S_{k+1}^{(3,3)}\right) S_{k+1}^{(3,2)} & Q_{k+1}^{(3 \rightarrow 2)}(2, 3, 3) &= 2 \cdot \left(1 - S_{k+1}^{(3,3)}\right) S_{k+1}^{(3,3)} \end{aligned}$$

$$\begin{aligned}
Q_{k+1}^{(3 \rightarrow 1)}(3, 3, 2) &= (1 - S_{k+1}^{(3,3)}) (1 - S_{k+1}^{(3,2)}) & Q_{k+1}^{(3 \rightarrow 1)}(2, 3, 3) &= (1 - S_{k+1}^{(3,3)})^2 \\
Q_{k+1}^{(3 \rightarrow 3)}(2, 2, 3) &= S_{k+1}^{(3,2)} \cdot S_{k+1}^{(3,3)} & Q_{k+1}^{(3 \rightarrow 3)}(2, 2, 2) &= (S_{k+1}^{(3,2)})^2 \\
Q_{k+1}^{(3 \rightarrow 2)}(2, 2, 3) &= (1 - S_{k+1}^{(3,2)}) S_{k+1}^{(3,3)} + S_{k+1}^{(3,2)} (1 - S_{k+1}^{(3,3)}) & Q_{k+1}^{(3 \rightarrow 2)}(2, 2, 2) &= 2 \cdot (1 - S_{k+1}^{(3,2)}) S_{k+1}^{(3,2)} \\
Q_{k+1}^{(3 \rightarrow 1)}(2, 2, 3) &= (1 - S_{k+1}^{(3,2)}) (1 - S_{k+1}^{(3,3)}) & Q_{k+1}^{(3 \rightarrow 1)}(2, 2, 2) &= (1 - S_{k+1}^{(3,2)})^2
\end{aligned}$$

Using the values of $Q_{k+1}^{(i)}$ and $Q_{k+1}^{(i \rightarrow i')}$, we can compute the values of $q_{k+1}^{(i')}$ for $i' \in \{1, 2, 3\}$. The denominator of the formula is the probability that the vertices u and v are white after $k+1$ rounds conditioned by the event that u and v are white after k rounds; the nominator is the probability that u and v are white and u has degree i' after $k+1$ rounds conditioned by the event that u and v are white after k rounds.

$$q_{k+1}^{(i')} = \frac{\sum_{\substack{i \geq 2 \\ i \geq i'}} \sum_{\vec{J} \in \mathcal{J}_i} q_k^{(i)} \cdot \prod_{j \in \vec{J}} q_k^{(j)} \cdot Q_{k+1}^{(i)}(\vec{J}) \cdot Q_{k+1}^{(i \rightarrow i')}(\vec{J})}{\sum_{i \geq 2} \sum_{\vec{J} \in \mathcal{J}_i} q_k^{(i)} \cdot \prod_{j \in \vec{J}} q_k^{(j)} \cdot Q_{k+1}^{(i)}(\vec{J})}$$

4.3 Solving the recurrences

The recurrences presented in this section were solved numerically using the Python program provided in the Appendix. The particular choice of parameters used in the program was $p_1 = p_2 = 10^{-5}$ and $K = 307449$. The choice of K was made in such a way that $w_K \leq 10^{-6}$. We also estimated the precision of our calculations based on the representation of float numbers to avoid rounding errors effecting the presented bound on significant digits. Solving the recurrences for the above choice of parameters we obtain that $r_K > 0.4352$.

5 High-girth graphs

In this section, we show how to modify the randomized procedure from the previous section so that it can be applied to cubic graphs with large girth. In order to use the analysis presented in the previous section, we have to cope with the dependence of some of the events caused by the presence of cycles in the graph. To do so, we introduce an additional parameter L which will

control the length of paths causing the dependencies. Then, we will be able to guarantee that the probability that a fixed vertex of a given cubic graph is red is at least $r_K - o(1)$ assuming the girth of the graph is at least $8KL + 2$. In particular, if L tends to infinity, we approach the same probability as for the infinite cubic tree.

5.1 Randomized procedure

We now describe how the randomized procedure is altered. Let G be the given cubic graph. We produce a sequence $G_0, \overline{G}_1, G_1, \dots, \overline{G}_K, G_K$ of vertex-colored subcubic graphs; the only vertices in these graphs that have less than three neighbors will always be assigned a new color—black. The graphs can also contain some additional vertices which do not correspond to the vertices of G ; such vertices will be called *virtual*. The graph G_0 is the cubic graph G with all vertices colored white. Assume that G_{k-1} is already defined. Let \overline{G}_k be the graph and its vertex coloring obtained from G_{k-1} using the randomized procedure for the k -th round exactly as described for the infinite tree.

Before we describe how the graph G_k is obtained from \overline{G}_k , we need to introduce additional notation. Let y_0, y_1, \dots, y_{2L} be a fixed path in the infinite cubic tree. Now consider restrictions to T_{y_1, y_0} of colorings of the tree after k rounds that satisfy that

1. the vertices y_0, y_1, \dots, y_{2L} are white after k rounds and
2. the vertices y_1, \dots, y_{2L-1} have degree two.

Let \mathcal{D}_k be the probability distribution on these restrictions such that the probability of each restriction is proportional to the probability of the original coloring after k rounds. In other words, we discard the colorings that do not satisfy the two constraints, restrict them to T_{y_1, y_0} and normalize the probabilities.

All graphs G_k will satisfy that all paths joining non-virtual vertices contain non-virtual vertices only. The graph G_k is obtained from \overline{G}_k by performing the following operation on every path P of type $1 \leftrightarrow 1$, $1 \leftrightarrow 3$ or $3 \leftrightarrow 3$ between vertices a and b that has length at least $2L$ and contains at least one non-virtual inner vertex. Let x_u be the non-virtual inner vertex of P that is the closest to a and x_v the one closest to b . Let P_x be the subpath between x_u and x_v (inclusively) in \overline{G}_k . Let u be the neighbor of x_u on P towards a , and

v the neighbor of x_v on P towards b . Observe that, since no path between non-virtual vertices contains a virtual vertex, if a is non-virtual, then x_u is the neighbor of a on P and $u = a$. Similarly, if b is non-virtual, then v is b .

We now modify the graph \overline{G}_k as follows. Color the vertices of P_x black and remove the edges $x_u u$ and $x_v v$ from the graph. Then attach to u and v rooted trees T_u and T_v , all of them fully comprised of virtual vertices, such that the colorings of T_u and T_v are randomly sampled according to the distribution \mathcal{D}_k . The roots of the trees will become adjacent to u or v , respectively. These trees are later referred to as *virtual trees*. Observe that we have created no path between two non-virtual vertices containing a virtual vertex.

After K rounds, the vertices of G receive colors of their counterparts in G_K . In this way, the vertices of G are colored white, blue, red and black and the red vertices form an independent set.

5.2 Refining the analysis

We argue that the analysis for the infinite cubic trees presented in Section 4 also applies to cubic graphs with large girth. We start with some additional definitions. Let u and v be two vertices of G_k . The vertex v is *reachable* from u if both u and v are white and there exists a path in G_k between u and v comprised of white vertices with all inner vertices having degree two. Clearly, the relation of being reachable is symmetric. The vertex v is *near* to u if both u and v are white and either v is reachable from u or v is a neighbor of a white vertex reachable from u . Note that the relation of being near is not symmetric in general. For a subset $X \subseteq V(G_k)$ of white vertices, $N_k(X) \subseteq V(G_k)$ is the set of white vertices that are near to a vertex of X in G_k .

We now prove the following theorem.

Theorem 7. *Let K be an integer, p_1 and p_2 positive reals, G a cubic graph and v a vertex of G . For every $\varepsilon > 0$ there exists an integer L such that if G has girth at least $8KL + 2$, then the probability that the vertex v will be red in G at the end of the randomized procedure is at least $r_K - \varepsilon$ where r_K is the probability that a fixed vertex of the infinite cubic tree is red after K rounds of the randomized procedure with parameters p_1 and p_2 .*

Proof. We keep the notation introduced in the description of the randomized procedure. As the first step in the proof, we establish the following two claims.

Claim 1. *Let k be a non-negative integer, u a vertex of G_k , c one of the colors, γ_1 and γ_2 two colorings of vertices of G_k such that u is white in both γ_1 and γ_2 . If the set of vertices near to u in γ_1 and γ_2 induce isomorphic trees rooted at u , then the probability that u has the color c in \overline{G}_{k+1} conditioned by the event that G_k is colored as in γ_1 , and the probability that u has the color c in \overline{G}_{k+1} conditioned by the event that G_k is colored as in γ_2 , are the same.*

Indeed, the color of u in \overline{G}_{k+1} is influenced only by the length and the types of the white paths in G_k containing u . All the vertices of these paths are near to u as well as the white neighbors of the other end-vertices of these paths (which are necessary to determine the types of the paths). By the assumption on the colorings γ_1 and γ_2 , the two probabilities from the statement of the claim are the same.

We now introduce yet another definition. For a subcubic graph H with vertices colored white, red, blue or black, a vertex $v \in H$ is said to be d -close to a white vertex $u \in H$ if v is white and there exists a path P from u to v comprised of white vertices such that

- the length of P is at most d , or
- P contains a vertex w such that w is at distance at most d from u on P and each of the first $(2L - 1)$ vertices following w (if they exist) has degree two.

Finally, a d -close tree of u is the subgraph comprised of all vertices d -close to u (for our choice of d , this subgraph will always be tree) that is rooted at u . By the definition, the d -close tree of u contains white vertices only.

Observe that if v is d -close to u , then it is also d' -close to u for every $d' > d$. Also observe that if a vertex v of a virtual subtree is d -close to a non-virtual vertex u , then all the white vertices lying in the same white component of the virtual subtree are also d -close to u : indeed, consider a path P witnessing that v is d -close to u and let P_0 be its subpath from u to the root of the virtual tree. The path P_0 witnesses that the root is d -close to u and P_0 can be prolonged by the path comprised of $2L - 1$ degree-two virtual vertices to a path to any white vertex v' of the same white component as v . The new path now witnesses that v' is also d -close to u .

Let us look at d -close sets in the infinite cubic trees.

Claim 2. *Let d be a non-negative integer, T an infinite cubic tree with vertices colored red, blue and white, and u a white vertex of T . If a vertex v is d -close to u in T , then every vertex v' that is near to v is $(d + 2L)$ -close to u .*

Let P be a path from u to v that witnesses that v is d -close to u . Assume first that the length of P is at most d . If v' lies on P or is a neighbor of a vertex of P , then v' is $(d + 1)$ -close to u . Otherwise, consider the path P' from u to v' ; observe that P is a subpath of P' and all the vertices following v on P with a possible exception of v' and the vertex immediately preceding it have degree two. If the length of P' is at most $d + 2L$, then v' is $(d + 2L)$ -close to u . Otherwise, v is followed by at least $2L - 1$ vertices of degree two and v' is again $(d + 2L)$ -close to u .

Assume now that the length of P is larger than d . Then, P contains a vertex w at distance at most d from u such that the first $2L - 1$ vertices following w (if they exist) have degree two. If v' lies on P or is adjacent to a vertex of P , then v' lies on P after w or is adjacent to w . In both cases, v' is $(d + 1)$ -close to u . In the remaining case, we again consider the path P' from u to v' which must be an extension of P (otherwise, v' would lie on P or it would be adjacent to a vertex on P). If there are at least $2L - 1$ vertices following w on P , then v' is d -close to u . Otherwise, either P' contains $2L - 1$ vertices of degree two following w or the length of P' is at most $d + 2L$. In both cases, v' is $(d + 2L)$ -close to u .

We are ready to prove our main claim.

Claim 3. *Let $k \leq K - 1$ be a positive integer, T a rooted subcubic tree such that its root is not contained in a path of degree-two vertices of length at least $2L$, u a vertex of the infinite cubic tree and v a non-virtual vertex of G_k . The probability that u is white and the $4(K - k)L$ -close tree of u in the infinite tree is isomorphic to T after k rounds is the same as the probability that v is white and the $4(K - k)L$ -close tree of v in G_k is isomorphic to T after k rounds.*

The proof proceeds by induction on k . For $k = 0$, both in the infinite tree and in G_0 , the probability is equal to one if T is the full rooted cubic tree of depth $4KL$ and it is zero, otherwise. Here, we use the girth assumption to derive that the subgraph of G induced by $4KL$ -close vertices to u is a tree (otherwise, G would contain a cycle of length at most $8KL + 1$).

Suppose $k > 0$. Let \tilde{T} be the $4(K - k + 1)L$ -close tree of v in G_{k-1} and W the set of the vertices that are $(4(K - k)L + 2L)$ -close to v in G_{k-1} . The degree and the color of each vertex in \overline{G}_k is determined by vertices that are near to it in G_{k-1} by Claim 1. In particular, the degrees and the colors of the vertices of W are determined by \tilde{T} by Claim 2. The induction assumption implies that every vertex of W is white and has a given degree i in \overline{G}_k with the same probability as its counterpart in the infinite cubic tree (assuming that the vertex u of the infinite tree does not lie on a path of degree-two vertices of length at least $2L$ after $k - 1$ rounds). If v lies on a path with at least $2L - 1$ degree-two vertices in \overline{G}_k , it becomes black.

The set W contains all vertices that are $4(K - k)L$ -close to v in G_k with the exception of the new virtual vertices that are $4(K - k)L$ -close to v . Since the colorings of newly added virtual trees have been sampled according to the distribution \mathcal{D}_k , Lemma 6 implies that the probability that v is white and the $4(K - k)L$ -close tree of v is equal to T is the same as the corresponding probability for u in the infinite cubic tree (conditioned by not being contained in a path of degree-two vertices of length at least $2L$).

Claim 4. *Let k be a non-negative integer and u a vertex of the infinite cubic tree. The probability that u is white and lies on a white path of length at least $2L$ after k rounds is at most $2 \cdot \left(q_k^{(2)}\right)^{L-1}$.*

If u lies on such a path, its degree must be two and the length of the path from u in one of the two directions is at least $L - 1$. The probability that this happens for each of the two possible directions from u is at most $\left(q_k^{(2)}\right)^{L-1}$. The claim now follows.

Let $p_0 = \sum_{k=1}^K 2 \left(q_k^{(2)}\right)^{L-1}$. Observe that $q_k^{(2)} < 1$. Indeed, if a vertex u of the infinite tree and all the vertices at distance at most $2K$ from u are white after the first round, then u and its three neighbors must have degree three after K rounds (all vertices at distance at most $2(K + 1 - k)$ from u are white after k rounds). Since this happens with non-zero probability, $q_k^{(3)} > 0$ and thus $q_k^{(2)} < 1$. This implies that p_0 tends to 0 with L approaching the infinity.

Fix a vertex v of G . By Claim 1, the probability that v is colored red in the k -th round is fully determined by the vertices that are near to v in G_{k-1} . All such vertices are also $2L$ -close to v by Claim 2. Consider a rooted

subcubic tree T . If the root of T does not lie on a path with inner vertices of degree two with length at least $2L$, then the probability that v is white and the $2L$ -close tree of v is isomorphic to T is the same as the analogous probability for a vertex of the infinite tree by Claim 3. Since the probability that v is white and it lies on a path of length at least $2L$ in its $2L$ -close tree at some point during the randomized procedure is at most p_0 by Claim 4, the probability that v is colored red in G_K is at least $r_K - p_0$. Since $p_0 < \varepsilon$ for L sufficiently large, the statement of the theorem follows. \square

6 Conclusion

The method we presented here, similarly to the method of Hoppen [5], can be applied to r -regular graphs for $r \geq 4$.

Another related question is whether the fractional chromatic number of cubic graphs is bounded away from 3 under a weaker assumption that the odd girth (the length of the shortest odd cycle) is large. This is indeed the case as we now show.

Theorem 8. *Let $g \geq 5$ be an odd integer. The fractional chromatic number of every subcubic graph G with odd girth at least g is at most $\frac{8}{3-6/(g+1)}$.*

Proof. Clearly, we can assume that G is bridgeless. If G contains two or more vertices of degree two, then we include G in a large cubic bridgeless graph with the same odd girth. Hence, we can assume that G contains at most one vertex of degree two. Consequently, G has a 2-factor F .

We now construct a probability distribution on the independent sets such that each vertex is included in the independent set chosen according to this distribution with probability at least $3(1 - 2/(g + 1))/8$. This implies the claim of the theorem.

Number the vertices of each cycle of F from 1 to ℓ where ℓ is the length of the cycle. Choose randomly a number k between 1 and $(g + 1)/2$ and let W be the set of all vertices with indices equal to k modulo $(g + 1)/2$. Hence, each vertex is not in W with probability $1 - 2/(g + 1)$.

Let V_1, \dots, V_m be the sets formed by the paths of $F \setminus W$. Since each set V_i , $i = 1, \dots, m$, contains at most $g - 1$ vertices, the subgraph $G[V_i]$ induced by V_i is bipartite. Choose randomly (and independently of the other subgraphs) one of its two color classes and color its vertices red. Observe that if an edge has both its end-points colored red, then it must be an edge of the matching

M complementary to F . If this happens, choose randomly one vertex of this edge and uncolor it.

The resulting set of red vertices is independent. We estimate the probability that a vertex v is red conditioned by $v \notin W$. With probability $1/2$, v is initially colored red. However, with probability at most $1/2$ its neighbor through an edge of M is also colored red (it can happen that this neighbor is in W). If this is the case, then the vertex v is uncolored with probability $1/2$. Consequently, the probability that v is red is at least $1/2 \cdot (1 - 1/4) = 3/8$. Multiplying by the probability that $v \notin W$, which is $1 - 2/(g + 1)$, we obtain that the vertex v is included in the independent set with probability at least $3(1 - 2/(g + 1))/8$ as claimed earlier. \square

Acknowledgments The authors would like to thank the anonymous referees for careful reading the manuscript and their valuable comments improving the presentation of the results.

References

- [1] W. Duckworh, M. Zito: Large independent sets in random regular graphs, *Theoretical Computer Science* 410 (2009), 5236–5243.
- [2] H. Hatami, X. Zhu: The fractional chromatic number of graphs of maximum degree at most three, *SIAM J. Discrete Math.* 23 (2009), 1762–1775.
- [3] A. M. Frieze, S. Suen: On the independence number of random cubic graphs, *Random Structures and Algorithms* 5 (1994), 649–664.
- [4] Jan Hladký: Bipartite subgraphs in a random cubic graph, Bachelor Thesis, Charles University (2006).
- [5] C. Hoppen: Properties of graphs with large girth, Ph.D. Thesis, University of Waterloo (2008).
- [6] B. D. McKay: Maximum bipartite subgraphs of regular graphs with large girth, *Proceedings of the 13th Southeastern Conf. on Combinatorics, Graph Theory and Computing*, Boca Raton, Florida (1982).

- [7] B. D. McKay: Independent sets in regular graphs of high girth, *Ars Combinatoria* 23A (1987), 179–185.
- [8] B. D. McKay: private communication, 2010.
- [9] J. Nešetřil: Aspects of structural combinatorics (graph homomorphisms and their use), *Taiwanese J. Math.* 3 (1999), 381–423.
- [10] N. C. Wormald: The asymptotic distribution of short cycles in random regular graphs, *J. Combin. Theory, Ser. B* 31 (1981), 168–182.
- [11] N. C. Wormald: Differential Equations for random processes and random graphs, *Annals of Applied Probability* 5 (1995), 1217–1235.
- [12] N. C. Wormald: private communication, 2009.
- [13] Ondřej Zýka: On the bipartite density of regular graphs with large girth, *Journal of Graph Theory* (1990), 631–634.

Appendix

```
W_THOLD = 1.0/1000000
p_1 = 0.00001
p_2 = 0.00001

def state():
    print("%d: r=%f b=%f w=%f q3=%f q2=%f q1=%f w3=%f w2=%f w1=%f w0=%f"
          % (k, p_r, p_b, p_w, q[3], q[2], q[1], w[3], w[2], w[1], w[0]))

def gen_C():
    S= {1: q[1], 2: q[2], 3: q[3]}
    for u in (1,2,3):
        A=(1,2,3)
        B=S if (u>1) else {"-":1}
        C=S if (u>2) else {"-":1}
        for a in A:
            for b in B:
                for c in C:
                    C3[(u,a,b,c)] = w[u]*q[a]*B[b]*C[c]
                    C2[(u,a,b,c)] = q[u]*q[a]*B[b]*C[c]

def init():
    global p_r, p_w, p_b

    t = (1-p_1)**2
    w[3] = t**3
    w[2] = 3*(t**2)*(1-t)
    w[1] = 3*t*(1-t)**2
    w[0] = (1-t)**3

    q[3] = t**2
    q[2] = 2*t*(1-t)
    q[1] = (1-t)**2

    p_b = 1-(1-p_1)**3
    p_r = p_1*(1-p_b)
    p_w = 1 - p_r - p_b

k=0
p_w = 1
p_b = p_r = 0

w = [0,0,0,3]
q = [0,0,0,1]
C3 = {}
C2 = {}

while (p_w > W_THOLD):
    k+=1

    if (k==1):
        init()
        state()
        continue
    o1 = q[1]/(1-q[2]**2)
```

```

e1 = q[2]*o1
p1 = o1 + e1

o3 = q[3]/(1-q[2]**2)
e3 = q[2]*o3
p3 = o3 + e3

p3_n = q[3]/(1-q[2]*(1-p_2))
o3_n = q[3]/(1-q[2]**2 * (1-p_2)**2)
e3_n = q[2]*(1-p_2)*o3_n

o3_y = (q[2]**2 *(1-(1-p_2)**2)*o3)/(1-q[2]**2 *(1-p_2)**2)
e3_y = q[2]*(p_2*o3 + (1-p_2)*o3_y)

Pr = [1,0,0,0]
Pb = [0,0,0,0]

Pr[1] = (e1 + o1*.5) + p3
Pb[1] = o1*.5

Pr[2] = e1**2 + e1*o1 + 2*e1*p3
Pr[2] += (1-p_2)*(o3_y**2 + 2*o3_y*o3_n+o3_y*e3_y+o3_n*e3_y+o3_y*e3_n)
Pr[2] += p_2*(o3**2 + o3*e3)

Pb[2] = o1**2 + e1*o1 + 2*o1*p3
Pb[2] += (1-p_2)*(e3_y**2 + 2*e3_y*e3_n+o3_y*e3_y+o3_n*e3_y+o3_y*e3_n)
Pb[2] += p_2*(e3**2+o3*e3)

Pb[3] = 1-(p3_n+e1+o3_y/2)**3

p_r += p_w*(w[0]+w[1]*Pr[1]+w[2]*Pr[2])
p_b += p_w*(w[1]*Pb[1]+w[2]*Pb[2]+w[3]*Pb[3])
p_w = 1-p_r-p_b

r32 = o1+(1-p_2)*(p3_n+.5*e3_y)+p_2*(.5*e3)
s33 = (p3_n+e1+o3_y/2)**2
s32 = (1-p_2)*p3_n / r32

gen_C()

T = [0,0,0,0]
sum = 0
for C in C3:
    deg1=False
    for x in C:
        if x==1:
            deg1=True
            break
    if deg1: continue

    N={}
    C_p = C3[C]

    if (C[0]==2):

```

```

    N[3] = {0:1}
    C_p*=1-p_2
    for i in (1,2):
        if (C[i]==3):
            N[i] = {0: s33 }
            N[i][1] = 1-N[i][0]
        elif (C[i]==2):
            C_p *= (1-p_2)*p3_n
            N[i] = {0: 1}

    elif (C[0]==3):
        for i in (1,2,3):
            if (C[i]==3):
                N[i] = {0: s33 }
            elif (C[i]==2):
                C_p *= r32
                N[i] = {0: s32 }

            N[i][1] = 1-N[i][0]

    sum += C_p
    for a in N[1]:
        for b in N[2]:
            for c in N[3]:
                T[C[0]-a-b-c] += C_p*N[1][a]*N[2][b]*N[3][c]

for i in (0,1,2,3): w[i] = T[i]/sum

T = [0,0,0,0]
sum = 0
for C in C2:
    deg1=False
    for x in C:
        if x==1:
            deg1=True
            break
    if deg1: continue

N={}
C_p = C2[C]

if (C[1]==3):
    C_p *= s33
elif (C[1]==2):
    C_p *= (1-p_2)*p3_n

if (C[0]==2):
    N[3] = {0:1}
    C_p*=1-p_2

    if (C[2]==3):
        N[2] = {0: s33 }
        N[2][1] = 1-N[2][0]
    elif (C[2]==2):
        C_p *= (1-p_2)*p3_n
        N[2] = {0: 1}

```



```

elif (C[0]==3):
    for i in (2,3):
        if (C[i]==3):
            N[i] = {0: s33 }
        elif (C[i]==2):
            C_p *= r32
            N[i] = {0: s32 }

            N[i][1] = 1-N[i][0]

    sum += C_p
    for b in N[2]:
        for c in N[3]:
            T[C[0]-b-c] += C_p*N[2][b]*N[3][c]

for i in (1,2,3): q[i] = T[i]/sum

state()

```